1 **AQMM: Enabling Absolute Quantification of Metagenome** 2 **and Metatranscriptome**

3 **Authors: Xiao-Tao Jiang, Ke Yu, Li-Guan Li, Xiao-Le, Yin, An-Dong, Li, Tong**
4 **Zhang***

5 **Affiliations: Environmental Biotechnology Laboratory, Department of Civil**
6 **Engineering, University of Hong Kong**

7 * Corresponding author Email: zhangt@hku.hk

8 **November 12, 2017**

9 **Abstract**

10 Metatranscriptome has become increasingly important along with the application of
11 next generation sequencing in the studies of microbial functional gene activity in
12 environmental samples. However, the quantification of target active gene is hindered
13 by the current relative quantification methods, especially when tracking the sharp
14 environmental change. Great needs are here for an easy-to-perform method to obtain
15 the absolute quantification. By borrowing information from the parallel metagenome,
16 an absolute quantification method for both metagenomic and metatranscriptomic data
17 to per gene/cell/volume/gram level was developed. The effectiveness of AQMM was
18 validated by simulated experiments and was demonstrated with a real experimental
19 design of comparing activated sludge with and without foaming. Our method provides
20 a novel bioinformatic approach to fast and accurately conduct absolute quantification
21 of metagenome and metatranscriptome in environmental samples. The AQMM can be
22 accessed from <https://github.com/biofuture/aqmm>.

23 **Keywords:** metagenome, metatranscriptome, absolute quantification, differential
24 expression genes

Background

Shotgun metatranscriptomics is a powerful tool in identifying the overall expression of microorganisms in an environment (Alexander et al. 2015, Gifford et al. 2011, Shi et al. 2009, Turner et al. 2013, Yu and Zhang 2012), shedding light on discovering how microbes respond to environmental changes or diseases status (Jorth et al. 2014, Mason et al. 2012) and capturing gene expression patterns for functionally important bacteria in engineering systems (Oyserman et al. 2015, Stark et al. 2014). For these applications, accurate quantification is required to detect the true variations or differential expression genes (DEGs).

Traditionally, the abundance of a transcript in RNA-sequencing (RNA-seq) is thought to be influenced by the library size and inherent dependence on the expression levels of other transcripts as described in a comprehensive review (Rapaport et al. 2013). Following this idea, transcripts in RNA-seq was generally quantified by within-sample normalization. One of the most common quantification methods was RPKM (Mortazavi et al. 2008) (reads per kilobase of exon model per million mapped reads) which considered factors of both the length of gene and library size. Another improved within-sample normalization method was TPM (transcript per million) (Wagner et al. 2012) which only considered the transcript rather than the whole library size and respected the invariance of relative molar RNA concentration (rmc). The TPM was thought to be better fitted in sample comparison due to its unit-free characteristics. The FPKM (substitute the reads with fragments in RPKM) was an adaption of RPKM to pair-end reads. These above methods are all relative quantification (RQ) and suffer from the ‘composition effects’ (the increase of one transcript will decrease other unrelated transcript). To relieve this problem, Robinson and Oshlack proposed a new normalization method “TMM” (trimmed mean of

M-values) to detect the DEGs under the hypothesis that most of the genes are not differentially expressed (Robinson and Oshlack 2010), which has been integrated into popular DEGs detection R software edgeR (Robinson et al. 2010). The scaling factor in edgeR for normalization is the TMM value. Another method was to compute the median of the ratio as the scaling factor and it could be conducted by R software DESeq/DESeq2 (Love et al. 2014). It is also based on the assumption that most genes are not DEGs and this method then calculates the scaling factor (median of ratios) associated with this sample to perform further normalization. In the two software, the negative binomial distribution was applied to adjust the distribution of transcript between different conditions to relieve the dispersion effects of deviation from standard poisson distribution (Rapaport et al. 2013). Although with these efforts in optimizing the normalization process, these indices were all still RQ based and the relationship could be distorted while performing comparative analysis across samples, especially when borrowing these methods from traditional Eukaryote RNA-seq to current Prokaryote metatranscriptome studies (Conesa et al. 2016). One feasible way to solve the problem was to get the absolute quantification (AQ) of expression level for each transcript. For example, the qRT-PCR has long been applied in RNA-seq or microarray data for AQ (Becker-André and Hahlbrock 1989, Whelan et al. 2003). In addition, there were methods by spiking in exterior/alien RNA in microarray to get the per cell absolute quantification (Kanno et al. 2006) and internal standard approach to estimate per liter expression in marine metatranscriptome (Gifford et al. 2011). However, the experiment to perform spiking internal standard was difficult due to its skill-demanding nature and for metatranscriptome data, factors like the time to add spike-in material, the type and the amount of alien RNA required still needed to be elaborately designed. Hence, it was not as popular as those RQ methods. The

quantification methods in the newly developed analyzing pipelines for metatranscriptome like IMP (Narayanasamy et al. 2016), MetaTrans (Martinez et al. 2016), COMAN (Ni et al. 2016) and SAMSA (Westreich et al. 2016) were still all based on RQ methods; this would result in accelerated spreading of the inaccurate quantification in many studies.

To solve the problem of RQ and get an accurate quantification without performing spike-in experiment, an AQ bioinformatics software package AQMM was developed by combining metagenome and metatranscriptome data to achieve the goal of accurate and comparable quantification. In this study, we firstly introduced the AQMM algorithm flow, and then compared and validated it with RQ methods by simulated metagenome and metatranscriptome data. Moreover, we further applied this algorithm to a real combination of metagenome and metatranscriptome dataset in quantifying genes and transcripts of resistome in six foaming activated sludge (FAS) and non-foaming activated sludge (NFAS) samples.

Results

Overall view of AQMM algorithm

The AQMM (**Fig. 1**) was designed to perform AQ of parallel metagenome and metatranscriptome dataset no matter whether spike-in experiment/internal standard was initially added or not. The major aims were to obtain the AQ of genes/transcripts/taxa in samples and to accurately detect DEGs in metatranscriptome data. The assumptions under the algorithm include: 1) with the known extraction ratio of DNA for a DNA extraction Kit for a type of sample, the total weight of DNA per volume of the sample could be calculated. The weight of the sequenced library of DNA could be estimated with the molecular weight and bases numbers of A, T, C and

G in the sample. Then, the ratio of sequenced DNA to total weight of DNA per volume of the sample could be calculated. In addition, by utilizing the universal single-copy phylogenetic marker genes (USCMGs), the number of cells for a metagenome library could be estimated accurately (Nayfach and Pollard 2015). With the above information, cells per volume could be calculated for a metagenome data; 2) Using the same volume of the sample contains the same number of cells for DNA and RNA extraction, the cell number per volume to extract RNA was the same as the parallel DNA sample; 3) With the known ratio of RNA extraction and the rRNA ratio of total RNA, by a similar process, the sequenced RNA weight ratio could be calculated, and then the equivalent cell numbers in a metatranscriptome could be deduced accordingly. With the cell numbers included in the metagenome and metatranscriptome data, the abundance of gene/transcript could be normalized to per cell level. Moreover, as the number of cells per volume is available, per cell quantification could be easily transformed into per volume quantification.

Comparing and validating AQMM with RQ methods using simulated metatranscriptome demo

To reveal the problem of RQ methods like RPKM, edgeR and DESeq2 and to assess the effectiveness of AQMM, simulated metatranscriptomic datasets comprised of known community structure and expression levels were generated (**Fig. 2; Details in methods**). The simulated data was with known ground truth absolute expression for each gene. For simplicity, to focus on the quantification of metatranscriptome in identifying DEGs, we assume the DNA content are not changed like what happens in a reactor with a stable biomass concentration, however the gene expression under condition A and B are significantly changed with fold of 2 or 16 in part of the bacteria like what happens under sharp environmental change. In order to focus only on the of

influence normalization methods, in generation of the simulated metatranscriptome, the base qualities of were all set with 50 and to eliminate the influence of mapping process, the mapping criteria of bowtie2 was set to exactly match without gap and mismatch allowed (bowtie2 parameters, -N 0 -L 31, --rdg 100,150 --rfg 100,150 --gbar 100,150). The result of DEGs detection was in **Table 1**. We can observe that compared with ground truth, the RQ methods detect quite a large portion of false positive higher gene expression under condition A. On the contrary, the AQMM method which aims to obtain the AQ has limited errors detection even with a given variance in RNA extraction efficiency (**Table 1**). Noticeably, in real combination of metagenome and metatranscriptome, the metagenome could also be totally different, and in this case, the AQMM is still applicable.

Case study: AQ of activate resistome in FAS and NFAS

The AQMM was applied in the six metagenome and metatranscriptome dataset of FAS and NFAS, the AQ of the sequenced cells generated by the pipeline were shown in **Table 2**. In detail, the metagenome contained 8 to 11.8 GBs data and metatranscriptome with a depth between 13 and 16 GBs for each sample. The “per cell/volume” quantifying values were the fundamental of normalizing to cells or volume in order to perform comparison among different studies. The cell number per milliliter in literature was at $3.3\text{E}+09$ using flow Cytometer to quantify (Foladori et al. 2010) and was from $2.1\text{E}+09$ to $5.5\text{E}+09$ using CFU and flow Cytometer (Manti et al. 2008) level for AS which was a bit lower than the obtained number in this study at the magnitude of $\text{E}+10$ cells per milliliter. Overall number of mRNA molecules per cell are 387.98 ± 102.86 and 235.21 ± 30.59 averagely for FAS and NFAS, respectively (**Table S1**), which is consistent with previous observation of coastal bacterioplankton by 142-238 mRNA molecules per Cell (Gifford et al. 2011, Moran et al. 2013).

149

150 As WWTPs become the hot-spot of antibiotic resistant genes (ARGs) to the receiving
 151 environment. Hence, the expressions of ARGs in the AS were in great concerns and
 152 further profiled. Overall, the abundance of ARGs per cell in FAS and NFAS were
 153 0.0517 ± 0.0034 and 0.0483 ± 0.0041 ; and the transcript of ARGs per cell were
 154 0.0140 ± 0.0039 and 0.0059 ± 0.0009 , respectively (**Table S2 & S3**). The overall
 155 transcription of ARGs was significantly higher in FAS compared with NFAS. At DNA
 156 level, only tetracycline resistance gene was higher in FAS and beta-lactam was higher
 157 in NFAS, other types were not significantly different. However, at transcript level, all
 158 the types were all significantly higher in FAS. Among the nine transcribed ARGs
 159 types, beta-lactam and sulfonamide resistance genes were the most abundant
 160 expressed ARG types in both FAS and NFAS. Per volume ARGs abundance and
 161 expression at type level were shown in **Fig. 3**. The overall ARGs abundance per
 162 milliliter AS in FAS and NFAS were $2.51\text{E}+09 \pm 2.44\text{E}+08$ and $2.66\text{E}+09 \pm$
 163 $5.63\text{E}+08$; and the transcript of ARGs per milliliter were $9.83\text{E}+09 \pm 3.82\text{E}+08$ and
 164 $4.49\text{E}+09 \pm 5.10\text{E}+08$, respectively. With the AQ results, the transcripts per copy gene
 165 (TPCG), which represents of the transcribe rate could be further derived. The
 166 unclassified, quinolone, multidrug and beta-lactam were more active in FAS
 167 compared with NFAS in terms of TPCG, (**Table S4**). For the detected ARGs, the host
 168 taxonomy was assigned by LCA algorithms using all the genes annotation in the same
 169 Contig. Thirteen orders were detected to carry ARGs and eleven of them were
 170 transcribed (**Fig. 3**). The most ARGs transcribed order was Enterobacteriales. The
 171 active ARGs in bacteria enclosed in foams of FAS posed potential threats for the
 172 public as ARGs carrying bacteria could spread into the air from the foams bubbles.

173

The co-expression of ARGs and MRGs was also studied to check whether there were co-expression effects at the RNA level. Using this dataset, we observed co-expression within ARGs, within MRGs, and between ARGs and MRGs (**Fig. 4**). Numerous types of MRGs were detected in the metagenome and metatranscriptome. The most abundant MRG was Cu resistant genes and for the ARGs, beta-lactam, tetracycline and aminoglycoside were the most expressed types. The highest number of co-expression within MRGs was Cr and Fe; while within ARGs was beta-lactam and tetracycline. The most MRG and ARGs co-expression was Cr, which co-expression with nine types of ARGs. This was the first transcript level evidence of the co-expression of ARGs and MRGs in AS.

Discussion

Metatranscriptome enabled the study of whole metabolic pathways expression of the system and many studies had already taken this advantage for different environments, such as in marine (Mason et al. 2012), rhizosphere of the plant (Turner et al. 2013), human oral disease (Jorth et al. 2014). Each study has specific method to integrate the metagenome and metatranscriptome information to understand the microbes and their activities in the system. The quantification of metatranscriptome was generally RQ based methods. The RQ methods are problematic as they may not be able to reflect the actual expression level of a population in the whole community. Due to the relative characteristics, the RQ methods are always suffer from the so-called composition effects, which indicates that the upgrade of one gene should definitely make other genes downgrade. Additionally, the RQ methods are just a relative portion rather than a value with biological implications. On the contrary, the AQ could be more biological meaningful at per cell/volume unit. Hence, it was necessary to conduct AQ to compare different samples. In this study, we proposed an AQ method

and developed a set of algorithms to conveniently calculate the absolute number of sequenced cells for each RNA library by borrowing cell numbers from a corresponding data set of DNA library of the same sample.

Noticeably, there were several hypotheses for the application of the proposed method. Firstly, the sample used to extract DNA and RNA should contain the same cell numbers per volume which could be easily met with sufficient mixing of samples. Secondly, the DNA and RNA extraction efficiency should be estimated, as well as the rRNA ratio in total RNA. This was likely difficult to achieve. However, for an environmental sample, generally literature based data could be used for the extraction kit, for example, to FastDNA SPIN Kit for Soil, the extract efficiency was estimated as 28.4% (Mumy and Findlay 2004). Most importantly, as the parallel samples were extracted under the same condition, the difference between samples was minimized **(DNA extraction data, unpublished)**. This AQMM method is capable of performing absolute quantification of both metagenome and metatranscriptome without the requirement to do complex spike-in experiments. Importantly, AQMM avoids the RQ problems of composition effects and able to detect accurate DEGs. Hence, the proposed AQMM is a method in between experimental spike-in based AQ methods and those improved RQ methods of TMM based edgeR.

With AQ, a number of indices with various biological meaning were proposed in this study (Methods), for example, the transcript per copy gene (TPCG) index is a reflection of the transcribe rate of the gene, which could never be delivered by RQ methods. It was demonstrated with simulating RNA-seq that the organism abundance (community structure) was important at normalizing metatranscriptome data in identifying DEGs (Klingenberg and Meinicke 2017). The gene per cell (GPC) and transcript per cell (TPC) in AQMM are global level normalization indices and the

scaling factor is the total number of cells in the DNA or RNA library. This global scaling factor could be easily transformed into taxa specific scaling factors with the relative quantification of different taxa with indices of transcript of taxon A per cell (TTPC). Hence, the normalization in AQMM is well fit for the factor of microbial abundance in metatranscriptome data.

AS is important biological wastewater treatment process and this system is considered as a hot spot for ARG dissemination into the receiving water. The foaming of AS would result in spreading of foams with AS bacteria into the surrounding environment. Understanding the active resistome and the host bacteria in foaming AS enables engineers understanding the risk of sludge foaming incurred to the surrounding environment. We observed a wide profile of active ARG types in the FAS, the identification of opportunity pathogen bacteria *Pseudomonas* carrying active ARGs alerts us the risk of spreading ARGs-carrying bacteria. Additionally, per cell mRNA molecules is an important indication of the activity of the cell, generally natural bacterial communities was observed to hold a lower inventory of transcripts (Moran et al. 2013); and the absolute quantification obtained with AQMM was well-fitted with previous observation.

Conclusions

In this study, we filled the gap of lacking a bioinformatic algorithm to perform AQ of metatranscriptomic data. The developed AQMM was demonstrated to gain enhanced performance at identifying DEGs compared with those RQ methods benchmarked with simulated metagenomic and metatranscriptomic data. Additionally, with the AQMM, the active resistome in foaming and normal activated sludge were quantified to per cell/volume level and even down to the transcription per copy gene. The active

ARG host were quantified and the co-expression of MRGs and ARGs was revealed for the first time in AS.

Materials and methods

Absolute quantification of gene abundance and transcript expression

We developed a package of scripts AQMM (absolute quantification of metagenome /metatranscriptome) to perform comparative analysis.

The formula for cells per mL:

$$C = N_c / \frac{L_{size} * 10^9 * \frac{(R_A * 313.2 + R_T * 304.2 + R_C * 289.18 + R_G * 329.21)}{6.022 * 10^{23}}}{X/\alpha} \quad (1)$$

C is value of cell numbers per mL AS

N_c is the estimated cell numbers for the sequenced DNA library with USCMGs

L_{size} is the sequencing depth

R_A , R_T , R_C and R_G are ratios of A, T, C and G

X is the overall extracted weight (ng) of DNA for 1 mL AS

α is DNA extraction efficiency, for FAST DNA Kit for Soil, α is estimated as 28.2%

(Mumy and Findlay 2004).

The sequenced cells for RNA sequencing, for a RNA-seq with library size of L_{size} after removing all ribosomal RNA, the equivalent sequenced cells for this sample is

$$E_c = C * \frac{L_{size} * 10^9 * \frac{R_A * 329.2 + R_U * 306.2 + R_C * 305.2 + R_G * 345.2}{6.022 * 10^{23}}}{Y * \gamma / \beta} \quad (2)$$

E_c is the estimated number of cells sequenced for this RNA library

C is value of cell numbers per mL AS

L_{size} is the sequencing depth

R_A , R_U , R_C and R_G are ratios of A, U, C and G, the value they multiplied are molecular weight

Y is the overall extracted weight (ng) of RNA for 1 mL AS

272 β is RNA extraction efficiency, the estimated β is about 7.5% as used in this study.

273 This value was deduced from AS empirical data of proportion of RNA biomass by
274 engineering perspective and the extracted RNA biomass.

275 γ is non-ribosomal RNA ratio, for AS the estimated γ is about 0.03.

276 Based on the two AQ numbers of cells for each sample, the gene or transcript
277 abundance matrix could be further normalized into the following indices.

278 **GPC** (Gene per Cell): an indication of the overall abundance of the gene in system.

$$279 \quad GPC = \frac{N_{read} * L_{read} / L_{gene}}{N_c} \quad (3)$$

280 **TPC** (Transcript per Cell): an indication of overall activity of the gene in system.

$$281 \quad TPC = \frac{N_{read} * L_{read} / L_{gene}}{E_c} \quad (4)$$

282 **TPCG** (Transcript per copy gene): an indication of the absolute activity of one copy
283 gene in the system, equivalent to transcribe rate for each gene.

$$284 \quad TPCG = TPC / GPC \quad (5)$$

285 **GTPC** (Gene of taxon A per Cell): an indication of the overall abundance of the taxon
286 in system averagely.

$$287 \quad GTPC = \sum_{i=1}^n GPC_i \quad (6)$$

288 **TTPC** (Transcript of taxon A per Cell): an indication of overall activity of the taxon
289 in system averagely.

$$290 \quad TTPC = \sum_{i=1}^n TPC_i \quad (7)$$

291 **ATCT** (Averagely transcript per copy gene of taxon A): indication of the averagely
292 absolute activity per copy expressed gene in taxon A

$$293 \quad ATCT = \frac{1}{n} \sum_{i=1}^n TPCG_i \quad (8)$$

294 N_c is the estimated cell numbers for the sequenced DNA library,

295 N_{read} is the number of reads or transcript mapping to the target gene

296 L_{read} is the length of reads
 297 L_{gene} is the length of the target gene
 298 n is the number of genes affiliated to taxa A.
 299 When the number of cells per mL was obtained, using the GPC, genes per mL could
 300 be calculated.

301 **Simulating metatranscriptome data**

302 To validate our method and comparing with those RQ methods in identifying the
 303 DEGs, simulated data was generated by workflow illustrated in **Fig. 2**. For simplicity,
 304 the DNA was set unchanged to mimic the activated sludge community composition
 305 with 16 strains from different phylogeny. The metatranscriptome data sets were
 306 generated for two conditions A and B, each with three biological duplications; for the
 307 condition A and B, there were part of the strains with folds of significantly changed
 308 expression (**Table S5**). To only focus on the quantification method, all the system
 309 errors caused by other factors like base qualities, cDNA synthesis, assembly, mapping
 310 parameters were not considered.

311 **Sampling**

312 AS samples were collected in Shatin wastewater treatment plant at three locations
 313 along the flow direction while serious foaming happened at 2016-04-08 and nearly no
 314 foaming happened at 2016-04-25. Samples were collected on site by storing in liquid
 315 nitrogen immediately and then transported to the laboratory for RNA extraction. The
 316 DNA samples were mixed with 1:1 100% ethanol and AS and then stored at -20 °C
 317 fridge. Totally six samples were collected for both DNA and RNA samples alongside
 318 the segment aeration tank in three locations as depicted in **Fig. 5**.

Whole DNA, total RNA extraction, removal of ribosomal RNA, cDNA synthesis and next generation sequencing

FAST DNA Kit was used to extract total DNA from 1 mL mixed AS samples. RNeasy Mini was used to extract the total RNA from 0.5 mL AS stored in liquid nitrogen. The extracted RNA was then processed by DNase I to eliminate the DNA in the RNA samples. Then both Illumina Ribo-Zero rRNA removal KIT (Bacteria) and Ribo-Zero rRNA removal KIT (Human/Mouse/Rat) was applied for each sample to remove rRNA from Prokaryote and Eukaryote respectively in order to get the total clean non-ribosomal RNA. Generally, metatranscriptome rRNA depletion was only used the Ribo-Zero for Bacteria, in this study, the addition of Eukaryote rRNA removal was due to a fact that by only using the Ribo-Zero Bacteria rRNA removal Kit for AS, there was still over half of RNA were rRNA from Eukaryote (our previous experiment, data unpublished). To get more non-rRNA, the Ribo-Zero rRNA Kit to remove Eukaryote was also used. RNA then was fragmented into 170 bps library and was reverse-transcribed to construct cDNA library for sequencing. The quality of DNA and RNA were assessed with Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA). All the samples was sent to sequence, considering the complexity of AS and the aims of this study to detect the expression of low abundance gene, we gave each sample a very deep sequencing depth which doubled the sequencing depth in previous studies. All the samples were sequenced with Hiseq 4000 in BGI-ShenZhen. DNA samples with PE-150 with library size of 300 bps. And RNA with PE101 of library size 170 bps.

Bioinformatics analysis

Quality filtering was firstly performed on DNA and RNA reads to keep only high quality reads using trimmomatic v1.04 (Bolger et al. 2014). DNA datasets were

344 pooled together and assembled by CLC Genomics Workbench 6.5.3 (CLC Bio,
345 Aarhus, Denmark, <https://www.qiagenbioinformatics.com/>) with default parameters.
346 Finally, 1,430,611 contigs with length over 100 bps (N50, 2,416 bps; 2,457,704,443
347 bps length in total) were obtained and 74.5% of reads could be mapped back to these
348 Contigs. All these contigs were sent to predict genes with Prodigal (version 1.5)
349 (Hyatt et al. 2010) using '-meta' parameter and finally 3,234,330 genes were obtained.
350 By removing exactly the same genes using USEARCH (version 8.0.1623) (Edgar
351 2010) unique command (parameters -fastx_uniques), 3,234,246 million genes were
352 kept; this set was defined as 'unique gene set'. Reads were mapped back to the contig
353 set and 'unique gene set' to obtain reads coverage matrixes for contigs and genes. The
354 matrix of genes was finally normalized to cell numbers. For metatranscriptome
355 samples, after quality filtering, the SortMeRNAv1.9 was used to remove all the
356 possible ribosomal RNA by aligning to six databases of bacteria, archaea and
357 eukaryotic small and large subunits (Kopylova et al. 2012). RNA reads for each
358 sample were then mapped back to the 'unique gene set' to get the transcript coverage
359 for each gene with CLC genomic workbench 6.5.3 using parameters of gap penalty 2,
360 gap extension 3, length fraction 0.8 and similarity at least 0.9.

361 Taxonomy composition of the metagenome was generated with MEGAN6 (Huson et
362 al. 2015). In detail, all genes were aligned to NCBI NR database (version 201603)
363 with diamondv1.09 (Buchfink et al. 2015) to find out the homology proteins. To each
364 gene, the local common ancestors (LCA) were applied using the taxonomy
365 information of the hit NR protein in NCBI taxonomy database (Acland et al. 2014)
366 and then this gene was annotated with the common ancestor taxonomy. We further
367 processed the NCBI taxonomy annotation results to remove those subdivisions and
368 subgroups to format the annotation to 7 levels from kingdom to species. Among total

3,234,246 unique genes predicted, 2,348,907 could be aligned to NR database. The remaining 885,339 (27.3%) genes could not be annotated with the NR database. The abundance of each taxon was a sum of all the annotated genes under that taxon in every sample. Antibiotic resistant genes (ARGs) were annotated with SARG database which contained a type-subtype structure annotation (Yang et al. 2016). Metal resistance genes (MRGs) were detected by aligning the “unique gene set” to the MRG database (Li et al. 2017). Absolute abundance and transcript was determined by AQMM.

Declarations

Data availability

The metagenome and metatranscriptome raw data were deposited in NCBI SRA under accession number XXX.

Analyzing document

The analyzing document for the whole data analysis and simulation process could be accessed from

https://github.com/biofuture/aqmm/blob/master/Analysing_document.txt

Conflict of interest

The authors declare no conflict of interest

Acknowledgements

The authors would like to thank GRF of Hong Kong for financial support (172099/14E). Xiaotao Jiang would like to thank The University of Hong Kong for the postgraduate scholarship, and Dr. Ke Yu, Dr. Li-Guan, and Dr. Andong, Li, would like to thank The University of Hong Kong for postdoc fellowship.

Contributions

T. Zhang and X.-T. Jiang design the study of quantification. X.-T. Jiang developed the software and performed the wet-lab and simulation experiments. X.-T. Jiang performed the bioinformatics analyses. X.-T. Jiang, A.D. Li and K. Y. did the DNA and RNA extraction experiment. L.-G. Li did the MRG analyses. T. Zhang and X.-T. Jiang wrote the manuscript. T. Zhang, X.-T. Jiang, A.D. Li, L.G. Li and X.L. Yin revised the manuscript.

Reference

- Acland, A., Agarwala, R., Barrett, T., Beck, J., Benson, D.A., Bollin, C., Bolton, E., Bryant, S.H., Canese, K. and Church, D.M. (2014) Database resources of the national center for biotechnology information. *Nucleic acids research* 42(Database issue), D7.
- Alexander, H., Jenkins, B.D., Rynearson, T.A. and Dyhrman, S.T. (2015) Metatranscriptome analyses indicate resource partitioning between diatoms in the field. *Proceedings of the National Academy of Sciences* 112(17), E2182-E2190.
- Becker-André, M. and Hahlbrock, K. (1989) Absolute mRNA quantification using the polymerase chain reaction (PCR). A novel approach by a PCR aided transcript titration assay (PATTY). *Nucleic acids research* 17(22), 9437-9446.
- Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, btu170.
- Buchfink, B., Xie, C. and Huson, D.H. (2015) Fast and sensitive protein alignment using DIAMOND. *Nature Methods* 12(1), 59-60.
- Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A., Szczesniak, M.W., Gaffney, D.J., Elo, L.L. and Zhang, X. (2016) A survey of best practices for RNA-seq data analysis. *Genome biology* 17(1), 13.
- Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26(19), 2460-2461.
- Foladori, P., Bruni, L., Tamburini, S. and Ziglio, G. (2010) Direct quantification of bacterial biomass in influent, effluent and activated sludge of wastewater treatment plants by using flow cytometry. *Water Research* 44(13), 3807-3818.
- Gifford, S.M., Sharma, S., Rinta-Kanto, J.M. and Moran, M.A. (2011) Quantitative analysis of a deeply sequenced marine microbial metatranscriptome. *The ISME journal* 5(3), 461-472.
- Huson, D., Beier, S., Buchfink, B., Flade, I., Górski, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.-J. and Tappu, R. (2015) MEGAN6-Microbiome analysis involving hundreds

of samples and billions of reads, preparation.

Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W. and Hauser, L.J. (2010) Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC bioinformatics* 11(1), 119.

Jorth, P., Turner, K.H., Gumus, P., Nizam, N., Buduneli, N. and Whiteley, M. (2014) Metatranscriptomics of the human oral microbiome during health and disease. *MBio* 5(2), e01012-01014.

Kanno, J., Aisaki, K.-i., Igarashi, K., Nakatsu, N., Ono, A., Kodama, Y. and Nagao, T. (2006) "Per cell" normalization method for mRNA measurement by quantitative PCR and microarrays. *BMC genomics* 7(1), 64.

Klingenberg, H. and Meinicke, P. (2017) How To Normalize Metatranscriptomic Count Data For Differential Expression Analysis. *bioRxiv*, 134650.

Kopylova, E., Noé, L. and Touzet, H. (2012) SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28(24), 3211-3217.

Li, L.G., Xia, Y. and Zhang, T. (2017) Co-occurrence of antibiotic and metal resistance genes revealed in complete genome collection. *ISME Journal* 11(3), 651-662.

Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* 15(12), 550.

Manti, A., Boi, P., Falcioni, T., Canonico, B., Ventura, A., Sisti, D., Pianetti, A., Balsamo, M. and Papa, S. (2008) Bacterial cell monitoring in wastewater treatment plants by flow cytometry. *Water Environment Research* 80(4), 346-354.

Martinez, X., Pozuelo, M., Pascal, V., Campos, D., Gut, I., Gut, M., Azpiroz, F., Guarner, F. and Manichanh, C. (2016) MetaTrans: an open-source pipeline for metatranscriptomics. *Scientific reports* 6, 26447.

Mason, O.U., Hazen, T.C., Borglin, S., Chain, P.S., Dubinsky, E.A., Fortney, J.L., Han, J., Holman, H.-Y.N., Hultman, J. and Lamendella, R. (2012) Metagenome, metatranscriptome and single-cell sequencing reveal microbial response to Deepwater Horizon oil spill. *The ISME journal* 6(9), 1715-1727.

Moran, M.A., Satinsky, B., Gifford, S.M., Luo, H., Rivers, A., Chan, L.-K., Meng, J., Durham, B.P., Shen, C. and Varaljay, V.A. (2013) Sizing up metatranscriptomics. *The ISME journal* 7(2), 237.

Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L. and Wold, B. (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods* 5(7), 621-628.

Mumy, K.L. and Findlay, R.H. (2004) Convenient determination of DNA extraction efficiency using an external DNA recovery standard and quantitative-competitive PCR. *Journal of Microbiological Methods* 57(2), 259-268.

Narayanasamy, S., Jarosz, Y., Muller, E.E., Heintz-Buschart, A., Herold, M., Kaysen, A., Laczny, C.C., Pinel, N., May, P. and Wilmes, P. (2016) IMP: a pipeline for reproducible

reference-independent integrated metagenomic and metatranscriptomic analyses. *Genome biology* 17(1), 260.

Nayfach, S. and Pollard, K.S. (2015) Average genome size estimation improves comparative metagenomics and sheds light on the functional ecology of the human microbiome. *Genome biology* 16(1), 51.

Ni, Y., Li, J. and Panagiotou, G. (2016) COMAN: a web server for comprehensive metatranscriptomics analysis. *BMC genomics* 17(1), 622.

Oyserman, B.O., Noguera, D.R., del Rio, T.G., Tringe, S.G. and McMahon, K.D. (2015) Metatranscriptomic insights on gene expression and regulatory controls in *Candidatus Accumulibacter phosphatis*. *The ISME journal*.

Rapaport, F., Khanin, R., Liang, Y., Pirun, M., Krek, A., Zumbo, P., Mason, C.E., Socci, N.D. and Betel, D. (2013) Comprehensive evaluation of differential gene expression analysis methods for RNA-seq data. *Genome biology* 14(9), 3158.

Robinson, M.D., McCarthy, D.J. and Smyth, G.K. (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1), 139-140.

Robinson, M.D. and Oshlack, A. (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome biology* 11(3), R25.

Shi, Y.M., Tyson, G.W. and DeLong, E.F. (2009) Metatranscriptomics reveals unique microbial small RNAs in the ocean's water column. *nature* 459(7244), 266-U154.

Stark, L., Giersch, T. and Wünschiers, R. (2014) Efficiency of RNA extraction from selected bacteria in the context of biogas production and metatranscriptomics. *Anaerobe* 29, 85-90.

Turner, T.R., Ramakrishnan, K., Walshaw, J., Heavens, D., Alston, M., Swarbreck, D., Osbourn, A., Grant, A. and Poole, P.S. (2013) Comparative metatranscriptomics reveals kingdom level changes in the rhizosphere microbiome of plants. *The ISME journal* 7(12), 2248-2258.

Wagner, G.P., Kin, K. and Lynch, V.J. (2012) Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. *Theory in Biosciences* 131(4), 281-285.

Westreich, S.T., Korf, I., Mills, D.A. and Lemay, D.G. (2016) SAMSA: a comprehensive metatranscriptome analysis pipeline. *BMC bioinformatics* 17(1), 399.

Whelan, J.A., Russell, N.B. and Whelan, M.A. (2003) A method for the absolute quantification of cDNA using real-time PCR. *Journal of immunological methods* 278(1), 261-269.

Yang, Y., Jiang, X.T., Chai, B.L., Ma, L.P., Li, B., Zhang, A.N., Cole, J.R., Tiedje, J.M. and Zhang, T. (2016) ARGs-OAP: online analysis pipeline for antibiotic resistance genes detection from metagenomic data using an integrated structured ARG-database. *Bioinformatics* 32(15), 2346-2351.

Yu, K. and Zhang, T. (2012) Metagenomic and metatranscriptomic analysis of microbial community structure and gene expression of activated sludge. Plos One 7(5), e38183.

Figures and legends

Fig. 1: Schematic flow diagram for absolute quantification of metagenome and metatranscriptome to cell/volume level.

509 **Fig. 2** Flowchart of the simulation datasets generation and analyzing process to get
510 the differential expression genes.

511
512 **Fig. 3:** Absolute quantification of type level ARGs abundance and transcription in
513 FAS and NFAS. ARGs-carry hosts abundance and expression. * represents significant
514 difference (P -value < 0.05).

515

523

524

525 **Fig. 5** Samples were collected for foaming activated sludge at 08/04/2016 and
 526 non-foaming activated sludge at 25/04/2016 alongside the bioreactor at Shatin
 527 wastewater treatment plant.

528

529 **Table 1** Comparing relative quantification methods with AQMM on detection of
530 DEGs for simulated metatranscriptome data.

	# of genes Higher expression in B	No expression difference	# of genes Higher expression in A
Theoretical Ground Truth	28524	36572	0
RPKM+t-Test (P < 0.05)	16477	11558	37062
edgeR	18278	20778	26040
AQMM-5%-variation	28744.72 ± 143.53	35807.52 ± 48.08	543.77 ± 129.72
AQMM-10%-variation	28740.83 ± 298.43	35801 ± 188.31	554.17 ± 256.81
AQMM-20%-variation	28549.48 ± 1007.17	35941.86 ± 919.86	604.66 ± 654.76
AQMM-50%-variation	16673.93 ± 9394.27	47694.99 ± 9600.33	727.08 ± 1775.09

531

532

533 **Table 2:** Summary of sequencing outputs and absolute quantification of each sample
534 at cell level with AQMM.

Sample ID	Type	Library (bps data)	size Total clean DNA (ng/mL)	extracted and RNA sequenced	Estimated cells *	Estimated cells per mL *
DNA1	Foaming AS	8,567,524,200	49,140		1,541	6.11E+10
DNA2	Foaming AS	11,786,228,700	54,600		2,179	6.98E+10
DNA3	Foaming AS	10,108,576,800	58,380		1,919	7.66E+10
DNA4	Normal AS	8,755,895,700	57,974		1,425	6.52E+10
DNA5	Normal AS	9,196,724,100	66,752		1,541	7.73E+10

DNA6	Normal AS	11,185,847,400	75,194	1,957	9.09E+10
RNA1	Foaming AS	14,894,959,100	12,270	98,936	
RNA2	Foaming AS	13,598,855,700	12,710	99,744	
RNA3	Foaming AS	15,551,044,400	20,290	78,449	
RNA4	Normal AS	15,376,790,700	8,350	160,343	
RNA5	Normal AS	16,156,607,900	8,790	189,776	
RNA6	Normal AS	13,700,741,100	10,735	154,925	

*: Estimated sequenced cells for DNA libraries was using MicrobeCensus and for RNA libraries using AQMM. The assumption for AQMM was that per ml sample used for DNA and RNA extraction contained the same number of cells.

535