Peer Community In Evolutionary Biology

Abu Awad D, Roze D. 2017. **Effects of partial selfing on the equilibrium genetic variance, mutation load and inbreeding depression under stabilizing selection.** *bioRxiv*, 180000,
<https://doi.org/10.1101/180000>

An article reviewed and recommended by *Peer Community In Evolutionary Biology*.

<http://dx.doi.org/10.24072/pci.evolbiol.100041>

Effects of partial selfing on the equilibrium genetic variance, mutation load and inbreeding depression under stabilizing selection

Diala Abu Awad* and Denis Roze^{†,‡}

* INRA, UMR AGAP, 34060 Montpellier, France

† CNRS, UMI 3614 Evolutionary Biology and Ecology of Algae, 29688 Roscoff,
France

‡ Sorbonne Universités, UPMC Université Paris VI, 29688 Roscoff, France

Running title: Selfing and stabilizing selection

Keywords: adaptive landscape, epistasis, evolutionary quantitative genetics, multi-locus population genetics, self-fertilization

Address for correspondence:

Denis Roze

Station Biologique de Roscoff

Place Georges Teissier, CS90074

29688 Roscoff Cedex

France

Phone: (+33) 2 56 45 21 39

Fax: (+33) 2 98 29 23 24

email: roze@sb-roscoff.fr

ABSTRACT

This preprint has been reviewed and recommended by Peer Community In Evolutionary Biology (<http://dx.doi.org/10.24072/pci.evolbiol.100041>).

The mating system of a species is expected to have important effects on its genetic diversity. In this paper, we explore the effects of partial selfing on the equilibrium genetic variance V_g , mutation load L and inbreeding depression δ under stabilizing selection acting on an arbitrary number n of quantitative traits coded by biallelic loci with additive effects. Overall, our model predicts a decrease in the equilibrium genetic variance with increasing selfing rates; however, the relationship between self-fertilization and the variables of interest depends on the strength of associations between loci, and three different regimes are observed. When the U/n ratio is low (where U is the total haploid mutation rate on selected traits) and effective recombination rates are sufficiently high, genetic associations between loci are negligible and the genetic variance, mutation load and inbreeding depression are well predicted by approximations based on single-locus models. For higher values of U/n and/or lower effective recombination, moderate genetic associations generated by epistasis tend to increase V_g , L and δ , this regime being well predicted by approximations including the effects of pairwise associations between loci. For yet higher values of U/n and/or lower effective recombination, a different regime is reached under which the maintenance of coadapted gene complexes reduces V_g , L and δ . Simulations indicate that the values of V_g , L and δ are little affected by assumptions regarding the number of possible alleles per locus.

INTRODUCTION

Genetic diversity maintained within populations plays an important role in defining their adaptive potential (for a species to evolve, there must be heritable phenotypic variation on which selection can act). The ultimate source of this diversity is mutation, with a substantial proportion of new mutations being of a slightly deleterious nature (Eyre-Walker and Keightley, 2007): hence, a corollary to the maintenance of genetic diversity is the existence of a mutation load, defined as the reduction in mean fitness of a population relative to the fitness of an optimal genotype (Haldane, 1937). Furthermore, the fact that most deleterious alleles are partially recessive causes inbred offspring to have a lower fitness (on average) than outbred ones, as they tend to carry higher numbers of homozygous mutations (inbreeding depression, Charlesworth and Charlesworth, 1987).

By affecting the average degree of homozygosity of individuals and the efficiency of recombination between loci, the reproductive system of a species is expected to have an important influence on the effect of selection against deleterious alleles, and thus on the mutation load, inbreeding depression and level of diversity maintained within populations. One mating system that has received considerable attention is self-fertilization, a reproductive strategy occurring at various rates in an important proportion of plant and animal species (Jarne and Auld, 2006; Goodwillie et al., 2005; Igic and Kohn, 2006). Self-fertilization, and inbreeding in general, may have different effects on genetic polymorphisms depending on the strength of selection acting on them (Glémin, 2007). When directional selection against deleterious alleles is sufficiently strong relative to drift ($N_e s \gg 1$), the increased homozygosity caused by inbreeding is expected

to improve the efficiency of selection against those alleles (purging), reducing the mutation load and inbreeding depression (Lande and Schmske, 1985; Charlesworth et al., 1990). At the other extreme, polymorphism at neutrally-behaving loci ($N_e s \ll 1$) should also be lowered by inbreeding, as the effective population size is reduced by identity-by-descent within loci (Pollak, 1987) and by stronger interference effects between loci — background selection, hitchhiking (Nordborg, 1997; Glémin and Ronfort, 2013; Roze, 2016). In intermediate regimes ($N_e s \sim 1$), however, the reduction in N_e due to inbreeding may cause an increased frequency of deleterious alleles (because selection is less effective), which may explain the higher π_N/π_S ratio observed in various selfing species compared with their outcrossing relatives (Brandvain et al., 2013; Burgarella et al., 2015, and other references listed in Table 1 of Hartfield, 2015).

Most classical results on the effects of selfing on genetic diversity, mutation load and inbreeding depression are based on single-locus models, and thus neglect the effects of linkage disequilibria and other forms of genetic associations among loci. Previous analytical and simulation models showed that intermediate selfing rates generate correlations in homozygosity between loci, termed “identity disequilibria” (Weir and Cockerham, 1973; Vitalis and Couvet, 2001), which tend to reduce the efficiency of purging when deleterious alleles are partially or fully recessive (an effect called “selective interference” by Lande et al., 1994). When the number of highly recessive mutations segregating within genomes is sufficiently high, these correlations in homozygosity may entirely suppress purging unless the selfing rate exceeds a given threshold (Lande et al., 1994; Scofield and Schultz, 2006; Kelly, 2007; Roze, 2015). Linkage disequilibrium corresponds to another form of association between loci that may also affect the efficiency of selection: in particular, selection may be strongly limited by Hill-Robertson

effects in highly selfing populations, due to the fact that selfing reduces the efficiency of recombination between loci — recombination having no effect when it occurs in homozygous individuals (Kamran-Disfani and Agrawal, 2014; Hartfield and Glémin, 2016). Epistatic interactions represent another possible source of linkage disequilibrium between selected loci. Charlesworth et al. (1991) considered a model in which epistasis between deleterious alleles is fixed and synergistic (the effects of mutations alone being smaller than when combined with others), and showed that the effect of the selfing rate on the load and inbreeding depression may be non-monotonic under this form of epistasis, with an increase in both variables above a (high) self-fertilization threshold. However, although models with fixed epistasis have lead to important insights, epistatic interactions are known to vary across pairs of loci, and this variation may have important evolutionary consequences (Phillips et al., 2000; Martin et al., 2007). Interestingly, several aspects of the complexity of epistatic interactions (such as possible compensatory effects between deleterious alleles, *i.e.*, reciprocal sign epistasis) are captured by models of stabilizing selection acting on quantitative traits, such as Fisher’s geometric model (Fisher, 1930). Furthermore, the distributions of epistasis generated by this type of model seem compatible with our empirical knowledge on epistasis (Martin et al., 2007).

Only a few models have explored the effect of self-fertilization on genetic variance for quantitative traits at equilibrium between mutation and stabilizing selection. Modeling a quantitative trait coded by additive loci, Wright (1951) showed that, in the absence of selection, the genetic variance for the trait is increased by a factor $1 + F$ (where F is the inbreeding coefficient), due to the increased homozygosity of the underlying loci. Selection will oppose this increase in variance, however, by elim-

inating genotypes that are too far from the optimum (purging). Stabilizing selection is also known to generate positive linkage disequilibrium between alleles at different loci having opposite effects on the trait (Bulmer, 1971), the immediate consequence of which is to reduce the genetic variance. These linkage disequilibria should also affect the efficiency of selection at each locus, and thus indirectly affect the genetic variance. Lande (1977) proposed a model of stabilizing selection acting on a single trait coded by additive loci in a partially selfing population, in which a Gaussian distribution of allelic effects is assumed to be maintained at each locus. He found that, as the increase in variance due to homozygosity is exactly compensated by the effect of purging, and the decrease in variance caused by linkage disequilibria is exactly compensated by the decreased efficiency of selection acting at each locus due to these linkage disequilibria, overall the equilibrium genetic variance is not affected by the selfing rate of the population. More recently, Lande and Porcher (2015) extended this model to multiple selected traits, and used a method developed by Kelly (2007) to take into account the effects of correlations in homozygosity across loci by splitting the population into selfing age classes (corresponding to classes of individuals having the same history of inbreeding), while assuming a Gaussian distribution of allelic effects at each locus within each class. Numerical iterations of the model showed that above a threshold selfing rate, a different regime is reached, in which strong compensatory associations between alleles at different loci reduce the genetic variance and may generate outbreeding depression (*i.e.*, lower fitness of outcrossed offspring relative to selfed offspring).

The hypothesis made by Lande (1977) and Lande and Porcher (2015) of a Gaussian distribution of allelic effects maintained at each locus (either in the whole population or in each selfing age class) has been criticized on the grounds that it

implicitly assumes an unrealistically high mutation rate per locus and/or very weak fitness effects of mutations (Turelli, 1984). Lande and Porcher (2015) also considered an infinitesimal model (in which traits are coded by an infinite number of loci, selection having a negligible effect on allele frequencies at each locus), and showed that a similar threshold pattern emerges, although the effect of selfing on the genetic variance and inbreeding depression above the threshold differs between the two models (in particular, outbreeding depression is not observed in the infinitesimal model). However, the effect of selfing on the genetic variance of quantitative traits under more general assumptions regarding the strength of selection at the underlying loci remains unclear.

In this paper, we introduce partial self-fertilization into previous models of stabilizing selection acting on quantitative traits coded by biallelic loci (Latter, 1960; Bulmer, 1972; Barton, 1986, 1989; Turelli and Barton, 1990; Roze and Blanckaert, 2014). Charlesworth and Charlesworth (1995) had extended such models to take complete selfing into account, and found that the genetic variance under stabilizing selection should be lower under complete selfing than under random mating. The present paper generalizes these results to arbitrary selfing rates and multiple selected traits. Assuming additive effects of alleles on phenotypes (no dominance or epistasis on phenotypic traits), we develop approximations incorporating the effects of pairwise associations between selected loci, and compare these approximations with results from individual-based simulations. Our results indicate that different regimes are possible depending on the effect of genetic associations on the genetic variance, this effect increasing as the overall mutation rate U and selfing rate σ increase, and decreasing as the genome map length R , mean fitness effect of mutations $\bar{s}$ and number of selected traits n increase. When U and σ are sufficiently low and R , $\bar{s}$ and n sufficiently high, the effect of associ-

ations is negligible and the mutation load, inbreeding depression and genetic variance are well predicted by classical expressions ignoring associations. As the strength of associations increases, a second regime is entered in which the overall effect of associations is to reduce purging, thereby increasing the genetic variance, mutation load and inbreeding depression; this “weak association” regime is generally well predicted by our approximations which include the effects of pairwise associations between loci. For yet higher U , σ and/or lower R , $\bar{s}$ or n , a third regime is reached in which strong associations between loci caused by compensatory effects among mutations reduce the genetic variance, load and inbreeding depression. Although our approximations break down in this “strong association” regime, the approximation proposed by Charlesworth and Charlesworth (1995) provides accurate results under complete selfing when the mutation rate is sufficiently high and the mean fitness effect of mutations sufficiently low.

MODEL

Genotype-phenotype map. The parameters and variables of our model are summarized in Table 1. We consider a diploid population of size N with discrete generations. Offspring are produced by self-fertilization with probability σ , and by random union of gametes with probability $1 - \sigma$. The fitness of an organism represents its overall relative fecundity (assumed very large for all individuals), and depends on the values of n quantitative phenotypic traits under stabilizing selection. In the following we use subscripts $\alpha, \beta, \gamma \dots$ to denote phenotypic traits, while subscripts $i, j, k \dots$ denote loci. The value of trait α in a given individual is denoted z_α , and can be decomposed into

164 a genetic and an environmental component:

$$z_{\alpha} = g_{\alpha} + e_{\alpha} \quad (1)$$

165 where the environmental component e_{α} is sampled from a Gaussian distribution with
 166 mean zero and variance V_e (the same for all traits). The genetic component g_{α} (“breed-
 167 ing value”) is controlled by a large number of biallelic loci with additive effects. The
 168 two alleles at each locus are denoted 0 and 1, while X_i^M and X_i^P are defined as indicator
 169 variables that equal zero if the individual carries allele 0 at locus i on its maternally
 170 (X_i^M) or paternally (X_i^P) inherited chromosome, while they equal 1 if allele 1 is present.
 171 We also assume that $g_{\alpha} = 0$ in an individual homozygous for allele 0 at all loci, so
 172 that:

$$g_{\alpha} = \sum_{i=1}^{\ell} r_{\alpha i} (X_i^M + X_i^P) \quad (2)$$

173 where ℓ is the number of loci affecting phenotypic traits, and $r_{\alpha i}$ the effect on phenotype
 174 α of changing the allelic state of one gene copy at locus i from 0 to 1 (note that $r_{\alpha i}$
 175 may be negative).

176 Following Chevin et al. (2010), Lourenço et al. (2011) and Roze and Blanckaert
 177 (2014), a parameter m measures the degree of pleiotropy of mutations: each locus
 178 affects a subset of m phenotypic traits, sampled randomly (and independently for
 179 each locus) among the n traits. Therefore, $m = 1$ means that each locus affects a
 180 single trait, while $m = n$ corresponds to full pleiotropy (each locus affecting all traits),
 181 as in Fisher’s geometric model (Fisher, 1930). We assume that the distribution of
 182 mutational effects $r_{\alpha i}$ over all loci affecting trait α has average zero and variance a^2
 183 (the same for all traits); if locus i does not affect trait α , then $r_{\alpha i} = 0$. For simplicity, we
 184 consider a fully isotropic model with no mutational covariance between traits. Finally,

u denotes the mutation rate from allele 0 to allele 1 and from allele 1 to allele 0 at each locus, while $U = u\ell$ is the haploid mutation rate over all loci (per generation).

From the previous definitions, and assuming that population size is sufficiently large, mean trait values are given by:

$$\overline{z_\alpha} \approx \overline{g_\alpha} = 2 \sum_{i=1}^{\ell} r_{\alpha i} p_i \quad (3)$$

where p_i is the frequency of allele 1 at locus i . As we assume no $G \times E$ interaction, the variance in trait α is given by:

$$V_{p,\alpha} = V_{g,\alpha} + V_e \quad (4)$$

where $V_{g,\alpha}$ is the variance in g_α (genetic variance). In the next subsection, we show how $V_{g,\alpha}$ can be expressed in terms of genetic associations within and between loci.

Genetic associations and decomposition of the genetic variance. Genetic associations are defined as in Kirkpatrick et al. (2002). In particular, the centered variables $\zeta_{i,\emptyset}$ and $\zeta_{\emptyset,i}$ are defined as:

$$\zeta_{i,\emptyset} = X_i^M - p_i, \quad \zeta_{\emptyset,i} = X_i^P - p_i. \quad (5)$$

Furthermore, products of $\zeta_{i,\emptyset}$, $\zeta_{\emptyset,i}$ variables are denoted:

$$\zeta_{\mathbb{U},\mathbb{V}} = \left(\prod_{i \in \mathbb{U}} \zeta_{i,\emptyset} \right) \left(\prod_{j \in \mathbb{V}} \zeta_{\emptyset,j} \right) \quad (6)$$

where $\mathbb{U}$ and $\mathbb{V}$ represent sets of loci. For example, for $\mathbb{U} = \mathbb{V} = \{i\}$, we have:

$$\zeta_{i,i} = (X_i^M - p_i) (X_i^P - p_i) \quad (7)$$

while for $\mathbb{U} = \{i, j\}$ and $\mathbb{V} = \{i\}$:

$$\zeta_{ij,i} = (X_i^M - p_i) (X_j^M - p_j) (X_i^P - p_i). \quad (8)$$

Finally, genetic associations $D_{\mathbb{U},\mathbb{V}}$ are defined as averages of $\zeta_{\mathbb{U},\mathbb{V}}$ variables over all individuals:

$$D_{\mathbb{U},\mathbb{V}} = \mathbb{E}[\zeta_{\mathbb{U},\mathbb{V}}] \quad (9)$$

where $\mathbb{E}$ stands for the average over all individuals in the population. We also define $\tilde{D}_{\mathbb{U},\mathbb{V}}$ as $(D_{\mathbb{U},\mathbb{V}} + D_{\mathbb{V},\mathbb{U}})/2$, and write $\tilde{D}_{\mathbb{U},\emptyset}$ as $\tilde{D}_{\mathbb{U}}$ (for simplicity). In particular, $D_{i,i}$ is a measure of excess homozygosity (due, for example, to non-random mating) at locus i ($D_{i,i} = 0$ at Hardy-Weinberg equilibrium). As shown in Supplementary File S1, it can be written as $D_{i,i} = F p_i q_i$, where F is the inbreeding coefficient (probability of identity by descent between two alleles present at the same locus in the same individual). The association $\tilde{D}_{ij}$ corresponds to the linkage disequilibrium between loci i and j (association between alleles present on the same haplotype), while $\tilde{D}_{i,j}$ is the association between alleles at loci i and j present on different haplotypes of the same individual. We will see that the association $D_{ij,ij}$ also appears in the computations, and can be expressed as $\phi_{ij} p_i q_i p_j q_j$, where ϕ_{ij} is the probability of joint identity by descent at loci i and j . The quantities ϕ_{ij} and F enter into the definition of the identity disequilibrium between loci i and j , given by $G_{ij} = \phi_{ij} - F^2$ (Weir and Cockerham, 1973), which will appear in some of our results.

From these definitions, and using equations 2 and 3, the genetic variance for trait α can be written as:

$$\begin{aligned} V_{g,\alpha} &= \mathbb{E}[(g_\alpha - \bar{g}_\alpha)^2] \\ &= \mathbb{E}\left[\sum_{i,j} r_{\alpha i} r_{\alpha j} (\zeta_{i,\emptyset} + \zeta_{\emptyset,i}) (\zeta_{j,\emptyset} + \zeta_{\emptyset,j})\right] \end{aligned} \quad (10)$$

where the last sum is over all i and j (including $i = j$). Using the fact that $\tilde{D}_{ii} = p_i q_i$,

one obtains from equation 9:

$$V_{g,\alpha} = 2 \sum_{i=1}^{\ell} r_{\alpha i}^2 (p_i q_i + D_{i,i}) + 2 \sum_{i,j \neq i} r_{\alpha i} r_{\alpha j} (\tilde{D}_{ij} + \tilde{D}_{i,j}). \quad (11)$$

Following previous usage (e.g., Bulmer, 1985), we will call *genic variance* (denoted $V_{g,\alpha}^0$) the quantity $2 \sum_{i=1}^{\ell} r_{\alpha i}^2 p_i q_i$, corresponding to the genetic variance in a population with the same allele frequencies, but in the absence of genetic association (within and between loci). As shown by equation 11, excess homozygosity tends to increase the genetic variance through the term in $D_{i,i}$. The second term of equation 11 (the effect of between-locus associations) tends to be negative under stabilizing selection, since the allele increasing the value of trait α at locus i tends to be associated with the allele decreasing its value at locus j (e.g., Bulmer, 1971, 1974; Lande, 1976; Turelli and Barton, 1990). However, below we show that that excess homozygosity and associations between loci also affect equilibrium allele frequencies, and thus the genic variance.

Fitness function. Most of the results derived in this paper assume an isotropic, Gaussian fitness function, the fitness of an individual being given by:

$$W = \exp \left[-\frac{d^2}{2\omega^2} \right] \quad (12)$$

where ω^2 measures the strength of selection, and $d = \sqrt{\sum_{\alpha=1}^n z_{\alpha}^2}$ is the Euclidean distance (in phenotypic space) between the individual's phenotype and the optimum, which we assume is located at $\mathbf{z} = (0, 0, \dots, 0)$. From equation 12, the fitness associated with a given genotype (obtained by averaging over environmental effects) is also Gaussian, and given by:

$$W_g = W_{g,\max} \exp \left[-\frac{d_g^2}{2V_s} \right] \quad (13)$$

with $V_s = \omega^2 + V_e$, $W_{g,\max} = (\omega^2/V_s)^{n/2}$ (the mean fitness of an optimal genotype), and
 $d_g = \sqrt{\sum_{\alpha=1}^n g_{\alpha}^2}$ (the Euclidean distance between the breeding value of the individual
and the optimum). Under our mutational model, the mean reduction in $\log W_g$ caused
by a heterozygous mutation present in an optimal genotype is:

$$\bar{s} = \frac{ma^2}{2V_s} \quad (14)$$

(e.g., Martin and Lenormand, 2006b). Under our assumption of additivity of pheno-
typic effects it is easy to show that the reduction in $\log W_g$ caused by a homozygous
deleterious allele (in an optimal genotype) is four times the reduction caused by the
same allele in the heterozygous state. Provided that most mutations have weak fitness
effects (so that $\log(1-s) \approx -s$), the dominance coefficient of deleterious alleles is
thus close to 0.25 at the fitness optimum (see Manna et al., 2011 for more general
results on dominance in Fisher’s geometric model).

The effect of the shape of the fitness peak will be explored using a generalized
version of equation 13 (e.g., Martin and Lenormand, 2006a; Tenaillon et al., 2007):

$$W_g = W_{g,\max} \exp \left[- \left(\frac{d_g}{\sqrt{2V_s}} \right)^Q \right]. \quad (15)$$

Gaussian fitness (equation 13) thus corresponds to $Q = 2$, while the fitness peak is
sharper around the optimum when $Q < 2$, and flatter when $Q > 2$. Importantly,
 Q affects the average dominance coefficient of deleterious alleles, making them more
dominant for $Q < 2$ and more recessive for $Q > 2$ (Manna et al., 2011), as well as
the average epistasis (on fitness) between alleles, positive for $Q < 2$, and negative
for $Q > 2$ (Gros et al., 2009). Approximations for the mutation load and inbreeding
depression can be derived for $Q \neq 2$ as long as the distribution of breeding values in
the population is approximately Gaussian.

Individual-based simulations. In order to verify the analytical results obtained, individual-based simulations were run using a C++ program described in Supplementary File S5 (and available from Dryad), in which the genome of each individual consists of two copies of a linear chromosome carrying ℓ equidistant biallelic loci affecting the n traits under selection. Another version of the program was used to consider a different genetic architecture, under which an infinite number of alleles are possible at each locus (see Supplementary File S5).

RESULTS

Neglecting associations between loci. In the following section we show that genetic associations between loci may be neglected when the haploid genomic mutation rate U is sufficiently low. In this case, equation 11 simplifies to:

$$V_{g,\alpha} \approx 2 \sum_{i=1}^{\ell} r_{\alpha i}^2 (p_i q_i + D_{i,i}). \quad (16)$$

Expressions for $p_i q_i$ and $D_{i,i}$ at equilibrium, assuming weak selection ($V_{g,\alpha} \ll V_s$) and neglecting associations among loci are derived in Supplementary File S1. To leading order, $D_{i,i} \approx F p_i q_i$ where $F = \sigma / (2 - \sigma)$ is the inbreeding coefficient. Neglecting associations between loci and assuming that mean phenotypes are at the optimum ($\bar{g}_\alpha = 0$), the effect of selection on $p_i q_i$ is given by:

$$\Delta_{\text{sel}} p_i q_i \approx -s_i (1 + 3F) (1 - 2p_i)^2 p_i q_i \quad (17)$$

where $s_i = \sum_{\alpha=1}^n r_{\alpha i}^2 / (2V_s)$ is the heterozygous effect of a mutation at locus i on log fitness in an optimal genotype. Furthermore, because mutation changes p_i to $p_i (1 - u) + u (1 - p_i)$, the change in $p_i q_i$ due to mutation is (to the first order in u):

$$\Delta_{\text{mut}} p_i q_i \approx u (1 - 2p_i)^2. \quad (18)$$

278 In regimes where genetic drift can be neglected, $\Delta_{\text{sel}} p_i q_i = -\Delta_{\text{mut}} p_i q_i$ at mutation-
279 selection balance, leading to either $p_i = 1/2$ or:

$$s_i p_i q_i \approx \frac{u}{1 + 3F}, \quad (19)$$

280 in agreement with results of previous biallelic models under random mating (e.g.,
281 Bulmer, 1972; Barton, 1986). A stability analysis indicates that the equilibrium given
282 by equation 19 is stable when $p_i q_i < 1/4$ (that is, when $s_i (1 + 3F) > 4u$), otherwise
283 $p_i = 1/2$ is stable. When all loci are at the equilibrium where $p_i q_i < 1/4$, summing
284 both sides of equation 19 over i yields, using $s_i = \sum_{\alpha=1}^n r_{\alpha i}^2 / (2V_s)$:

$$\sum_{\alpha=1}^n V_{g,\alpha}^0 \approx \frac{4V_s U}{1 + 3F} \quad (20)$$

285 where again $V_{g,\alpha}^0 = 2 \sum_{\alpha=1}^n r_{\alpha i}^2 p_i q_i$ is the genic variance. By symmetry the equilibrium
286 genic variance should be the same for all traits, and thus:

$$V_{g,\alpha}^0 \approx \frac{4V_s U}{n(1 + 3F)} = \frac{2V_s U}{n} \frac{2 - \sigma}{1 + \sigma}. \quad (21)$$

287 From equations 16 and 21, the equilibrium genetic variance is:

$$V_{g,\alpha} \approx V_{g,\alpha}^0 (1 + F) \approx \frac{4V_s U}{n(1 + \sigma)}. \quad (22)$$

288 When $\sigma = 0$ and $n = 1$, equation 22 is equivalent to the result of previous biallelic
289 models (e.g., Latter, 1960; Bulmer, 1972) and to Turelli's house-of-cards approximation
290 (Turelli, 1984).

291 Assuming that the variance in log-fitness is small, mean fitness is approximately
292 $\bar{W} \approx e^{\overline{\log W_g}}$. Defining the mutation load L as the reduction in $\bar{W}$ relative to the average
293 fitness of an optimal genotype, one obtains from equation 13:

$$\begin{aligned} L &= 1 - \frac{\bar{W}}{W_{g,\text{max}}} \\ &\approx 1 - \exp \left[-\frac{\sum_{\alpha=1}^n V_{g,\alpha}}{2V_s} \right]. \end{aligned} \quad (23)$$

Equations 22 and 23 yield:

$$L \approx 1 - \exp \left[-\frac{2U}{1 + \sigma} \right]. \quad (24)$$

Inbreeding depression δ measures the mean fitness of selfed offspring, relative to the mean fitness of outcrossed offspring. Under the same assumptions, it is given by:

$$\begin{aligned} \delta &= 1 - \frac{\overline{W}_{\text{self}}}{\overline{W}_{\text{out}}} \\ &\approx 1 - \exp \left[-\frac{\sum_{\alpha=1}^n (V_{g,\alpha}^{\text{self}} - V_{g,\alpha}^{\text{out}})}{2V_s} \right] \end{aligned} \quad (25)$$

where $V_{g,\alpha}^{\text{self}}$ and $V_{g,\alpha}^{\text{out}}$ are the genetic variances for trait α among selfed and outcrossed offspring, respectively (e.g., Lande and Schemske, 1985). The intralocus association $D_{i,i}$ among selfed offspring is $D_{i,i}^{\text{self}} = \frac{1}{2} (p_i q_i + D_{i,i})$ and therefore $V_{g,\alpha}^{\text{self}} = V_{g,\alpha}^0 \left[1 + \frac{1}{2} (1 + F) \right]$, while $V_{g,\alpha}^{\text{out}} = V_{g,\alpha}^0$, yielding (using equation 21):

$$\delta \approx 1 - \exp \left[-\frac{\sum_{\alpha=1}^n V_{g,\alpha}}{4V_s} \right] \approx 1 - \exp \left[-\frac{U}{1 + \sigma} \right]. \quad (26)$$

Equations 24 and 26 are equivalent to the classical expressions obtained for the load and inbreeding depression at mutation-selection balance when the dominance coefficient h of deleterious alleles is set to 0.25 (e.g., Charlesworth and Charlesworth, 1987), in agreement with the fact that $h \approx 0.25$ under Gaussian stabilizing selection when mutations have additive effects on phenotypes (see previous section).

Figure 1 shows that the mutation load is well predicted by equation 24 when $N_e \bar{s}$ is sufficiently large (for $U = 0.1$ and $n = 50$), and generally decreases as selfing increases — results for different numbers of loci ℓ are shown in Supplementary Figure S1, while Supplementary Figures S2 and S3 show that the genetic variance and inbreeding depression follow similar patterns. Drift may have significant effects on genetic variation, however, when $N_e \bar{s}$ is ≈ 1 or lower. Following Bulmer (1972), a

diffusion model can be used to compute the expected value of $p_i q_i$ under selection, mutation and drift, provided that the effects of associations between loci are neglected. As explained in Supplementary File S2, the result can then be integrated over the distribution of s_i across loci to obtain the equilibrium genetic variance, inbreeding depression and mutation load. Figures 1 and S1 – S3 show that drift increases V_g , L and δ in regimes where $p_i q_i$ tends to stay small at most loci at the deterministic equilibrium ($\bar{s} = 10^{-2}, 10^{-3}$ in Figure 1), and has the opposite effect in regimes where $p_i q_i$ is high ($\bar{s} = 10^{-4}$ in Figure 1). Simple approximations can be obtained when the effect of selection is negligible at most loci (see Supplementary File S2), which provide accurate predictions when $N_e \bar{s}$ is sufficiently low, or when $\bar{s} \ll u$ so that $p_i = 1/2$ at most loci at the deterministic equilibrium (Figures S1 – S4). In this mutation-drift regime, V_g , L and δ are nearly independent of σ when $N_e u \ll 1$ (the increase in variance caused by excess homozygosity being exactly compensated by the reduction in variance caused by the lower effective population size), or increase with σ for larger values of $N_e u$. The discrepancies between analytical and simulation results observed in Figure 1 at high selfing rates are partly due to the reduction in effective population size N_e caused by background selection, which is not accounted for in the diffusion model. An estimation of N_e using the equilibrium diversity at a neutral locus (with an infinite number of possible alleles) at the mid-point of the chromosome (as in Roze, 2016) yielded an N_e of approximately 740, 300 and 200 for $\bar{s} = 10^{-4}$, 10^{-3} and 10^{-2} (respectively) for $N = 5,000$ and $\sigma = 1$ (right-most points in Figure 1B). Replacing N by $N_e (1 + F)$ in the diffusion model provides predictions that closely match the simulation results for $\bar{s} = 10^{-4}$ and 10^{-3} , suggesting that the initial discrepancy was indeed caused by background selection reducing N_e (results not shown). However, for

$\bar{s} = 10^{-2}$, the diffusion model still performs poorly despite the corrected N_e . This implies that the discrepancy between analytical and simulation results is more likely due to interactions among loci, and possibly also to deviations of mean phenotypes from the optimum caused by genetic drift (that are not taken into account in the analysis).

In Supplementary File S3, we derive expressions for the genetic variance, mutation load and inbreeding depression (for both the mutation-selection and the mutation-drift regimes) under the generalized fitness function given by equation 15. In the mutation-selection regime ($s_i \gg 1/N_e$, u at most loci), one obtains:

$$\frac{V_g}{V_s} \approx \left[\frac{4U}{Q(1+\sigma)} \frac{\Gamma\left(\frac{n}{2}\right)}{\Gamma\left(\frac{Q+n}{2}\right)} \right]^{\frac{2}{Q}} \quad (27)$$

(where Γ is Euler's Gamma function), while

$$L \approx 1 - \exp \left[-\frac{4U}{Q(1+\sigma)} \right] \quad (28)$$

$$\delta \approx 1 - \exp \left[-\frac{4U}{Q(1+\sigma)} \left[\left(\frac{3-\sigma}{2} \right)^{\frac{Q}{2}} - \left(\frac{2-\sigma}{2} \right)^{\frac{Q}{2}} \right] \right], \quad (29)$$

these equations being equivalent to equations 22, 24 and 26 when $Q = 2$. As shown by Figure 2, equations 27 – 29 provide good predictions of the simulation results when the population size and number of loci are sufficiently large (and selfing is not too high). As Q increases, the fitness peak becomes flatter around the optimum, and the equilibrium genetic variance increases (Figure 2B). However, despite increasing the genetic variance, higher values of Q lead to lower mutation loads due to the fact that deleterious alleles are more often eliminated when present in combination within the same genome: this corresponds to the classical result that negative epistasis reduces the mutation load in sexually reproducing populations (e.g., Kimura and Maruyama, 1966; Kondrashov and Crow, 1988). Indeed, the average epistasis between deleterious

alleles equals zero for $Q = 2$, but becomes negative when $Q > 2$, and positive when $Q < 2$ (Gros et al., 2009). By contrast, inbreeding depression is less affected by Q , δ slightly increasing or decreasing as Q increases, depending on the selfing rate.

Figure 3 shows the effects of the parameters m and n (for $Q = 2$). The degree of pleiotropy m of mutations affects their distribution of fitness effects (e.g., Lourenço et al., 2011). In Figure 3A, $\bar{s}$ is kept constant by decreasing the variance of mutational effects a^2 as m increases (see equation 14). Increasing m (while keeping $\bar{s}$ constant) decreases the variance in fitness effects of mutations: indeed, one can show that the variance of mutational effects on log fitness (at the optimum) is given by $2\bar{s}^2/m$. Figure 3A shows that $m = 5$ and $m = 50$ yield very close results when $\bar{s} = 10^{-4}$, as selection has a negligible effect at most loci for both values of m (for the parameter values used here), and the genetic variance does not depend on m at mutation-drift equilibrium (see equation B8 in Supplementary File S2). When $\bar{s} = 10^{-2}$, most loci are at mutation-selection balance ($s_i \gg 1/N, u$) for both $m = 5$ and $m = 50$, and the genetic variance is again not affected by m (see equation 22). Slightly different results are obtained for $m = 1$, due to the higher variance in fitness effects of mutations, causing a larger fraction of loci to be substantially affected by both selection and drift (this effect being captured by the diffusion model). Similarly, the effect of m is more pronounced when $\bar{s} = 10^{-2}$ and $\sigma = 1$, as N_e is greatly reduced by background selection when selfing is high, causing higher proportions of loci to be substantially affected by drift.

As shown by Figure 3B, the number of selected traits n has only little effect on the load in the mutation-drift regime ($\bar{s} = 10^{-4}$), in agreement with equation B9 in Supplementary File S2. However, while the diffusion model also predicts very little effect of n in the mutation-selection regime ($\bar{s} = 10^{-2}$), larger effects are observed in the

simulations, with larger deviations from the analytical predictions (and higher load) for lower values of n . These deviations are caused by associations between loci (which are neglected in equation 24 and in the diffusion model). In the next subsection, we show that the relative effect of these associations is indeed stronger when n is lower, and derive an approximation including the effect of pairwise genetic associations that better matches the simulation results.

Effects of associations between loci. In Supplementary File S1, we derive approximations for the effects of associations between pairs of loci on the genetic variance at mutation-selection balance, under a Gaussian fitness function ($Q = 2$). For this, we assume that these associations remain weak, and neglect the effects of all associations involving more than two loci. As shown by equation 11, associations $\tilde{D}_{ij}$, $\tilde{D}_{i,j}$ (between alleles at loci i and j , either on the same or on different haplotypes) directly affect the genetic variance. At equilibrium, these associations are approximately given by:

$$\tilde{D}_{ij} \approx \frac{1}{1-F} \left(\frac{1}{\rho_{ij}} + 2F \right) \Delta_{\text{sel}} \tilde{D}_{ij} \quad (30)$$

$$\tilde{D}_{i,j} \approx \frac{F}{1-F} \left(\frac{1}{\rho_{ij}} + 2 \right) \Delta_{\text{sel}} \tilde{D}_{i,j}, \quad (31)$$

where again $F = \sigma / (2 - \sigma)$, ρ_{ij} is the recombination rate between loci i and j , and $\Delta_{\text{sel}} \tilde{D}_{ij}$ is the change in $\tilde{D}_{ij}$ and $\tilde{D}_{i,j}$ due to selection:

$$\Delta_{\text{sel}} \tilde{D}_{ij} \approx - \frac{\sum_{\alpha=1}^n r_{\alpha i} r_{\alpha j}}{V_s} [(1+F)^2 + G_{ij}] p_i q_i p_j q_j. \quad (32)$$

The term G_{ij} in equation 32 represents the identity disequilibrium between loci i and j : the correlation in identity by descent between loci, generating a correlation in homozygosity (Weir and Cockerham, 1973, Supplementary File S1). Equation 32 shows that stabilizing selection generates a positive association between alleles at loci

401 i and j that tends to displace phenotypes in opposite directions (allele 1 with allele
402 1, and allele 0 with allele 0 if $\sum_{\alpha=1}^n r_{\alpha i} r_{\alpha j} < 0$): the effect of the deleterious allele at
403 locus i is then partially compensated by its associated allele at locus j (e.g., Bulmer,
404 1974; Lande, 1976; Turelli and Barton, 1990). This effect of selection is strengthened
405 by homozygosity (and correlations in homozygosity between loci) caused by selfing.
406 As may be seen from equations 30 and 31, $\tilde{D}_{i,j} \approx F \tilde{D}_{ij}$ when loci are tightly linked
407 ($\rho_{ij} \ll 1$), as expected from separation of timescales arguments (e.g., Nordborg, 1997;
408 Roze, 2016). However, our approximations diverge as recombination tends to zero (or
409 as the selfing rate tends to 1), due to the assumption that genetic associations remain
410 weak.

411 We show in Supplementary File S1 how $\tilde{D}_{ij}$ and $\tilde{D}_{i,j}$ can be summed over all
412 pairs of loci in order to compute their overall direct effect on the genetic variance (sec-
413 ond term of equation 11). These associations depend on recombination rates through
414 the terms in $1/\rho_{ij}$ in equations 30 and 31, and also through the identity disequilibrium
415 G_{ij} in equation 32. However, because G_{ij} only weakly depends on the recombination
416 rate, its average over all pairs of loci is often very close to the value obtained under
417 free recombination, provided that the genome map length is not too small (see Sup-
418 plementary Figure S5). In the following, we thus approximate G_{ij} by its expression
419 for freely recombining loci, denoted G :

$$G = \frac{4\sigma(1-\sigma)}{(4-\sigma)(2-\sigma)^2}. \quad (33)$$

420 By contrast, linkage has more effect on the average of $1/\rho_{ij}$ over all pairs of loci,
421 corresponding to the inverse of the harmonic mean recombination rate between all
422 pairs of loci (denoted ρ_H thereafter). Assuming that the number of loci is large,

one obtains for the direct effect of linkage disequilibria on the genetic variance (see Supplementary File S1):

$$2 \sum_{i,j \neq i} r_{\alpha i} r_{\alpha j} (\tilde{D}_{ij} + \tilde{D}_{i,j}) \approx -\frac{2}{V_s} \frac{2 + \sigma}{(1 - \sigma)(2 - \sigma)(4 - \sigma)} \left[\frac{1}{\rho_H} + 2\sigma \right] (V_{g,\alpha}^0)^2 \quad (34)$$

where the genic variance $V_{g,\alpha}^0$ may be replaced by its expression to leading order, given by equation 21. Equation 34 shows that the immediate effect of associations between alleles with compensatory phenotypic effects is to reduce the genetic variance (since this term is negative). The fraction in equation 34 is an increasing function of σ , which implies that self-fertilization increases the strength of associations, thus decreasing $V_{g,\alpha}$. However, because the genic variance is expected to decrease with σ (equation 21), the direct effect of linkage disequilibria on $V_{g,\alpha}$ may remain approximately constant (or even slightly decrease) as σ increases from zero.

Associations between loci do not only affect $V_{g,\alpha}$ through equation 34, however, but also affect the equilibrium allele frequencies and the excess homozygosity $D_{i,i}$ at each locus. The effect on $D_{i,i}$ is mainly driven by identity disequilibria: indeed, neglecting associations between 3 or more loci, one obtains (see Supplementary File S1):

$$D_{i,i} \approx F \left[1 - 2 \sum_{j \neq i} s_j G_{ij} p_j q_j \right] p_i q_i. \quad (35)$$

Equation 35 is equivalent to equation 5 in Roze (2015) (which is expressed to the first order in p_j), as can be noted by replacing s and h in Roze (2015) by $4s_j$ and $1/4$. It shows that identity disequilibria reduce the excess homozygosity at each locus: this is due to the fact that homozygotes at locus i are more likely to be also homozygous at locus j , while homozygotes at locus j have a lower fitness than heterozygotes when deleterious alleles are partially recessive. Identity disequilibria thus tend to reduce

the genetic variance through this effect on $D_{i,i}$, by an amount corresponding to the sum of the term in G_{ij} in equation 35 over all pairs of loci. Approximating G_{ij} by its expression for freely recombining loci (equation 33), one obtains that this effect reduces $V_{g,\alpha}$ by approximately $-n F G (V_{g,\alpha}^0)^2 / (2V_s)$, where again $V_{g,\alpha}^0$ may be replaced by the expression given by equation 21, to leading order (see Supplementary File S1).

Finally, associations between loci affect equilibrium allele frequencies ($p_i q_i$) at each locus. As shown in Supplementary File S1, both the linkage disequilibria generated by epistasis and the identity disequilibria caused by partial selfing reduce the efficiency of purging, thereby increasing $p_i q_i$ and thus the genic variance. Indeed, an expression for the effect of selection on $p_i q_i$ that includes the effects of pairwise associations is, to leading order:

$$\begin{aligned} \Delta_{\text{sel}} p_i q_i \approx & -s_i (1 - 2p_i)^2 \left[p_i q_i + 3D_{i,i} - 6(1 + F) \sum_{j \neq i} s_j G_{ij} p_i q_i p_j q_j \right] \\ & + 2(1 - 2p_i)^2 \sum_{j \neq i} a_{ij} \left[(1 + 2F) \tilde{D}_{ij} + \tilde{D}_{i,j} \right. \\ & \left. + a_{ij} \frac{16(1 + \sigma)(2 + \sigma)}{(2 - \sigma)^2(4 - \sigma)} p_i q_i p_j q_j \right] \end{aligned} \quad (36)$$

with $a_{ij} = -\sum_{\alpha} r_{\alpha i} r_{\alpha j} / (2V_s)$, and where $\tilde{D}_{ij}$, $\tilde{D}_{i,j}$ and $D_{i,i}$ are given by equations 30, 31 and 35. The first line of equation 36 is equivalent to equation 6 in Roze (2015), showing that identity disequilibria reduce the efficiency of purging by decreasing the excess homozygosity ($D_{i,i}$), and by two additional effects represented by the term in $6(1 + F)$ (see Roze (2015) for interpretation of these effects). The term on the second and third lines (proportional to a_{ij}^2) represents the effect of epistasis between loci: this term also reduces purging, since selection against deleterious alleles is less efficient when these alleles are partially compensated by alleles at other loci.

An expression for the genic variance at mutation-selection balance is given by

equation A65 in Supplementary File S1. From this, one obtains for the genetic variance:

$$V_{g,\alpha} = \frac{4V_s U}{n(1+\sigma)} \left[1 + 2U \frac{\sigma(1-\sigma)[6+\sigma(2-\sigma)]}{(2-\sigma)(4-\sigma)(1+\sigma)^2} + \frac{4U}{n} \frac{(2-\sigma)(2+\sigma)}{(1-\sigma)(4-\sigma)(1+\sigma)} \left(\frac{1}{2\rho_H} + \frac{2+\sigma(1-\sigma)(2-\sigma)}{2+\sigma(1-\sigma)} \right) \right] \quad (37)$$

where the terms in U between the brackets correspond to the effect of between-locus associations. The first of these terms (on the first line of equation 37) represents the effect of identity disequilibria, while the term in U/n on the second line represents the effect of epistasis (compensatory effects between alleles at different loci). Both terms are positive, indicating that the overall effect of interactions between loci is to increase the genetic variance, due to the fact that correlations in homozygosity and compensatory effects between mutations both reduce the efficiency of purging (equation 36). Furthermore, while the effect of identity disequilibria scales with U , the effect of epistasis scales with U/n : indeed, it becomes less and less likely that alleles at different loci have compensatory effects on all of the traits as the dimensionality of the fitness landscape increases. Finally, the effect of epistasis is more strongly affected by linkage between loci (through the term in $1/\rho_H$); the effect of linkage on the term in U representing the effect of linkage disequilibria is weaker, and has been neglected in equation 37. Under random mating ($\sigma = 0$), equation 37 simplifies to:

$$V_{g,\alpha} = \frac{4V_s U}{n} \left[1 + \frac{2U}{n} \left(\frac{1}{\rho_H} + 2 \right) \right] \quad (38)$$

which takes a similar form as equation 4.16 in Turelli and Barton (1990) in the case of a single selected trait ($n = 1$). Supplementary Figure S6 shows how the equilibrium genetic variance and its different components vary with the selfing rate, in a regime where both identity disequilibria and epistasis have significant effects ($U = 1$).

Assuming that $\overline{W} \approx e^{\overline{\log W_g}}$, an approximation for the load at mutation-selection balance is $1 - \exp[-n V_g / (2V_s)]$, where V_g is the genetic variance given by equation 37 (the same for all traits). A slightly better approximation can be obtained by using $\overline{W} \approx e^{\overline{\log W_g}} (1 + \frac{1}{2} \text{Var}[\log W_g])$, where $\text{Var}[\log W_g]$ is the variance in log fitness in the population (Roze, 2015). To leading order, it is given by (see Supplementary File S1):

$$\text{Var}[\log W_g] \approx 2\bar{s}U \frac{1+3\sigma}{1+\sigma} + 4U^2 \frac{\sigma(1-\sigma)}{(4-\sigma)(1+\sigma)^2} + \frac{8U^2}{n} \frac{(2-\sigma)(2+\sigma)}{(4-\sigma)(1+\sigma)^2}, \quad (39)$$

simplifying to $2U(\bar{s} + 4U/n)$ in the absence of selfing. The first term of equation 39 represents the sum of single-locus contributions to the variance in log fitness, while the second and third term correspond to the effects of identity disequilibria and epistasis (respectively), both increasing the variance in fitness. The mutation load is then given by:

$$L \approx 1 - \left(1 + \frac{1}{2} \text{Var}[\log W_g]\right) \exp\left[-\frac{nV_g}{2V_s}\right]. \quad (40)$$

Similarly, we show in Supplementary File S1 that an expression for inbreeding depression including the effects of pairwise associations between loci is:

$$\delta \approx 1 - \left(1 + \frac{1}{2} \Delta \text{Var}[\log W_g]\right) \exp\left[-\frac{nV_g}{4V_s}\right] \quad (41)$$

where $\Delta \text{Var}[\log W_g]$ is the difference in variance in log fitness between selfed and outcrossed offspring, given by:

$$\Delta \text{Var}[\log W_g] \approx \frac{7\bar{s}U}{1+\sigma} + U^2 \frac{\sigma(1-\sigma)}{(4-\sigma)(1+\sigma)^2} + \frac{2U^2}{n} \frac{(10-\sigma)(2-\sigma)}{(4-\sigma)(1+\sigma)^2} \quad (42)$$

(the terms in $\text{Var}[\log W_g]$ and $\Delta \text{Var}[\log W_g]$ in equations 40 and 41 are often small, however, and may thus be neglected). After replacing V_g , $\text{Var}[\log W_g]$ and $\Delta \text{Var}[\log W_g]$ by the expressions given by equations 37, 39 and 42, the approximations obtained for

the load and inbreeding depression include terms in U^2 representing the effect of identity disequilibria, and terms in U^2/n representing the effect of epistasis between loci. The terms in U^2 are identical to the terms representing the effect of identity disequilibria in a model with purely multiplicative selection against deleterious alleles (no epistasis) when setting the dominance coefficient h of deleterious alleles to $1/4$ (equations 11 and 14 in Roze, 2015). The novelty here thus corresponds to the effect of epistasis (compensatory effects between deleterious alleles), that tends to increase V_g , L , δ by reducing the efficiency of purging.

Figure 3B shows that equations 37, 39 and 40 capture the increase in load observed in the simulations as the number of traits n decreases (see Supplementary Figure S7 for the genetic variance and inbreeding depression). Note that the harmonic mean recombination rate ρ_H between pairs of loci under our simulated genetic architecture (linear chromosome with equally spaced loci) can be obtained from:

$$\frac{1}{\rho_H} = \frac{2}{\ell(\ell-1)} \sum_{i=1}^{\ell-1} \frac{2(\ell-i)}{1 - \exp\left[-2i\frac{R}{\ell-1}\right]} \quad (43)$$

(see Appendix 2 in Roze and Blanckaert, 2014), yielding $\rho_H \approx 0.42$ for $\ell = 1,000$ and $R = 20$. Figure 4 shows that for low or moderate selfing rates, decreasing the genome map length from $R = 20$ to $R = 1$ increases the mutation load, by increasing the strength of linkage disequilibria caused by epistasis, that in turn reduce the efficiency of purging. In this regime, equations 37, 39 and 40 provide an accurate prediction for the load (see Supplementary Figure S8 for genetic variance and inbreeding depression). At high selfing rates, however, a different regime is entered, in which the assumption of weak genetic associations breaks down. As can be seen in Figure 4, in this regime (which spans a broader parameter range under tighter linkage) the load decreases more

rapidly as σ increases. Increasing linkage tends to reduce the mutation load when the selfing rate is high, although the effect of R vanishes when $\sigma = 1$. When linkage is extremely tight, the approximations given above break down for all values of σ : as shown by Figures 4 and S8, decreasing R has a non-monotonic effect on the genetic variance, load and inbreeding depression when selfing is small to moderate, the lowest values of V_g , L and δ being reached when $R = 0$ (in which case selfing has no effect). An approximation for the genetic variance under complete linkage can be obtained by treating the whole genome as a single locus with a very large number of possible alleles, and assuming a Gaussian distribution of allelic effects in the population (Lande, 1977; Supplementary File S4). This yields:

$$\frac{V_g}{2V_s} \approx 2\sqrt{\frac{\bar{s}U}{2n} \frac{2}{2 + \sigma(1 - \sigma)}}, \quad (44)$$

$$L \approx 1 - \exp\left[-\frac{nV_g}{2V_s}\right], \quad \delta \approx 1 - \exp\left[-\frac{nV_g}{4V_s}\right]. \quad (45)$$

Note that equation 44 is equivalent to equation 3A in Charlesworth and Charlesworth (1995) when $\sigma = 1$ and $n = 1$. As shown by Figures 4 and S8, equations 44 and 45 only slightly overestimate V_g , L and δ when $\sigma = 1$ and/or $R = 0$. As shown below, better predictions are observed for higher values of U/n and lower values of $\bar{s}$.

The effects of identity disequilibria between loci remain negligible for the parameter values used in Figures 3 and 4. As shown by Figure 5, identity disequilibria become more important for higher values of the mutation rate U . Indeed, the relative effects of identity disequilibria on the load can be deduced from the differences between the three curves in each panel of Figure 5, the red curves showing the predicted mutation load in the absence of epistasis, but taking into account identity disequilibria (obtained by removing the terms in U^2/n from equations 37, 39 and 40, leading to an

expression equivalent to equation 11 in Roze, 2015). The difference between the black and red curves thus represents the predicted effect of identity disequilibria on the load, while the difference between the red and green curves corresponds to the additional effect of epistasis. Simulations indicate that the change in regime observed above a threshold selfing rate (around $\sigma = 0.5$ for $U = 1$ in Figure 5) is due to epistasis, since this threshold is not observed in simulations without epistasis (red dots). Supplementary Figure S9 shows that this threshold pattern is little affected by population size N , as long as the effects of drift remain small. Similarly, the results only weakly depend on the number of loci ℓ , as long as the mutation rate per locus $u = U/\ell$ is small enough so that $p_i q_i < 1/4$ at most loci (see Supplementary Figure S10 for distributions of allele frequencies in simulations with $\ell = 1,000$ and $\ell = 10,000$). Figure S9 also shows that the results are little affected by the degree of pleiotropy of mutations m , as long as $\bar{s}$ remains constant. However, $\bar{s}$ does affect V_g , L and δ in the regime where our approximations break down. As shown by Figure 6, decreasing $\bar{s}$ lowers the threshold selfing rate above which our approximations are not valid and results in lower equilibrium mutation loads (see Supplementary Figure S11 for results on V_g and δ). Figures 6 and S11 also show that, when $\bar{s}$ is sufficiently small, the single-locus, Gaussian model (equations 44 and 45, dotted curves on the figures) provides accurate predictions for V_g , L and δ under complete selfing ($\sigma = 1$).

In Figure 7 we show that decreasing the number of traits under selection n decreases the threshold selfing rate above which our approximations break down (see Supplementary Figure S12 for inbreeding depression and scaled genetic variance). Below the threshold, the mutation load decreases as n increases, as predicted by our analytical results (although our approximations become less precise for low n and high

568 U), while n has the opposite effect above the threshold. Overall, we observe that in
 569 this second regime (in which interactions between loci have important effects), the
 570 mutation load generally increases with the number of selected traits, the fitness effects
 571 of mutations $\bar{s}$, the mutation rate U and recombination rate (through the parameter
 572 R). However, Figure 8 shows that the effects of these parameters on inbreeding de-
 573 pression are more complicated. In particular, outbreeding depression (negative δ) may
 574 occur in regimes where the effects of epistasis are particularly strong (high U , low n)
 575 and when the selfing rate is moderate to high (above 0.5 but below 1), outbreeding
 576 depression becoming stronger when $\bar{s}$, U and R increase (the approximation derived
 577 from equation 37 fails for all values of σ for the parameter values used in Figure 8,
 578 and is not shown here). Supplementary Figures S13 and S14 show that for the same
 579 parameter values, V_g and L always increase when $\bar{s}$, U and R increase.

580 DISCUSSION

581 The response of a population to environmental change depends critically on its
 582 genetic diversity. Our results predict that the level of genetic variation maintained at
 583 equilibrium under stabilizing selection acting on quantitative traits is generally lower
 584 in more highly selfing population, due to more efficient purging (although increasing
 585 selfing may sometimes increase genetic variation, for example when mutations have
 586 weak fitness effects, as shown by Figure S3). This finding agrees with Charlesworth
 587 and Charlesworth's (1995) theoretical prediction that fully selfing populations should
 588 maintain lower genetic variance for quantitative traits under stabilizing selection than
 589 fully outcrossing ones, and with several empirical studies comparing levels of genetic

variation for morphological traits in closely related pairs of plant species with con-
trasted mating systems (Charlesworth and Charlesworth, 1995; Geber and Griffen,
2003; Bartkowska and Johnston, 2009 and references therein). We also show that the
lower level of variation present in more highly selfing populations is associated with
lower values of the mutation load and inbreeding depression. The meta-analysis carried
out by Winn et al. (2011) showed that inbreeding depression is indeed lower in highly
selfing plant species compared to species with lower selfing rates, while no significant
difference is observed between species with low vs. intermediate selfing rates. It has
been put forth that correlations in homozygosity between selected loci may suppress
purging at moderate selfing rates (“selective interference”, Lande et al., 1994; Winn et
al., 2011); this, however, would imply that a large number of segregating deleterious al-
leles have very low dominance coefficients, generating very high inbreeding depression
(Kelly, 2007; Roze, 2015), which seems unlikely. Another possible explanation for the
lack of purging at intermediate selfing rates involves epistasis (compensatory effects
between mutations coding for the same quantitative trait, Lande and Porcher, 2015).
Our analysis of the effects of epistasis (under assumptions that differ from those made
in Lande and Porcher’s model) shows that different regimes are possible, and outlines
how the parameters affect transitions between these regimes.

In our model, the effect of epistasis on the equilibrium genetic variance V_g is
inversely proportional to effective recombination rates between selected loci, and scales
with U/n (where n is the number of selected traits and U the total mutation rate on
those traits). Indeed, U/n determines the number of segregating “interacting” mu-
tations, that is, mutations with epistatic fitness effects. As n tends to infinity, all
mutations become orthogonal in phenotypic space (with independent fitness effects),

and our results converge to the results from previous population genetics models without epistasis (e.g., Charlesworth and Charlesworth, 1987; Roze, 2015). When U/n is small and map length R is sufficiently large, associations between loci have little effect. Under Gaussian stabilizing selection ($Q = 2$), the average coefficient of epistasis between mutations (on fitness) is zero (Martin and Lenormand, 2006b) while the dominance coefficient of deleterious alleles (in an optimal genotype) is close to 0.25 under the assumption of additive effects on phenotypes. In this case, we found that classical deterministic expressions based on single-locus models (hence neglecting the variance in epistatic interactions) provide accurate predictions for the mutation load L and inbreeding depression δ . Simple approximations are also obtained under the more general fitness function given by equation 15, confirming that the mutation load is an increasing function of the average coefficient of epistasis between mutations (Kimura and Maruyama, 1966; Kondrashov and Crow, 1988; Phillips et al., 2000; Roze and Blanckaert, 2014). Neglecting the effect of associations between loci also allowed us to explore the effects of drift using diffusion methods. As in previous studies (e.g., Charlesworth, 2013; Roze and Blanckaert, 2014), we found that drift may lower the mutation load by reducing V_g . However, this result probably strongly depends on the assumption that mutations may increase or decrease phenotypic traits with the same probability (no mutational bias): indeed, previous works showed that drift may increase the load in the presence of a mutational bias by displacing mean phenotypes away from the optimum (Zhang and Hill, 2008; Charlesworth, 2013). Given that partial selfing reduces effective population size, it would be of interest to study the combined effects of drift and mutational bias in models with selfing.

The variance in epistasis has stronger effects as the U/n ratio increases and/or as

the effective recombination rate decreases (*i.e.* due to selfing). Our results showed that two different regimes are possible. (1) When genetic associations (linkage disequilibria) generated by epistasis stay moderate, the overall effect of epistasis is to increase V_g , L and δ by decreasing the efficiency of selection against deleterious alleles. This regime is generally well described by our model taking into account the effects of associations between pairs of loci. This result bears some similarity with the result obtained by Phillips et al. (2000), showing that the variance in epistasis between deleterious alleles increases the mutation load. Equation 2.1 in Phillips et al. (2000) is not fully equivalent to our expression for the load under random mating, however, possibly due to different assumptions on the relative orders of magnitude of s_i , s_j and e_{ij} (where s_i and s_j are the strength of selection at loci i and j and e_{ij} is epistasis between those loci) and how they covary. Nevertheless, it is interesting to note that both results become equivalent if $\text{Var}[e_{ij}/(s_i s_j)]$ in Phillips et al. (2000) is replaced by $\text{Var}[e_{ij}/\bar{s}^2]$ (where Var stands for the variance across all pairs of loci), using the fact that $\text{Var}[e_{ij}] = 4\bar{s}^2/n$ in our model (with a Gaussian fitness function). (2) Increasing the value of U/n and/or reducing effective recombination rates or $\bar{s}$ generates a transition to a different regime in which the effect of the variance in epistasis switches, reducing V_g , L and δ . Because our analytical approach fails in this regime (presumably due to higher-order associations between loci), it is more difficult to obtain an intuitive understanding of the selective mechanisms involved. However, it is likely that selection operates on multilocus genotypes (comprising combinations of alleles with compensatory effects) that can be maintained over many generations due to high selfing rates and/or low recombination. A similar transition from genic to genotypic selection as recombination decreases was described by Neher and Shraiman (2009), using a haploid model in which

epistasis is randomly assigned to genotypes.

Although our results show some qualitative similarities with those obtained by Lande and Porcher (2015) — e.g., the same transition between regimes occurs in both models as selfing increases — several differences can be observed. In particular, Lande and Porcher’s model predict little or no effect of selfing on V_g below the threshold selfing rate corresponding to the change in regime, and an abrupt change in V_g at the threshold (except in their infinitesimal model). A step change such as this is never observed in our model, even for parameter values at which the effect of drift should be negligible at most loci. These differences between the models are not due to the different genetic architectures considered (biallelic vs. multiallelic): indeed, Supplementary Figures S15 and S16 show that assuming biallelic loci or an infinite number of possible alleles per locus in our individual-based simulations yields very similar results (for $\ell = 1,000$ and $\ell = 10,000$). Rather, they must be due to Lande and Porcher’s assumption of a Gaussian distribution of allelic effects maintained at each locus in each selfing age class, implicitly assuming a sufficiently high mutation rate per locus u and low fitness effect of mutations $\bar{s}$ (Turelli, 1984). In our multiallelic simulations (with $u = 10^{-5}$ to 10^{-3} and $\bar{s} = 0.01$), the number of alleles maintained at each locus is not sufficiently large to generate a Gaussian distribution of segregating allelic effects (see Figures S15 and S16). One may also note that the effect of the number of selected traits n seems different in both models (compare Lande and Porcher’s Figure 5 and 6 to our Figure 7), but this is due to the fact that the overall mutation rate U is proportional to n in Lande and Porcher’s model (while U is fixed in Figure 7). Increasing both n and U in order to maintain a constant U/n ratio, we indeed observed that the transition between regimes occurs at lower selfing rates when n is larger, as in Lande and Porcher’s Figure

5 and 6 (results not shown). In general, whether U should scale with n depends on the degree of pleiotropy of mutations (Lande and Porcher assume no pleiotropy). Our model allowed us to explore the effects of pleiotropy through the parameter m , showing that pleiotropy mostly affects the results through its effect on $\bar{s}$ (equation 14). The equilibrium genetic variance thus depends on m in regimes where V_g is affected by $\bar{s}$, in particular when $N_e \bar{s} \approx 1$ or lower (Figures 1, S1 – S4), and when genetic associations are strong (Figure 6). However, pleiotropy may have stronger effects under different assumptions regarding the genetic architecture of traits, for example when different sets of traits are affected by different sets of loci (modular pleiotropy, Welch and Waxman, 2003). The effects of selective or mutational covariance among traits would also be interesting to explore: indeed, such covariances decrease the effective number of selected traits (Martin and Lenormand, 2006b), potentially increasing the importance of associations between loci.

In the regime where genetic associations generated by epistasis reduce V_g (regime (2) mentioned above), outbreeding depression may occur due to the lower fitness of recombinants between selfing lineages maintaining coadapted gene complexes (Figure 8), a result shared with Lande and Porcher’s (2015) Gaussian Allele Model. In our additive model of phenotypic effects, outbreeding depression should only be expressed in F2 individuals (that is, among the offspring of an individual produced by a cross between different selfing lineages), once recombination has disrupted compensatory associations between alleles at different loci. This explains why outbreeding depression is not observed under complete (or nearly complete) selfing in Figure 8, as all outcrossed individuals are F1 hybrids between selfing lineages. Outbreeding depression between lineages collected from the same geographical location has been observed in highly

selfing plants (Parker, 1992; Volis et al., 2011) and *Caenorhabditis* nematodes (Dolgin et al., 2007; Gimond et al., 2013). In all cases, estimated selfing rates are higher than those leading to $\delta < 0$ in our simulations, however, and outbreeding depression was observed in F1 offspring of crosses between inbred lines of nematodes. The occurrence of outbreeding depression at higher selfing rates may be partly explained by the fact that experimental crosses were often performed between genetically different lines; by contrast, in our simulations the parents of an outcrossed individual may share the same genotype (in particular when the number of genetically different selfing lineages is reduced due to the low effective size of highly selfing populations), reducing the magnitude of outbreeding depression. However, the occurrence of outbreeding depression in F1 individuals must involve dominance effects which are absent from our model. Exploring the effects of dominance/recessivity of mutations on phenotypic traits would be an interesting extension of this work.

Due to the lower genetic diversity of self-fertilizing populations, it has been suggested that they should be less able to adapt to a changing environment (e.g., Stebbins, 1957; Williams, 1992; Takebayashi and Morrell, 2001). In the absence of epistasis, existing models indeed predict that selfing populations should have lower rates of adaptation than outcrossing ones (Glémin and Ronfort, 2013; Hartfield and Glémin, 2016). When compensatory effects between mutations are possible, however, a substantial amount of genetic variance may be hidden by genetic associations between loci in highly selfing populations (Lande and Porcher, 2015, the present study). After a change in environment, this variance may be liberated by rare outcrossing events, increasing the short-term evolutionary response of highly (but not fully) selfing populations. Exploring how selfing affects adaptation under directional selection, and more

generally how the variability of epistatic interactions between loci may influence the evolution of mating systems represents a natural next step of this work.

Acknowledgements. We thank Aneil Agrawal, Patrice David, Sylvain Glémin, Frédéric Guillaume, Thomas Lenormand, Emmanuelle Porcher, Ophélie Ronce, Joëlle Ronfort and an anonymous reviewer for helpful discussions and comments, and the bioinformatics and computing service of Roscoff’s Biological Station (Abims platform) for computing time. This work was supported by the French Agence Nationale de la Recherche (project SEAD, ANR-13-ADAP-0011 and project SexChange, ANR-14-CE02-0001).

LITERATURE CITED

- 744
745 Bartkowska, M. P. and M. O. Johnston. 2009. Quantitative genetic variation in pop-
746 ulations of *Amsinckia spectabilis* that differ in rate of self-fertilization. *Evolution*
747 63:1103–1117.
- 748 Barton, N. H. 1986. The maintenance of polygenic variation through a balance between
749 mutation and stabilizing selection. *Genet. Res.* 47:209–216.
- 750 ———. 1989. The divergence of a polygenic system subject to stabilizing selection,
751 mutation and drift. *Genet. Res.* 54:59–77.
- 752 Brandvain, Y., T. Slotte, K. M. Hazzouri, S. I. Wright, and G. Coop. 2013. Ge-
753 nomic identification of founding haplotypes reveals the history of the selfing species
754 *Capsella rubella*. *PLoS Genetics* 9:e1003754.
- 755 Bulmer, M. G. 1971. The effect of selection on genetic variability. *Am. Nat.* 105:201–
756 211.
- 757 ———. 1972. The genetic variability of polygenic characters under optimizing selec-
758 tion, mutation and drift. *Genet. Res.* 19:17–25.
- 759 ———. 1974. Linkage disequilibrium and genetic variability. *Genet. Res.* 23:281–289.
- 760 ———. 1985. *The Mathematical Theory of Quantitative Genetics*, 2nd edition. Oxford
761 University Press, Oxford.
- 762 Burgarella, C., P. Gayral, M. Ballenghien, A. Bernard, P. David, P. Jarne, A. Correa,
763 S. Hurtrez-Boussès, J. Escobar, N. Galtier, and S. Glémin. 2015. Molecular evolution
764 of freshwater snails with contrasting mating systems. *Mol. Biol. Evol.* 32:2403–2416.

- 765 Charlesworth, B. 2013. Why we are not dead one hundred times over. *Evolution*
766 67:3354–3361.
- 767 Charlesworth, B., M. T. Morgan, and B. Charlesworth. 1990. Inbreeding depression,
768 genetic load, and the evolution of outcrossing rates in a multilocus system with no
769 linkage. *Evolution* 44:1469–1489.
- 770 Charlesworth, B., M. T. Morgan, and D. Charlesworth. 1991. Multilocus models of
771 inbreeding depression with synergistic selection and partial self-fertilization. *Genet.*
772 *Res.* 57:177–194.
- 773 Charlesworth, D. and B. Charlesworth. 1987. Inbreeding depression and its evolution-
774 ary consequences. *Ann. Rev. Ecol. Syst.* 18:237–268.
- 775 ———. 1995. Quantitative genetics in plants: the effect of the breeding system on
776 genetic variability. *Evolution* 49:911–920.
- 777 Chevin, L.-M., G. Martin, and T. Lenormand. 2010. Fisher’s model and the genomics
778 of adaptation: restricted pleiotropy, heterogenous mutation, and parallel evolution.
779 *Evolution* 64:3213–3221.
- 780 Dolgin, E. S., B. Charlesworth, S. E. Baird, and A. D. Cutter. 2007. Inbreeding and
781 outbreeding depression in *Caenorhabditis* nematodes. *Evolution* 61:1339–1352.
- 782 Eyre-Walker, A. and P. D. Keightley. 2007. The ditribution of fitness effects of new
783 mutations. *Nat. Rev. Genet.* 8:610–618.
- 784 Fisher, R. A. 1930. The genetical theory of natural selection. Clarendon Press, Oxford.

- 785 Geber, M. A. and L. R. Griffen. 2003. Inheritance and natural selection on functional
786 traits. *Int. J. Plant Sci.* 164:S21–S42.
- 787 Gimond, C., R. Jovelin, S. Han, C. Ferrari, A. D. Cutter, and C. Braendle. 2013. Out-
788 breeding depression with low genetic variation in selfing *Caenorhabditis* nematodes.
789 *Evolution* 67:3087–3101.
- 790 Glémin, S. 2007. Mating systems and the efficacy of selection at the molecular level.
791 *Genetics* 177:905–916.
- 792 Glémin, S. and J. Ronfort. 2013. Adaptation and maladaptation in selfing in outcross-
793 ing species: new mutations versus standing variation. *Evolution* 67:225–240.
- 794 Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed
795 mating systems in plants: occurrence, theoretical explanations, and empirical evi-
796 dence. *Ann. Rev. Ecol. Evol. Syst.* 36:47–79.
- 797 Gros, P.-A., H. Le Nagard, and O. Tenaillon. 2009. The evolution of epistasis and
798 its links with genetic robustness, complexity and drift in a phenotypic model of
799 adaptation. *Genetics* 182:277–293.
- 800 Haldane, J. B. S. 1937. The effect of variation on fitness. *Am. Nat.* 71:337–349.
- 801 Hartfield, M. 2015. Evolutionary genetic consequences of facultative sex and outcross-
802 ing. *J. Evol. Biol.* 29:5–22.
- 803 Hartfield, M. and S. Glémin. 2016. Limits to adaptation in partially selfing species.
804 *Genetics* 203:959–974.

- 805 Igic, B. and J. R. Kohn. 2006. The distribution of plant mating systems: study bias
806 against obligately outcrossing species. *Evolution* 60:1098–1103.
- 807 Jarne, P. and J. R. Auld. 2006. Animals mix it up too: the distribution of self-
808 fertilization among hermaphroditic animals. *Evolution* 60:1816–1824.
- 809 Kamran-Disfani, A. and A. F. Agrawal. 2014. Selfing, adaptation and background
810 selection in finite populations. *J. Evol. Biol.* 27:1360–1371.
- 811 Kelly, J. K. 2007. Mutation-selection balance in mixed mating populations. *J. Theor.*
812 *Biol.* 246:355–365.
- 813 Kimura, M. and T. Maruyama. 1966. The mutational load with epistatic gene inter-
814 actions in fitness. *Genetics* 54:1337–1351.
- 815 Kirkpatrick, M., T. Johnson, and N. H. Barton. 2002. General models of multilocus
816 evolution. *Genetics* 161:1727–1750.
- 817 Kondrashov, A. S. and J. F. Crow. 1988. King’s formula for the mutation load with
818 epistasis. *Genetics* 120:853–856.
- 819 Lande, R. 1976. The maintenance of genetic variability by mutation in a polygenic
820 character with linked loci. *Genet. Res.* 26:221–235.
- 821 ———. 1977. The influence of the mating system on the maintenance of genetic
822 variability in polygenic characters. *Genetics* 86:485–498.
- 823 Lande, R. and E. Porcher. 2015. Maintenance of quantitative genetic variance un-
824 der partial self-fertilization, with implications for the evolution of selfing. *Genetics*
825 200:891–906.

- 826 Lande, R. and D. W. Schemske. 1985. The evolution of self-fertilization and inbreeding
827 depression in plants. I. Genetic models. *Evolution* 39:24–40.
- 828 Lande, R., D. W. Schemske, and S. T. Schultz. 1994. High inbreeding depression,
829 selective interference among loci, and the threshold selfing rate for purging recessive
830 lethal mutations. *Evolution* 48:965–978.
- 831 Latter, B. D. H. 1960. Natural selection for an intermediate optimum. *Aust. J. Biol.*
832 *Sci.* 13:30–35.
- 833 Lourenço, J., N. Galtier, and S. Glémin. 2011. Complexity, pleiotropy and the fitness
834 effect of mutations. *Evolution* 65:1559–1571.
- 835 Manna, F., G. Martin, and T. Lenormand. 2011. Fitness landscapes: an alternative
836 theory for the dominance of mutation. *Genetics* 189:923–937.
- 837 Martin, G., S. F. Elena, and T. Lenormand. 2007. Distributions of epistasis in microbes
838 fit predictions from a fitness landscape model. *Nat. Genet.* 39:555–560.
- 839 Martin, G. and T. Lenormand. 2006a. The fitness effect of mutations across environ-
840 ments: a survey in light of fitness landscape models. *Evolution* 60:2413–2427.
- 841 ———. 2006b. A general multivariate extension of Fisher’s geometrical model and the
842 distribution of mutation fitness effects across species. *Evolution* 60:893–907.
- 843 Neher, R. A. and B. I. Shraiman. 2009. Competition between recombination and
844 epistasis can cause a transition from allele to genotype selection. *Proc. Natl. Acad.*
845 *Sci. U. S. A.* 106:6866–6871.

- 846 Nordborg, M. 1997. Structured coalescent processes on different time scales. *Genetics*
847 146:1501–1514.
- 848 Parker, M. A. 1992. Outbreeding depression in a selfing annual. *Evolution* 46:837–841.
- 849 Phillips, P. C., S. P. Otto, and M. C. Whitlock. 2000. Beyond the average: the evolu-
850 tionary importance of gene interactions and variability of epistatic effects. Pp. 20–38
851 *in* J. B. Wolf, E. D. Brodie, and M. J. Wade, eds. *Epistasis and the Evolutionary*
852 *Process*. Oxford University Press, New York.
- 853 Pollak, E. 1987. On the theory of partially inbreeding finite populations. I. Partial
854 selfing. *Genetics* 117:353–360.
- 855 Roze, D. 2015. Effects of interference between selected loci on the mutation load,
856 inbreeding depression and heterosis. *Genetics* 201:745–757.
- 857 ———. 2016. Background selection in partially selfing populations. *Genetics* 203:937–
858 957.
- 859 Roze, D. and A. Blanckaert. 2014. Epistasis, pleiotropy and the mutation load in
860 sexual and asexual populations. *Evolution* 68:137–149.
- 861 Scofield, D. G. and S. T. Schultz. 2006. Mitosis, stature and evolution of plant mating
862 systems: low- Φ and high- Φ plants. *Proc. Roy. Soc. (Lond.) B* 273:275–282.
- 863 Stebbins, G. L. 1957. Self fertilization and population variability in higher plants. *Am.*
864 *Nat.* 91:337–354.
- 865 Takebayashi, N. and P. L. Morrell. 2001. Is self-fertilization an evolutionary dead

866 end? Revisiting an old hypothesis with genetic theories and a macroevolutionary
867 approach. *Am. J. Bot.* 88:1143–1150.

868 Tenaillon, O., O. K. Silander, J.-P. Uzan, and L. Chao. 2007. Quantifying organismal
869 complexity using a population genetic approach. *PLoS One* 2:e217.

870 Turelli, M. 1984. Heritable genetic variation via mutation-selection balance: Lerch’s
871 zeta meets the abdominal bristle. *Theor. Popul. Biol.* 25:138–193.

872 Turelli, M. and N. H. Barton. 1990. Dynamics of polygenic characters under selection.
873 *Theor. Popul. Biol.* 38:1–57.

874 Vitalis, R. and D. Couvet. 2001. Two-locus identity probabilities and identity disequi-
875 librium in a partially selfing subdivided population. *Genet. Res.* 77:67–81.

876 Volis, S., I. Shulgina, M. Zaretsky, and O. Koren. 2011. Epistasis in natural populations
877 of a predominantly selfing plant. *Heredity* 106:300–309.

878 Weir, B. S. and C. C. Cockerham. 1973. Mixed self and random mating at two loci.
879 *Genet. Res.* 21:247–262.

880 Welch, J. J. and D. Waxman. 2003. Modularity and the cost of complexity. *Evolution*
881 57:1723–1734.

882 Williams, G. C. 1992. *Natural Selection: Domains, Levels, and Challenges*. Oxford
883 University Press, New York, NY.

884 Winn, A. A., E. Elle, S. Kalisz, P.-O. Cheptou, C. G. Eckert, C. Goodwillie, M. O.
885 Johnston, D. A. Moeller, R. H. Ree, R. D. Sargent, and M. Vallejo-Marín. 2011.

886 Analysis of inbreeding depression in mixed-mating plants provides evidence for se-
 887 lective interference and stable mating systems. *Evolution* 65:3339–3359.

888 Wright, S. 1951. The genetical structure of populations. *Ann. Eugen.* 15:323–354.

889 Zhang, X.-S. and W. G. Hill. 2008. The anomalous effects of biased mutation revis-
 890 ited: mean-optimum deviation and apparent directional selection under stabilizing
 891 selection. *Genetics* 179:1135–1141.

Table 1: Parameters and variables of the model.

N	Population size
σ	Selfing rate
n	Number of selected traits
m	Degree of pleiotropy of mutations
a^2	Variance of mutational effects on selected traits
V_e	Environmental variance (on selected traits)
ω^2	Strength of stabilizing selection on phenotypic traits
$V_s = \omega^2 + V_e$	Strength of stabilizing selection on breeding values g_α
Q	Shape of the fitness peak (equation 15)
ℓ	Number of loci affecting selected traits
u	Mutation rate per locus
$U = u \ell$	Mutation rate (per haploid genome) on loci affecting selected traits
R	Genome map length
ρ_H	Harmonic mean recombination rate between pairs of loci affecting selected traits
$\bar{s}$	Average heterozygous effect of mutations on log fitness (in an optimal genotype)
z_α	Value of phenotypic trait α (in a given individual)
g_α, e_α	Genetic and environmental components of trait α

895

$r_{\alpha i}$	Effect of allele 1 at locus i on trait α
p_i, q_i	Frequencies of alleles 1 and 0 at locus i
$D_{i,i}$	Excess homozygosity at locus i
$\tilde{D}_{ij}$	Association between alleles 1 at loci i and j on the same haplotype (linkage disequilibrium)
$\tilde{D}_{i,j}$	Association between alleles 1 at loci i and j on different haplotypes
$V_{g,\alpha}$	Genetic variance for trait α (variance of g_α)
$V_{g,\alpha}^0$	Genic variance for trait α ($2 \sum_i r_{\alpha i}^2 p_i q_i$)
F	Inbreeding coefficient
G_{ij}	Identity disequilibrium between loci i and j
G	Identity disequilibrium between freely recombining loci
L	Mutation load
δ	Inbreeding depression

896

897

898 **Figure 1.** Mutation load L as a function of the selfing rate σ . Black curve: approx-
 899 imation for mutation-selection regime neglecting genetic associations (equation 24).
 900 The different colors correspond to different values of $\bar{s}$ as shown in A. Colored solid
 901 curves: results from the diffusion model (Supplementary File S2). Dots correspond to
 902 simulation results; in this and the following figures, error bars (computed by splitting
 903 the last 70,000 generations into 7 batches of 10,000 generations and calculating the
 904 standard error over batches) are smaller than the size of symbols in most cases. Other
 905 parameter values are $U = 0.1$, $R = 20$, $n = 50$, $m = 5$, $\ell = 10,000$.

906

907 **Figure 2.** A: fitness as a function of the (scaled) distance from the optimum, for
 908 different values of the parameter Q (from equation 15). B, C, D: scaled genetic vari-
 909 ance, mutation load and inbreeding depression as a function of the selfing rate σ , for
 910 different values of Q . The curves represent the analytical results (neglecting associa-
 911 tions between loci) at mutation-selection balance (equations 27 – 29), while the dots
 912 correspond to simulation results. Parameter values: $N = 50,000$, $\ell = 10,000$, $U = 0.1$,
 913 $R = 20$, $n = 50$, $m = 5$, $a^2 / (2V_s) = 0.0002$ (yielding $\bar{s} = 0.001$ for $Q = 2$).

914

915 **Figure 3.** Mutation load L as a function of the selfing rate σ for $\bar{s} = 10^{-2}$ (top, filled
916 circles) and $\bar{s} = 10^{-4}$ (bottom, filled squares). A: the different colors correspond to
917 different values of m (degree of pleiotropy of mutations); B: the different colors corre-
918 spond to different values of n (number of selected traits). Black curve: approximation
919 for mutation-selection regime, neglecting genetic associations (equation 24). Colored
920 solid curves: results from the diffusion model (Supplementary File S2). Colored dashed
921 curves (in B): approximation including the effect of pairwise interactions among loci
922 (equations 37, 39 and 40). Other parameter values are $U = 0.1$, $R = 20$, $N = 5,000$,
923 $\ell = 1,000$, $n = 50$ (in A), $m = 5$ (in B).

924

925 **Figure 4.** Mutation load L as a function of the selfing rate σ , for $\bar{s} = 0.01$, $n =$
926 $m = 5$ and different values of the genome map length R , yielding (using equation 43)
927 $\rho_H \approx 0.07, 0.13$ and 0.42 for $R = 1, 2, 20$ (respectively). Dots: simulation results;
928 black curve: approximation for mutation-selection regime neglecting genetic associ-
929 ations (equation 24); colored curves: approximation including the effect of pairwise
930 interactions among loci (equations 37, 39 and 40); dashed grey curve: single-locus
931 model with many alleles, assuming a Gaussian distribution of allelic values (equations
932 44 and 45). Other parameter values are as in Figure 3.

933

934 **Figure 5.** Mutation load L as a function of the selfing rate σ , for $\bar{s} = 0.01$, $n = 50$, $m =$
935 5 , $U = 0.5$ (left) and 1 (right). The black curves correspond to the approximation for
936 mutation-selection regime, neglecting genetic associations (equation 24). Green curves:
937 approximation including the effect of pairwise interactions among loci (equations 37,
938 39 and 40); red curves: approximation including the effects of identity disequilibria
939 between loci, but not the effects of epistasis (obtained by removing the terms in U^2/n
940 from equations 37, 39 and 40, equivalent to equation 11 in Roze, 2015). Green dots:
941 simulation results; red dots: results from the simulation program used in Roze (2015)
942 representing multiplicative selection (no epistasis), with $s = 0.04$ and $h = 0.25$. Other
943 parameter values are $N = 5,000$, $\ell = 10,000$, $R = 20$ (yielding $\rho_H \approx 0.38$).

944

945 **Figure 6.** Mutation load L as a function of the selfing rate σ , for different values of the
 946 mutation rate U and average heterozygous effect of mutations $\bar{s}$; other parameter values
 947 are as in Figure 5. Dots: simulation results; black curves: approximation for mutation-
 948 selection regime, neglecting genetic associations (equation 24); solid colored curves:
 949 approximation including the effect of pairwise interactions among loci (equations 37,
 950 39 and 40); dotted colored curves: single-locus model with many alleles, assuming a
 951 Gaussian distribution of allelic values (equations 44 and 45).

952

953 **Figure 7.** Mutation load L as a function of the selfing rate σ , for different values
 954 of the mutation rate U and number of selected traits n ; other parameter values are
 955 as in Figure 5. Dots: simulation results; black curves: approximation for mutation-
 956 selection regime, neglecting genetic associations (equation 24); solid colored curves:
 957 approximation including the effect of pairwise interactions among loci (equations 37,
 958 39 and 40); dotted colored curves: single-locus model with many alleles, assuming a
 959 Gaussian distribution of allelic values (equations 44 and 45).

960

961 **Figure 8.** Inbreeding depression δ as a function of the selfing rate σ , for $n = m = 5$
 962 and different values of $\bar{s}$ (A), U (B) and R (C). Dots: simulation results; dotted curves:
 963 single-locus model with many alleles, assuming a Gaussian distribution of allelic values
 964 (equations 44 and 45). Other parameter values are $N = 5,000$, $\ell = 10,000$, $R = 20$,
 965 $U = 1$ and $\bar{s} = 0.01$.