Instructive starPEG-Heparin biohybrid 3D cultures for modeling human neural stem cell plasticity, neurogenesis, and neurodegeneration

4

5 Christos Papadimitriou^{1,2}, Mehmet I. Cosacak^{1,2}, Violeta Mashkaryan^{1,2},
6 Hilal Celikkaya^{1,2}, Laura Bray^{3,4}, Prabesh Bhattarai^{1,2}, Heike Hollak^{1,2}, Xin
7 Chen⁵, Shuijin He⁵, Christopher L. Antos^{5,6}, Alvin K. Thomas⁷, Jens
8 Friedrichs^{2,3}, Andreas Dahl⁷, Yixin Zhang⁸, Uwe Freudenberg^{2,3}, Carsten
9 Werner^{2,3}, Caghan Kizil^{1,2,*}

10

11 ¹ German Center for Neurodegenerative Diseases (DZNE) Dresden,
12 Helmholtz Association, Arnoldstr. 18, 01307, Dresden, Germany.

13 ² Center for Regenerative Therapies (CRTD), Technische Universität
14 Dresden, Fetscherstr. 105, 01307, Dresden, Germany.

15 ³ Leibniz Institute of Polymer Research Dresden, Max Bergmann Center of
16 Biomaterials Dresden, Hohe Str. 6, 01069, Dresden, Germany.

17 ⁴ Institute of Health Biomedical Innovation (IHBI), Queensland University
18 of Technology, 60 Musk Avenue, Kelvin Grove, 4059, Australia.

19 ⁵ School of Life Sciences and Technology, ShanghaiTech University,
20 Shanghai 201210, People's Republic of China.

21 ⁶ Institut für Pharmakologie und Toxikologie, Technische Universität
22 Dresden Medizinische Fakultät, Fetscherstr. 74, 01307, Dresden,
23 Germany.

24 ⁷ Deep Sequencing Group, Biotechnology Center, TU Dresden, Fetscherstr.
25 105, 01307, Dresden, Germany.

26 ⁸ B CUBE, Center for Molecular Bioengineering, TU Dresden, Arnoldstr. 18,
27 10307, Dresden, Germany.

28

29 *Correspondence should be addressed to: C.K. (caghan.kizil@dzne.de).
30 Fetscherstr. 105, 01307, Dresden, Germany.

31

Keywords:

Human neural stem cell, plasticity, Amyloid-beta42, whole transcriptome sequencing, 3D culture, neurogenesis

Abstract

Three-dimensional models of human neural development and neurodegeneration are crucial when exploring stem-cell-based regenerative therapies in a tissue-mimetic manner. However, existing 3D culture systems are not sufficient to model the inherent plasticity of NSCs due to their ill-defined composition and lack of controllability of the physical properties. Adapting a glycosaminoglycan-based, cell-responsive hydrogel platform, we stimulated primary and induced human neural stem cells (NSCs) to manifest neurogenic plasticity and form extensive neuronal networks *in vitro*. The 3D cultures exhibited neurotransmitter responsiveness, electrophysiological activity, and tissue-specific extracellular matrix (ECM) deposition. By whole transcriptome sequencing, we identified that 3D cultures express mature neuronal markers, and reflect the *in vivo* make-up of mature cortical neurons compared to 2D cultures. Thus, our data suggest that our established 3D hydrogel culture supports the tissue-mimetic maturation of human neurons. We also exemplarily modeled neurodegenerative conditions by treating the cultures with A β 42 peptide and observed the known human pathological effects of Alzheimer's disease including reduced NSC proliferation, impaired neuronal network formation, synaptic loss and failure in ECM deposition as well as elevated Tau hyperphosphorylation and formation of neurofibrillary tangles. We determined the changes in transcriptomes of primary and induced NSC-derived neurons after A β 42, providing a useful resource for further studies. Thus, our hydrogel-based human cortical 3D cell culture is a powerful platform for studying various aspects of neural development and neurodegeneration, as exemplified for A β 42 toxicity and neurogenic stem cell plasticity.

Significance

Neural stem cells (NSC) are reservoir for new neurons in human brains, yet they fail to form neurons after neurodegeneration. Therefore, understanding the potential use of NSCs for stem cell-based regenerative therapies requires tissue-mimetic humanized experimental systems. We report the adaptation of a 3D bio-instructive hydrogel culture system where human NSCs form neurons that later form networks in a controlled microenvironment. We also modeled neurodegenerative toxicity by using Amyloid-beta4 peptide, a hallmark of Alzheimer's disease, observed phenotypes reminiscent of human brains, and determined the global gene expression changes during development and degeneration of neurons. Thus, our reductionist humanized culture model will be an important tool to address NSC plasticity, neurogenicity, and network formation in health and disease.

Text

Introduction

Human brain development and neuronal diseases cannot be modeled adequately by current animal models (LaFerla and Green, 2012); therefore, the development of novel humanized systems that manifest neurogenic plasticity is necessary. The brain's plasticity provides an endogenous reservoir of cells that could be harnessed to physiologically enhance brain capacity or for neuronal repair (Gage and Temple, 2013; Wyss-Coray, 2016). Therefore, it is fundamentally important to understand how neurons develop to form a hard-wired network, how new networks are generated when newly formed neurons are incorporated, and how stem cells contribute to these processes. Animal models and cell culture experiments examining how mammalian brains develop and elucidating the molecular programs regulating this process have been invaluable (Molyneaux et al., 2007). However, the human brain might exhibit differences compared with the model systems that might not be discernable with existing tools.

The human brain cannot repair the loss of neurons caused by neurodegenerative diseases (ND), in part due to reduced stem cell proliferation and neurogenesis (Tincer et al., 2016). These combinatorial effects exacerbate the manifestation of the disease. In ND states in humans, overall plasticity is severely decreased (Heneka et al., 2015; Nalbantoglu et al., 1997; Selkoe, 2002), however, we know little about how to circumvent this reduction due to lack of appropriate experimental systems. Alzheimer's disease (AD) is the most common ND, and one of its hallmarks is the aggregation of amyloid protein cleavage products – mainly amyloid- β -42 (A β 42) peptides (Esler and Wolfe, 2001; Haass and Selkoe, 2007; Nalbantoglu et al., 1997). The disease state produces an inhospitable environment lacking homeostasis in which stem cells are unable to form neurons and new cells do not survive and successfully integrate into existing circuitry. Therefore, understanding stem cell plasticity and neuronal behavior in disease-related settings is critical to determine if a stem cell can regain proliferative and neurogenic function or whether a newborn neuron can survive and integrate into the remaining circuitry despite prevalent amyloid toxicity in the brain.

The overall plasticity of the human brain requires neural stem cell (NSC) proliferation, neurogenesis and neuronal network formation (Alvarez-Buylla et al., 2002; Gage, 2000). However, although NSCs in human brain possess the plasticity to fulfill all these steps, 2D culture conditions are insufficient to generate the connected arbors and long-term behaviors observed in the brain (Haycock, 2011; Justice et al., 2009). Culture conditions that drive the generation of neurons that retain a mature neuronal morphology and form synapses and 3D patterns are needed to address how the entire spectrum of plasticity manifests in the human brain. In addition, paracrine effects from the interplay between glia and neurons, the nascent cellular microenvironment, and the extracellular matrix (ECM) regulate cell fate decisions (Lutolf et al., 2009). In particular, the modulation of mechanical cues, the degradability of the matrix, and the administration of soluble effectors is known to control stem cell fate (Discher et al., 2009). Therefore, we require advanced customized and controllable assay systems for human neural stem cells.

Animal models of AD are unable to recapitulate the entire human disease spectrum (LaFerla and Green, 2012), suggesting that human cells might have a different physiological response than animal cells in response to NDs. Additionally, revolutionary 3D technologies are useful tools for addressing specific questions; however, they require highly complex culture conditions, are difficult to establish and reproduce, and the content of the scaffolds often have unintended consequences on the encapsulated cells. Therefore, *in vitro* systems utilizing human cells to model neural development or neurodegeneration in an *in vivo*-like 3D environment that is amenable to manipulation and monitoring would be highly beneficial. Therefore, in this study, we adapted a highly tunable and defined glycosaminoglycan (GAG)-based 3D matrix system (Chwalek et al., 2014; Maitz et al., 2013) to culture primary and induced human neural stem cells, which manifest their plasticity and neurogenic capacity. This system allowed us to systematically vary the local cellular environment in terms of stiffness, degradability and presentation of GAG-affine signaling molecules (Capilla and Linhardt, 2002) to identify optimal conditions that reproducibly form networks of mature neurons and glia that serve as neural stem cells. We also recapitulated important aspects of AD by modeling amyloid toxicity. A β 42 treatment impaired network formation and progenitor cell proliferation and induced human pathological hallmarks, including Tau hyperphosphorylation and neurofibrillary tangle formation, suggesting that our biohybrid 3D culture system can be used to address questions regarding neural stem cell proliferation, neurodevelopment and neurodegenerative diseases in a reductionist and tissue mimetic pre-clinical setting.

Results

To generate a culture system that would allow NSCs to manifest their plasticity and neurogenic capacity in a tissue-representative manner, we applied a modular biohybrid material based on star-shaped poly(ethylene glycol) (starPEG) and GAG heparin (HEP) that can be used to independently tune mechanical cues and biomolecular functionalization (Freudenberg et al., 2012; Tsurkan et al., 2013) when embedding primary human neural stem cells (human cortical astrocytes, from here on primary

NSCs) derived from fetal tissue at gestation week 21 (Fig. 1A). We systematically varied the biohybrid matrix in terms of stiffness, cell-responsive remodeling potential, and the presence of soluble effector mediating GAGs to induce the cellular morphology reminiscent of *in vivo* (Supplementary Figure 1). After varying the stiffness (Young's modulus) of the hydrogels from 0.5 kPa to 3 kPa (Fig. 1B,B'), 1.2 kPa was the optimal stiffness to promote the formation of extended neuronal networks by primary NSCs (Fig. 1B'). Although soft gels (0.5 kPa) disintegrated within 1 week (data not shown), stiff gels (3 kPa) resulted in round non-proliferative cells that do not grow into the matrix or form a network (Fig. 1B').

Tissue characteristics and patterning can be influenced by the cell-responsive remodeling of biohybrid hydrogels, which can be achieved by the incorporation of matrix-metalloprotease-cleavable linkers (Chwalek et al., 2011; Tsurkan et al., 2010). Using MMP-cleavable hydrogels (starPEG-MMP-HEP) (Fig. 1C,C'), directly linked starPEG-HEP gels (Fig. 1D,D') and non-degradable controls with a scrambled (MMP-insensitive) peptide sequence (starPEG-scr-HEP) (Fig. 1E,E'), we found that only MMP-responsive hydrogels induced the formation of a neuronal network (Fig. 1C). Given that the optimal stiffness was 1200 Pa for all of these gels (Fig. 1F), these results clearly point to enzymatic remodeling involving MMPs (Agrawal et al., 2008) as a crucial process for axodendritic outgrowth and the formation of mature arbors. Moreover, if an enzymatically degradable hydrogel of inert starPEG was formed, the cells did not proliferate and did not display axodendritic extensions, highlighting the importance of a factor regulating GAG-heparin interaction within the matrices (data not shown). Because heparin can bind multiple insoluble matrix proteins (e.g., laminin, fibronectin) and other soluble growth factors (Capila and Linhardt, 2002; Garg et al., 2011), it might mediate the activity and presentation of cell-secreted molecules and thus indirectly control cell fate processes. Sulfated GAGs, particularly heparan sulfate and chondroitin sulfate, are well known to be a major component of neuronal ECMs and are crucial for developmental processes and axon guidance (Lau et al., 2013). Thus, the MMP-cleavable heparin hydrogel matrix provided optimal

conditions with respect to initial stiffness and the presence of sulfated GAGs and promoted remodeling in a well-orchestrated and timely way via MMP-sensitive peptide linkers.

Since the development of 3D cultures requires MMP-cleavage, we hypothesized that the observed maturation and patterning might follow a replacement of the initial scaffold with the cells' own matrix. Therefore, we immunostained the cultures for fibronectin and laminin to test this hypothesis and found that the 3D gels generate fibronectin (Fig. 1G,H) and laminin (Fig. 1I,J) *de novo*, suggesting that neural stem cells and neurons remodel the matrix allowing to generate a liberal stem cell niche according to the needs of the NSCs. To investigate whether this remodeling would alter the stiffness of the gel matrix, we performed atomic force microscopy analyses of the gels at different time points and observed that the initial elastic modulus of 1.3 kPa is reduced gradually to 0.3 kPa after 14 days (Fig. 1K), indicating that the development of starPEG-Heparin 3D cultures generate human brain-mimetic physical properties. This is particularly important because widely used 3D scaffolds (e.g.: Matrigel) cannot be regulated in their stiffness by the encapsulated cells, and therefore they deviate from tissue-mimetic physical features. Indeed, when we compared the network forming ability of primary human NSCs in identical culture conditions in starPEG-Heparin and Matrigel, we found that the neurogenic capacity and network-formation ability of primary NSCs manifest significantly better in starPEG-Heparin gels (Supplementary Figure 2), indicating that starPEG-Heparin composition favors an unprecedented tissue-mimetic environment for primary NSCs.

Based on our cell-matrix interaction results, we hypothesized that the maturation of neuronal networks (Fig. 1L) might resemble human neurodevelopment. We analyzed the gels at various time points to address this hypothesis (Fig. 1M-P). One week after seeding, the gel contained sparsely distributed GFAP-positive glia with a 3D arborized morphology (Fig. 1M). After two weeks of culture, we started to observe acetylated tubulin-positive neurons extending processes and organizing themselves into clusters (Fig. 1N; Suppl. Video 1). At 3 weeks, the cultures produced an extensive and elaborate network of neurons with

interspersed glia (Fig. 1O) in close association with neurons (Fig. 1P). We also determined that the primary NSC cultures expressed neural stem cell markers such as SOX2 and GFAP, and neural fate determinants such as NEUROD in spatially overlapping, but distinct domains (Fig. 1Q, X-axis view; Fig. 1R, Z-axis view), indicating that neural stem cell plasticity programs and neurogenic activity drive the development of the neuronal networks, and the stem cell compartments may pattern in a 3D topology similar to human brains.

Based on our findings, we hypothesized that 3D cultures would provide a superior 3D topological environment to pNSCs, and therefore would instruct a gene expression profile that would be closer to *in vivo*. To investigate how 3D cultures would molecularly differ from 2D cultures, we cultured primary NSCs in 2D and 3D using identical culture conditions for three weeks. After isolating total mRNA, we performed whole transcriptome sequencing, and observed that a considerable number of genes are differentially expressed between 2D and 3D cultures (Fig. 2A, Supplementary Dataset 1). In order to identify the pathways and molecular programs represented better in 3D cultures, we performed pathway and enrichment analyses (Fig. 2B). We found that 3D cultures express genes related to several pathways that characterize mature neuronal physiology such as focal adhesion, ECM-receptor interaction, axon guidance, and various signaling pathways (Fig. 2B, Suppl. Dataset 2). Additionally, cellular component analyses indicate that the 3D cultures express the genes, the protein products of which are related to various mature neuronal processes such as synapses and axons (Fig. 2B). These results indicate that our hydrogel cultures provide the 3D topology and instructive environment to generate neuronal networks from primary human NSCs in a tissue-mimetic manner, which is not the case for 2D cultures.

To verify that the 3D cultures of primary human NSCs generate mature neurons resembling the *in vivo* conditions, we immunostained 3D cultures for the synaptic marker Synaptophysin (Fig. 2C), which clusters at neuronal junctions (Fig. 2D) and boutons (Fig. 2E), and generate pre- and post-synaptic termini (Fig. 2F), indicating that neurons in 3D cultures

develop enough to form synaptic connections, which we did not observe in 2D conditions (data not shown). We found that the neurons formed in the 3D starPEG-HEP-based hydrogels were also expressing neurotransmitter receptors such as VGLUT1 (Fig. 2G), and are responsive to neurotransmitters such as glutamate, as shown by increased intracellular calcium levels (Fig. 2H-J; Suppl. Video 2) following the transfection of plasmids expressing the GCaMP6f calcium sensor driven by the CMV promoter. We also performed electrophysiology experiments assessing sodium and potassium channel activity to determine whether the cells from the matrices were functional neurons. We observed potassium and sodium channel activity as well as spontaneous firing of neurons in whole cell patch clamp recordings (Fig. 2K), indicating that cells cultured within the hydrogels differentiated into functionally active neurons. These results indicate that starPEG-HEP-hydrogel-based 3D cultures of primary human NSCs are able to generate an elaborate and mature network of neurons and glia in a three dimensional organization.

To determine the cortical subtypes produced in 3D and 2D conditions, we compared the expression levels of a selected set of cortical marker genes in relation to their spatial confinement to different layers (Fig. 2L; from (Molyneaux et al., 2007)). We observed that 3D conditions favor for expression of a larger and more complete set of cortical marker genes especially associated with layers IV, V and VI, compared to 2D (Fig. 2L). Similarly, although 2D and 3D conditions allow expression of NSC markers at high levels, the major cortical pro-neural gene ASCL1 is expressed in rather low levels in 2D conditions while 3D cultures allow significantly higher levels of ASCL1 expression (Fig. 2M). To validate our deep sequencing results and heat map analyses, we performed immunohistochemistry for pro-neural determinant ASCL1 and early neuronal marker DCX, cortical layer markers SATB2, CTIP2, Reelin (RELN), DBX1, FOXO1, FOXP2, CRYM, and mature neuronal markers Neurofilament-200 and NeuN (Fig. 2N). We found that 3D cultures allow expression of pro-neural fate determinants, and production of different lineage subtypes of cortical neurons (Fig. 2N).

To test whether we could use our system with other types of neural stem cells, we cultured iPSC-derived human neural stem cells (iNSCs) in culture under conditions identical to those of primary NSCs (pNSCs), and observed extensive neurogenesis and network formation similar to pNSCs at 3 weeks of cultures (Fig. 3A,B, Suppl. Movie 3), suggesting that our instructive hydrogel composition can support development of networks from various types of neural stem cell sources.

To compare the molecular expression profiles of pNSC and iNSC cultures, we isolated total RNA and performed whole transcriptome sequencing on iNSCs and compared the reads to those of the cultures with pNSCs (Fig. 3C, Suppl. Dataset 3). We found that although both iNSCs and pNSCs can generate neuronal networks in 3D cultures, these two cell types differ significantly in their gene expression patterns (Fig. 3C; Suppl. Dataset 4). Interestingly, we found that compared to iNSCs, pNSC cultures express genes related to extracellular matrix and plasma membrane more (Fig. 3D; Suppl. Dataset 5), and these differences enrich pathways such as various signaling pathways, axon guidance, neuroactive ligand-receptor interaction and metabolism (Fig. 3D; Suppl. Dataset 6). Furthermore, when we compared cortical layer marker expression in iNSCs and pNSCs, we found that compared to pNSCs, iNSC cultures cannot form a subset of neuronal lineages especially for layers II, III and IV in our particular culture conditions (Fig. 3E). These findings suggest that our 3D culture system can be used to dissect the properties and neurogenic capacities of different progenitor types in particular culture settings and under certain physical parameters, and may serve as a suitable tool for investigating the physiological differences between induced and primary NSC populations.

Since our culture system can form cortical neurons, the development of which relies on the plasticity and neurogenic ability of human NSCs, we hypothesized that we could model disease conditions that lead to impaired NSC plasticity and neurogenic output. In the human brain, A β 42 aggregation impairs neuronal network formation and neuronal connectivity due to death of existing neurons as well as loss of neurogenic ability and neural stem cell plasticity (Hardy and Selkoe, 2002; Kienlen-

Campard et al., 2002; LaFerla et al., 2007; Selkoe, 2002; Tincer et al., 2016). A β 42 was shown to negatively affect the plasticity and neurogenesis of NSCs in mouse models of AD (Ermini et al., 2008; Haughey et al., 2002; Heo et al., 2007), and the same effect is one of the principle limitations of current neuro-regenerative approaches to the treatment of AD in humans (Cosacak et al., 2015; Demars et al., 2010; Tincer et al., 2016). However, the mechanisms underlying the impact of A β 42 on NSC plasticity are still largely unknown and cannot be elucidated analytically in human brains. We therefore extended our above-described 3D cultures by treating primary NSCs with A β 42, a major hallmark of AD pathology, before embedding them in the biohybrid hydrogels (Figure 4A). We used an A β 42 form that we previously found to be causing pathological outcomes in vertebrate brains (Bhattacharai et al., 2016; Bhattacharai et al., 2017b). In control gels, the neurons formed highly connected networks (Fig. 4B) that were disrupted upon A β 42 treatment (Fig. 4C). We developed an algorithm to trace the connected neuronal paths as skeletonized arbors and quantified the extent of the neuronal connections, and observed that neuronal networks reduce significantly after Amyloid toxicity (Fig. 4D-F). Similar to human brains, A β 42 resulted in dystrophic axons (Fig. 4G-G''; Supplementary Video 5), impaired ECM composition and stiffness (Figure 4H-J), Tau hyperphosphorylation (Figure 4K-N, Supplementary Movie 5), neurofibrillary tangle formation as observed by Gallyas silver impregnation (Fig. 4m, Suppl. Fig. 3) and Thioflavin S staining (Fig. 4N), microtubule disassembly (Figure 4O), and amyloid aggregation and autophagy (Figure 4P,Q).

Amyloid toxicity not only impairs neurogenesis and neuronal survival but also reduces synaptic plasticity (Selkoe, 2002). Amyloid load prevents the formation of new synapses, and newly added cells cannot integrate into the circuitry, rendering exogenous stem cell therapy inefficient (Lilja et al., 2015; Tong et al., 2015). We developed a transplantation paradigm with cultured gels to test whether our 3D culture model could be used to address questions regarding the neurogenic potential and capacity of transplanted cells to integrate into existing networks (Fig. 4R). We labeled all pNSCs with a nuclear stain, and injected them into another hydrogel

that had been pre-cultured with embedded pNSCs for 1 week (Fig. 4R). At 1 week after transplantation, the injected cells formed neurons with arbors (red nuclei, Fig. 4R; Supplementary Video 7) and connected to pre-existing cells in the control gels (Fig. 4R,R'). In contrast, cells injected into A β 42-containing gels did not acquire an arborized morphology or connect to the existing cells that were also not arborized (Fig. 4S,S'; Supplementary Video 8). Combined with the findings that A β 42 impairs the synaptic connections overall (Fig. 4T), these results suggest that 3D cultures can be used for analyzing how synaptic connections can be regenerated and how new neurons can be forced to integrate into the existing circuitry upon A β 42 toxicity.

A β 42 reduces the NSC plasticity and neurogenic capacity in human brains, and stem cell-based regenerative therapies would require therapeutic activation of NSCs (Tincer et al., 2016). To investigate whether A β 42 reduced NSC plasticity in our 3D cultures, we determined the proliferative capacity and prevalence of pNSCs after A β 42 by BrdU/EdU treatment and immunohistochemical stainings for NSC markers SOX2 and GFAP (Fig. 4U-X). We treated the cultures with BrdU at 1 week of development and with EdU at 2 weeks of development, and analyzed the presence of BrdU and EdU positive cells at 3 weeks of cultures where double-positive cells would indicate constitutively proliferating stem cells (Fig. 4U,V). We found that A β 42 reduced BrdU-EdU incorporation and the number of constitutively proliferating cells (Fig. 4Y) as well as reducing numbers of GFAP and SOX2-positive NSCs (Fig. 4Y), indicating that our 3D culture system can be used for modeling A β 42-induced impairment of NSC plasticity. Thus, our 3D cultures of NSCs can also successfully serve as a novel *in vitro* sporadic AD model of A β 42 toxicity on human NSC plasticity and neurogenesis (Figure 4Z). Importantly, we found that all those effects of A β 42 observed in 3D cultures are specific because scrambled A β 42 or other A β species such as A β 38 do not show any phenotypes above (Suppl. Figure 4).

To determine if A β 42 would affect the neuronal network formation and NSC plasticity in iPSC-derived cultures, we treated the iNSCs with A β 42, encapsulated in 3D gels. To determine the neuronal network formation,

NSC prevalence, and proliferative capacity, we performed immunostainings for GFAP and TUBB3 (Fig. 5A,B), SOX2 and GFAP (Fig. 5C,D), and GFAP and BrdU (Fig. 5E,F) together with A β 42 detection. Compared to controls, A β 42 reduces the total number of newborn cells and NSCs (GFAP/SOX2-positive) (Fig. 5G), average number of networks (Fig. 5H), average number of branches per network (Fig. 5I), average branch length per network (Fig. 5J), length of longest connected path (Fig. 5K), and maximum branch length (Fig. 5L) in cultures started with iNSCs. These results show that A β 42 impairs the plasticity, neurogenic ability and network forming capacity of primary and human NSC cultures in our 3D hydrogel matrix, and our system can also be used for iPSC-derived cell types.

To determine the gene expression changes exerted by A β 42 in 3D cultures, we performed whole transcriptome sequencing on control and A β 42-treated cultures initiated with human iNSCs (Fig. 5M, Suppl. Dataset 7) and pNSCs (Fig. 5N, Suppl. Dataset 8). Cellular component analyses and KEGG pathway enrichment analyses in iNSC-derived (Fig. 5O, Suppl. Dataset 9, 10) and primary (Fig. 5P, Suppl. Dataset 11, 12) cultures showed that divergent pathways are affected by A β 42 in 3D cultures. Hierarchical clustering (Fig. 5Q) and multivariate analyses (Fig. 5R) indicated that primary and induced NSC cultures have their own molecular signatures of gene expression, which are affected by A β 42. By plotting a heat map of gene expression changes, we also found that several cortical marker genes are differentially expressed after A β 42 in induced and primary human NSC-based 3D cultures (Fig. 5S), which is suggestive of the alterations in the cortical neuronal subtypes (for instance, POU3F2, CRYM, and FOXO1). Our results indicate that although A β 42 causes impaired neural stem cell plasticity, neurogenesis and network formation in both iNSC- and pNSC-derived cultures (Fig. 4, Fig. 5), the molecular programs it alters in these cultures do differ (Fig. 5O, P, S). This finding is consistent with previous documentations that primary and iPSC-derived neural stem cells have profound physiological differences that might affect subsequent global gene expression, neuronal maturation capacity, and resilience to disease conditions (Kim et al., 2011; Kim et al., 2010;

Verpelli et al., 2013; Xia et al., 2016). Therefore, our results suggest that starPEG-Heparin 3D culture system can also be used to dissect the effects of A β 42 on different stem cell and neuronal populations. Furthermore, our 3D cultures may help refining the overall toxicity of A β 42 to distinct physiological states and cellular characteristics of experimental cellular systems. Since our cultures do not contain inflammatory cells, this system will also help to investigate the direct roles of A β 42 on stem cells and neurons in a reductionist and dissection manner.

Discussion

Neurodegenerative diseases such as AD present with a perplexing set of impairments, including neuronal death, synaptic degeneration, and the inability of stem cells to produce neurons to replace lost neurons (Kienlen-Campard et al., 2002; LaFerla et al., 2007; Lindvall and Kokaia, 2006; Selkoe, 2002, 2003). Thus, designing effective regenerative therapies for patients with AD requires assay systems that address the parameters of neurodegenerative pathology individually and in combination in a reductionist way using human cells. Rodent models of neurodegeneration cannot recapitulate various aspects of human pathology (Gotz and Ittner, 2008; LaFerla and Green, 2012). Thus, we used primary and iPSC-derived human neural stem cells (iNSCs) in 3D cultures to better reflect neurodevelopmental paradigms and neurodegenerative processes in a reductionist manner. 3D cultures using a well-defined biohybrid hydrogel system based on star-shaped PEG and heparin allowed the generation of extensive neuronal networks that could be used to address neurogenesis and connectivity-based questions.

Existing 3D culture systems, including MatrigelTM-based cultures, are chemically undefined and heterogeneous in composition and cannot be modified for various parameters, such as stiffness, scaffold composition or bioresponsiveness (Ravi et al., 2015). Thus, the interpretation of often quite variable results is difficult, and it is rarely possible to dissect the influences of different exogenous and paracrine signals on cellular development in an isolated and controllable experimental setup. Our biohybrid hydrogel system based on heparin and star-shaped PEG provides valuable advantages by enabling the independent and ongoing

adjustment of biophysical and biomolecular matrix signals. Indeed, we determined that starPEG-Heparin gels allow faster and more elaborate network formation compared to Matrigel upon identical culture conditions (Suppl. Fig 2). Moreover, the method we use to generate the gels does not cause cell death or DNA mutation that occurs in other matrices through the formation of free radicals upon polymer network formation.

Various 3D systems, including organoids, cannot form reproducibly sized or formed structures (Fatehullah et al., 2016). However, MMP-sensitive, starPEG-heparin-based 3D cultures can be customized for all of these parameters, providing better-defined conditions. Compared to previous reports modeling AD in 3D cultures using Matrigel (Choi et al., 2014; Smith et al., 2015), despite differences in the initial cell source, starPEG-heparin 3D gels enabled significantly faster development of neuronal networks that offer various advantages such as high-throughput screening approaches that are suitable for drug discovery.

The composition and architecture of the ECM is an integral parameter governing stem cell activity and tissue modeling. However, due to the complex interplay between multiple ECM-derived signals and their pleiotropic effects, *in vivo* assays pose a challenge to identifying the roles of exogenous cues in tissue patterning. Previously described methods for the formation of three-dimensional neuronal networks (Choi et al., 2014; Fatehullah et al., 2016; Tang-Schomer et al., 2014; Zhang et al., 2014) lack a defined composition and the ability to tune the artificial extracellular matrix. Our novel 3D matrix platform would be advantageous to dissect the roles of matrix properties in stem cell activity and differentiation, because cells can dynamically interact with the scaffold to generate their “own” cell-secreted ECM. Additionally, our hydrogels can be covalently functionalized with different matrix-derived peptides or could be used for the effective administration of GAG-affine soluble signal molecules. By doing so, the effects of exogenous cues could be individually tested on human neural stem/progenitor cell proliferation and neuronal network formation. Similarly, cells in 3D cultures can be transfected with plasmids for misexpression studies, a powerful tool for rapid investigation of gene function.

Our new 3D culture platform provides a novel and improved model to study the plasticity of human neural stem cells and disease states in real time. The application of customized A β 42 peptides for optimized cellular uptake within the hydrogel-based 3D cultures indicates that our advanced culture method can recapitulate the major pathophysiology of human A β 42 toxicity. Additionally, our gels can be used to experimentally investigate how new cells are incorporated into diseased brains, and how different sources of neural stem cells would molecularly affect the outcomes of A β 42 toxicity (e.g.: primary versus induced neural stem cells). Individual differences between cell types and their response to disease stimuli can also be measured in a tissue-mimetic composition using our gel system. In overall, starPEG-heparin-hydrogel-based 3D human neural stem cell culture is a novel and comprehensive method for analyzing various stages of neural development and disease of the human cortex, from stem cell proliferation to neurogenesis and from neuronal maturation to integration into the circuitry, in a highly defined and controllable method in an *in vitro* environment. Our system can be expanded for examining embryonic stem cells, organoids or adult-derived cortical cells from humans, and analyze their stem cell properties or neurogenic capacity in a comparative manner to complement previous studies (Choi et al., 2014; Koutsopoulos and Zhang, 2013; Zhang et al., 2014). Beyond that, our established 3D hydrogel cultures can be expected to enable personalized medicine approaches targeting brain diseases or drug efficacy tests.

Author contributions: C.P. and C.K. conceived and designed the experiments. M.I.C. analyzed the next generation sequencing data and performed the bioinformatics analyses. L.B., U.F., and C.W. provided the gel materials, J.F. performed AFM studies. C.P. and H.C. performed cell cultures, imaging and quantifications. P.B., H.H. and V.M helped the cell cultures. C.P. optimized the culture conditions for iPSCs and Matrigel. A.K.T. and Y.Z. provided the Amyloid peptides. X.C., S.H. and C.L.A. performed whole-cell patch clamping. C.K. wrote the manuscript, C.K., C.P., U.F., and C.W. revised the manuscript.

Acknowledgements

This work was supported by DZNE and Helmholtz Association (VH-NG-1021, C.K.), DFG (KI1524/6, C.K.), (AN797/4-1, C.L.A.) and (CRC TR 67, CRC SFB 655, FOR/EXC999, C.W.); and BMBF (PRECIMATRIX-FKZ-03XP0083-310117, C.W.). We wish to thank M. Wagner for electrophysiology, R. Szech for image rendering, and T. Kurth for electron microscopy.

References

- Agrawal, S.M., Lau, L., and Yong, V.W. (2008). MMPs in the central nervous system: where the good guys go bad. *Semin Cell Dev Biol* 19, 42-51.
- Alexa, A., Rahnenfuhrer, J., and Lengauer, T. (2006). Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. *Bioinformatics* 22, 1600-1607.
- Alvarez-Buylla, A., Seri, B., and Doetsch, F. (2002). Identification of neural stem cells in the adult vertebrate brain. *Brain Res Bull* 57, 751-758.
- Bhattacharai, P., Thomas, A.K., Cosacak, M.I., Papadimitriou, C., Mashkaryan, V., Zhang, Y., and Kizil, C. (2017a). Modeling Amyloid- β 42 Toxicity and Neurodegeneration in Adult Zebrafish Brain. *Journal of Visualized Experiments* 128.
- Bhattacharai, P., Thomas, A.K., Papadimitriou, C., Cosacak, M.I., Mashkaryan, V., Froc, C., Kurth, T., Dahl, A., Zhang, Y., and Kizil, C. (2016). IL4/STAT6 signaling activates neural stem cell proliferation and neurogenesis upon Amyloid- β 42 aggregation in adult zebrafish brain. *Cell Reports* 17, 941-948.
- Bhattacharai, P., Thomas, A.K., Zhang, Y., and Kizil, C. (2017b). The effects of aging on Amyloid- β 42-induced neurodegeneration and regeneration in adult zebrafish brain. *Neurogenesis*.
- Bray, L.J., Binner, M., Holzheu, A., Friedrichs, J., Freudenberg, U., Hutmacher, D.W., and Werner, C. (2015). Multi-parametric hydrogels support 3D in vitro bioengineered microenvironment models of tumour angiogenesis. *Biomaterials* 53, 609-620.
- Capila, I., and Linhardt, R.J. (2002). Heparin-protein interactions. *Angew Chem Int Ed Engl* 41, 391-412.
- Capilla, I., and Linhardt, R.J. (2002). Heparin-protein interactions,. *Angew Chem Int Ed Engl* 41, 391-412.
- Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., *et al.* (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499, 295-300.
- Choi, S.H., Kim, Y.H., Hebisch, M., Sliwinski, C., Lee, S., D'Avanzo, C., Chen, H., Hooli, B., Asselin, C., Muffat, J., *et al.* (2014). A three-dimensional human neural cell culture model of Alzheimer's disease. *Nature* 515, 274-278.
- Chwalek, K., Bray, L.J., and Werner, C. (2014). Tissue-engineered 3D tumor angiogenesis models: Potential technologies for anti-cancer drug discovery. *Adv Drug Deliv Rev*.
- Chwalek, K., Levental, K.R., Tsurkan, M.V., Zieris, A., Freudenberg, U., and Werner, C. (2011). Two-tier hydrogel degradation to boost endothelial cell morphogenesis. *Biomaterials* 32, 9649-9657.

583 Cosacak, M.I., Papadimitriou, C., and Kizil, C. (2015). Regeneration, Plasticity, and
584 Induced Molecular Programs in Adult Zebrafish Brain. *Biomed Res Int* 2015:769763.

585 Demars, M., Hu, Y.S., Gadadhar, A., and Lazarov, O. (2010). Impaired neurogenesis
586 is an early event in the etiology of familial Alzheimer's disease in transgenic mice. *J*
587 *Neurosci Res* 88, 2103-2117.

588 Discher, D.E., Mooney, D.J., and Zandstra, P.W. (2009). Growth factors, matrices,
589 and forces combine and control stem cells. *Science* 324, 1673-1677.

590 Ermini, F.V., Grathwohl, S., Radde, R., Yamaguchi, M., Staufenbiel, M., Palmer,
591 T.D., and Jucker, M. (2008). Neurogenesis and alterations of neural stem cells in
592 mouse models of cerebral amyloidosis. *Am J Pathol* 172, 1520-1528.

593 Esler, W.P., and Wolfe, M.S. (2001). A portrait of Alzheimer secretases--new features
594 and familiar faces. *Science* 293, 1449-1454.

595 Fabig, G., Kretschmar, S., Weiche, S., Eberle, D., Ader, M., and Kurth, T. (2012).
596 Labeling of ultrathin resin sections for correlative light and electron microscopy.
597 *Methods Cell Biol* 111, 75-93.

598 Falcon, S., and Gentleman, R. (2007). Using GOstats to test gene lists for GO term
599 association. *Bioinformatics* 23, 257-258.

600 Fatehullah, A., Tan, S.H., and Barker, N. (2016). Organoids as an in vitro model of
601 human development and disease. *Nat Cell Biol* 18, 246-254.

602 Freudenberg, U., Sommer, J.U., Levental, K.R., Welzel, P.B., Zieris, A., Chwalek, K.,
603 Schneider, K., Prokoph, S., Prewitz, M., Dockhorn, R., *et al.* (2012). Using Mean
604 Field Theory to Guide Biofunctional Materials Design. *Adv Funct Mater* 22, 1391-
605 1398.

606 Gage, F.H. (2000). Mammalian neural stem cells. *Science* 287, 1433-1438.

607 Gage, F.H., and Temple, S. (2013). Neural stem cells: generating and regenerating the
608 brain. *Neuron* 80, 588-601.

609 Garg, H.G., Linhardt, R.J., and Hales, C.A. (2011). Chemistry and Biology of
610 Heparin and Heparan Sulfate (Elsevier).

611 Gotz, J., and Ittner, L.M. (2008). Animal models of Alzheimer's disease and
612 frontotemporal dementia. *Nat Rev Neurosci* 9, 532-544.

613 Haass, C., and Selkoe, D.J. (2007). Soluble protein oligomers in neurodegeneration:
614 lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8, 101-
615 112.

616 Hardy, J., and Selkoe, D.J. (2002). The amyloid hypothesis of Alzheimer's disease:
617 progress and problems on the road to therapeutics. *Science* 297, 353-356.

618 Haughey, N.J., Liu, D., Nath, a., Borchard, a.C., and Mattson, M.P. (2002).
619 Disruption of neurogenesis in the subventricular zone of adult mice, and in human
620 cortical neuronal precursor cells in culture, by amyloid beta-peptide: implications for
621 the pathogenesis of Alzheimer's disease. *Neuromolecular Med* 1, 125-135.

622 Haycock, J.W. (2011). 3D cell culture: a review of current approaches and techniques.
623 *Methods Mol Biol* 695, 1-15.

624 Heneka, M.T., Carson, M.J., El Khoury, J., Landreth, G.E., Brosseron, F., Feinstein,
625 D.L., Jacobs, A.H., Wyss-Coray, T., Vitorica, J., Ransohoff, R.M., *et al.* (2015).
626 Neuroinflammation in Alzheimer's disease. *The Lancet Neurology* 14, 388-405.

627 Heo, C., Chang, K.-A., Choi, H.S., Kim, H.-S., Kim, S., Liew, H., Kim, J.a., Yu, E.,
628 Ma, J., and Suh, Y.-H. (2007). Effects of the monomeric, oligomeric, and fibrillar
629 A β 42 peptides on the proliferation and differentiation of adult neural stem cells from
630 subventricular zone. *Journal of Neurochemistry* 102, 493-500.

631 Justice, B.A., Badr, N.A., and Felder, R.A. (2009). 3D cell culture opens new
632 dimensions in cell-based assays. *Drug Discov Today* 14, 102-107.

633 Kienlen-Campard, P., Miolet, S., Tasiaux, B., and Octave, J.N. (2002). Intracellular
634 amyloid-beta 1-42, but not extracellular soluble amyloid-beta peptides, induces
635 neuronal apoptosis. *J Biol Chem* 277, 15666-15670.

636 Kim, J., Efe, J.A., Zhu, S., Talantova, M., Yuan, X., Wang, S., Lipton, S.A., Zhang,
637 K., and Ding, S. (2011). Direct reprogramming of mouse fibroblasts to neural
638 progenitors. *Proc Natl Acad Sci U S A* 108, 7838-7843.

639 Kim, K., Doi, A., Wen, B., Ng, K., Zhao, R., Cahan, P., Kim, J., Aryee, M.J., Ji, H.,
640 Ehrlich, L.I., *et al.* (2010). Epigenetic memory in induced pluripotent stem cells.
641 *Nature* 467, 285-290.

642 Kizil, C., Iltzsche, A., Kuriakose, A., Bhattarai, P., Zhang, Y., and Brand, M. (2015).
643 Efficient cargo delivery using a short cell-penetrating peptide in vertebrate brains.
644 *PLoS One* 10, e0124073.

645 Koutsopoulos, S., and Zhang, S. (2013). Long-term three-dimensional neural tissue
646 cultures in functionalized self-assembling peptide hydrogels, matrigel and collagen I.
647 *Acta Biomater* 9, 5162-5169.

648 LaFerla, F.M., and Green, K.N. (2012). Animal models of Alzheimer disease. *Cold*
649 *Spring Harbor perspectives in medicine* 2.

650 LaFerla, F.M., Green, K.N., and Oddo, S. (2007). Intracellular amyloid-beta in
651 Alzheimer's disease. *Nat Rev Neurosci* 8, 499-509.

652 Lau, L.W., Cua, R., Keough, M.B., Haylock-Jacobs, S., and Yong, V.W. (2013).
653 Pathophysiology of the brain extracellular matrix: a new target for remyelination. *Nat*
654 *Rev Neurosci* 14, 722-729.

655 Liao, Y., Smyth, G.K., and Shi, W. (2013). The Subread aligner: fast, accurate and
656 scalable read mapping by seed-and-vote. *Nucleic Acids Res* 41, e108.

657 Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general
658 purpose program for assigning sequence reads to genomic features. *Bioinformatics*
659 30, 923-930.

660 Lilja, A.M., Malmsten, L., Röjdner, J., Voytenko, L., Verkhatsky, A., Ögren, S.O.,
661 Nordberg, A., and Marutle, A. (2015). Neural Stem Cell Transplant-Induced Effect on
662 Neurogenesis and Cognition in Alzheimer Tg2576 Mice Is Inhibited by Concomitant
663 Treatment with Amyloid-Lowering or Cholinergic $\alpha 7$ Nicotinic Receptor Drugs.
664 *Neural Plast* 2015, 370432.

665 Lindvall, O., and Kokaia, Z. (2006). Stem cells for the treatment of neurological
666 disorders. *Nature* 441, 1094-1096.

667 Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
668 and dispersion for RNA-seq data with DESeq2. *Genome biology* 15, 550.

669 Luo, W., and Brouwer, C. (2013). Pathview: an R/Bioconductor package for pathway-
670 based data integration and visualization. *Bioinformatics* 29, 1830-1831.

671 Lutolf, M.P., Gilbert, P.M., and Blau, H.M. (2009). Designing materials to direct
672 stem-cell fate. *Nature* 462, 433-441.

673 Maitz, M.F., Freudenberg, U., Tsurkan, M.V., Fischer, M., Beyrich, T., and Werner,
674 C. (2013). Bio-responsive polymer hydrogels homeostatically regulate blood
675 coagulation. *Nat Commun* 4, 2168.

676 Molyneaux, B.J., Arlotta, P., Menezes, J.R., and Macklis, J.D. (2007). Neuronal
677 subtype specification in the cerebral cortex. *Nat Rev Neurosci* 8, 427-437.

678 Nalbantoglu, J., Tirado-Santiago, G., Lahsaini, A., Poirier, J., Goncalves, O., Verge,
679 G., Momoli, F., Welner, S.A., Massicotte, G., Julien, J.P., *et al.* (1997). Impaired
680 learning and LTP in mice expressing the carboxy terminus of the Alzheimer amyloid
681 precursor protein. *Nature* 387, 500-505.

682 Ravi, M., Paramesh, V., Kaviya, S.R., Anuradha, E., and Solomon, F.D. (2015). 3D
683 cell culture systems: advantages and applications. *J Cell Physiol* 230, 16-26.

684 Selkoe, D.J. (2002). Alzheimer's disease is a synaptic failure. *Science* 298, 789-791.

685 Selkoe, D.J. (2003). Folding proteins in fatal ways. *Nature* 426, 900-904.

686 Smith, I., Silveirinha, V., Stein, J.L., de la Torre-Ubieta, L., Farrimond, J.A.,
687 Williamson, E.M., and Whalley, B.J. (2015). Human neural stem cell-derived cultures
688 in three-dimensional substrates form spontaneously functional neuronal networks. *J*
689 *Tissue Eng Regen Med*.

690 Tang-Schomer, M.D., White, J.D., Tien, L.W., Schmitt, L.I., Valentin, T.M.,
691 Graziano, D.J., Hopkins, A.M., Omenetto, F.G., Haydon, P.G., and Kaplan, D.L.
692 (2014). Bioengineered functional brain-like cortical tissue. *Proc Natl Acad Sci U S A*
693 111, 13811-13816.

694 Tincer, G., Mashkaryan, V., Bhattarai, P., and Kizil, C. (2016). Neural
695 stem/progenitor cells in Alzheimer's disease. *Yale J Biol Med* 89, 23-35.

696 Tong, L.M., Fong, H., and Huang, Y. (2015). Stem cell therapy for Alzheimer's
697 disease and related disorders: current status and future perspectives. *Experimental &*
698 *molecular medicine* 47, e151.

699 Tsurkan, M.V., Chwalek, K., Levental, K.R., Freudenberg, U., and Werner, C.
700 (2011). Modular StarPEG-Heparin Gels with Bifunctional Peptide Linkers. *Macromol*
701 *Rapid Commun* 31, 1529-1533.

702 Tsurkan, M.V., Chwalek, K., Prokoph, S., Zieris, A., Levental, K.R., Freudenberg, U.,
703 and Werner, C. (2013). Defined polymer-peptide conjugates to form cell-instructive
704 starPEG-heparin matrices in situ. *Adv Mater* 25, 2606-2610.

705 Tsurkan, M.V., Levental, K.R., Freudenberg, U., and Werner, C. (2010).
706 Enzymatically degradable heparin-polyethylene glycol gels with controlled
707 mechanical properties. *Chem Commun (Camb)* 46, 1141-1143.

708 Venable, J.H., and Coggeshall, R. (1965). A Simplified Lead Citrate Stain for Use in
709 Electron Microscopy. *J Cell Biol* 25, 407-408.

710 Verpelli, C., Carlessi, L., Bechi, G., Fusar Poli, E., Orellana, D., Heise, C.,
711 Franceschetti, S., Mantegazza, R., Mantegazza, M., Delia, D., *et al.* (2013).
712 Comparative neuronal differentiation of self-renewing neural progenitor cell lines
713 obtained from human induced pluripotent stem cells. *Front Cell Neurosci* 7, 175.

714 Wieduwild, R., Tsurkan, M., Chwalek, K., Murawala, P., Nowak, M., Freudenberg,
715 U., Neinhuis, C., Werner, C., and Zhang, Y. (2013). Minimal peptide motif for non-
716 covalent peptide-heparin hydrogels. *J Am Chem Soc* 135, 2919-2922.

717 Wu, T.D., Reeder, J., Lawrence, M., Becker, G., and Brauer, M.J. (2016). GMAP and
718 GSNAP for Genomic Sequence Alignment: Enhancements to Speed, Accuracy, and
719 Functionality. *Methods Mol Biol* 1418, 283-334.

720 Wyss-Coray, T. (2016). Ageing, neurodegeneration and brain rejuvenation. *Nature*
721 539, 180-186.

722 Xia, N., Zhang, P., Fang, F., Wang, Z., Rothstein, M., Angulo, B., Chiang, R., Taylor,
723 J., and Reijo Pera, R.A. (2016). Transcriptional comparison of human induced and
724 primary midbrain dopaminergic neurons. *Sci Rep* 6, 20270.

725 Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology
726 analysis for RNA-seq: accounting for selection bias. *Genome biology* 11, R14.

727 Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for
728 comparing biological themes among gene clusters. *OMICS* 16, 284-287.

729 Zhang, D., Pekkanen-Mattila, M., Shahsavani, M., Falk, A., Teixeira, A.I., and
730 Herland, A. (2014). A 3D Alzheimer's disease culture model and the induction of

731 P21-activated kinase mediated sensing in iPSC derived neurons. *Biomaterials* 35,
732 1420-1428.
733 Zhang, Y., Fussel, S., Reimer, U., Schutkowski, M., and Fischer, G. (2002).
734 Substrate-based design of reversible Pin1 inhibitors. *Biochemistry* 41, 11868-11877.
735
736

Main figures and legends

Figure 1

Physical properties and dynamic remodeling of 3D starPEG-Heparin hydrogels and tissue-mimetic behavior of primary human neural stem/progenitor cells (NSPCs).

(A) Simplified composition and preparation scheme of glycosaminoglycan (GAG)-based minimalist multifunctional hydrogels.

(B, B') Bright field gel image (BFI) at 1-kPa (B) and 3-kPa (B') stiffness.

(C-E') BFI (C,D,E) and immunostaining for acetylated-tubulin (aTub) and GFAP (C',D',E') in a 3-week-old gel at 1 kPa stiffness with (C, C'), without (D, D') and with scrambled MMP cleavage sites (E, E').

(F) Rheological measurements of various gel types.

(G) Acetylated-tubulin and fibronectin.

(B) KEGG pathway analyses pie chart showing significantly enriched molecular pathways.

(C-E) Synaptophysin and Acetylated tubulin immunostaining in 3D cultures.

(F) Synaptophysin, PSD95 and Acetylated tubulin immunostaining in 3D cultures.

(G) VGLUT1 and Acetylated tubulin immunostaining in 3D cultures.

(H) GCaMP6 signal without Glutamate treatment.

(I) GCaMP6 signal with Glutamate treatment.

(J) Fluorescence intensity histogram. Note the peak at the time of Glutamate treatment.

(K) Patch clamp recordings for Na⁺ (left) and K⁺ (middle) currents. Detection of spontaneous firing of neurons (right).

(L) Heat map for expression levels of cortical marker genes for 2D and 3D cultures. Asterisks indicate significant difference between two samples. Genes are denoted with their respective cortical layer expression with color codes (left to the heat map).

(M) Heat map for expression levels of neural stem cell markers.

(N) Immunostaining for ASCL1 (proneural determinant), DCX (early neuronal marker), SATB2, CTIP2, Reelin (RELN), DBX1, FOXP2, CRYM, FOXO1 (cortical markers), and Neurofilament (NF200) and NeuN (mature neuronal markers).

Scale bars: 10 μ m for C-I; 25 μ m for N. Gels: 3 weeks of culture. See also Supplementary Datasets 1, 2 and 3.

Figure 3

Culture of iPSC-derived neural stem cells in 3D, and comparison of whole transcriptome profile to primary human neural stem cell-derived cultures.

(A) Immunostaining for GFAP (red) and TUBB3 (green) on 3 weeks of 3D cultures with primary human cortical astrocytes (neural stem cells). Insets below the panel are individual fluorescence channels including DAPI (white).

(B) Immunostaining for GFAP (red) and TUBB3 (cyan) on 3 weeks of 3D cultures with iPSC-derived human neural stem cells. Insets below the panel are individual fluorescence channels including DAPI (white).

(C) MA-plot for differentially expressed genes in primary over iPSC-derived NSC cultures. Red: over-represented in primary cultures, green: underrepresented in primary cultures compared to iPSC cultures.

(D) KEGG pathway analyses pie chart showing significantly enriched molecular pathways in primary cultures compared to iPSC cultures.

(E) Heat map for expression levels of cortical marker genes in iPSC and primary cultures. Asterisks indicate significant difference between two samples. Genes are denoted with their respective cortical layer expression with color codes (left to the heat map).

Scale bars: 100 μ m. Gels: 3 weeks of culture.

See also Supplementary Datasets 4, 5 and 6.

Figure 4

3D hydrogel cultures of primary human NSCs as an Aβ42 toxicity model.

(A) Simplified gel preparation/Aβ42 administration scheme.

(B,C) X-axis views for Aβ42, Acetylated-tubulin, GFAP in control (B) and Aβ42 gels (C). Double stainings under the panels.

(D,E) Maximum projection of the skeletonized-connected neuronal pathways in control (D) and Aβ42 gels (E).

(F) Quantification.

(G-G'') 3D-reconstruction of TUBB3/Aβ42 showing dystrophic axons.

(H,I) Fibronectin/laminin in control (H) and Aβ42 gels (I).

(J) Time-course atomic force microscopy measurements of stiffness.

(K,L) Phospho-Tau (AT180) in control (K) and Aβ42 gels (L).

(M) Gallyas silver staining.

843 (N) Thioflavin-S staining in A β 42 gels.
 844 (O) EM images of microtubules in control (left) and A β 42 gels (right).
 845 (P,Q) A β 42 depositions (P), and autophagic vacuoles (Q).
 846 (R-S') Acetylated-tubulin and A β 42 in control (R) and A β 42 gels (S) after
 847 transplantation. Transplanted cell nuclei labeled with TOPRO3 (red). 3D-
 848 reconstruction of R (R') and S (S').
 849 (T) Acetylated-tubulin, synaptophysin, A β 42 in control (upper) and A β 42
 850 gels (lower).
 851 (U,V) BrdU and EdU in control (U) and A β 42 gels (V). Arrows: double-
 852 positive cells.
 853 (W,X) SOX2, GFAP, A β 42 in control (W) and A β 42 gels (X).
 854 (Y) Quantification of U-X.
 855 (Z) Schematics for effects of A β 42.
 856 Scale bars: 10 μ m for G, M, N, T, R, and S; 200 nm for O-Q; 50 μ m for
 857 the other figures. Gels: 3 weeks of culture.
 858 See also Supplementary Movies 4-8.

859

860

861 **Figure 5**

862 A β 42 toxicity model with iPSC-derived neural stem cells in 3D hydrogels,
 863 analysis of transcriptional changes, and comparison to pNSC cultures.

864

865 (A) Immunostaining for A β 42 (green), TUBB3 (cyan) and GFAP (red) on
866 control iPSC-derived NSC cultures.

867 (B) Immunostaining for A β 42 (green), TUBB3 (cyan) and GFAP (red) on
868 A β 42-treated iPSC-derived NSC cultures.

869 (C) Immunostaining for A β 42 (green), SOX2 (cyan) and GFAP (red) on
870 control iPSC-derived NSC cultures.

871 (D) Immunostaining for A β 42 (green), SOX2 (cyan) and GFAP (red) on
872 A β 42-treated iPSC-derived NSC cultures.

873 (E) Immunostaining for A β 42 (green), BrdU (cyan) and GFAP (red) on
874 control iPSC-derived NSC cultures. BrdU is given at 1 week of culture.

875 (F) Immunostaining for A β 42 (green), TUBB3 (cyan) and GFAP (red) on
876 A β 42-treated iPSC-derived NSC cultures. BrdU is given at 1 week of
877 culture.

878 (G) Quantification graph for number of cells in control and A β 42-treated
879 cultures.

880 (H-L) Quantification of average number of networks (H), average number
881 of branches per network (I), average branch length per network (J),
882 length of longest connected path (K), and maximum branch length (L).

883 (M) MA-plot for differentially expressed genes in iPSC-derived NSC
884 cultures after A β 42 treatment. Red: upregulated, green: downregulated.

885 (N) MA-plot for differentially expressed genes in primary NSC cultures
886 after A β 42 treatment. Red: upregulated, green: downregulated.

887 (O) KEGG pathway analyses pie chart showing significantly enriched
888 molecular pathways in iPSC cultures after A β 42 treatment. .

889 (P) KEGG pathway analyses pie chart showing significantly enriched
890 molecular pathways in primary cultures after A β 42 treatment.

891 (Q) Heat map for changes in expression levels of cortical marker genes in
892 iPSC and primary cultures after A β 42 treatment. Genes are denoted with
893 their respective cortical layer expression with color codes (above the heat
894 map).

895 Scale bars: 100 μ m. Gels: 3 weeks of culture.

896 See also Supplementary Datasets 7-12.

897

898

Materials and Methods

Cell Culture

Astrocytes isolated from the cerebral cortex at gestation week 21 were obtained from ScienCell Research Laboratory (SRL, Catalog Number 1800, Carlsbad, CA, USA) at passage one and delivered as frozen stocks. The cells (from here on primary human neural stem cells, pNSC) are certified to be negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi. PHCCs were seeded on conventional T75 flasks or 24-well plates and cultured in complete astrocyte medium (AM) composed of Astrocyte medium (SRL, Catalog Number 1801) supplemented with 2% fetal bovine serum (SRL, Catalog Number 0010), 1% astrocyte growth supplement (SRL, Catalog Number 1852) and 1% penicillin/streptomycin solution (SRL, Catalog Number 0503) in an incubator with a 5% CO₂/95% air atmosphere at 37 °C. Human induced neural stem cells (HIP, BC1 line, Amsbio, catalog number GSC-4311, from here on iNSC) were seeded in Geltrex pre-coated cultureware in complete AM as described in the above paragraph.

Generation of starPEG-Heparin hydrogels and cell encapsulation

StarPEG-heparin hydrogels were generated as previously described (Maitz et al., 2013; Wieduwild et al., 2013) with the following modifications: pNSCs or iNSCs were collected from culture flasks using Accutase (Invitrogen, CA, USA). After centrifugation (271 g for 10 minutes), cells were resuspended in PBS at a concentration of 8×10^6 cells per ml. For each hydrogel, we first resuspended the cells in 5 µl of PBS, then added 5 µl Heparin maleimide conjugate solution (90 µg/µl in PBS) and 10 µl starPEG-MMP-peptide conjugate solution (Tsurkan et al., 2011) to a final volume of 20 µl and a cell density of 2×10^6 cells/ml. Next, the 20 µl droplet was placed on a Parafilm sheet for approximately two minutes until it began to gelate. The gels were placed in 24-well plates, and each well contained 1 ml of astrocyte culture medium (AM) (Supplementary Movie 8). The gels were cultured and incubated in 5% CO₂/95% air at 37 °C until the desired time points (1 week, 2 weeks, and 3 weeks).

Generation of Matrigel cultures

For the generation of Matrigel cultures, we used BD Matrigel (catalog number: 356234). Prior to any cell culture work and use of the Matrigel, pipette tips and Eppendorf tubes has been frozen at -20 °C according to manufacturer's instruction following the "thick gel method". The BD Matrigel has been thawed overnight on ice at 4 °C. pNSCs were collected from culture flasks using Accutase (Invitrogen). After centrifugation (271 g for 10 minutes), pNSCs were re-suspended in BD Matrigel in concentration 2×10^6 cells per ml. Droplets of the cell/Matrigel mix were placed in the bottom of culture and then waiting to solidify at 37 °C. Then, cell medium (SRL, Catalog Number 1801) added and the gels cultured for 3 weeks (Supplementary Movie 8). Cell medium changed the day after the generation of gels and then every other day.

Synthesis of Amyloid peptides

The peptides were synthesized as previously described (Bhattacharai et al., 2017a; Bhattacharai et al., 2016; Bhattacharai et al., 2017b; Kizil et al., 2015; Wieduwild et al., 2013). For peptide synthesis, all of the required chemicals were purchased from IRIS Biotech GmbH (Marktredwitz, Germany). Acetonitrile (for UPLC/LCMS), dichloromethane (DCM), diethylether, dimethyl sulfoxide (DMSO), formic acid (FA), trifluoroacetic acid (TFA), and triisopropylsilane (TIS) were purchased from MERCK KGaA (Darmstadt, Germany). Acetic anhydride and N-methylmorpholine (NMM) were purchased from Sigma-Aldrich Co. LLC (St. Louis, MO, USA). Dithiothreitol (DTT) was obtained from Prolab VWR International, LCC (Radnor, PA, USA). Acetonitrile (for HPLC) was purchased from TH Geyer (Renningen, Germany). 5(6)-Carboxyfluorescein was purchased from Acros Organics (Fisher Scientific Company LLC). The TentaGel S RAM Fmoc rink amide resin was purchased from RappPolymere GmbH (Tübingen, Germany). The peptide synthesis columns and syringes with included filters were purchased from Intavis AG (Cologne, Germany). Water was obtained from a Milli-Q water purifier (Milli-Q Advantage A10, EMD Millipore Corporation, Billerica, MA, USA) with a LCPAK0001 Milli-Q filter. The polytetrafluoroethylene (PTFE) filter, polyvinylidene fluoride (PVDF) syringe filter, and filter holder were purchased from Sartorius Stedtim (Aubagne, France).

A β 42 peptides were prepared using standard 9-fluorenylmethoxycarbonyl (Fmoc) chemistry with 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) activation on an automated solid-phase peptide synthesizer (ResPep SL, Intavis)(Wieduwild et al., 2013; Zhang et al., 2002). Each amino acid was coupled twice at 5-fold excess followed by capping the non-reacted amino groups with acetic anhydride to achieve high quality synthesis. Upon completion of the peptide synthesis, 5(6)-carboxyfluorescein was coupled to the N-terminus using HBTU as the coupling reagent. The peptide was then cleaved from the resin with TFA/TIS/water/DTT (90(v/v):5(v/v):2.5(v/v):2.5(m/v)) for 2 hours. The product was precipitated and washed with ice-cold diethyl ether.

The peptide was dissolved in Milli-Q water, and peptide purification was performed via reverse-phase high-pressure liquid chromatography (HPLC) on a semi-preparative HPLC (Waters) equipped with a semi-preparative column (PolymerX RP-1, 250 x 10 mm, Phenomenex). The peptide was eluted from the column by applying a gradient of 5% to 100% solvent B over 30 min at 20 ml/min, in which solvent A is 0.1% TFA in water and solvent B is 0.1% TFA and 5% water in acetonitrile.

Purity was confirmed on an analytical reverse phase ultra-high pressure liquid chromatograph (UPLC Aquity with UV Detector) equipped with an analytical C18 column (Acquity UPLC BEH C18, bead size 1.7 μ m, 50 x 2.1 mm) using an isocratic gradient and electrospray ionization mass spectrometry (ESI-MS) (Acquity TQ Detector).

A β 42 treatment

The A β 42 treatment was performed 24 h post-thaw for a period of 48 h at 10 μ M final concentration for pNSCs and iNSCs in 3D cultures. Amyloid plaques remain in the culture attached to the cells throughout the cultures. The medium was removed and the cells were washed with PBS after 48 h A β 42 treatment. Then, the cells were collected using Accutase, counted and centrifuged at 271 g for 10 min. The cell pellet was resuspended in PBS to obtain at 8×10^6 cells/ml. This cell suspension was mixed with an equal volume of 6.00 mM heparin maleimide conjugate in

PBS to obtain a 3.00 mM heparin maleimide conjugate-cell suspension mix at 4.0×10^6 cells/ml. 10 μ l of this mix was combined with 10 μ l of 2.25 mM starPEG-MMP-peptide conjugate solution in PBS (Tsurkan et al., 2011), quickly triturated a few times, and the resulting 20 μ l volume was pipetted onto a Parafilm sheet forming a droplet. This droplet was allowed to gelate for about 2 min resulting in formation of a 20 μ l hydrogel containing 40,000 cells (2.0×10^6 cells/ml) at a final concentration of 1.50 mM heparin maleimide conjugate and 1.12 mM starPEG-MMP-peptide conjugate. The gels were then placed in 0.75 ml of AM culture medium per well in 24-well plates, and incubated in 5% CO₂/95% air at 37 °C until the desired time points (1 week, 2 weeks, and 3 weeks). The medium was changed 3 times a week throughout the incubation period.

2D pNSC cultures were treated with 2 μ M A β 42 in the culture medium for 48 hours between second and fourth days of culture. The medium is removed and cells are washed with cell culture medium twice. Amyloid plaques remain in the culture attached to the cells throughout the cultures. New medium is added and cultures were continued.

Transplantation

Primary human neural stem cells (pNSCs) were cultured for two days and treated with TOPRO-3 (Thermo Scientific) for 1 hour before transplantation. After a washing step, the cells were harvested with Accutase and re-suspended in astrocyte cell culture medium at 3×10^5 cells per ml. With a pipette tip, 2 μ l of the cell suspension was injected into the center of the hydrogel. After a week of culture, the hydrogels were fixed and processed for immunocytochemistry.

Immunocytochemistry

All of the hydrogels were fixed with ice-cold 4% paraformaldehyde and incubated for 1.5 hours at room temperature followed by washing in PBS overnight at 4 °C. For immunocytochemistry, the hydrogels were blocked and permeabilized in blocking solution for 4 hours at room temperature. For BrdU-treatment, the gels were incubated with 2 M HCl for 20 minutes at 37 °C followed by three washes in PBS (2 hours each). EdU staining was performed according to the manufacturer's protocol (Life

Technologies, C10638) using a 1-hour incubation step. The hydrogels were incubated with primary antibodies (Supplementary Table 1) in blocking solution overnight at 4 °C. The gels were washed for two subsequent days at 4 °C, with occasional changes of the PBS. After washing, the gels were incubated with the secondary antibodies (1:500 in blocking solution) at room temperature for 6 hours. After 3 washing steps of 2 hours each, DAPI staining was performed (1:3000 in PBS, 2 hours at room temperature). Immunostaining for SOX2 (Santa Cruz Biotechnology, 1:100), SATB2 (Abcam, 1:300), ASCL1 (Neuromics, 1:100), CTIP2 (Abcam, 1:100), EEA1 (Abcam, 1:500), neurofilament (NF-M+H+L) (Life Technologies, 1:500), TUBB3 (R&D Systems, 1:500), CASP3 (Santa Cruz Biotechnology, 1:500), MKI67 (Abcam, 1:1000), acetylated tubulin (Sigma, 1:500), SYN (Millipore, 1:500), GFAP (Novex, 1:500), DCX (Novex, 1:300), MAPT (Abcam, 1:500), A β 42 (Cell Signaling Technology, host: Rabbit, 1:500), Reelin (Abcam, 1:500), FOXO1 (Thermo Scientific, 1:500), FOXP2 (R&D, 1:200), CRYM (Thermo Fischer, 1:50), PSD95 (Thermo Fischer, 1:300), DBX1 (Abcam, 1:300), VGLUT1 (Thermo Fischer, 1:500), BrdU (AdB Serotec, 1:500) was performed. All of the secondary antibodies were conjugated to AlexaFluor dyes (Life Technologies).

Fluorescent imaging

For the hydrogels, fluorescent imaging was performed using a Leica SP5 inverted Laser Scanning Confocal microscope. The hydrogels were placed in glass bottom Petri dishes. Sixty microliters of PBS were added on top of the hydrogels to avoid desiccation. The Z-stacks were captured using a 25x water immersion lens. Every Z-stack had a z-distance of 500 μ m. Monolayers were imaged using an inverted Zeiss Apotome 2 microscope.

Histological analyses

For Gallyas silver staining, the 3D hydrogels were cryo-frozen and sequentially incubated in 5% periodic acid (5 minutes), an alkaline silver iodide solution (1 minute), acetic acid (3 minutes), 0.1% gold chloride (5 minutes), 1% sodium thiosulfate (5 minutes), and 2.5% aluminum sulfate (1 minutes), with intermittent washes with distilled water. For Thioflavin S

staining, samples were incubated in 1% Thioflavin S (8 minutes), absolute ethanol (3 minutes), and DAPI (10 minutes).

Transfection with GCaMP6f plasmids and calcium imaging

TurboFectin 8.0 reagent (OriGene, Cat# TF81001) was used to transfect adherent (2D cultures) and encapsulated (3D cultures) cells with 700 µg plasmid per reaction in 1 ml of cell growth medium. The pGP-CMV-GCaMP6f plasmid was a gift from Douglas Kim (Addgene plasmid # 40755)(Chen et al., 2013). The images were captured using a Leica SP5 inverted Laser Scanning Confocal microscope in resonant scanner mode with photon counting. Images were acquired every 100 milliseconds. Analysis of the calcium image spectrum was performed with the Leica LAS AF software by using region of interests (ROIs) and photon counting.

Patch clamp recordings

Single neurons were recorded in Artificial Cerebrospinal Fluid (ACSF) (119 mM NaCl, 2.5 mM KCl, 2 mM CaCl₂, 1.3 mM MgCl₂, 1 mM NaH₂PO₄, and 10 mM glucose, pH 7.3) and patched with nerve solution (125 mM K⁺-gluconate, 0.1 mM CaCl₂, 0.6 mM MgCl₂, 8 mM NaCl, 1 mM EGTA, 0.01 mM HEPES, and 4 mM Na-ATP, pH 7.23) The Whole-cell patch recordings were assessed using a HEKA set up and Pulse program. Membrane voltage resistance was held at -80 mV with the pipette resistance of 4-6 MOhms. For measurements of K⁺ and Na⁺ currents, test pulses were applied in 80 ms durations from -80 mV to 30 mV every 2 s. All experiments were done at 20-23 °C.

Electron microscopy

For electron microscopy, the hydrogel-embedded cells were fixed in modified Karnovsky's fixative (2% glutaraldehyde + 2% paraformaldehyde in 50 mM HEPES) at least overnight at 4 °C. The samples were washed 2x in 100 mM HEPES and 2x in water and postfixed in a 2% aqueous OsO₄ solution containing 1.5% potassium ferrocyanide and 2 mM CaCl₂ for 30 min on ice. Next, washes in water, 1% thiocarbohydrazide in water (20 minutes at room temperature), water, and a second osmium contrasting step in 2% OsO₄/water (30 minutes on ice). After several washes in water, the samples were *en bloc* contrasted

with 1% uranyl acetate/water for 2 hours on ice, washed again in water, dehydrated in a graded series of ethanol/water up to 100% ethanol, and infiltrated with Epon 812 (Epon/ethanol mixtures: 1:3, 1:1, 3:1 for 1.5 hours each, pure Epon overnight, and pure Epon for 5 hours). The samples were embedded in flat embedding molds and cured overnight at 65 °C. Ultrathin sections were prepared with a Leica UC6 ultramicrotome (Leica Microsystems, Vienna, Austria), collected on Formvar-coated slot grids and stained with lead citrate and uranyl acetate as previously described (Venable and Coggeshall, 1965).

For CLEM, cells that were embedded in the hydrogels with the fluorescein-labeled peptide were fixed with 4% paraformaldehyde in 100 mM phosphate buffer (PB). After several washes in water, the samples were dehydrated in 50% (15 minutes at 4 °C), 70%, 90%, and 100% acetone (45 minutes each at -25 °C) and incubated with LR Gold (London Resin Company, Reading, UK) solutions of 33% and 66% LR Gold/acetone, pure LR Gold (1 hour each at -25 °C), and LR Gold + 0.1% benzil (1 hour, overnight at -25 °C). Finally, the samples were transferred to LR Gold-containing 1% benzil and polymerized using the UV lamp of the Leica AFS2 freeze substitution unit (Leica Microsystems, Vienna, Austria) for 48 hours at -25 °C. Ultrathin sections were mounted on Formvar-coated EM grids, stained with DAPI, imaged with a wide field fluorescence microscope, washed, and contrasted with 1% uranyl acetate for EM as previously described (Fabig et al., 2012). Contrasted ultrathin sections were analyzed on a FEI Morgagni D268 (FEI, Eindhoven, The Netherlands) or a Jeol JEM1400 Plus at 80 kV acceleration voltage.

Atomic force microscopy

Atomic force microscopy (AFM) was performed to determine the mechanical properties of the gels. Briefly, AFM measurements were collected at 37 °C using a Nanowizard II AFM (JPK Instruments, Berlin, Germany). Tipless silicon nitride cantilevers with a nominal spring constant of 80 mN*m⁻¹ (PNP-TR-TL-Au; Nanoworld) were used. The cantilevers were modified with silica beads (Æ10 µm, Kisker Biotec GmbH), as previously described (Bray et al., 2015). Force-distance curves were acquired in closed loop, constant height mode using a 3 nN contact

force and a 5 $\mu\text{m/s}$ approach/retract velocity. Each data set was generated by probing a minimum of 70 different spots on each sample. The data processing software provided by the AFM manufacturer (JPK Instruments) was used to extract the Young's Modulus E from the approach force–distance curves.

RNA Isolation

RNA isolation from 2D cell culture was performed by TriZol (Invitrogen). Total RNA isolation from 3D gels was performed by Norgen Total RNA isolation Kit (Cat#17200). 5 gels were lysed in 1ml RL buffer with 10 μL β -mercaptoethanol, and after centrifugation at 12,000g for 7 min at room temperature, the supernatant was collected in a new Eppendorf and mixed with absolute Ethanol. The remaining steps performed as previously described (Bhattarai et al., 2016).

Next generation sequencing of whole transcriptome

cDNA libraries were prepared by following the protocol for NEBNext® Ultra I Directional RNA Library Prep Kit. This involves the following steps: mRNA isolation via poly(A)+ selection and fragmentation, first strand and second strand cDNA synthesis, purification using the Agencourt® AMPure® Kit and end repair/dA-tailing of cDNA. Adapters were ligated to the dA-tailed cDNA, followed by an size selection using AMPure XP Beads. Indexing of the library constructs was done with illumina® index primer during the following PCR amplification using NEBNext® Q5 2X PCR Master Mix. Lastly, libraries were purified using the Agencourt® AMPure® Kit. Libraries were pooled and sequenced on an illumina® NextSeq 500 system, resulting in ca. 27 – 38 million 75 bp single-end reads. All protocols are performed according to the manufacturers' instructions.

Data Analysis

The reads in fastq files were aligned to the human genome (hg19/GRCh38) with gsnap (version 2016-09-23) (Wu et al., 2016), and featureCounts (v1.5.3) (Liao et al., 2013, 2014) was used to assign reads to each gene using Ensembl version 90 [Homo_sapiens.GRCh38.90.gtf]. DESeq2 (1.18.0) (Love et al., 2014) was used to normalize the reads, calculate fold changes and p-values. 2-fold change and padj value of less

than 0.1 were used to identify differentially expressed genes. For KEGG pathway analysis GStats (2.44.0) (Falcon and Gentleman, 2007), GSeq (1.30.0) (Young et al., 2010), and clusterProfiler (3.6.0) (Yu et al., 2012) were used. topGO (2.30.0) (Alexa et al., 2006) was used for GO analysis and pathview (1.18.0) (Luo and Brouwer, 2013) was used for drawing KEGG pathway. All data analysis pipeline scripts were written in R in our lab, and are available upon request.

Sequencing datasets

All deep sequencing experiments (pNSC 2D, pNSC 3D, iNSC 3D control and Amyloid-beta42-treated) can be found under the GEO accession number GSE78117.

Image analysis and statistics

The 3D reconstructions of hydrogel images and videos were generated using Arivis 4D software. Images from monolayers were processed using Zeiss ZEN software. The statistical analyses were performed using GraphPad Prism and two-tailed Student's t-tests. The levels of significance were *: $p \leq 0.05$, **: $p \leq 0.01$, and ***: $p \leq 0.001$. In all graphs, means $\pm$ standard deviations are shown.

The effect size was calculated using G-Power, and the sample size was estimated with n-Query. The data conforms to normal distribution as determined by Pearson's chi-squared test. The variations between the samples are similar as determined by variance estimation using Microsoft Excel software. For 3D gels, 9 gels were used for quantifications (3 technical replicates in every experiment, and 3 experiments as biological replicates). All experiments were replicated many times in the laboratory and results were confirmed independently (80-120 gels were qualitatively analyzed to check the consistency of the results for every individual experiment).

Generation of skeletonized networks and quantification

To examine the axons of neural cells, the length and branching were obtained by thinning binary images to a skeleton, which was performed in all three dimensions. In detail, the raw images were processed with a Gaussian filter and then with the tubeness filter to enhance linear

1198 structures. Then, an automatic threshold was applied, followed by several
1199 morphological operations to facilitate the skeletonization. Fiji software
1200 (www.fiji.sc) was used for image processing. Skeletons were quantified
1201 using KNIME freeware (Supplementary Figure 4).

Supplementary Information

Supplementary Figure 1-5

Supplementary Movies 1-8

Supplementary Dataset 1-12

Supplementary figures

Supplementary Figure 1

Formation of 3D topology and arborizations in starPEG-Heparin gels

(A) Culture of primary human NSCs in 2D.

(B,C) Culture of primary human NSCs in 3D. Note the arborized morphology and cellular processes reminiscent of in vivo (white arrows).

Scale bars 25 μ m. Related to Figure 1.

Supplementary Figure 2

Comparison of Matrigel and starPEG-Heparin cultures

(A) Immunostaining for GFAP and SOX2 in Matrigel cultures.

(B) Immunostaining GFAP and Acetylated tubulin in Matrigel cultures Z-view.

(C) Immunostaining GFAP and Acetylated tubulin in Matrigel cultures X-view.

(D) Immunostaining for GFAP and SOX2 in starPEG-Heparin cultures.

(E) Immunostaining GFAP and Acetylated tubulin in starPEG-Heparin cultures Z-view.

(F) Immunostaining GFAP and Acetylated tubulin in starPEG-Heparin cultures X-view.

Scale bars: 20 μ m. All gels are 3 weeks of culture.

Supplementary Figure 3

Formation of neurofibrillary tangles and senile plaques in 3D cultures of primary human NSCs.

Image shows Gallyas silver impregnation staining for neurofibrillary tangles in 3D cultures. Inset shows the senile plaques (white arrow).

Supplementary Figure 4

Amyloid aggregation dynamics

(A) Immunostaining for acetylated tubulin and early endosomes (EEA1) in control cultures.

(A', A'') Individual fluorescence channels for EEA1 and acetylated tubulin in control cultures.

(B) Immunostaining for acetylated tubulin and early endosomes (EEA1) in Aβ42-treated cultures.

(B', B'') Individual fluorescence channels for EEA1 and acetylated tubulin in Aβ42-treated PHCCs.

(C, C') Co-localization of EEA1 and Aβ42.

(D) Quantification of the average number of early endosomes per cell and the average size of early endosomes per cell in control and Aβ42-treated cultures.

(E) Immunostaining of Aβ42-treated cells for acetylated tubulin (red), Aβ42 (green) and caspase-3 (white).

(E') Individual fluorescence channel for acetylated tubulin.

(E'') Fluorescence channels for Aβ42 and Caspase-3.

1261 (F) Quantification of Caspase-3-positive cells in control and A β 42-treated
 1262 cultures.
 1263 (G) Bright field image of control cultures.
 1264 (G') Bright field image of A β 42-treated cultures.
 1265 (G'') Bright field image of A β 38-treated cultures.
 1266 (H) Confocal image of acetylated tubulin immunostaining on A β 38-treated
 1267 cultures.
 1268 (H') Close-up of a region from H showing cells treated with A β 38.
 1269 (I) Quantification of the relative number of untreated, A β 42-treated and
 1270 A β 38-treated cells immunoreactive for GFAP, acetylated tubulin or BrdU.
 1271 (J) Confocal image over the z-axis and orthogonal section over the y-axis
 1272 of A β 42-treated human-derived fibroblasts. F-actin was stained with
 1273 phalloidin, and DNA was stained with DAPI.
 1274 (K) Confocal image over the z-axis and orthogonal section over the y-axis
 1275 of A β 38-treated human-derived fibroblasts. F-actin was stained with
 1276 phalloidin, and DNA was stained with DAPI.
 1277 (L) Quantification of the relative total number of fibroblasts in control,
 1278 A β 42-treated and A β 38-treated samples.
 1279 Scale bars 20 μ m. Related to Figure 4.
 1280

```

1 title = getTitle();
2 dir = getDirectory("image");
3 getVoxelSize(width, height, depth, unit);
4 run("Duplicate...", "title=Duplicate duplicate channels=2");
5 selectWindow(title);
6 close();
7 selectWindow("Duplicate");
8 run("Gaussian Blur...", "sigma=0.85 stack");
9 run("Tubeness", "sigma=3");
10 setAutoThreshold("Triangle dark stack");
11 setOption("BlackBackground", false);
12 run("Convert to Mask", "method=Triangle background=Dark black");
13 run("Invert LUT");
14 run("Options...", "iterations=6 count=3 pad do=Close");
15 run("Erode (3D)", "iso=255");
16 run("Dilate (3D)", "iso=255");
17 run("Dilate (3D)", "iso=255");
18 run("Erode (3D)", "iso=255");
19 run("Skeletonize (2D/3D)");
20 setVoxelSize(width, height, depth, unit);
21 run("Analyze Skeleton (2D/3D)", "prune=none calculate");
22 newtitle = substring(title,0,lastIndexOf(title, ".lif"));
23 saveAs("results", dir+newtitle+".xls");
24 selectWindow("Tagged skeleton");
25 saveAs("Tiff", dir+newtitle+"_skel.tif");
26 //run("Close All");

```

Supplementary Figure 5

Script for KNIME software

Supplementary Movies

Supplementary Movie 1

Neuronal network in pNSC cultures

3D representation of a 0.18 mm³ portion of a control hydrogel containing networks of neurons as assessed by acetylated tubulin staining (cyan). Related to Figure 1.

Supplementary Movie 2

Neuronal network in iNSC cultures

3D reconstruction of iPSC-derived NSC cultures in PEG-Heparin gels. Cyan: acetylated tubulin, white: DAPI, red: GFAP. Related to Figure 1.

Supplementary Movie 3

Calcium activity

Encapsulated primary human cortical astrocytes in PEG-HEP gels transfected with the GCaMP6f Calcium sensor exhibit green fluorescence in response to calcium influx after treatment with glutamate. Related to Figure 1.

Supplementary Movie 4

Dystrophic axons

A TUBB3-positive neuron (cyan) with retracted processes, dystrophic axonal ends, and A β 42 aggregation (green) is shown in a 3D reconstruction. DAPI is shown in white. Related to Figure 2.

Supplementary Movie 5

3D reconstruction of a neuron with hyperphosphorylated Tau

A 3D reconstruction of primary human cortical astrocyte cultures after 3 weeks in A β 42-treated gels and staining for A β 42 (green), DAPI (white)

and phosphorylated Tau (AT180, red). Note that neurons exhibit staining for hyperphosphorylated Tau. Related to Figure 2.

Supplementary Movie 6

Transplantation in control gels

3D reconstruction of a labeled neuron (red nuclei, transplanted) that established connections to the existing neurons in the control gels. Cyan: acetylated tubulin, white: DAPI, red: TOPRO3. Related to Figure 2.

Supplementary Movie 7

Transplantation in Amyloid-containing gel

3D reconstruction of a labeled neuron (red nuclei, transplanted) that was unable to make connections with the existing, already dystrophic, neurons in amyloid-containing gels. Green: A β 42, cyan: acetylated tubulin, white: DAPI, red: TOPRO3. Related to Figure 2.

Supplementary Movie 8

Comparison of gel preparation

Comparison of preparation of starPEG-Heparin and Matrigel cultures of primary human astrocytes. Related to Figure 2.

Supplementary Datasets

Supplementary Dataset 1

Differential expression analyses between pNSC 3D cultures and pNSC 2D cultures.

Supplementary Dataset 2

Elim Fischer cellular component analyses of differentially expressed genes between pNSC 3D cultures and pNSC 2D cultures.

Supplementary Dataset 3

KEGG pathway analysis of differentially expressed genes between pNSC 3D cultures and pNSC 2D cultures.

Supplementary Dataset 4

Differential expression analyses between iNSC 3D cultures and pNSC 3D cultures.

Supplementary Dataset 5

Elim Fischer cellular component analyses of differentially expressed genes between iNSC 3D cultures and pNSC 3D cultures.

Supplementary Dataset 6

KEGG pathway analysis of differentially expressed genes between iNSC 3D cultures and pNSC 3D cultures.

Supplementary Dataset 7

Differential expression analyses upon A β 42 in 3D iNSC cultures.

Supplementary Dataset 8

Differential expression analyses upon A β 42 in 3D pNSC cultures.

Supplementary Dataset 9

Elim Fischer cellular component analyses of differentially expressed genes in A β 42-treated 3D iNSC cultures.

Supplementary Dataset 10

KEGG pathway analysis of differentially expressed genes in A β 42-treated 3D iNSC cultures.

Supplementary Dataset 11

Elim Fischer cellular component analyses of differentially expressed genes in A β 42-treated 3D pNSC cultures.

Supplementary Dataset 12

KEGG pathway analysis of differentially expressed genes in A β 42-treated 3D pNSC cultures.