1 Measuring the net outcome of conditional mutualism: a case study with oaks and scatterhoarding
2 rodents

3

4

5 Michał Bogdziewicz^{1,2}, Elizabeth E. Crone³ & Rafał Zwolak^{1,4}

6 ¹ Department of Systematic Zoology, Faculty of Biology, Adam Mickiewicz University, Poznań,
7 Poland

8 ² CREAM, Cerdanyola del Valles, Catalonia, 08193 Spain

9 ³ Department of Biology, Tufts University, 163 Packard Ave, Medford, MA 02155, USA

10 ⁴ Department of Environmental Science and Policy, University of California at Davis, One Shields
11 Avenue, Davis, CA 95616

12

13 *corresponding author: michalbogdziewicz@gmail.com

14

15 **Authors contributions**

16 MB, RZ, and ECC conceived and designed the study, MB collected the data, MB, RZ, and ECC
17 analyzed the data, MB and RZ wrote the manuscript, all authors contributed substantially to the
18 revising of the paper.

19

20 **Data for the paper will be archived at Dryad repository.**

21 **Abstract**

22 Numerous interactions between plants and animals vary in their outcome between antagonism and
 23 mutualism, but it has proven to be difficult to quantify their final outcome. Interactions between plants
 24 and scatterhoarding animals provide a prime example of this phenomenon. Scatterhoarders consume
 25 large quantities of seeds (potentially reducing plant establishment), yet also disperse seeds and bury
 26 them in shallow caches (potentially improving recruitment). However, it has been rarely determined
 27 which role prevails for particular plant species. We demonstrate how plant-scatterhoarder interactions
 28 can be placed at the antagonism-mutualism continuum, with interactions between rodents and two
 29 oaks species (sessile oak *Quercus petraea*, and red oak *Q. rubra*) as an empirical example. Our
 30 approach consists of quantifying the net outcome of the interaction through assembling different vital
 31 rates (e.g. probability of seedling recruitment with and without rodents; near and far from conspecific
 32 trees; with and without seed pilferage) piecewise with a simple mathematical model. Our results
 33 indicate that during the period of the study, interactions between scatterhoarding rodents and both
 34 focal oaks were antagonistic. Even though acorn burial increased the likelihood of seedling
 35 establishment, this effect was not strong enough to compensate for the costs of seed predation.
 36 Furthermore, we found no evidence that the short-distance transportation that is usually provided by
 37 small mammals benefited early oak recruitment. Our study demonstrates how readily accessible field
 38 data can be used to gauge the outcomes in conditional mutualisms.

39

40 **Key words:** antagonism, biotic invasion, conditional mutualism, scatterhoarding, seed dispersal

41

42

43 Introduction

44 Numerous interactions between plants and animals vary in their outcome between antagonism and
45 mutualism (Bronstein, 1994; Palmer *et al.*, 2010). Interactions between plants and scatterhoarding
46 animals, such as rodents or corvids, are a prime example of this phenomenon because scatterhoarders
47 play a dual role in plant regeneration. On the one hand, they consume large quantities of seeds and
48 reduce plant establishment (Howe & Brown, 2001; Zwolak *et al.*, 2010; Larios *et al.*, 2017). On the
49 other hand, however, they disperse seeds and bury them in shallow caches, which for some plant
50 species provide the only means of successful recruitment (Vander Wall, 1992; Asquith *et al.*, 1999;
51 Muñoz & Bonal, 2011; Pesendorfer *et al.*, 2016). Determining which role prevails for particular plant
52 species can help guide nature conservation, forest management, and control of invasive species
53 because management strategies will depend on the interaction outcome. Yet, these outcomes have
54 been rarely quantified.

55 Whether scatterhoarding granivores are beneficial or detrimental for plant population depends
56 on whether recruitment with granivores is greater or less than recruitment without seed caching
57 (Jansen & Forget, 2001; Theimer, 2005; Schupp *et al.*, 2010; Zwolak & Crone, 2012). Quantifying
58 this has proven challenging, but one approach is to build up the net outcome from separately-measured
59 components (Zwolak & Crone, 2012). This kind of approach is similar to predicting population
60 dynamics from separately-measured vital rates (e.g., Morris & Doak 2002), but has been used less
61 often to evaluate species interactions. Intuitively, when the benefits from seed caching are high, plants
62 can bear higher costs in the form of seed consumption. Following on this notion, the interaction is
63 mutualistic when the probability of caching and not retrieving cached seeds exceeds the ratio of
64 seedling emergence from surface to seedling emergence from caches.

65 To briefly review our past use of this approach (Zwolak & Crone, 2012), we started from the
66 premise that granivores are beneficial when plant recruitment in the presence of granivores is greater
67 than plant recruitment in the absence of granivores. This inequality is written in mathematical terms as
68 follows:

$$69 \quad e_s < p_H p_C e_C + (1 - p_H) e_s \quad (0)$$

Where e_S is seedling emergence from surface, p_H is the proportion of seeds harvested by granivores, p_C is the probability that seeds will be cached and left uneaten, and e_C is the seedling emergence from caches. In studies of plant-granivore interactions, it is much easier and more common to measure the emergence rates (e_S and e_C) than the caching rates (p_H and p_C ; see Zwolak and Crone 2012). Therefore, in order to compare the role of granivores across studies, Zwolak and Crone (2012) rearranged the equation to calculate the minimum value of p_C that would be necessary for granivores to increase plant recruitment:

$$p_C < \frac{e_S}{e_C} \quad (0)$$

Thus, granivores help plant recruitment when the proportion of cached and uneaten seeds exceeds a threshold value (hereafter p_C , after Zwolak and Crone 2012) determined by the seedling emergence ratio, i.e. the benefits of seed burial.

Scatterhoarders not only bury the seeds in the topsoil, but also move them away from the parent plant. This also modifies recruitment probability, with effects that are usually believed to be positive, due to colonization of ephemeral microsites or escape from distance- and density dependent mortality (Jansen *et al.*, 2008; Comita *et al.*, 2010; Johnson *et al.*, 2012; Fricke *et al.*, 2014). However, the effects can also be negative, e.g. when habitat quality is autocorrelated, it often declines with distance from maternal plants (John *et al.*, 2007; Condit *et al.*, 2013). Furthermore, the distance to the seed source may alter rodent foraging activity and seed pilferage rates through changes in local seed availability (Stapanian & Smith, 1984; Gálvez *et al.*, 2009). Nonetheless, even though factors shaping dispersal distance by scatterhoarders, especially by rodents, are intensively studied (Jansen *et al.*, 2004; Xiao *et al.*, 2005; Moore *et al.*, 2007; Sunyer *et al.*, 2014; Lichti *et al.*, 2017), the actual influence of dispersal distance on recruitment probability is seldom quantified. The intertwined escape-related (transportation distance) and condition-related (burial) benefits constrain our ability to understand mechanisms that drive the ecological interactions between plants and scatterhoarders.

Here, we use empirical data to illustrate an approach for separating burial- and distance-dependent benefits of rodent seed dispersal. We used two oak species as model system: sessile oak (*Quercus petraea*), and northern red oak (*Q. rubra*). The sessile oak is the dominant native oak in

Central European forests. The northern red oak was introduced to European forests from North America in the 17th century as an ornamental species (Woziwoda *et al.*, 2014b). Currently, it is one of the most frequent foreign deciduous species in the region (Woziwoda *et al.*, 2014b), and its occurrence proved to be troublesome as it suppresses abundance and richness of numerous native species (Chmura, 2013; Woziwoda *et al.*, 2014a). For both oaks, the primary means of reproduction is thought to be abandonment of seed caches made by scatter-hoarding rodents and birds (den Ouden *et al.*, 2005; Steele, 2008; Myczko *et al.*, 2014). Nonetheless, whether caching benefits exceeds the costs imposed by seed predation has never been experimentally evaluated.

Drawing on past studies, we evaluated the following hypotheses (Table 1): (1) Benefits of seed caching should be larger than transportation benefits. Seed burial plays a critical role in regeneration of many plant species because it protects seeds from desiccation and strict seed predators (Haas & Heske, 2005; Vander Wall, 2010; Zwolak *et al.*, 2016). On the other hand, rodents move seeds over rather short distances (within population: den Ouden *et al.* 2005), which is unlikely to result in strong distance-dependent effects. (2) The distance-related benefits should be smaller in red oak than in sessile oak. This result might occur because invasive species often escape from the inhibitory effects of soil biota in their alien range (Agrawal *et al.*, 2005; Maron *et al.*, 2014). (3) Seed removal by rodents should increase with distance from adult plants. This result might occur because dense seed shadows under the tree may create local satiation effects (Xiao *et al.*, 2015). (4) Pilferage of cached acorns should be more frequent in red than in sessile oaks. This result might occur because heavier acorns are more likely to be pilfered (Perea *et al.*, 2016) and red oak acorns are over twice heavier than sessile oak acorns (Bogdziewicz *et al.*, 2017a). (5) Benefits of seed dispersal and burial will be large enough to compensate predation, leading to mutualistic relationship in which small mammals will aid recruitment of focal oaks. This prediction is based on results of a meta-analysis of previous studies, which suggested that outcomes of plant-rodent interactions tend to be weakly mutualistic (Zwolak and Crone 2012).

We used the estimates of seedling establishment probability obtained in field experiments to place the focal interactions at the antagonism-mutualism continuum, using the modeling framework developed by Zwolak and Crone (2012). However, the original model did not include the potential

changes in caching benefits driven by seed pilferage (Steele *et al.*, 2014; Sunyer *et al.*, 2015; Zwolak *et al.*, 2016). Therefore, as part of this paper, we extend the logic of the p_c calculation to include pilferage by naïve foragers.

Materials and Methods

We established four study sites in sessile – red oak mixed forests in Drawieńska Forest, western Poland (spaced 1 – 15 km from each other). This area is located in the temperate climate zone, with average annual precipitation of 592 mm and means monthly temperature ranging from 17°C in July to -2°C in January. These mixed forests comprise almost exclusively of the three oak species (*Q. petraea*, *Q. robur*, and *Q. rubra*), with single trees of common hornbeam (*Carpinus betulus*) and European beech (*Fagus sylvatica*). The understory is poorly developed, with some patches of raspberry (*Rubus* sp.) and common nettle (*Urtica dioica*), and seedlings of oaks and beech. Oak acorns are relatively large (average weight; *Q. petraea* 1.26 g., *Q. rubra*: 2.85 g.), and readily dispersed and eaten by small mammals (Bogdziewicz *et al.* 2017b, Bogdziewicz, unpublished). As revealed by camera traps, small mammals at our study sites were dominated by *Apodemus* sp., most likely *A. flavicollis*, a seed specialist (Selva *et al.*, 2012, Gasperini *et al.*, 2017).

To quantify the effects of acorn burial on seedling emergence we conducted seed addition experiments. We randomly chose 12 *Q. petraea* trees (3 per site) and 12 *Q. rubra* trees (3 per site). We added acorns of the focal species in 20 x 20 x 20 cm wire mesh cages (5 acorns per cage). Cages were buried 10 cm into the ground in sets of four. In half of the cages we buried acorns 1-2 cm into the ground and in the other half we placed acorns at the top of the litter layer and covered them with leaves to mimic autumn leaf fall. This treatment was crossed with rodent exclusion: in half of the cages we cut 8 x 8 cm holes to allow rodent access, and the other half remained closed to exclude rodent foraging. A comparison of seedling recruitment from acorns that were buried vs. placed on surface allowed us to estimate burial-dependent benefits of rodent seed dispersal. Rodent exclusion allowed us to estimate seed pilferage by comparing recruitment of buried acorns in open vs. closed cages.

To address the distance-related effects of rodent seed dispersal, the above-described cage sets were placed along transects. Under each tree, we established a transect along one cardinal direction, aiming to maximize the distance of the transect to other conspecifics. This was done assuming that rodents tend to carry and cache seeds towards areas of lower conspecific seed density (Stapanian & Smith, 1984; Hirsch *et al.*, 2012; Steele *et al.*, 2014; Yang *et al.*, 2016). Thus, our estimates of distance-dependent effects may be overly positive, if such directed dispersal does not occur in our system. We placed five sets of cages at each transect. We used tree crown as a reference point and buried one set of cages directly underneath the crown border, another set 5 m towards the tree trunk (underneath the crown), and the remaining 3 sets every 5 m in the opposite direction (i.e. away from the tree trunk). We used 25 m as the maximum evaluated distance because acorn-tracking experiments report that vast majority of acorns transported by rodents are cached within that radius (den Ouden *et al.*, 2005; Xiao *et al.*, 2005; Muñoz & Bonal, 2011; Bogdziewicz *et al.*, 2017b). We set up experimental cages in October 2016 and quantified seedling establishment in August 2017. The overall sample size equaled 2400 acorns (480 seedling cages).

Statistical analysis

To test how acorn burial, distance from the tree, and rodent foraging affect the seedling establishment, we built a separate generalized linear mixed model (GLMM) for each oak species. We used nested random effects of cage set, tree, and study site, logit link, and binomial family error distribution, and implemented the models via lme4 package in R (Bates et al. 2015). In each model, we used proportion of established seedlings as the response variable, and burial (surface vs. sowed), rodent access (excluded vs. allowed), and distance from the tree as fixed effects. We also included all possible 2-way interaction terms between fixed effects, and the 3-way interaction (which was removed when non-significant). We calculated marginal (i.e. the proportion of variance explained by fixed effects) and conditional (i.e. the proportion of variance explained by fixed and random effects) R^2 for GLMMs using the MuMIn package (Nakagawa & Schielzeth, 2013; Bartoń, 2016).

Calculating the p_c threshold and the effects of seed pilferage

We evaluated how the interactions between rodents and oaks are placed along the antagonism – mutualism continuum (Zwolak & Crone, 2012). The p_c threshold was calculated as a ratio of emergence from seeds sown on surface vs. emergence from buried seeds, with rodents excluded. Seed pilferage was gauged as the ratio of seedling recruitment from buried seeds in open vs. closed cages. Implicitly, the original definition of the proportion of seeds cached and uneaten (p_c) combined three processes (Zwolak & Crone, 2012): the probability that a seed is cached, the probability it is eaten by the cache owner, and the probability it is pilfered:

$$p_c = p_{cached} \times (1 - p_{eaten\ by\ cache\ owner}) \times (1 - p_{pilfered}) \quad (1)$$

where p_{cached} is the probability a seed is cached, p_{eaten} by cache owner is the probability of retrieval by individuals responsible for seed burial and $p_{pilfered}$ is the probability of retrieval by pilferers. The p_c threshold is the minimum value of when the benefits for plants balance the costs of seed consumption. Thus, if the threshold is determined by the proportion of seedling emergence from surface, e_s (estimated with data on seedling emergence from seeds sown on surface in closed cages), to emergence from caches, e_c (i.e. by benefits of burial, estimated with data on seedling emergence from seeds buried in closed cages), i.e.:

$$\tilde{p}_c = \frac{e_s}{e_c}$$

then the equation can be expanded to show the effects of pilferage:

$$p_{cached} \times (1 - p_{eaten\ by\ cache\ owner}) \times (1 - p_{pilfered}) = \frac{e_s}{e_c}$$

and rearranged to include only the unknown proportion of seeds cached and uneaten by the cache owner:

$$\tilde{p}_{CO} = p_{cached} \times (1 - p_{eaten\ by\ cache\ owner}) = \frac{e_s}{e_c (1 - p_{pilfered})} \quad (2)$$

In the above equation, p_{CO} is the minimum beneficial proportion of seeds cached and uneaten by *cache owner*, and all other parameters are as defined above. In other words, p_{co} threshold defines the burial benefits while taking into account cache pilferage by naïve forages.

Confidence intervals for these parameters (e_s , e_c , and p_c) were obtained with parametric bootstrapping, i.e. sampling from the distributions defined by the mean and standard error of each coefficient to obtain a joint distribution for the derived variables. We repeated the calculations of p_c and p_{co} for both near the conspecific probability of germination (i.e. germination rate estimated at the distance 0 m, and far i.e. germination rate estimated at distance 25m). Probabilities of germination were derived from the above-described GLMMs.

The empirical p_{co} value (the ultimate probability that the acorn will be cached and not retrieved, accounting for retrieval by the whole granivore community) for both oak species was derived from a parallel study investigating rodent seed dispersal of the focal oaks, i.e. 17% for red oaks and 2% for sessile oaks (Bogdziewicz et al. unpublished).

Results

In accordance with hypothesis (1), benefits of acorn burial were higher than benefits of acorn transportation away from adult trees. Acorn burial enhanced seedling establishment in both species, mainly through reducing seed predation. In *Q. petraea*, when rodents had access, acorn burial increased establishment probability 2-fold (open cages, surface vs. buried acorns: 18% vs. 39%). This effect was considerably weaker when rodents were excluded (closed cages, 57% for acorns on the surface vs. 65% for buried acorns; rodent exclusion $\times$ burial interaction in Table 2a, Fig. 1). Similarly, burial increased establishment probability almost 3-fold in *Q. rubra* (open cages: 25% vs. 71%, Fig. 1). This effect was weaker when rodents were excluded (50% vs. 73%; rodent exclusion $\times$ burial interaction in Table 2b).

We did not detect any distance-related benefits of seed dispersal and contrary to hypothesis (2), the oak species did not differ in this regard. In fact, seedling establishment probability decreased with the distance from the focal tree (Table 2, Fig. 2 and 3). This phenomenon was caused by an increase in acorn removal, which supports hypothesis (3). In *Q. petraea* increased removal was apparent only when acorns were sown on surface (Table 2a, Fig. 2). Pilferage of cached acorns did not differ with distance from the tree in this species. In *Q. rubra*, this effect occurred both for acorns that were buried and those that were left on surface.

Acorn burial was more beneficial for *Q. rubra* than for *Q. petraea* (see above, and Fig. 1). In contrast to hypothesis (4), pilferage rates were higher in *Q. petraea* than in *Q. rubra*. In *Q. petraea* seedling establishment from buried acorns was 1.5 times higher when rodents were excluded (39% in open vs. 65% in closed cages). In *Q. rubra*, burial provided almost complete protection from pilferage (open vs. closed cages: 71% vs. 73%). Note that the percentages estimates are the model intercepts, and decrease with distance from the tree in some treatments (see below).

Estimated p_{CO} values (the minimum beneficial proportion of seeds cached and uneaten by cache owner, i.e. those taking into account cache pilferage) equaled 1.21 (95% CI: 0.86-1.79) for *Q. petraea* (both near and far from the tree), and 0.69 (95% CI: 0.46-0.97) in *Q. rubra* near, and 1.24 (95% CI: 0.77-1.99) in *Q. rubra* far from the tree (Fig. 4). Note that the p_{CO} value for *Q. petraea* does not differ with distance because the pilferage rates were distance-independent (Fig. 2a). These values are either impossible to reach (when they exceed 1) or would require almost all cached acorns to be never retrieved to approach the mutualism parameter space of the interaction (Fig. 4). Thus, hypothesis (5) was not supported: interactions between scatterhoarders and oaks in our study system were clearly antagonistic.

A fraction of removed acorns could be recached rather than consumed. If we assume that pilfered seeds are as likely to be eaten or re-cached as seeds collected for the first time (Jansen *et al.*, 2004), then the consequences of burial depend on the number of rounds of recaching (Online Supplement 1). Nonetheless, for the parameters observed in our system, recaching of pilfered acorns would not affect the conclusion that scatterhoarders acted antagonistically in their interactions with oaks (Online Supplement 1).

Discussion

Our study is the first to separate and directly quantify two most important services provided to plants by their rodent partners: seed transportation away from parent plants and seed burial in topsoil. Our results suggest that widely-accepted benefits of transportation might in many ecological systems be smaller than expected. Moreover, they demonstrate that even relatively large improvements in seedling establishment after seed burial do not necessarily render the plant-scatterhoarder interaction

mutualistic. Finally, and most generally, our study illustrates a straightforward empirical approach that can be used to evaluate the ambiguous role of scatterhoarding granivores in plant regeneration.

This approach consists of quantifying the net outcome of an interaction through assembling different parameters piecewise with a simple mathematical model. This method, although used here to a specific plant-scatterhoarder dataset, is very general: in fact, it is analogous to building population models with separately-measured vital rates (e.g. Caswell 2001, Morris and Doak 2002). Similarly, just like other ecological models, it is a simplification, that can be made more realistic by adding additional information (see, e.g., Maron and Crone 2006, Ehrlén et al. 2016 for examples in a demographic context). Here we used data on probability of seedling recruitment with and without rodents; near and far from adult conspecifics; with and without seed pilferage. Possible future extensions of our seed-scatterhoarder model include e.g. effects of temporal variability or effects of directed dispersal of seeds into particularly favorable microsites (see below).

In addition, this case study provides a method to compare the importance of different types of benefits or costs of interspecific interactions. Animal dispersers often provide several different services that are not of equal value for their partner plants, but benefits provided by each are rarely separated. Yet, a few case studies that have done so demonstrated that it may change the way we think about particular interactions. For example, birds that disperse chili peppers (*Capsicum chacoense*) remove pathogens from the dispersed seeds (condition-related benefit) and transport seeds far from parent plants (escape-related benefit). However, only gut passage enhances seed survival (Fricke *et al.*, 2013). A similar situation was reported for Iberian pears (*Pyrus bourgaeana*), where pulp removal was more important than transportation distance for plant recruitment (Fedriani *et al.*, 2012). By untwining the role of caching (condition-related benefit) from transportation distance in oak-scatterhoarders interactions, we demonstrated that acorn burial was the main benefit of this interaction (hypothesis 1 in Table 1), at least during the period of the study. The lack of distance-dependent benefits was unsurprising because in locally common plant species, the benefits should disappear when the species become so common that their predator and pathogen communities become functionally uniform across the landscape (Janzen, 1971; Schupp, 1992; Fricke *et al.*, 2013; Garzon-Lopez *et al.*, 2015). This is likely the case in our system because sessile and red oaks dominate forest stands. Our results suggest

that the generally greater establishment of some species far from parent plants may be due distance-independent benefits of burial rather than distance-dependence *per se*. In other words, intertwined caching and transportation may create false positive effect of transportation, while it is only caching that helps recruitment (as in our system). This calls for increased attention on condition-dependent benefits of seed dispersal, which have been often overlooked as researchers focused on dispersal distance and final location of seeds (Fricke *et al.*, 2013).

The patterns of seedling establishment suggest that rodent foraging is a strong filter of oaks spatial recruitment in our system. When rodents were excluded, distance from the tree did not change the probability of seedling establishment in either species (contrary to hypothesis 2 in Table 1). However, when rodent foraging was allowed, seedling establishment rate was highest near the trees, indicating that locally dense seed shadows may allow higher proportion of undispersed seeds to survive and germinate (hypothesis 3 in Table 1). On the other hand, distance to adult trees influenced pilferage of buried acorns in a species-specific manner. In *Q. petraea*, the pilferage did not change with distance from the tree, while it decreased with distance in *Q. rubra*. It suggests that ambient seed density has a stronger effect on cache pilferage rates in *Q. rubra* than in *Q. petraea* (Gálvez *et al.*, 2009). However, we did not measure seed shadows, and thus this pattern might have resulted from larger crop size of *Q. rubra* at the time of our experiments.

Q. rubra acorns were generally better protected from seed predation once buried than acorns of *Q. petraea* (contrary to hypothesis 4 in Table 1). The pilferage rates of buried seeds are related to their release of odorant molecules (Vander Wall, 1998; Vander Wall & Joyner, 1998; Yi *et al.*, 2016). *Q. rubra* appears to rely more on rodent dispersal in their native range than does *Q. petraea* in Europe (den Ouden *et al.*, 2005; Steele, 2008). Thus, the tighter coevolution between small mammals and *Q. rubra* in the species native range (Steele, 2008), may led to reduced emission of volatile compounds in this species, reducing the overall pilferage rates. In consequence, lower pilferage rates in *Q. rubra* increased overall burial benefits, moving the interaction closer to the mutualism.

Nonetheless, in contrast to our fifth and final hypothesis (Table 1), scatterhoarding rodents reduced recruitment of focal oak species during the period of our study. As anticipated, acorn burial increased the likelihood of seedling establishment. However, the seedling establishment of unburied

acorns in the absence of small mammal foraging was high for both species. As a final result, burial benefits were too small to override the costs of seed predation (Zwolak & Crone, 2012). In fact, even if the probability of seedling establishment from rodent caches would elevate to 100%, the p_{CO} value for *Q. petraea* would equal 0.57 while for *Q. rubra* 0.50. This level of survival of cached acorns appears unlikely, as the reported values range from 1 to 20% (Vander Wall & Joyner, 1998; Vander Wall, 2002; Gomez *et al.*, 2008; Xiao *et al.*, 2013; Bogdziewicz *et al.*, 2017b; Wróbel & Zwolak, 2017). Therefore, while it is still best for an acorn to be cached and then forgotten, presence of small mammals did not help recruitment of oaks in our system.

We note, however, that the balance of benefits and costs in conditional mutualisms typically changes over time (Theimer 2005, Klinger and Rejmánek 2010, Zwolak *et al.* 2016) and our study was conducted during relatively short time frame (1 year). It is possible that fluctuating environmental conditions (e.g. years with droughts or severe winters) can increase the benefits of acorn caching and shift the oak-rodent relationship towards mutualism. This is particularly likely given that burial benefits in our study were remarkably low: e.g. Kollmann & Schill (1996), García *et al.* (2002), and Xia *et al.*, (2016) reported higher benefits in oaks. In this situation, the role of small mammals in oak recruitment would change over time. Moreover, both focal oaks species exhibit mast years (Sork *et al.*, 1993; Bogdziewicz *et al.*, 2017c), which in turn drives fluctuations in small mammal population abundance and may create satiation effects, both at the seed source and after caching (Kelly, 1994; Xiao *et al.*, 2013; Zwolak *et al.*, 2016). Yet, the potential increase in cache survival caused by the phenomenon is unlikely to counterbalance the costs of seed predation – unless it is accompanied by environmental changes that increase the benefits of seed caching. Thus, an interesting venue for future studies would be to quantify temporal variation in this interaction.

As a final caveat to this study, Janzen-Connell effects are stronger at the seedling than seed-to-seedling stage (Comita *et al.*, 2014). Therefore, the benefits of transportation may appear at later stages of plants life cycle. However, several studies that evaluated distance-dependent survival rates at seedling stage in temperate oaks did not found the effect (Reinhart *et al.*, 2012; Comita *et al.*, 2014). Furthermore, directed dispersal increases the likelihood of colonization of microhabitats that are favorable for germination and establishment (Steele *et al.*, 2013; Yi *et al.*, 2013). Although such

effects were never reported for focal oaks, our experimental design could underestimate these effects and thus, benefits of acorn transportation by scatterhoarding rodents.

To conclude, we presented simple means by which outcomes of conditional plant-scatterhoarder interactions can be classified. The strength of our approach lies in its versatility: it uses mathematics to combine different types of data and can be easily modified to incorporate new information when data on other parameters becomes accessible. Our empirical results demonstrated that certain common assumptions (e.g. that scatterhoarding by rodents invariably improves plant recruitment; that improved germination after seed burial is sufficient to make plant-scatterhoarder interactions mutualistic; that transportation away from maternal plants is highly beneficial) do not always hold and should be tested rather than taken for granted.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgements

We thank Nadleśnictwo Krzyż Wlkp. for the permission to conduct the study. Kinga Stępnia, Agnieszka Amborska-Bogdziewicz, and Jan Bogdziewicz helped during the field work, and Anna and Bogusław Bogdziewicz provided logistical assistance. Dawid Manyś provided invaluable assistance during field site selection. The study was supported by the (Polish) National Science Centre grant ‘Preludium’ no. 2015/17/N/NZ8/01565.

References

- Agrawal AA, Kotanen PM, Mitchell CE, Power AG, Godsoe W, Klironomos J (2005) Enemy release? An experiment with congeneric plant pairs and diverse above- and belowground enemies. *Ecology* 86: 2979-2989.
- Asquith NM, Terborgh J, Arnold AE, Riveros CM (1999) The fruits the agouti ate: *Hymenaea courbaril* seed fate when its disperser is absent. *Journal of Tropical Ecology* 15: 229-235.
- Bartoń K (2016) MuMIn: Multi-Model Inference. R package version 1.15.6. <https://CRAN.R-project.org/package=MuMIn>
- Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67: 1-48.
- Bogdziewicz M, Bonal R, Espelta JM, Kalembe E, Steele MA, Zwolak R (2017)a Invasive oaks escape pre-dispersal insect seed predation and trap enemies in their seeds. *Integrative Zoology* DOI: 10.1111/1749-4877.12285
- Bogdziewicz M, Crone EE, Steele MA, Zwolak R (2017)b Effects of nitrogen deposition on reproduction in a masting tree: benefits of higher seed production are trumped by negative biotic interactions. *Journal of Ecology* 105: 310-320.
- Bogdziewicz M, Szymkowiak J, Kasprzyk I, Grewling L, Borowski Z, Borycka K, Kantorowicz W, Myszkowska D, Piotrowicz K, Ziemianin M, et al. (2017)c Masting in wind-pollinated trees: the role of weather and pollination dynamics in driving seed production. *Ecology*. 98: 615–2625
- Bronstein JL (1994) Conditional outcomes in mutualistic interactions. *Trends in Ecology & Evolution* 9: 214-217.
- Caswell H (2001) Matrix population models. John Wiley & Sons, Ltd.
- Chmura D (2013) Impact of alien tree species *Quercus rubra* L. on understorey environment and flora: a study of the silesian upland (southern Poland). *Polish Journal of Ecology* 61: 431-441.
- Comita LS, Muller-Landau HC, Aguilar S, Hubbell SP (2010) Asymmetric density dependence shapes species abundances in a tropical tree community. *Science* 329: 330-332.

- Comita LS, Queenborough SA, Murphy SJ, Eck JL, Xu K, Krishnadas M, Beckman N, Zhu Y (2014) Testing predictions of the Janzen–Connell hypothesis: a meta-analysis of experimental evidence for distance-and density-dependent seed and seedling survival. *Journal of Ecology* 102: 845-856.
- Condit R, Engelbrecht BMJ, Pino D, Perez R, Turner BL (2013) Species distributions in response to individual soil nutrients and seasonal drought across a community of tropical trees. *Proceedings of the National Academy of Sciences of the United States of America* 110: 5064-5068.
- den Ouden J, Jansen PA, Smit R (2005) Jays, Mice and Oaks: Predation and Dispersal of *Quercus robur* and *Q. petraea* in North-western Europe. Seed Fate: Predation, dispersal, and seedling establishment: 223.
- Ehrlén J, Morris WF, von Euler T, Dahlgren JP (2016) Advancing environmentally explicit structured population models of plants. *Journal of Ecology* 104:292–305.
- Fedriani JM, Zywiec M, Delibes M (2012) Thieves or mutualists? Pulp feeders enhance endozoochore local recruitment. *Ecology* 93: 575-587.
- Fricke EC, Simon MJ, Reagan KM, Levey DJ, Riffell JA, Carlo TA, Tewksbury JJ (2013) When condition trumps location: seed consumption by fruit-eating birds removes pathogens and predator attractants. *Ecology Letters* 16: 1031-1036.
- Fricke EC, Tewksbury JJ, Rogers HS (2014) Multiple natural enemies cause distance-dependent mortality at the seed-to-seedling transition. *Ecology Letters* 17: 593-598.
- García D, Bañuelos M-J, Houle G (2002) Differential effects of acorn burial and litter cover on *Quercus rubra* recruitment at the limit of its range in eastern North America. *Canadian Journal of Botany* 80: 1115-1120.
- Garzon-Lopez CX, Ballesteros-Mejia L, Ordonez A, Bohlman SA, Olff H, Jansen PA (2015) Indirect interactions among tropical tree species through shared rodent seed predators: a novel mechanism of tree species coexistence. *Ecology Letters* 18: 752-760.

- Gasperini S, Bonacchi A, Bartolommei P, Manzo E, Cozzolino R (2017) Seasonal cravings: plant food preferences of syntopic small mammals. *Ethology Ecology & Evolution* 14:1-4.
- Gomez JM, Puerta-Pinero C, Schupp EW (2008) Effectiveness of rodents as local seed dispersers of Holm oaks. *Oecologia* 155: 529-537.
- Gálvez D, Kranstauber B, Kays RW, Jansen PA (2009) Scatter hoarding by the Central American agouti: a test of optimal cache spacing theory. *Animal Behaviour* 78: 1327-1333.
- Haas JP, Heske EJ (2005) Experimental study of the effects of mammalian acorn predators on red oak acorn survival and germination. *Journal of Mammalogy* 86: 1015-1021.
- Hirsch BT, Kays R, Pereira VE, Jansen PA (2012) Directed seed dispersal towards areas with low conspecific tree density by a scatter-hoarding rodent. *Ecology Letters* 15: 1423-1429.
- Howe H, Brown J (2001) The ghost of granivory past. *Ecology Letters* 4: 371-378.
- Jansen PA, Bongers F, Hemerik L (2004) Seed mass and mast seeding enhance dispersal by a neotropical scatter-hoarding rodent. *Ecological Monographs* 74: 569-589.
- Jansen PA, Bongers F, Van Der Meer PJ (2008) Is farther seed dispersal better? Spatial patterns of offspring mortality in three rainforest tree species with different dispersal abilities. *Ecography* 31: 43-52.
- Jansen PA, Forget P-M (2001) Scatterhoarding rodents and tree regeneration. *Nouragues: Springer*, 275-288.
- Janzen DH (1971) Seed predation by animals. *Annual review of ecology and systematics*: 465-492.
- John R, Dalling JW, Harms KE, Yavitt JB, Stallard RF, Mirabello M, Hubbell SP, Valencia R, Navarrete H, Vallejo M, et al. (2007) Soil nutrients influence spatial distributions of tropical tree species. *Proceedings of the National Academy of Sciences of the United States of America* 104: 864-869.
- Johnson DJ, Beaulieu WT, Bever JD, Clay K (2012) Conspecific negative density dependence and forest diversity. *Science* 336: 904-907.
- Kelly D (1994) The evolutionary ecology of mast seeding. *Trends in Ecology & Evolution* 9: 465-470.

- Kollmann J, Schill HP (1996) Spatial patterns of dispersal, seed predation and germination during colonization of abandoned grassland by *Quercus petraea* and *Corylus avellana*. *Vegetatio* 125: 193-205.
- Klinger R, Rejmánek M (2010) A strong conditional mutualism limits and enhances seed dispersal and germination of a tropical palm. *Oecologia* 162: 951-63.
- Larios L, Pearson DE, Maron JL (2017) Incorporating the effects of generalist seed predators into plant community theory. *Functional Ecology* 31: 1856-1867.
- Lichti NI, Steele MA, Swihart RK (2017) Seed fate and decision-making processes in scatter-hoarding rodents. *Biological Reviews* 92: 473:504
- Maron JL, Crone E (2006) Herbivory: effects on plant abundance, distribution and population growth. *Proceedings of the Royal Society of London B: Biological Sciences*. 273:2575-84.
- Maron JL, Klironomos J, Waller L, Callaway RM (2014) Invasive plants escape from suppressive soil biota at regional scales. *Journal of Ecology* 102: 19-27.
- Moore JE, McEuen AB, Swihart RK, Contreras TA, Steele MA (2007) Determinants of seed removal distance by scatter-hoarding rodents in deciduous forests. *Ecology* 88: 2529-2540.
- Morris WF, Doak DF (2002) Quantitative conservation biology. Sinauer, Sunderland, Massachusetts, USA.
- Muñoz A, Bonal R (2011) Linking seed dispersal to cache protection strategies. *Journal of Ecology* 99: 1016-1025.
- Myczko Ł, Dylewski Ł, Zduniak P, Sparks TH, Tryjanowski P (2014) Predation and dispersal of acorns by European Jay (*Garrulus glandarius*) differs between a native (Pedunculate Oak *Quercus robur*) and an introduced oak species (Northern Red Oak *Quercus rubra*) in Europe. *Forest Ecology and Management* 331: 35-39.
- Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods in Ecology and Evolution* 4: 133-142.
- Palmer TM, Doak DF, Stanton ML, Bronstein JL, Kiers ET, Young TP, Goheen JR, Pringle RM (2010) Synergy of multiple partners, including freeloaders, increases host fitness in a multispecies mutualism. *Proceedings of the National Academy of Sciences* 107: 17234-17239.

- Perea R, Dirzo R, San Miguel A, Gil L (2016) Post-dispersal seed recovery by animals: is it a plant- or an animal-driven process? *Oikos* 125: 1203-1210.
- Pesendorfer MB, Sillett TS, Koenig WD, Morrison SA (2016) Scatter-hoarding corvids as seed dispersers for oaks and pines: A review of a widely distributed mutualism and its utility to habitat restoration. *The Condor* 118: 215-237.
- Reinhart KO, Johnson D, Clay K (2012) Effects of trees on their recruits in the southern Appalachians, USA. *Forest Ecology and Management* 263: 268-274.
- Schupp EW (1992) The Janzen-Connell model for tropical tree diversity - population implications and the importance of spatial scale. *American Naturalist* 140: 526-530.
- Schupp EW, Jordano P, Gómez JM (2010) Seed dispersal effectiveness revisited: a conceptual review. *New Phytologist* 188: 333-353.
- Selva N, Hobson KA, Cortés-Avizanda A, Zalewski A, Donázar JA (2012) Mast pulses shape trophic interactions between fluctuating rodent populations in a primeval forest. *PloS One* 7: e51267.
- Sork VL, Bramble J, Sexton O (1993) Ecology of mast-fruiting in three species of North American deciduous oaks. *Ecology* 74: 528-541.
- Stapanian MA, Smith CC (1984) Density-dependent survival of scatterhoarded Nuts: an experimental approach. *Ecology* 1387-1396.
- Steele MA (2008) Evolutionary interactions between tree squirrels and trees: a review and synthesis. *Current Science* 95: 871-876.
- Steele MA, Contreras TA, Hadj-Chikh LZ, Agosta SJ, Smallwood PD, Tomlinson CN (2014) Do scatter hoarders trade off increased predation risks for lower rates of cache pilferage? *Behavioral Ecology* 25: 206-215.
- Sunyer P, Boixadera E, Munoz A, Bonal R, Espelta JM (2015) The Interplay among Acorn Abundance and Rodent Behavior Drives the Spatial Pattern of Seedling Recruitment in Mature Mediterranean Oak Forests. *Plos One* 10(6).
- Sunyer P, Espelta JM, Bonal R, Munoz A (2014) Seeding phenology influences wood mouse seed choices: the overlooked role of timing in the foraging decisions by seed-dispersing rodents. *Behavioral Ecology and Sociobiology* 68: 1205-1213.

- Theimer TC (2005) Rodent Scatterhoarders as Conditional Mutualists. Seed Fate: Predation, Dispersal, and Seedling Establishment: 283.
- Vander Wall SB (1992) The role of animals in dispersing a "wind-dispersed" pine. *Ecology*: 614-621.
- Vander Wall SB (1998) Foraging success of granivorous rodents: effects of variation in seed and soil water on olfaction. *Ecology* 79: 233-241.
- Vander Wall SB (2002) Masting in animal-dispersed pines facilitates seed dispersal. *Ecology* 83(12): 3508-3516.
- Vander Wall SB (2010) How plants manipulate the scatter-hoarding behaviour of seed-dispersing animals. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 365(1542): 989-997.
- Vander Wall SB, Joyner JW (1998) Recaching of Jeffrey pine (*Pinus jeffreyi*) seeds by yellow pine chipmunks (*Tamias amoenus*): potential effects on plant reproductive success. *Canadian Journal of Zoology* 76: 154-162.
- Woziwoda B, Kopec D, Witkowski J (2014)a. The negative impact of intentionally introduced *Quercus rubra* L. on a forest community. *Acta Societatis Botanicorum Poloniae* 83: 39-49.
- Woziwoda B, Potocki M, Sagan J, Michal Z, Tomusiak R, Wilczynski S. (2014)b. Commercial Forestry as a Vector of Alien Tree Species - the Case of *Quercus rubra* L. Introduction in Poland. *Baltic Forestry* 20(1): 131-141.
- Wróbel A, Zwolak R (2017) Deciphering the effects of disperser assemblages and seed mass on patterns of seed dispersal in a rodent community. *Integrative Zoology*. 10.1111/1749-4877.12265
- Xia K, Tan HY, Turkington R, Hu JJ, Zhou ZK (2016) Desiccation and post-dispersal infestation of acorns of *Quercus schottkyana*, a dominant evergreen oak in SW China. *Plant Ecology* 217: 1369-1378.
- Xiao Z, Zhang Z, Krebs CJ (2013) Long-term seed survival and dispersal dynamics in a rodent-dispersed tree: testing the predator satiation hypothesis and the predator dispersal hypothesis. *Journal of Ecology* 101: 1256-1264.

- Xiao Z, Zhang Z, Wang Y (2005) Effects of seed size on dispersal distance in five rodent-dispersed fagaceous species. *Acta Oecologica* 28: 221-229.
- Xiao ZS, Zhang ZB, Krebs CJ (2015) Seed size and number make contrasting predictions on seed survival and dispersal dynamics: A case study from oil tea *Camellia oleifera*. *Forest Ecology and Management* 343: 1-8.
- Yang YQ, Zhang MM, Yi XF (2016) Small rodents trading off forest gaps for scatter-hoarding differs between seed species. *Forest Ecology and Management* 379: 226-231.
- Yi XF, Liu GQ, Steele MA, Shen Z, Liu CQ (2013) Directed seed dispersal by a scatter-hoarding rodent: the effects of soil water content. *Animal Behaviour* 86: 851-857.
- Yi XF, Wang ZY, Zhang HM, Zhang ZB (2016) Weak olfaction increases seed scatter-hoarding by Siberian chipmunks: implication in shaping plant-animal interactions. *Oikos* 125: 1712-1718.
- Zwolak R, Crone EE (2012) Quantifying the outcome of plant–granivore interactions. *Oikos*.
- Zwolak R, Pearson DE, Ortega YK, Crone EE (2010) Fire and mice: Seed predation moderates fire's influence on conifer recruitment. *Ecology* 91: 1124-1131.
- Zwolak R, Bogdziewicz M, Wrobel A, Crone EE (2016) Advantages of masting in European beech: timing of granivore satiation and benefits of seed caching support the predator dispersal hypothesis. *Oecologia* 180: 749-758.

Table 1. Summary of hypotheses and results.

Hypothesis	Result
(1) Benefits of acorn caching exceed benefits of acorn transportation.	Confirmed.
(2) Benefits of acorn transportation is lower in <i>Q. rubra</i> than in <i>Q. petraea</i> .	Not confirmed: in both oak species, seedling establishment did not change with distance of acorn dispersal.
(3) Seed removal and pilferage increases with distance from the mother tree.	Confirmed.
(4) Pilferage rate is higher in <i>Q. rubra</i> than in <i>Q. petraea</i> .	Not confirmed: pilferage rate was lower in <i>Q. rubra</i> , perhaps due to lower emission of volatile compounds.
(5) Interactions between scatterhoarders and both species of oaks are mutualistic.	Not confirmed: due to relatively low burial benefits and no distance-related benefits, interaction were antagonistic in both species of oaks.

Table 2. Effects of burial, distance from the parent tree, and rodent access on the germination probability of focal oaks. The marginal R^2 of the model for *Q. petraea* equaled 0.28, while conditional 0.34. For *Q. rubra*, the marginal R^2 of the model equaled 0.33, while conditional 0.38.

A) Sessile oak *Quercus petraea*

Fixed effect	Estimate	SE	z	p
Intercept (rodent access allowed, burial: surface)	-1.49	0.35	-4.23	<0.001
Rodents excluded	1.77	0.33	5.31	<0.001
Burial: cached	1.04	0.33	3.10	0.001
Distance	-0.06	0.02	-2.16	0.03
Rodents excluded × Burial: cached	-0.67	0.31	-2.19	0.03
Rodent excluded × Distance	0.04	0.02	1.82	0.07
Burial: cached × Distance	0.04	0.02	2.03	0.04

B) Northern red oak *Quercus rubra*

Fixed effect	Estimate	SE	z	p
Intercept (rodent access allowed, burial: surface)	-1.06	0.39	-2.71	0.006
Rodents excluded	1.09	0.37	2.95	0.003
Burial: cached	1.97	0.36	5.36	<0.001
Distance	-0.10	0.03	-3.75	<0.001
Rodents excluded × Burial: cached	-0.97	0.33	-2.94	0.003
Rodent excluded × Distance	0.06	0.02	2.88	0.004
Burial: cached × Distance	0.03	0.02	1.21	0.22

Figure 1. Probability of seedling establishment of *Q. petraea* and *Q. rubra* in closed (rodent access excluded), and open (rodent access allowed) cages. Sowed indicate acorns sowed in 1-2 cm in the soil, while surface indicate acorns sown on soil surface and covered with leaves to mimic autumn leaf fall. Whiskers indicate standard error.

Figure 2. Probability of seedling establishment of sessile oaks (*Quercus petraea*) as a function of distance from the mother tree. Open cages indicate rodent access, while closed rodent exclusion; red lines acorns buried in topsoil, blue sowed on soil surface. Solid line indicates relationship significantly different from 0, while dashed line non-significant relationship. Shaded areas indicate 95% confidence intervals.

Figure 3. Probability of seedling establishment of red oaks (*Quercus rubra*) as a function of distance from the mother tree. Open cages indicate rodent access, while closed rodent exclusion; red lines acorns buried in topsoil, blue sowed on soil surface. Solid line indicates relationship significantly different from 0, while dashed line non-significant relationship. Shaded areas indicate 95% confidence intervals.

Figure 4. Classification of oak–granivore interactions based on the probability of caching and not retrieving seeds, and the ratio of seedling emergence from the ground to emergence from caches. The net effect of granivores is beneficial at any point above the dotted gray line and antagonistic at any point below. The ‘far’ and ‘near’ categories indicate the establishment ratio calculated based on germination rate estimated at the distance 0 m (near), and 25 m (far). For sessile oaks, the ratio components did not differ with the distance from the seed source tree (see Fig. 2). The values on x-axis (proportion of acorns cached and not retrieved) are derived from parallel study (Bogdziewicz et al. unpublished).

Figure 1.

Figure 2.

Figure 3.

Figure 4.