1 **The brown alga *Saccharina japonica* features distinct**
2 **vanadium-dependent bromoperoxidases and iodoperoxidases**

3

4 **Shan Chi^{1,2}, Tao Liu^{1*}, Hongxin Yin¹, Xin Xu¹, Weiming Zhu¹, Yi Wang¹, Cong Wang¹, Hui**
5 **Lv¹**

6 ¹ Ocean University of China, Qingdao, P. R. China,

7 ² Qingdao Haida BlueTek Biotechnology Co., Ltd, Qingdao, P. R. China.

8

9 ***Corresponding author: liutao@ouc.edu.cn (Tao Liu)**

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30 Abstract

31 Marine algae have an extraordinary ability to absorb halogens which provide algae with an
 32 inorganic antioxidant impacting atmospheric chemistry. Although brown algal Laminariales
 33 species are the most efficient iodine accumulators among all living systems, and *Saccharina*
 34 *japonica* is the primary material used for iodine extraction, the functions and regulatory
 35 mechanisms of these species have not been fully documented. In this study, a functional genomics
 36 analysis of the algal vanadium-dependent haloperoxidase (*vHPO*) gene family was conducted;
 37 there genes can introduce halogen atoms into organic compounds. The comprehensive analyses
 38 regarding the bioinformatics and phylogenetics of novel genomic and transcriptomic sequencing
 39 data of 21 Rhodophyta and 19 Ochrophyta marine algal species revealed that brown algal *vHPOs*
 40 have two gene types, vanadium-dependent bromoperoxidase (*vBPO*) and vanadium-dependent
 41 iodoperoxidase (*vIPO*), with secondary endosymbiotic host origin. The enzyme activity of *S.*
 42 *japonica* *vBPO* and *vIPO* were verified for the first time and were quite stable in a wide range of
 43 temperature and pH values. However, the specific activity and optimal conditions were
 44 considerably different between *vBPO* and *vIPO*. The transcript expression analysis in different *S.*
 45 *japonica* tissues (including rhizoids), generations (sporophytes and gametophytes), sexes (male
 46 and female), and stress conditions (hyposaline and hyperthermia) also showed great differences
 47 between *vBPOs* and *vIPOs*. Most of the *vBPOs* were constitutively expressed with higher
 48 expression dose, which may be responsible for basal halogen metabolism. On the contrary, *vIPOs*
 49 mainly showed specific expression, which may be involved in tissue differentiation, generation
 50 differentiation, sex differentiation, and stress regulation. Comprehensive analysis of gene family
 51 evolution, enzyme biochemical characteristics, and complex transcriptional mechanisms were
 52 conducive to the environmental adaptation and sophisticated system evolution of Laminariales.
 53 The successful bromination of small-molecule compound substrate by *SjavBPO* provided high
 54 activity and efficient enzymatic tools for artificial synthesis of halogenated compounds.

55

56 **Key words:** *Saccharina japonica*, vanadium-dependent haloperoxidases, Laminariales, halogen,
 57 enzyme activity, RNA sequencing

58

59

60 Introduction

61 A halogen substituent is often an essential structural feature of natural products, drugs, or
 62 signaling molecules. At present, over 5,000 halogenated compounds have been isolated, including
 63 halogenated hydrocarbons, halogenated acetylenes, halogenated phenols, halogenated tyrosine,
 64 halogenated fatty acids, and halogenated terpenes (Frank, *et al.*, 2016). They are mainly derived
 65 through marine algal biosynthesis, which has important biological functions (*e.g.*, signaling
 66 molecules and defense compounds) and ecological and atmospheric significance (La Barre, *et al.*,
 67 2010). Brown algae are widely distributed in temperate and subtropical zones and are an important
 68 component of subtidal and intertidal ecosystems (Charrier, *et al.*, 2012). The order Laminariales is
 69 the most efficient iodine accumulator among all living systems, with an average content of 1.0%
 70 dry weight in most *Laminaria* and *Saccharina* species, representing approximately a 30,000-fold
 71 accumulation of this element from seawater (Saenko, *et al.*, 1978; Leblanc, *et al.*, 2006). This is
 72 the reason *Laminaria* was previously used in Europe as raw material for iodine extraction (Lüning,
 73 1985) and *Saccharina* in China as a dietary iodine supplement to prevent goiter (Brinkhuis, *et al.*,
 74 1987).

75 Nature has evolved exquisite methods to introduce halogen atoms into organic compounds
 76 using halogenating enzymes. The vanadium-dependent haloperoxidase (vHPO) is one of the most
 77 studied types among these enzymes due to its biocatalytic properties, including an unusual
 78 stability and tolerance for heat and organic solvents (Coupe, *et al.*, 2007; Fernández-Fueyo, *et al.*,
 79 2015; Sabuzi, *et al.*, 2015; Weichold, *et al.*, 2016). It can be divided into vanadium-dependent
 80 chloroperoxidase (vCPO), vanadium-dependent bromoperoxidase (vBPO), and
 81 vanadium-dependent iodoperoxidase (vIPO), depending on the oxidation ability of the halogen
 82 ions. The first vHPO to be isolated and characterized was the vBPO from the brown alga
 83 *Ascophyllum nodosum* (Vitler, 1984). To date, vHPOs have been characterized from all major
 84 classes of marine algae, such as brown algae (Vreeland, *et al.*, 1998; Weyand, *et al.*, 1999), red
 85 algae (Itoh, *et al.*, 1986, 1987; Shimonishi, *et al.*, 1998; Isupov, *et al.*, 2000), green algae (Itoh,
 86 1985; Wever, *et al.*, 1985; de Boer, *et al.*, 1986; Sheffield, 1993; Ohshiro, 1999; Manley, 2001;
 87 Colin, 2003; Suthiphongchai, 2008), as well as terrestrial lichens (Plat, *et al.*, 1987), fungi (van
 88 Schijndel, *et al.*, 1993; Barnett, *et al.*, 1998), and cyanobacteria (Frank, *et al.*, 2016). These three
 89 enzymes, in terms of the gene structure, have the same conservative metal ion binding sites; the

histidine residues of imidazole rings, the region for vanadium ions, and amino acid series have a high degree of homology (Littlechild, *et al.*, 2002). Overall, *vCPO* gene is mainly found in terrestrial organisms, whereas *vBPO* is mainly found in marine organisms, and *vIPO* is only found in a small number of marine brown algae (Gribble, 2004).

There are few studies on the metabolism of halogens in marine algae. It has been shown that in some brown algae (Wever, *et al.*, 1991) and a green macroalga (Manley, 2001), the vBPOs were located at or near the surface of the seaweed. The possible role of the extracellular enzymatic system may be to control the colonization of the surfaces of the seaweed by generating HOBr, which is directly bactericidal (Hansen, *et al.*, 2003; Renirie, *et al.*, 2008). Until recently, the mechanism of their accumulation, metabolism, and transportation needed further study. In the present study, the first functional *vBPO* and *vIPO* from *S. japonica* were identified. In addition to phylogenetic analysis and transcriptional regulation, this may be worthy of an in-depth study regarding physiological adaptations and relationships with ecological systems and the atmospheric environment. Such research might also provide a platform for diverse protein engineering efforts, and thus an opportunity to establish a new chemoenzymatic halogenation tool in the future.

Materials and methods

Algal sample collection

Preserved *S. japonica* haploid gametophytes (male and female) were available as laboratory cultures and were obtained from our Laboratory of Genetics and Breeding of Marine Organisms. Fresh samples of the *Saccharina* sporophytes (rhizoids, stipe, blade tip, blade pleat, blade base, and blade fascia) were collected from east China (Rongcheng, Shandong Province, 37°8'53"N, 122°34'33"E). These samples were used for RNA Sequencing analysis. Various gametophyte samples (male and female) and tissues of sporophytes (rhizoids, stipe, blade tip, blade pleat, blade base, and blade fascia) were collected to analysis relative gene expressions. To detect the influences of abiotic factors, the female gametophytes and blade base of sporophytes were cultured under different temperatures (Control: 8°C; Hyperthermia: 18°C), salinities (Control: 30‰; Hyposaline: 12‰), and circadian rhythms (Control: 30 μ mol photons/m²·s for 12 h; Darkness: no irradiance for 12 h).

120 **Sequence analysis**

121 In the present study, genes were identified by analyzing transcriptomic and genomic sequencing
122 data of *S. japonica* (Tao Liu, unpublished data), as well as of the species whose genomes and
123 transcriptomes were sequenced and published in OneKP (www.onekp.com) or NCBI. Matching
124 sequences were manually checked for accuracy with the corresponding known cDNA sequences.
125 The unigenes related to *vHPO* were re-verified using the BLASTX algorithm
126 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Sequence identities were calculated using the Clustal
127 Omega tool (<http://www.ebi.ac.uk/Tools/msa/clustalo/>).

128

129 **Phylogenetic tree construction**

130 All downloaded sequences are listed in Supplementary Fig. S1. The sequences were aligned using
131 ClustalX 1.83 software (Thompson, *et al.*, 1997). The amino acid phylogenetic trees were
132 constructed using MrBayes 3.1.2 software (Ronquist and Huelsenbeck, 2003). The posteriori
133 probability was based on the Metropolis-Hastings-Green algorithm through four chains (Markov
134 Chain Monte Carlo, MCMC) with the temperature set to 0.2°C. The chains were run for
135 10,000,000 cycles (Posada and Crandall, 1998; Ronquist and Huelsenbeck, 2003). Random trees
136 were constructed in the MCMC analysis, and one tree in every 1,000 generations was saved. After
137 discarding the aging 25% of all tree samples, the residual samples were used to construct a
138 consensus tree; the tree was rendered using Tree View v.1.6.5 software (Page, 1996).

139

140 **Purification of recombinant proteins expressed in *Escherichia coli***

141 One *vBPO* (*SjavBPO*) and one *vIPO* (*SjavIPO*) from *S. japonica* were synthesized (Shanghai
142 Xuguan Biotechnological Development Co. LTD) to construct recombinant plasmids (NCBI
143 accession number MG195954 and MG195955). *SjavBPO* and *SjavIPO* were cloned in pET32a.
144 These recombinant plasmids were transformed into *E. coli* BL21 (DE3) cells. Isopropyl
145 β-D-1-thiogalactopyranoside (IPTG) was added at a concentration of 0.5 mM to induce the
146 over-expression of the target proteins, and the bacterial cultures were incubated for 16 h at 20°C.
147 His•Bind Resin and GST•Bind Resin were used according to the manufacturer's instructions
148 (www.yuekebio.com). The proteins were stored at -80°C.

149

150 *Assays for enzyme kinetics*

151 The vHPO activity of the purified enzymes was detected using previously described methods
152 (Coupe, *et al.*, 2007). For enzymatic characterization, Cl⁻, Br⁻, and I⁻, which are considered
153 potential substrates, were tested. The effect of pH on the enzymatic activities of the purified
154 proteins was determined in the range of 6.0 to 10.0 for SjavBPO1 and 2.5 to 6.5 for SjavIPO1.
155 The effect of temperature on these enzymes was determined over a range of 10°C to 60°C. Four
156 replicates were analyzed for each condition to ensure the consistency of the experimental results.
157 In each case, boiled purified recombinant enzymes were used as the negative control. All data
158 were subjected to a one-way analysis of variance (one-way ANOVA) followed by a Student's
159 *t*-test.

160 The small molecular compound HD-ZWM-163 used in the halogen addition experiment was
161 provided by Professor Zhu Weiming from Ocean University of China. It is a type of alkaloid
162 staurosporine extracted from actinomycetes *Streptomyces fradiae* 007 var. M315. Its molecular
163 weight is 466 g/mol, with a purity of more than 98% (Supplementary Fig. S2). HD-ZWM-163
164 replaced MCD in the above enzyme assay at 20°C for 12 h. The reactants were extracted three
165 times using ethyl acetate and were dissolved in chromatographic methanol. The HPLC-UV test
166 was performed with 75% methanol (v/v, 1.5% TFA; flow rate: 1 mL/min), and the peak samples
167 were collected for the determination of the HPLC-MS.

168

169 *Transcriptome sequencing*

170 Total RNA was extracted using an improved CTAB method (Gareth, *et al.*, 2006). A total amount
171 of 3 µg RNA per sample was used as input material for the RNA sample preparations. Sequencing
172 libraries were generated using NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (NEB,
173 USA) following the manufacturer's recommendations, and index codes were added to attribute
174 sequences to each sample. The clustering of the index-coded samples was performed on a cBot
175 Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the
176 manufacturer's instructions. After cluster generation, the library preparations were sequenced on
177 an Illumina HiSeq platform, and 125 bp/150 bp paired-end reads were generated. HTSeq v0.6.1
178 was used to count the reads numbers mapped to each gene. The FPKM of each gene was then
179 calculated based on the length of the gene and the reads count mapped to this gene.

180

181 **Results**

182 ***Phylogenetic analysis of algal vHPO genes***

183 Novel transcriptomic sequencing data were obtained for 21 Rhodophyta and 19 Ochrophyta
184 marine algal species (OneKP database), and re-sequencing genomic data for *S. japonica* were used
185 to identify *vHPO* genes. Additionally, 104 new full-length candidate genes from 10 brown algal
186 species and 18 genes from 12 red algal species were detected (Supplementary Fig. S1). There
187 were 21 *vBPOs* and 68 *vIPOs* isolated from genomic and transcriptomic data for *S. japonica*. The
188 homology comparison and structure prediction confirmed that these sequences all belonged to the
189 *vHPO* superfamily.

190 Phylogenetic trees based on full-length amino acid sequences of *vHPO*-related genes from
191 archaeal taxa, bacteria, fungi, and eukaryotic algae were constructed using Bayesian methods
192 (only representative candidates were included). Based on this tree, all *vHPO* genes formed a
193 monophyletic group sharing a common ancestor with the *vCPO* genes in fungi, after which they
194 evolved independently in red and brown algae (Fig. 1). Red algae only have *vBPO* genes.
195 However, brown algae contain *vBPOs* and *vIPOs*, which both have secondary endosymbiotic host
196 origins. These two types of *vHPOs* were paralogues resulting from an ancestral gene duplication.
197 Interestingly, brown alga *Ectocarpus siliculosus* has only one *vBPO* gene in its genome, whereas
198 the closely related species *S. japonica* has 89 *vHPOs*. Based on the consensus tree, 21 *S. japonica*
199 *vBPOs* were clustered into three groups (I–III), and 68 *vIPOs* were clustered into five groups
200 (I–V). The large number of family members are expected to have been derived from recent
201 tandem duplication events, which occurred after the differentiation of *Saccharina* and *Ectocarpus*.

202

203 ***Characterization and confirmation of the functions of vHPO genes from S. japonica***

204 In this study, one *vBPO* (*SjavBPO1*) and one *vIPO* (*SjavIPO1*) were chosen to be over-expressed
205 in *E. coli* to verify their encoding enzyme activities. This was the first functional analysis of these
206 enzymes in *Saccharina*. Specificity of *SjavHPOs* was determined by assaying its activity in the
207 presence of different potential substrates. The oxidation ability of halogen ions Cl^- , Br^- , and I^-
208 were tested. These experiments demonstrated that the *SjavBPO1* enzyme exhibited oxidation
209 activity for both Br^- and I^- , whereas *SjavIPO1* could only oxidize I^- (Table 1). Purified *SjavBPO1*

210 had a specific activity of 5861.1 U/mg and 2342.88 U/mg for Br⁻ and I⁻, respectively. SjavIPO1
211 had a specific activity of 333.6 U/mg for I⁻. These activities were almost the highest of those
212 measured for algal vHPOs (Table 1). In addition, over 90% of the enzymatic activity was detected
213 in SjavHPOs after having been stored at 4°C for 72 h, suggesting that the recombinant proteins
214 were stable under the purification conditions tested.

215

216 Table 1. The protein activity of vHPOs in different organisms.

Gene	Species	Iron	Specific activity (U/mg)	Reference
<i>SjavBPO</i>	<i>Saccharina japonica</i>	Br ⁻	5861.1	This study
<i>SjavBPO</i>	<i>Saccharina japonica</i>	I ⁻	2342.9	This study
<i>SjavBPO</i>	<i>Saccharina japonica</i>	I ⁻	333.6	This study
<i>vBPO</i>	<i>Corallina officinalis</i>	Br ⁻	9.9-469	Zhang, <i>et al.</i> , 2011
<i>vBPO</i>	<i>Corallina officinalis</i>	Br ⁻	450-5000	Coupe, <i>et al.</i> , 2007
<i>vBPO</i>	<i>Corallina pilulifera</i>	Br ⁻	2.32-286	Shimonishi, <i>et al.</i> , 1998
<i>vBPO</i>	<i>Ascophyllum. nodosum</i>	Br ⁻	50-250	Hartung, <i>et al.</i> , 2008
<i>vBPO</i>	<i>Laminaria digitata</i>	Br ⁻	42	Colin, <i>et al.</i> , 2003
<i>vBPO</i>	<i>Laminaria digitata</i>	I ⁻	62	Colin, <i>et al.</i> , 2003
<i>vIPO</i>	<i>Laminaria digitata</i>	I ⁻	310	Colin, <i>et al.</i> , 2003

217

218 The activities of the two purified proteins under different temperatures and pH values were
219 determined to elucidate their biological characteristics. Based on these experiments, the maximum
220 activity of SjavBPO1 for Br⁻ was at 20°C, whereas the activity was 68% and 93% of the maximum
221 activity at 10°C and 25°C, respectively. The optimum temperature for I⁻ was 25°C, with 78% and
222 72% of the residual activity at 20°C and 30°C, respectively (Fig. 2A). For SjavIPO1, the optimum
223 temperature was much higher (50°C), with 68% and 81% of the residual activity at 40°C and 60°C,
224 respectively (Fig. 2B). The optimum pH for SjavBPO1 for both Br⁻ and I⁻ was 6.5, with the
225 activity being 71% to 94% of the maximum activity at other pH values (6.0, 7.0, 8.0, 9.0, 10.0)
226 (Fig. 2C). The optimum pH for SjavIPO1 was determined to be 3.0, and 58% to 99% of the

227 activity remained intact at pH values from 2.5 to 6.5 (Fig. 2D).

228

229 ***Halogen addition of SjavBPO to small-molecule compound***

230 A small-molecule compound (HD-ZWM-163) was used to validate the halogen addition activity
231 of SjavBPO1. In Fig. 3A, a peak of a single compound appears, which is the unmodified
232 compound HD-ZWM-163 (MW = 466 g/mol). The first peak in Fig. 3B and 3C was
233 2-(N-Morpholino)ethanesulfonic acid (MES) in the reaction Buffer, and peak A and peak B were
234 the derivatives of the halogen addition reaction. Peak B2 was approximately four times that of B1,
235 confirming the effect of the SjavBPO1 protein on the HD-ZWM-163 enzymatic reaction. The
236 samples of peaks A1, A2, B1, and B2 were then detected by mass spectrometry analysis.
237 Compounds A1 and A2 were identified as monobromo-HD-ZWM-163 products (Supplementary
238 Fig. S3), whereas B1 and B2 were dibromo-HD-ZWM-163 products (MW = 623 g/mol, Fig. 3D).
239 In addition, the abundance ratio of peak 623:625:627 is approximately 1:2:1 and also confirms
240 that compound B contain two Br atoms. After adding the SjavBPO1 protein, the production of B2
241 was more obvious than that of B1, indicating that SjavBPO1 played a role in the compound
242 HD-ZWM-163 dibromination reaction.

243

244 ***Expression differences of Saccharina vHPOs***

245 The expression of 89 *S. japonica* vHPO genes was determined in different generations
246 (sporophytes and gametophytes), tissue (rhizoids, stipe, blade tip, blade pleat, blade base, and
247 blade fascia), sexes (male and female gametophytes), and stress conditions (hyperthermia,
248 hyposaline) (Fig. 4). In all normal and stress conditions, 26.1% (6/23) and 50% (33/66) of vBPOs
249 and vIPOs were not detected, respectively. However, there were many constitutive vHPOs
250 expressed in different tissues (14 vBPOs and four vIPOs), generations (nine vBPOs and eight
251 vIPOs), sexes (six vBPOs and four vIPOs), and stress conditions (seven vBPOs and two vIPOs). It
252 was determined that vHPOs were widely involved throughout the process of *S. japonica* growth,
253 development, and environmental adaptation (Fig. 5).

254 The vBPOs and vIPOs identified in *S. japonica* showed diverse patterns of expression. The
255 expression of vIPOs was more specific than was the expression of vBPOs (Fig. 6). There were 27
256 vIPO and 16 vBPO genes expressed in sporophyte tissues. The specific expression ratio of vIPOs

(23/27, 85.2%) was 6.8 times higher than that of *vBPOs* (2/16, 12.5%). Sixteen *vBPOs* and 30 *vIPOs* were expressed in different generations, and the specific expression ratios of *vIPOs* and *vBPOs* were 73.3% (22/30) and 43.8% (7/16), respectively. There were nine *vBPOs* and 11 *vIPOs* expressed in male and female gametophytes. The specific expression ratios of *vIPOs* and *vBPOs* were 63.6% (7/11) and 33.3% (3/9), respectively. Female gametophytes displayed a similar response under hyperthermia. The specific expression ratio of *vIPOs* (8/15, 53.3%) was higher than that of *vBPOs* (2/9, 22.2%). On the contrary, the gene number of constitutive expressed *vBPOs* was more than that of *vIPOs*, and expressed constitutive *vBPOs* exhibited higher expression levels than *vIPOs* (Fig. 5). For example, the total expression dose of *vBPOs* in different tissues was about 10^2 – 10^3 times (27.1–520.1 times) that of *vIPOs*.

Considering the different generations, the average expression of sporophyte *vBPOs* (FPKM value = 3307.6) was much higher than that of gametophyte *vBPOs* (FPKM value = 12.7) ($P = 0.001$), whereas there was no significant difference in *vIPOs* between the two groups (FPKM value = 34.1 vs 45.2) (Fig. 7A). There were 26 *vHPOs* (7 *vBPOs* and 19 *vIPOs*) that were only expressed in sporophytes and three *vIPOs* that were only expressed in gametophytes. Among the sporophytes, some genes were only expressed in unique tissues, such as *vIPO55* in the rhizoids, *vIPO36* in the stipes, *vIPO62* in the blade bases, *vIPO7* in the blade tips, *vIPO11* in the blade pleats, and *vIPO60* in the blade fascia. The expression levels were higher in the blade tips than other sporophyte samples, followed by the blade fascia, rhizoids, blade bases, blade pleats, and stipes (Fig. 7B).

The expression of *vHPOs* differed between male and female gametophytes. Few genes were expressed during the gametophyte stage, including nine *vBPOs* and 11 *vIPOs*. One *vBPO* and three *vIPOs* were specifically expressed in male gametophytes, whereas two *vBPOs* and four *vIPOs* were specifically expressed in female gametophytes. Some of the constitutively expressed genes were highly expressed in female gametophytes (*vBPO18*, $P = 0.02$), and some were highly expressed in male gametophytes (*vIPO37*, $P < 0.01$) (Fig. 8).

The expression of *vHPOs* under abiotic stress was significantly upregulated in female gametophytes. Compared to gametophyte samples cultured under normal condition, three *vBPOs* (*vBPO6*, *vBPO7*, and *vBPO21*) and three *vIPOs* (*vIPO38*, *vIPO55*, and *vIPO18*) were expressed under hyposaline induction; two *vBPOs* (*vBPO12*, *vBPO21*) and eight *vIPOs* (*vIPO9*, *vIPO13*,

287 *vIPO15*, *vIPO17*, *vIPO18*, *vIPO19*, *vIPO49*, and *vIPO58*) expressed under hyperthermia (Fig. 9).
 288 However, the expression of the *vHPOs* in gametophytes did not change significantly under the
 289 normal circadian rhythm (12 h irradiation vs. 12 h darkness) or in sporophytes under hyposaline or
 290 hyperthermia stresses.

291

292 **Discussion**

293 *The origin and evolution of vHPOs contributed to the differentiation of brown algal vBPO and* 294 *vIPO genes*

295 Our new data allow for the identification of *vHPO* genes widely distributed among different red
 296 (such as Gigartinales, Gracilariales, and Halymeniales) and brown (such as Laminariales,
 297 Desmarestiales, Ishigeales, and Fucales) algal taxa (Fig. 1). Among the *vHPO* gene family, *vCPO*
 298 appeared first in fungi, followed by *vBPO* and *vIPO* with functional specialization. The eukaryotic
 299 algal *vHPOs* clustered with the *vCPO* genes of fungi, indicating its endosymbiotic host origin. The
 300 differentiation time of *vBPOs* and *vIPOs* could be later than that of the *vCPOs*, which might have
 301 occurred because of variations in the earth's environment. Previous results have demonstrated the
 302 independent evolution of *vHPOs* in red algae and brown algae (Ye, *et al.*, 2015). This study
 303 obtained several *vHPO* genes from several algal species and performed a comprehensive
 304 phylogenetic analysis using a wide range of algal species, clarifying their evolutionary
 305 relationships.

306 Furthermore, there are diverse gene duplications among most red and brown algal taxa, as they
 307 have undergone gene duplications subsequently at different evolutionary time scales. Because
 308 many brown algae were included in the analysis, we confirmed that gene duplications occurred
 309 both before (different groups of *SjavBPOs* and *SjavIPOs*) and after species differentiation (such as
 310 *SjavBPO1* and *SjavBPO2*, *SjavIPO1* and *SjavIPO2*). This suggested that the *vHPO* gene family in
 311 brown algae recently underwent, and perhaps is still undergoing, rapid evolution. Through
 312 genomic sequencing analysis, *E. siliculosus* has only one *vBPO*. However, in Laminariales species,
 313 the gene family expanded dramatically, especially in the genera *Saccharina* and *Laminaria*. As we
 314 know, the iodine content is much higher in these two genera. Vinogradov (1953) demonstrated that
 315 the iodine content in *L. digitata* on the west coast of France was as high as 1.7%. Ji (1963) showed
 316 that the iodine content in *S. japonica* was also greater than 1%. The appearance of functional

317 *vHPOs* indicated the more efficient immobilization of halogen.

318

319 ***The diverse enzyme characteristics of SjavBPO1 and SjavIPO1 were conducive to better***
 320 ***environmental adaption***

321 The enzyme activities of *S. japonica* vBPO1 and vIPO1 were confirmed (Fig. 2), verifying the
 322 authenticity of the pathways as determined through a bioinformatical approach. The oxidation
 323 activity of SjavBPO1 is now known as the highest activity of vBPO and vIPO proteins, compared
 324 to those measured for algal vHPOs listed in Table 1. This also suggested the massive accumulation
 325 of halogen in *S. japonica*. Under the same experimental conditions, the SjavBPO1 protein
 326 exhibited no activity with Cl⁻, and the specific activity with Br⁻ was three times that of the activity
 327 with I⁻. Meanwhile, the SjavIPO1 protein only had the activity with I⁻. This demonstrated the
 328 functional differentiation and specialization between SjavBPO and SjavIPO.

329 The biochemical characteristics of SjavBPO1 and SjavIPO1 differed considerably. The optimal
 330 pH value for SjavBPO1 (pH = 6.5) was similar to that of brown alga *L. digitata* vBPO and vIPO
 331 (optimal pH = 5.5; Colin, *et al.*, 2003) and red alga *Corallina officinalis* vBPO (optimal pH =
 332 7.0–7.5; Coupe, *et al.*, 2007); SjavIPO1 activity was lower (optimal pH = 3.0) (Fig. 2). In contrast,
 333 the SjavIPO1 protein was active up to 60°C, which was the optimal temperature for maximum
 334 activity, similar to *L. digitata* vIPO (optimal temperature = 50°C; Colin, *et al.*, 2003) and *C.*
 335 *officinalis* vBPO (optimal temperature 65°C–70°C; Coupe, *et al.*, 2007). The SjavBPO1 maximum
 336 activity was considerably lower at 20°C and 25°C for Br⁻ and I⁻, respectively; this is similar to the
 337 optimal temperature for *L. digitata* vBPO (30°C) for I⁻ (Colin, *et al.*, 2003). The variation of
 338 optimum conditions for different proteins indicates that they may play a role under different
 339 environmental conditions. Additionally, more than 40% of the maximum activity remained intact
 340 in the temperature range of 20°C–50°C for both SjavBPO1 and SjavIPO1 enzymes. For pH
 341 stability, over 70% of the maximum activity was in the range of 6.0–10.0 for SjavBPO1, and over
 342 57% of the maximum activity was in the range of 2.5–6.5 for SjavIPO1. These results suggested
 343 the high stability of SjavHPO enzymes, which remain active under a wide range of variable
 344 conditions. The iodine content of kelp increases with seawater depth (Saenko, *et al.*, 1978) and
 345 differs between growing seasons (Ar Gall, *et al.*, 2004). In the present study, only two
 346 representative SjavHPOs were analyzed. The dozens of remaining proteins in *S. japonica* might

347 exhibit different biochemical characteristics. These characteristics could help this species to use
348 halogen and facilitate signal transduction and self-defense under different environmental
349 conditions.

350

351 ***The complex regulation of vHPOs was a driving force for the sophisticated system evolution in***
352 ***brown algae***

353 The *SjavHPO* gene expression showed significant differences in various tissues, during different
354 developmental stages, and under different types of stress (Fig. 4). Firstly, a large number of genes
355 were not transcribed in the tested samples, which may be redundant gene backup, or were only
356 expressed under special conditions. In contrast, the constitutively expressed genes maintained
357 basic halogen metabolism under different conditions (Fig. 5). Halogen compounds have important
358 ecological and physiological value for algae, such as free-oxygen cleaning and antimicrobial
359 defense (Renirie, *et al.*, 2008; Hansen, *et al.*, 2003). They form an important source of bromine to
360 the troposphere and lower stratosphere, and they contribute significantly to the global budget of
361 halogenated hydrocarbons (Wever and van der Horst 2013). Additionally, constitutively expressed
362 vHPOs have great physiological and ecological significance.

363 The expression of *vIPOs* was more specific than that of *vBPOs* in different tissues, generations,
364 sexes, and under hyperthermia stress (Fig. 6). However, the expression doses of *vBPOs* were far
365 higher than those of *vIPOs* in sporophytes (Fig. 4). With respect to the higher specific activity of
366 *SjavBPO1* than that of *SjavIPO1* (Table 1), *vBPOs* were supposed to be the core of *S. japonica*
367 halogen metabolism, and responsible for the basal halogen accumulation. In contrast, *vIPOs* are
368 mainly involved in generation differentiation, tissue differentiation, sex differentiation and stress
369 regulation.

370 *S. japonica* possess heteromorphic haploid-diploid life cycles with a macroscopic thallus
371 sporophyte and microscopic gametophyte generation. The gametophyte is similar to some
372 filamentous brown algae, with an isomorphic haploid-diploid filamentous generation (Cock, *et al.*,
373 2014; Bartsch, *et al.*, 2008). Therefore, the comparison between *S. japonica* sporophytes and
374 gametophytes might provide an explanation for the evolution of brown algae from unicellular to
375 multicellular organisms (Chi, *et al.*, 2017). The expression level was higher in sporophytes than in
376 gametophytes (Fig. 7A). This differs from previous studies (Ye, *et al.*, 2015), likely because we

377 annotated more *Saccharina vHPO* genes (89 *vHPOs* in our study vs. 76 *vHPOs* in Ye, *et al.*, 2015),
 378 and the materials were derived from different *Saccharina* strains. The higher expression in the
 379 blade tips (Fig. 7B) is consistent with the higher content of iodine elements in Laminariales distal
 380 blades (Shaw, 1962; Ar Gall, *et al.*, 2004; Küpper, *et al.*, 2008). The blade tips are more sensitive
 381 to environmental stresses and pathogen infections than are the regions beneath the thalli (Küpper,
 382 *et al.*, 1998). This increased gene expression led to greater halogen accumulation. The reserved
 383 material might move toward the halogen-requiring basal meristems through the highly specialized
 384 elongated sieve elements (Amat and Srivastava, 1985). The differences in gene expression and
 385 synthetic product translocation are conducive to the functional differentiation of tissues and
 386 necessary for the supply of halo-containing compounds in brown macroalgae. In addition,
 387 *SjavHPO* expression was significantly upregulated in the gametophytes under stress, such as
 388 hyperthermia and hyposaline conditions (Fig. 9), indicating that *vHPOs* present in *S. japonica*
 389 were involved in stress resistance. However, the expression level in sporophytes did not change
 390 under the same abiotic stresses. One possible explanation is that the gametophyte stage is more
 391 vulnerable to external stress than is the sporophyte stage.

392

393 ***Algal vHPOs have important industrial and pharmaceutical values***

394 Halo-containing compounds, such as acetogenins (anti-microbial activity), and indoles
 395 (anti-inflammatory and anti-cancer activities) could have medical applications (Butler, 1998).
 396 However, conventional chemical manufactures generate potentially harmful byproducts. For
 397 example, synthetic bromination typically yields approximately 50% of the remaining bromine in
 398 the form of waste compounds. Currently, biohalogenation approaches are particularly valuable as
 399 an alternative, which could markedly reduce the amount of halogen pollutants produced. For
 400 example, using a biotransformation approach employing haloperoxidases to produce drugs such as
 401 Vancomycin, Maracen A, and cryptophycins (all halogen-containing compounds) could markedly
 402 reduce toxic levels in wastewater (Coupe, *et al.*, 2007). Therefore, the halogenation of *SjavBPO*
 403 on small molecule compounds (Fig. 3) could provide a more efficient and convenient method for
 404 biohalogenation, which has important economic value and environmental significance.

405

406 In conclusion, the huge diversities in gene family members, enzyme catalytic activities,

407 biochemical characteristics, and gene expression patterns of brown algal *vBPOs* and *vIPOs*
 408 exhibited great differences between large individual parenchyma (sporophyte, 2n) and single-row
 409 filamentous cells (gametophyte, n) regarding biochemistry, physiology, and ecology. The
 410 Phaeophyceae-exclusive *vIPOs*, with a large number of gene expansions and differential
 411 expression, and function in scavenging free oxygen, play a crucial role in the evolution of
 412 unicellular organisms (such as filamentous *S. japonica*) into multicellular organisms (such as *S.*
 413 *japonica* thalli). The deep resolution of the *vHPO* gene family in brown algae has important
 414 biological, ecological, and economic significance.

415

416 **Figure legends**

417 Figure 1. Bayesian phylogenetic tree based on the translated amino acids of *vHPO* with bootstrap
 418 values (when >50%) indicated at the nodes. Brown algae *vHPOs* originated from eukaryotic hosts,
 419 and underwent gene duplication in their common ancestor. Brown algal *vBPOs* were clustered into
 420 3 Group (I, II and III), and *vIPOs* were clustered into 5 Group (I to V). All *vHPO* sequences were
 421 obtained from GenBank or OneKP databases (Supplementary Fig. S1).

422

423 Figure 2. Effects of temperature and pH on activity of SjavBPO1 and SjavIPO1. A. Enzyme
 424 activity of SjavBPO1 at 20°C for Br⁻ and 25°C for I⁻ were set to 100%, respectively. B. Enzyme
 425 activity of SjavBPO1 at pH 6.5 for both Br⁻ and I⁻ were set to 100%. C. Enzyme activity of
 426 SjavIPO1 at 50°C for I⁻ was set to 100%. D. Enzyme activity of SjavIPO1 at pH 3.0 for I⁻ was set
 427 to 100%. Data represent means ± SD of four independent experiments.

428

429 Figure 3. Halogen addition of SjavBPO1 to HD-ZWM-163. (A) The HPLC result of
 430 HD-ZWM-163 (flow rate: 1 mL/min; tR = 3.70 min). (B) The HPLC result of HD-ZWM-163 in
 431 reaction buffer without SjavBPO1 (tA1 = 4.91 min, tB1 = 7.51 min). (C) The HPLC result of
 432 HD-ZWM-163 in reaction buffer with SjavBPO1 (tA2 = 4.91 min, tB2 = 7.51 min). Peaks A1, B1,
 433 A2, and B2 were the derivatives of the halogen addition reaction. (D) The ESI-MS result of B1 in
 434 (B) and B2 in (C). Compounds B1 and B2 were dibromo-HD-ZWM-163 products.

435

436 Figure 4. The transcriptional expression of 89 *S. japonica vHPO* genes in different generations

(sporophytes and gametophytes), tissue (rhizoids, stipe, blade tip, blade pleat, blade base, and blade fascia), sexes (male and female gametophytes) and stress conditions (hyperthermia, hyposaline).

440

Figure 5. Constitutive expressed *SjavHPOs* in different tissues (A), sexes (B) and stress conditions (C).

443

Figure 6. The specific expression gene ratios of *vBPOs* and *vIPOs* in different tissues, generations, sexes and under hyperthermia condition. The expression of *vIPOs* was more specific than that of *vBPOs*.

447

Figure 7. The transcriptional expression of *vBPOs* and *vIPOs* in different sexes (A) and tissues (B). A. The average expression dose of sporophyte *vBPOs* was much higher than that of gametophyte *vBPOs*. The difference between *vIPO* from sporophyte and gametophyte was not significant. B. The highest total expression dose of *vHPOs* was in blade tip, followed by blade fascia, rhizoids, blade base, blade pleat, and stipe.

453

Figure 8. The transcriptional expression of nine *vBPOs* and 11 *vIPOs* in female and male gametophytes. Three *vBPOs* and four *vIPOs* genes were highly expressed in female gametophytes, and one *vBPOs* and four *vIPOs* genes were highly expressed in male gametophytes.

457

Figure 9. The up-regulated and down-regulated gene ratio of *vBPOs* and *vIPOs* under hyposaline and hyperthermia stresses in female gametophytes.

460

461 **Supplementary data**

Supplemental Figure S1. List of *vHPO* sequences used for phylogenetic analysis.

Supplemental Figure S2. The chemical structure of small-compound HD-ZWM-163.

Supplemental Figure S3. The ESI-MS result of peak A1 in Figure 3B and peak A2 in Figure 3C.

465

466 **Acknowledgements**

467 This work was supported by the National Natural Science Foundation of China (NSFC No.
468 41376143), Leading Talents Program in Taishan Industry of Shandong Province, Leading Talents
469 Program in Entrepreneurship and Innovation of Qingdao, China-ASEAN Maritime Cooperation
470 Fund “China-ASEAN Center for Joint Research and Promotion of Marine Aquaculture
471 Technology”, and China Agriculture Research System (CARS-50).

472

473 **References**

474 **Amat MA, Srivastava LM.** 1985. Translocation of iodine in *Laminaria saccharina* (Phaeophyta).
475 J Phycol **21**, 330–333.

476 **Ar Gall E, Küpper FC, Kloareg B.** 2004. A survey of iodine contents in *Laminaria digitata*. Bot
477 Mar **47**, 30–37.

478 **Barnett P, Hemrika W, Dekker HL, Muijsers AO, Renirie R, and Wever R.** 1998. Isolation,
479 characterization, and primary structure of the vanadium chloroperoxidase from the fungus
480 *Embellisia didymospora*. J Biol Chem **273**, 23381–23387.

481 **Brinkhuis BH, Levine HG, Schlenk CG, Tobin S.** 1987. *Laminaria* cultivation in the Far East
482 and North America. In Seaweed Cultivation for Renewable Resources (Bird, K.T. & Benson,
483 P.H., editors), 107–146. Elsevier, Amsterdam, The Netherlands.

484 **Butler A.** 1998. Vanadium haloperoxidases. Current Opinion in Chemical Biology **2(2)**, 279–285.

485 **Charrier B, Le Bail A, de Reviers B.** 2012. Plant Proteus: brown algal morphological plasticity
486 and underlying developmental mechanisms. Trends Plant Sci **17(8)**, 468–477.

487 **Chi S, Liu T, Wang X, Wang R, Wang S, Wang G, Shan G, Liu C.** 2017. Functional genomics
488 analysis reveals the biosynthesis pathways of important cellular components (alginate and
489 fucoidan) of *Saccharina*. Curr Genet doi: 10.1007/s00294-017-0733-4.

490 **Colin C, Leblanc C, Michel G, Wagner E, Leize-Wagner E, van Dorsselaer A, Potin P.** 2005.
491 Vanadium-dependent iodoperoxidases in *Laminaria digitata*, a novel biochemical function
492 diverging from brown algal bromoperoxidases, J Biol Inorg Chem **10**, 156–166.

493 **Colin C, Leblanc C, Wagner E, Delage L, Leize-Wagner E, Van Dorsselaer A, Kloareg**
494 **B, Potin P.** 2003. The brown algal kelp *Laminaria digitata* features distinct bromoperoxidase
495 and iodoperoxidase activities. J Biol Chem **278(26)**, 23545–23552.

496 **Coupe EE, Smyth MG, Fosberry AP.** 2007. The dodecameric vanadium-dependent

497 haloperoxidase from the marine algae *Corallina officinalis*: Cloning, expression, and refolding
 498 of the recombinant enzyme. Protein Expression & Purification **52**(2), 265–272.

499 **de Boer E, Tromp MGM, Plat H, Krenn BE, Wever R.** 1986. Vanadium (V) as an essential
 500 element for haloperoxidase activity in marine brown algae: Purification and characterization of
 501 a vanadium (V)-containing bromoperoxidase from *Laminaria saccharina*. Biochim Biophys
 502 Acta **872**, 104–115.

503 **Fernández-Fueyo E, van Wingerden M, Renirie R, Wever R, Ni Y, Holtmann D, Hollmann F.**
 504 2015. Chemoenzymatic Halogenation of Phenols by using the Haloperoxidase from *Curvularia*
 505 *inaequalis*. Chemcatchem **7**(24), 4035–4038.

506 **Frank A, Seel CJ, Groll M, Gulder T.** 2016. Characterization of a Cyanobacterial
 507 Haloperoxidase and Evaluation of its Biocatalytic Halogenation Potential. Chembiochem
 508 **17**(21), 2028–2032.

509 **Gribble GW.** 2004. Natural organohalogens : a new frontier for medicinal agents. Journal of
 510 Chemical Education **81**(10), 1441.

511 **Hansen EH, Albertsen L, Schäfer T, Johansen C, Frisvad JC, Molin S, Gram L.** 2003.
 512 *Curvularia* haloperoxidase: antimicrobial activity and potential application as a surface
 513 disinfectant. Applied & Environmental Microbiology **69**(8), 4611–4617.

514 **Isupov MN, Dalby AR, Brindley AA, Izumi Y, Tanabe T, Murshudov GN, Littlechild JA.**
 515 2000. Crystal structure of dodecameric vanadium-dependent bromoperoxidase from the red
 516 algae *Corallina officinalis*. J Mol Biol **299**, 1035–1049.

517 **Itoh N.** 1986. Characterization of nonheme type bromoperoxidase in *Corallina pilulifera*. J Biol
 518 Chem **261**, 5194–5200.

519 **Itoh N.** 1987a. Characterization of nonheme iron and reaction mechanism of bromoperoxidase in
 520 *Corallina pilulifera*. J Biol Chem **262**, 11982–11987.

521 **Itoh N, Izumi Y, Yamada H.** 1985. Purification of bromoperoxidase from *Corallina pilulifera*.
 522 Biochem Biophys Res Commun **131**, 428–435.

523 **Ji MH.** 1963. Studies on the chemical composition of the Chinese economic brown seaweeds - II.
 524 Seasonal variations in the main chemical components of *Laminaria japonica*, *Sargassum*
 525 *pallidum* and *Sargassum kjellmanianum* from the north China (in Chinese). Oceanologia et
 526 limnologia sinica **5**(1), 1–10.

527 **Küpper FC, Carpenter L, McFiggans GB, et al.** 2008. Iodide accumulation provides kelp with
528 an inorganic antioxidant impacting atmospheric chemistry. *Proc Natl Acad Sci USA* **105**,
529 6954–6958.

530 **Küpper FC, Schweiger N, Ar Gall E, Legendre J-M, Vilter H, Kloareg B.** 1998. Iodine uptake
531 in Laminariales involves extracellular, haloperoxidase-mediated oxidation of iodide. *Planta* **207**,
532 163–171.

533 **La Barre S, Potin P, Leblanc C, Delage L.** 2010. The halogenated metabolism of brown algae
534 (Phaeophyta), its biological importance and its environmental significance. *Mar Drugs* **8(4)**,
535 988–1010.

536 **Leblanc C, Colin C, Cosse A, et al.** 2006. Iodine transfers in the coastal marine environment:
537 the key role of brown algae and of their vanadium-dependent haloperoxidases. *Biochimie* **88(11)**,
538 1773.

539 **Littlechild J, Garcíarodríguez E, Dalby A, et al.** 2002. Structural and functional comparisons
540 between vanadium haloperoxidase and acid phosphatase enzymes. *Journal of Molecular*
541 *Recognition* **15(5)**, 291–296.

542 **Lüning K.** 1985. *Meeresbotanik: verbreitung, ökophysiologie und nutzung der marinen*
543 *makroalgen*. Thieme Verlag, Stuttgart, Germany.

544 **Manley SL, Barbero PE.** 2001. Physiological constraints on bromoform (CHBr₃) production by
545 *Ulva lactuca* (Chlorophyta). *Limnology & Oceanography* **46(6)**, 1392–1399.

546 **Ohshiro T, Nakano S, Takahashi Y, Suzuki M, Izumi Y.** 1999. Occurrence of bromoperoxidase
547 in the marine green macro-alga, *Ulva lactuca*, and emission of volatile brominated methane by
548 the enzyme. *Phytochemistry* **52(7)**, 1211–1215.

549 **Ohshiro T, Littlechild J, Garcia-Rodríguez E, Isupov MN, Iida Y, Kobayashi T, Izumi Y.**
550 2004. Modification of halogen specificity of a vanadium-dependent bromoperoxidase. *Protein*
551 *Science* **13**, 1566–1571.

552 **Page RD.** 1996. TREEVIEW: An application to display phylogenetic trees on personal computers.
553 *Computer Applications in the Biosciences* **12**, 357–358.

554 **Plat H, Krenn BE, Wever R.** 1987. The bromoperoxidase from the lichen *Xanthoria parietina* is
555 a novel vanadium enzyme. *Biochemical Journal* **248(1)**, 277–279.

556 **Posada D, Crandall KA.** 1998. MODELTEST: testing the model of DNA substitution.

- 557 Bioinformatics **14**, 817–818.
- 558 **Renirie R, Dewilde A, Pierlot C, Wever R, Hober D, Aubry JM.** 2008. Bactericidal and
- 559 virucidal activity of the alkalophilic P395D/L241V/T343A mutant of vanadium
- 560 chloroperoxidase. Journal of Applied Microbiology **105(1)**, 264–270.
- 561 **Ronquist F, Huelsenbeck JP.** 2003. Mrbayes 3: Bayesian phylogenetic inference under mixed
- 562 models. Bioinformatics **19(12)**, 1572–1574.
- 563 **Sabuzi F, Churakova E, Galloni P, Wever R, Hollmann F, Floris B, Conte V.** 2015. Thymol
- 564 Bromination–A Comparison between Enzymatic and Chemical Catalysis. European Journal of
- 565 Inorganic Chemistry **21**, 3519–3525.
- 566 **Saenko GN, Kravtsova YY, Ivaneneko VV, Sheludko SI.** 1978. Concentration of Iodine and
- 567 Bromine by Plants in the Seas of Japan and Okhotsk, Marine Biology **47**, 243–250.
- 568 **Shaw TI.** 1962. Halogens. In Lewin RA (ed.). Physiology and Biochemistry of Algae, Academic
- 569 Press, New York and London pp247–253.
- 570 **Sheffield DJ, Harry T, Smith AJ, Rogers LJ.** 1993. Purification and characterization of the
- 571 vanadium bromoperoxidase from the macroalga *Corallina officinalis*. Phytochemistry **32(1)**,
- 572 21–26.
- 573 **Shimonishi M, Kuwamoto S, Inoue H, Wever R, Ohshiro T, Izumi Y, Tanabe T.** 1998. Cloning
- 574 and expression of the gene for a vanadium-dependent bromoperoxidase from a marine
- 575 macro-alga, *Corallina pilulifera*. Febs Letters **428(1-2)**, 105–110.
- 576 **Suthiphongchai T, Boonsiri P, Panijpan B.** 2008. Vanadium-dependent bromoperoxidases from
- 577 *Gracilaria* algae. Journal of Applied Phycology **20(3)**, 271–278.
- 578 **Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG.** 1997. The ClustalX
- 579 windows interface: flexible strategies for multiple sequence alignment aided by quality analysis
- 580 tools. Nucleic Acids Research **24**, 4876–4882.
- 581 **van Schijndel JWPM, Vollenbroek EGM, Wever R.** 1993. The chloroperoxidase from the
- 582 fungus *Curvularia inaequalis*, a novel vanadium enzyme. Biochim Biophys Acta **1161**,
- 583 249–256.
- 584 **Vilter H.** 1984. Peroxidases from phaeophyceae: A vanadium(V)-dependent peroxidase from
- 585 *Ascomyllum nodosum*. Phytochemistry **23(7)**, 1387–1390.
- 586 **Vinogradov AP.** 1953. The elementary chemical composition of marine organisms. Sears

587 Foundation for Marine Research, Yale University, 647pp.

588 **Vreeland V, Ng KL, Epstein L.** 1998. cDNA sequence and active recombinant vanadium
589 bromoperoxidase from *Fucus* embryos. *Mol BiolCell* **9**, 1043.

590 **Weichold V, Milbredt D, van Pée KH.** 2016. Specific enzymatic halogenation-from the
591 discovery of halogenated enzymes to their applications in vitro and in vivo. *Angew Chem Int*
592 *Ed Engl* **55(22)**, 6374–6389.

593 **Wever R, Plat H, de Boer E.** 1985. Isolation procedure and some properties of the
594 bromoperoxidase from the seaweed *Ascophyllum nodosum*. *Biochim Biophys Acta* **830**,
595 181–186.

596 **Wever R, Tromp MGM, Krenn BE, Marjani A, Tol MV.** 1991. Brominating activity of the
597 seaweed *Ascophyllum nodosum*: impact on the biosphere. *Environmental Science and*
598 *Technology (USA)* **25(3)**, 446–449.

599 **Wever R, van der Horst MA.** 2013. The role of vanadium haloperoxidases in the formation of
600 volatile brominated compounds and their impact on the environment. *Dalton Trans* **42(33)**,
601 11778–11786.

602 **Weyand M, Hecht HJ, Kieß M, Liaud MF, Vilter H, Schomburg D.** 1999. X-ray structure
603 determination of a vanadium-dependent haloperoxidase from *Ascophyllum nodosum* at 2.0 Å
604 resolution. *J Mol Biol* **293**, 595–611.

605 **Ye N, Zhang X, Miao M, et al.** 2015. *Saccharina* genomes provide novel insight into kelp
606 biology. *Nat Commun* **6**, 6986.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9