The strong grip of childhood conditions in older Europeans

Gindo Tampubolon¹

Maria Fajarini²

**1 Manchester Institute for Collaborative Research on Ageing
University of Manchester**

**2 Evidence & Analytics
Manchester**

* tampubolon@manchester.ac.uk

Abstract

Among older Europeans grip strength has been found to be marked by a disadvantaged adulthood. Across the Channel, among older Britons gait speed as another measure of physical function has been found to be marked by disadvantaged childhood. Using the Survey of Health, Ageing, and Retirement in Europe (2004-2013), we studied whether childhood poverty led to Europeans aged 50 to 104 years having a weaker grip. We then drew their trajectories of repeatedly measured grip strength to discern a steeper decline among the childhood poor. Retrospective childhood poverty some four to nine decades in the past was treated as a

latent construct following the above literature; attrition during repeated measurements is handled using inverse proportional to attrition weighting. The data showed the childhood poor to have a weaker grip for half a century in later life. However, they do not show a steeper decline. Most important, by contributing to levels of grip strength in later life, adult condition holds the potential to shape the strong and long arm of childhood condition. The results are another impetus to eliminate childhood poverty to ensure healthy ageing Europeans.

Keywords: Survey of Health, Ageing, and Retirement in Europe; childhood; grip strength

Introduction

Physical functioning is a key driver for the wellbeing of older people, with grip strength and gait speed its two important and well-characterised markers [1]. Therefore, maintaining high levels of physical function throughout later life is singled out as an objective for public health in response to the ageing population challenge [2]. It is also important because with life expectancy at 60 years extending secularly, the welfare implication of impaired physical function in later life is considerable. Long term care of physical disabilities in later life is costly. In the Netherlands and Sweden in 2011, it costs more than 3.5% of their gross domestic products [2]. We therefore aimed to draw trajectories of physical function of Europeans aged 50 to 104 over an extended period.

Grip strength has been repeatedly shown to predict incident disability, morbidity, and mortality [3,4]. Thus a recent study explored its predictors among Europeans aged 65 to 90 using the Survey of Health, Ageing, and

Retirement in Europe (SHARE, 2004-2013), examining the roles of
parental occupation and individual occupation at midlife [5]. The authors
found that grip weakens linearly with age with steeper slopes among men
than women; similar results had been found earlier in a cross-section of
Europeans aged 50 and older [6] and in Danes aged 46 to 102 [7]. In
addition, men with elementary or lower occupation at midlife had a weaker
grip at ages beyond 65, though there was no evidence that they
experienced a steeper decline. Beyond age and sex variations, an earlier
study of SHARE found that grip strength varied considerably across
individual height and geographic region (north – south), advising that
these variables should be adjusted for [6].

Another report used gait speed as a measure of physical function in its
sister study, the English Longitudinal Study of Ageing (ELSA), appraising
the role of an even earlier stage in life course: childhood condition [8]. The
author found that material poverty during childhood associates with slower
gait in Britons aged 50 to 90 years, with material poverty indicated by lack
of essential facilities, overcrowding, and number of books in the childhood
home, as well as financial hardship during childhood. Notably, childhood
information was elicited retrospectively, collecting potentially inaccurate
information [9], and requiring new methods based on latent construct to
deal with inaccurate information [8]. Examined with the new methods, the
data showed that childhood poverty was associated with lower levels of
health status in later life overall: slower gait, poorer memory, and more
depression. The mechanism invoked to link the childhood condition and
later life emphasised the broader effects early life adversity can have. The
results evinced the long arm of childhood condition across the spectrum of

health from physical to mental health.

To understand more about the long arm of childhood condition [10], four improvements can be made. First, most empirical studies are content with explaining levels of health, effectively associating a stage in childhood and a time in later life. The study of older Britons above for example explained the levels of gait speed, episodic memory, and depression by childhood condition. No attempts was made at explaining their rates of change. But surely it is more fruitful to understand whether childhood poverty puts people onto a trajectory of steeper decline. So far, the limited evidence shows no steeper decline among those with disadvantaged childhood or midlife [5].

Second, with some exceptions [5,8], most empirical studies stopped at adulthood. No doubt, this is a function of available data. Although theory suggests that a disadvantaged childhood can be compensated for in adulthood and midlife such that later life health is freed from childhood condition [11], very little evidence is furnished about older people and their childhood. On the other hand, epigenetic change in early life is posited to have a stable effect well into later life [12]. Once biological imprinting has transpired through DNA methylation and histone modification, the effect of childhood condition can persist. Therefore, more empirical investigation is necessary to examine whether childhood condition reach into health trajectories in later life.

Third, information about childhood condition of the oldest old [13] is rarely available in prospective survey. This lack is felt more strongly if a nationally representative sample is required. ELSA collected rich information about people aged 50 years and over prospectively, except

when it comes to information about their childhood which was collected
retrospectively. This information may not be entirely accurate. For
instance, among 50 year old Britons (who had been prospectively followed
since birth), when asked about the numbers of people and bedrooms in
their childhood home, only one in three got both right [9]. Fortunately,
new methods to work with such inaccurate retrospective information have
been proposed and subsequently shown to work with these kinds of data;
such methods need to be applied more often [8].

Lastly, studying older people over time to draw their trajectories of
physical function inevitably faces attrition problems since older people
tend to attrite from a longitudinal study due to worse health
function [14,15]. Recently, a number of solutions have been proposed
including joint modelling and weighting [16,17]; inverse proportional to
attrition weighting is applied here.

We therefore aimed to distinguish the roles of childhood poverty and
adult condition in explaining the trajectories of grip strength of Europeans
aged 50 to 104 years. To tie the four strands together, three questions are
raised. Do those with a poor childhood enter later life with a weaker grip
and remain so throughout? Are their grip strength trajectories also
steeper? Lastly, does good condition in adulthood render negligible any
disadvantage identified earlier?

In answering these questions, this report contributes three ideas to the
literature. The arm of childhood condition is long and strong in predicting
grip strength much later in life. Childhood condition can be recovered
retrospectively and should be considered when explaining health outcomes
of people above 50. Epigenetic changes imprinted by poverty early in life

may lie behind this long and strong arm of childhood condition.

Materials and methods

The Survey of Health, Ageing, and Retirement in Europe (SHARE) is an ongoing longitudinal study of ageing in 20 countries so far [18]. Our use of this anonymised secondary research data has been approved for exemption by the ethical board of the University of Manchester.

As [6] we studied 11 countries, repeatedly surveyed and grouped into two regions: northern-continental (Austria, Denmark, France, Germany, the Netherlands, Sweden, Switzerland) and southern (Greece, Italy, and Spain). We used all waves (2004-2013) matched with the life course survey in 2008 following [5]. The matched sample differed from the rest in the following ways: the participants are older (67.0 vs 66.0 year, $t = 17.5, p < 0.001$) and somewhat weaker (33.5 vs 34.2 kg, $t = 12.1, p < 0.001$). There is a higher proportion of women to men in the analytic sample than in the excluded sample ($\chi^2_1 = 47.7, p < 0.001$).

The outcome variable is objectively measured as the maximum grip strength of the dominant hand obtained using a dynamometer (Smedley, S Dynamometer, TTM, Tokyo, 100 kg) [6]. In contrast, childhood condition as the key exposure was retrospectively obtained. The condition concerned situation at ten years of age i.e. some four to nine decades in the past, indicating lack of the following: indoor toilet, hot and cold running water, central heating, fixed bath; plus overcrowding (more people than bedrooms) as well as number of books in the house, following [8].

It is tempting to use the information unmodified, but this should be resisted. A latent construct solution to obtaining poverty status when its

indicators were inaccurate has been proposed [19]; a particular application 119
has been fruitfully used on ELSA [8] as well as on the China Health and 120
Retirement Longitudinal Study [20], sister studies of SHARE. Following 121
this we built using latent class analysis a childhood poverty status giving 122
poor versus non-poor class based on the indicators above. Beyond dealing 123
with measurement error, this latent construct approach offers substantive 124
advantages that we shall revisit in the discussion. 125

The literature on longitudinal ageing studies is keenly aware that 126
participants tend to attrite non-randomly, hence a number of approaches 127
have been proposed including pattern mixture [5], joint model [16, 21], 128
multilevel multiple imputation [22, 23], and weighting [17, 24–26]. We 129
joined the last stream to apply inverse proportional to attrition weighting. 130
Specifically following [26] in their study of cognition in Atherosclerosis Risk 131
in Community study, the attrition model includes age, sex, smoking, 132
cognition, education, hypertension, cardiovascular disease, diabetes, and 133
retirement status; stabilised weights were then computed with a base 134
model including age, sex, and education. 135

The trajectories are derived using mixed model, also known as latent 136
growth or random coefficients model, which has been used for this 137
sample [5]. We included random intercepts only because there were no 138
meaningful variations in the random age slopes nor extensive discussion of 139
this in the literature [5, 7], retaining the virtue of parsimony [27]. Instead 140
of positing that, *ceteris paribus*, the trajectories change randomly as age 141
unfolds, we posited that they change systematically i.e. the childhood poor 142
have a steeper decline. 143

We explored new factors unexamined in previous work on longitudinal 144

trajectories of grip strength in SHARE. Social inequality in morbidities in later life is well documented, and this suggests inclusion of markers of socioeconomic position and marital status. Log of household income with purchasing power parity exchange rate, education (ISCED three levels: less than high school as reference, high school, and college or higher), occupation (ISCO three levels: elementary as reference, managerial or professional, and others), and marital status (fourfold: never married as reference, married or in partnership, separated or divorced, widowed). We included two markers of disadvantaged adult condition: following [5], adult occupational position (elementary occupation or not), and following [8], adult illness period.

Poverty class as derived above is one of the covariates. Because this is a derived latent class instead of an observed variable, adjustment to standard errors was made following a new method proposed by Vermunt and colleagues [28–30].

To answer the research questions, we built four models separately for men and women following [5–7]. The level model showed childhood poverty association with levels of grip strength, the slope model additionally showed association with the slope of annual decline by interacting poverty with age, while the alternative adult model showed, instead of the interaction term, additional adult condition associations. Lastly, the complete model includes them all. Modelling is done in Latent GOLD 5.1 [31] with model fit judged using Bayes-Schwarz information criterion of the smallest being best.

Results

Women made up the majority of the sample (37,756, 55.6%) which has the average age of 66.9 years (standard deviation [SD]: 9.7 years). Women have weaker maximum grip: 26.0 kg (SD: 7.1 kg) compared to men: 42.6 kg (SD: 10.4 kg). The average of the maximum grip strength among northern-continental Europeans is (34.8 kg, SD: 12.0) while among southern Europeans is (31.2 kg, SD: 11.7 kg). The sample is further summarised in Table 1.

The latent class analysis of childhood poverty revealed that 45.9% of the participants had a poor childhood at age ten (Table 2). The indicators of childhood condition showed plausible loadings. For instance, lacking more facilities is positively loaded on being a poor child while having more books is negatively loaded on being a poor child.

As women are found to have lower levels of grip strength than men, we presented their grip strength trajectories separately. The best model based on BIC is the adult model for both. Information criteria and key coefficients for both models are put together in Table 3 and discussed each in turn, putting complete information criteria (Supplement Table 4) and complete coefficients (Supplement Table 5) in the Supplement.

As has been widely documented, men have stronger grip (higher intercepts) but have steeper annual decline (418 versus 303 gram). Both slopes are significant ($p < 0.001$) and gentler than the estimates for the Danes [7]. Individual height and geographic region are also significant, in accordance with the literature [6]. Adult illness, a marker of adult condition, inversely associates with grip strength throughout; and so is being employed as elementary worker, which accords with [5]. Despite the

Table 1. Description of the analytic sample (SHARE 2002-2013)

Variable	Women		Men	
	Mean or <i>N</i>	Std. dev. or %	Mean or <i>N</i>	Std. dev. or %
Grip strength	25.996	7.059	42.625	10.375
Age	66.836	10.029	67.042	9.338
Height	162.218	6.611	173.684	7.465
Married	24,361	64.52	24,511	81.22
Single	2,112	5.59	1,827	6.05
Sep/divorced	3,252	8.61	1,949	6.46
Widowed	8,031	21.27	1,892	6.27
Elementary occ.	996	2.64	539	1.79
Intermediate	36,068	95.53	28,574	94.68
Professional	692	1.83	1,066	3.53
Primary or less	14,185	37.57	9,284	30.76
High School	17,031	45.11	14,155	46.90
College	6,540	17.32	6,740	22.33
Household income	21,820	60,656	25,629	47,118
Adult illness: none	30,181	80.28	24,839	82.52
One	5,052	13.44	3,849	12.79
Two	1,022	2.72	734	2.44
Three	378	1.01	198	0.66
More than three	577	1.53	252	0.84
Most adulthood	384	1.02	230	0.76
Adult elementary occ.: no	27,644	73.22	21,060	69.78
Yes	10,112	26.78	9,119	30.22
North	23,259	61.60	18,525	61.38
South	14,497	38.40	11,654	38.62
Lack facility: none	6,580	17.43	5,466	18.11
1	3,999	10.59	3,012	9.98
2	3,475	9.20	2,708	8.97
3	5,691	15.07	4,560	15.11
4	7,786	20.62	6,270	20.78
Lack all five	10,225	27.08	8,163	27.05
Num. books: none/very few	17,103	45.78	13,794	46.13
One shelf	7,910	21.17	6,230	20.83
One bookcase	7,419	19.86	5,994	20.04
Two bookcases	2,483	6.65	1,870	6.25
Three or more	2,442	6.54	2,016	6.74
Over-crowded: no	10,103	26.76	8,312	27.54
Yes	27,653	73.24	21,867	72.46

Table 2. Latent classes of poor and non-poor childhood.

Indicator	Non-poor	Poor
Size	54.1%	45.9%
Over-crowded		
No	0.4384	0.0597
Yes	0.5616	0.9403
Lack facility		
None	0.2418	0.0116
1	0.1782	0.0228
2	0.1709	0.0567
3	0.1686	0.1398
4	0.1334	0.2673
Lack all five	0.1071	0.5019
Number of books		
None/very few	0.2156	0.7091
One shelf	0.2377	0.2104
One bookcase	0.3245	0.0729
Two bookcases	0.1093	0.0061
Three or more	0.1129	0.0016

Table 3. Mixed models of trajectories of grip strength (adjusting for household income, occupation, education, and marital status); SE: standard error. Source: SHARE 2004-2013.

Covariate	Women			Men		
	coef	SE	<i>p</i>	coef	SE	<i>p</i>
Intercept	28.1459	3.7292	< 0.001	53.7396	3.3773	< 0.001
Age	-0.3028	0.0100	< 0.001	-0.4184	0.0112	< 0.001
Height	0.0926	0.0212	< 0.001	0.1303	0.0188	< 0.001
Southern Europe	-2.7182	0.2662	< 0.001	-2.5454	0.2862	< 0.001
Adult illness	-0.3664	0.0780	< 0.001	-0.6170	0.1445	< 0.001
Adult elementary occupation	-0.3857	0.0934	< 0.001	-0.7460	0.0801	< 0.001
Poor childhood	-1.1932	0.2370	< 0.001	-2.1917	0.2189	< 0.001
BIC	359347			1990012		

strong effects of adult condition, the childhood-poor Europeans still have weaker grip in their later lives: men by 2.19 kg and women by 1.19 kg. In summary, the coefficients are plotted in Figure 1 to help in making comparison, and the trajectories of predicted grip strength for men and women who were childhood-poor and otherwise are drawn in Figure 2.

Figure 1. Plots of key coefficients for women (left pane) and men (right pane).

Discussion

Maintaining higher levels of grip strength is key, since it is a core component to avoid frailty and sarcopenia and ensure healthy ageing and wellbeing of older people. Here is the first evidence that being poor in life's first decade goes with weaker grip in life's last five decades.

Beyond covering a more extended age group than recent studies [5, 8], our study confirmed that adult condition (elementary occupation or ill health in adulthood) is associated with a weaker grip. These results are

Figure 2. Predicted trajectories of grip strength, distinguished by childhood poverty status for women (left pane) and men (right pane).

robust to inaccuracies in the measurement of childhood condition and to the attrition so common in longitudinal ageing studies. The results on the associations of childhood and adulthood conditions are strengthened because other factors have been accounted for including household income, education, occupational class, and marital status [32,33]. In short, excepting the question about a steeper decline, the results supplied affirmative answers to all our questions: both childhood poverty and adulthood disadvantage go hand in hand with a weaker grip in later life.

Such long range results can be underpinned by a biosocial mechanism, especially with chronic inflammation playing a major role [34]. Older age is often marked by chronic or low grade inflammation which can impair muscle function. In turn, inflammation itself can be upregulated as a result of childhood adversity. The mechanism therefore has two major steps: [i]

childhood adversity to lifetime inflammation, and [ii] inflammation
disrupting myogenic processes of regeneration and functioning. We take
each in turn.

Epigenetic literature has been accumulating evidence using animal
models, such as mice, rats, and macaques, to examine whether early life
adversity imprints epigenetic changes (DNA methylation and histone
modification) to otherwise similar genotypes, resulting in different
phenotypic response [34,35]. Childhood poverty, pointing to broader early
life adversity, entails more than just material lack but includes social
deprivation when parents' nurturing is compromised due to their time
being absorbed in providing for household members and making ends meet.

Therefore animal models capable of reflecting some of the complexity of
material and social deprivation are uniquely revealing, especially macaques
studies. They have been used in a randomised design (of parental caring of
frequent versus infrequent licking or grooming) to study the causal effect of
early life deprivation on DNA methylation [36,37]. The study found stable
and organised epigenetic changes, involving genes in the pathways of the
immune system and the hypothalamic pituitary adrenal (HPA) axis
responsible for responding to stress. The peripheral immune system
interacts with the HPA axis and has a role in brain function; evidence
consistent with this interaction has been shown in this sample in our
previous work [38].

A key gene for regulating the HPA axis function, the glucocorticoid
receptor (*NR3C1*), is activated in the hypothalamus in response to stress
and releases glucocorticoid. Glucocorticoid receptor is differentially
expressed according to the experience of social deprivation, by epigenetic

programming through histone acetylation and DNA methylation of the
 exon **1_F**. This epigenetic programming differentiates similar DNA
 sequences phenotypically, resulting in blunted feedback by glucocorticoid
 and heightened stress response and demodulated immune system response,
 a pattern that is stable throughout the life course. The bidirectional
 interaction between the HPA axis and the immune system facilitates the
 imprinting of childhood adversity through epigenetic changes. This can
 lead to chronic inflammation that is stable through later life as reflected in
 higher levels of circulating tumour necrosis factor- α (TNF- α).

By discussing the role of inflammatory cytokines such as TNF- α , the
 literature on muscle regeneration and muscle function has provided
 evidence to complete the mechanism. Inflammation is known to impair
 both muscle regeneration and muscle functioning. In normal activities of
 daily living which involve muscle exertion, some minute damage to muscle
 tissue may occur [39]. In these circumstances, the pluripotent myosatellite
 cells respond by proliferating and differentiating to form muscle fibres and
 cover the damaged tissue. But circulating inflammatory cytokines such as
 TNF- α have been shown to impair this process of regeneration in two ways:
 apoptosis of myoblasts [40] and inhibition of the differentiation stage,
 leaving proliferated cells unable to differentiate and replace the damaged
 tissue [41]. Beyond impairing the myogenesis process in common minute
 damage, inflammation also impairs functioning by reducing the power of
 the single permeable fibre [42]. So in mice, TNF- α rapidly reduces the
 force generating capacity or specific tension of muscle fibres independent of
 loss of muscle volume [43].

In short, inflammation impairs muscle functioning in older people at

least along three points: it encourages myosatellite cell deaths [44], it
interrupts the step of differentiation into myonuclei and muscle fibres;
lastly, even if muscle fibres were successfully regenerated, inflammation
reduces the febrile tensile strength. Childhood poverty, through
upregulating inflammation, impairs grip strength in later life.

This study has a number of weaknesses. First, by matching only
individuals with childhood information with those with longitudinal
observations, inevitably some unmatched observations were set aside. It is
impossible to measure the direction of possible bias this might entail.
Second, although epigenetic changes are posited to be the mechanism,
there is no direct evidence of the extent of DNA methylation in the sample.
This is a potentially rectifiable weakness. Despite these weaknesses, this
study has some strengths. First, the sample is designed to represent the
countries and not only some clinical groups or cities, hence facilitating
generalisation. Finally, this study is also the first to link broad childhood
condition (subject to recall error) with later life trajectories (subject to
attrition), reinforcing sustained links across the life course.

As alluded to above, besides uncovering the strong results on childhood
poverty, the method with which childhood poverty is constructed i.e. as a
latent class of poverty, holds potential to advance research work on the life
course and health. It is useful to call to mind that childhood information,
such as a lack of the five facilities above, can be used alternatively as (i)
indicators of a latent factor in factor analysis or (ii) five additional
covariates. Now the use of a latent class of poverty facilitates discussion,
for instance when presenting whether the childhood poor (compared to the
non-poor) show better health outcomes in later life. On the other hand,

with the latent factor we have to compare those on one standard deviation away from the mean against those on the mean of the latent factor. This is hardly intuitive. Or with five additional covariates, we are led to scrutinise each effect which makes discussion potentially unwieldy.

Second, a latent class is also easier to use when testing a hypothesis of a steeper decline; it simply needs an interaction term (of poverty class and age). The interpretation will be similarly intuitive: the childhood poor declined more steeply by a certain kg (the coefficient) per year if the interaction term was found significantly negative. On the other hand, although with the latent factor a similar interaction can be used, there remains the attendant difficulty of interpretation. Or with five additional covariates, we are required to use five interaction terms. Depending on the choice, interpretation may be hindered.

Most important, a latent class enables cross-country comparison. It is conceivable that the long arm of childhood condition hypothesis may be tested in other countries, for example in equatorial developing countries where lack of running hot water or central heating may not hold similar salience. Without these indicators, nevertheless, poverty class can still be constructed with this method and the hypothesis tested. In this way, latent class of childhood poverty based on retrospective information should always be considered in life course and ageing investigation anywhere in the world.

In conclusion, although childhood and adulthood conditions last a lifetime [12], there is a potential role for interventions in adulthood both in the labour market and the health sector. On the basis of evidence uncovered here, childhood could be a critical period to stave costly long

term care. It is never too early to invest in later life.

325

Acknowledgments

This work was supported by grants from the Medical Research Council and Economic & Social Research Council (No. G1001375/1), the National Institute of Health Research (No. ES/L001772/1) and the European Union's Horizon 2020 Research and Innovation Programme (No. 668648) to Gindo Tampubolon.

This paper used data from SHARE Waves 1, 2, 3 (SHARELIFE), 4 and 5 (DOIs: 10.6103/SHARE.w1.500, 10.6103/SHARE.w2.500, 10.6103/SHARE.w3.500, 10.6103/SHARE.w4.500, 10.6103/SHARE.w5.500), see Börsch-Supan et al. (2013) for methodological details. The SHARE data collection has been primarily funded by the European Commission through the FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812) and FP7 (SHARE-PREP: No. 211909, SHARE-LEAP: No. 227822, SHARE M4: No. 261982). Additional funding from the German Ministry of Education and Research, the U.S. National Institute on Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGHA_04-064) and from various national funding sources is gratefully acknowledged (see www.share-project.org)

Table 4. Supplement Table 4 Model comparisons.

Model	BIC
Male: level	2012538
Male: slope	2008427
Male: adult condition	1990012
Male: all	2007423
Female: level	376341
Female: slope	369201
Female: adult condition	359347
Female: all	363772

Table 5. Supplement Table 5 Models with adult condition for women (left pane) and men (right pane)

Covariate	Women			Men		
	coef	SE	<i>p</i>	coef	SE	<i>p</i>
Intercept	28.1459	3.7292	< 0.001	53.7396	3.3773	< 0.001
Poor childhood	-1.1932	0.2370	< 0.001	-2.1917	0.2189	< 0.001
Age	-0.3028	0.0100	< 0.001	-0.4184	0.0112	< 0.001
Log household income	0.1375	0.0026	< 0.001	0.1633	0.0148	< 0.001
High school	2.2142	0.2716	< 0.001	1.9144	0.2723	< 0.001
College	5.5721	0.3256	< 0.001	10.2978	0.5578	< 0.001
Intermediate	1.7347	0.1905	< 0.001	-1.8704	0.3893	< 0.001
Managerial	3.7451	0.3295	< 0.001	0.2778	0.5475	0.61
Married	1.9782	0.3128	< 0.001	-2.1077	0.4301	< 0.001
Sep/divorced	4.7904	0.7347	< 0.001	4.3090	0.3546	< 0.001
Widowed	2.0289	0.3310	< 0.001	-3.5030	0.5218	< 0.001
Height	0.0926	0.0212	< 0.001	0.1303	0.0188	< 0.001
Southern Europe	-2.7182	0.2662	< 0.001	-2.5454	0.2862	< 0.001
Adult illness period	-0.3664	0.0780	< 0.001	-0.6170	0.1445	< 0.001
Adult elementary work	-0.3857	0.0934	< 0.001	-0.7460	0.0801	< 0.001
σ^2	4.1348	0.0805	< 0.001	4.1061	0.0940	< 0.001

Supplement Table 4 and 5

References

1. Dodds RM, Syddall HE, Cooper R, Benzeval M, Deary IJ, Dennison EM, et al. Grip strength across the life course: Normative data from twelve British studies. PLoS ONE. 2014 12;9(12):1–15.

2. WHO. World Report on Ageing and Health. Geneve: WHO; 2015.
Available from:
www.who.int/ageing/publications/world-report-2015/.
3. Sasaki H, Kasagi F, Yamada M, Fujita S. Grip strength predicts cause-specific mortality in middle-aged and elderly persons. *The American Journal of Medicine*. 2007;120(4):337 – 342.
4. Cooper R, Kuh D, Cooper C, Gale CR, Lawlor DA, Matthews F, et al. Objective measures of physical capability and subsequent health: a systematic review. *Age and Ageing*. 2011;40(1):14–23.
5. Kröger H, Fritzell J, Hoffmann R. The association of levels of and decline in grip strength in old age with trajectories of life course occupational position. *PLoS ONE*. 2016;p. e0155954.
6. Andersen-Ranberg K, Petersen I, Frederiksen H, Mackenbach JP, Christensen K. Cross-national differences in grip strength among 50+ year-old Europeans: results from the SHARE study. *European Journal of Ageing*. 2009;6(3):227–236.
7. Frederiksen H, Hjelmberg J, Mortensen J, McGue M, Vaupel JW, Christensen K. Age trajectories of grip strength: Cross-sectional and longitudinal data among 8,342 Danes aged 46 to 102. *Annals of Epidemiology*. 2006;16(7):554–562. Available from:
<http://dx.doi.org/10.1016/j.annepidem.2005.10.006>.
8. Tampubolon G. Growing up in poverty, growing old in infirmity: The long arm of childhood conditions in Great Britain. *PLoS ONE*. 2015 12;10(12):1–16.

9. Brown M. Assessing recall of early life circumstances: evidence from the National Child Development Study. *Longitudinal and Life Course Studies*. 2014;5(1):64–78.
10. Hayward MD, Gorman BK. The long arm of childhood: The influence of early-life social conditions on men’s mortality. *Demography*. 2004;41(1):87–107.
11. Shanahan MJ, Mortimer JT, Johnson MK, editors. *Handbook of the Life Course*. New York: Springer; 2016.
12. Szyf M. How do environments talk to genes? *Nat Neurosci*. 2013 Jan;16(1):2–4. Available from: <http://dx.doi.org/10.1038/nn.3286>.
13. Suzman RM, Willis DP, Manton KG, editors. *The Oldest Old*. Oxford: Oxford University Press; 1992.
14. Newman AB. An overview of the design, implementation, and analyses of longitudinal studies on aging. *Journal of the American Geriatrics Society*. 2010;58(S2):S287–S291.
15. Hardy SE, Allore H, Studenski SA. Missing data: A special challenge in aging research. *Journal of the American Geriatrics Society*. 2009;57(4):722–729.
16. Tampubolon G. Cognitive ageing in Great Britain in the new century: Cohort differences in episodic memory. *PLoS ONE*. 2016 12;10(12):1–17.

17. Tampubolon G. Trajectories of the healthy ageing phenotype among middle-aged and older Britons, 2004-2013. *Maturitas*. 2016;88(7):9 – 15.
18. Börsch-Supan A, Brandt M, Hunkler C, Kneip T, Korbmacher J, Malter F, et al. Data Resource Profile: The Survey of Health, Ageing and Retirement in Europe (SHARE). *International Journal of Epidemiology*. 2013; Available from: <http://ije.oxfordjournals.org/content/early/2013/06/18/ije.dyt088.abstract>.
19. Breen R, Moisisio P. Poverty dynamics corrected for measurement error. *The Journal of Economic Inequality*. 2004;2(3):171–191. Available from: <http://dx.doi.org/10.1007/s10888-004-3227-9>.
20. Yang F, Lou V. Childhood adversities, urbanisation and depressive symptoms among middle-aged and older adults: evidence from a national survey in China. *Ageing and Society*. 2016;36(5):1031–1051.
21. Tampubolon G. Delineating the Third age: Joint models of older people's quality of life and attrition in Britain 2002-2010. *Aging & Mental Health*. 2015;19:576–583. Available from: <http://dx.doi.org/10.1080/13607863.2014.1003279>.
22. Romaniuk H, Patton GC, Carlin JB. Multiple imputation in a longitudinal cohort study: A case study of sensitivity to imputation methods. *American Journal of Epidemiology*. 2014;.

23. Kalaycioglu O, Copas A, King M, Omar RZ. A comparison of multiple-imputation methods for handling missing data in repeated measurements observational studies. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*. 2016;179(3):683–706.
Available from: <http://dx.doi.org/10.1111/rssa.12140>.
24. Cole SR, Hernán MA. Constructing inverse probability weights for marginal structural models. *American Journal of Epidemiology*. 2008;168(6):656–664.
25. Hernán MA, Brumback B, Robins JM. Marginal structural models to estimate the causal effect of zidovudine on the survival of HIV-positive men. *Epidemiology*. 2000;11(5):561–570.
26. Gottesman RF, Rawlings AM, Sharrett AR, Albert M, Alonso A, Bandeen-Roche K, et al. Impact of differential attrition on the association of education with cognitive change over 20 years of follow-up: The ARIC Neurocognitive Study. *American Journal of Epidemiology*. 2014;179(8):956–966.
27. Box GEP. Some problems of statistics and everyday life. *Journal of the American Statistical Association*. 1979;74(365):1–4.
28. Vermunt JK. Latent class modeling with covariates: Two improved three-step approaches. *Political Analysis*. 2010;18(4):450–469.
29. Bakk Z, Tekle FB, Vermunt JK. Estimating the association between latent class membership and external variables using bias-adjusted three-step approaches. *Sociological Methodology*. 2013;43(1):272–311.

30. Bakk Z, Oberski DL, Vermunt JK. Relating latent class assignments to external variables: Standard errors for correct inference. *Political Analysis*. 2014;22(4):520–540.
31. Vermunt JK, Magidson J. Upgrade Manual for Latent GOLD 5.1. Belmont, MA: Statistical Innovations; 2016.
32. Deaton A. The Great Escape: health, wealth, and the origins of inequality. Princeton, NJ: Princeton University Press; 2013.
33. Marmot M. The Health Gap: The Challenge of an Unequal World. London: Bloomsbury; 2015.
34. McGowan PO, Sasaki A, D'Alessio AC, Dymov S, Labonte B, Szyf M, et al. Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nat Neurosci*. 2009 Mar;12(3):342–348. Available from: <http://dx.doi.org/10.1038/nn.2270>.
35. Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, et al. Epigenetic programming by maternal behavior. *Nature Neuroscience*. 2004;7(8):847–854.
36. Liu D, Diorio J, Tannenbaum B, Caldji C, Francis D, Freedman A, et al. Maternal care, hippocampal glucocorticoid receptors, and Hypothalamic-Pituitary-Adrenal responses to stress. *Science*. 1997;277(5332):1659–1662.
37. Szyf M. The early-life social environment and DNA methylation. *Clinical Genetics*. 2012;81(4):341–349. Available from: <http://dx.doi.org/10.1111/j.1399-0004.2012.01843.x>.

38. Tampubolon G. Repeated systemic inflammation was associated with cognitive deficits in older Britons. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*. 2016;3:1 – 6.
39. Degens H. The role of systemic inflammation in age-related muscle weakness and wasting. *Scandinavian Journal of Medicine & Science in Sports*. 2010;20(1):28–38.
40. Saini A, Al-Shanti N, Faulkner SH, Stewart CE. Pro- and anti-apoptotic roles for IGF-I in TNF-alpha-induced apoptosis: A MAP kinase mediated mechanism. *Growth Factors*. 2008;26(5):239–253.
41. Langen RCJ, Schols AMWJ, Kelders MCJM, van der Velden JLJ, Wouters EFM, Janssen-Heininger YMW. Muscle wasting and impaired muscle regeneration in a murine model of chronic pulmonary inflammation. *American Journal of Respiratory Cell and Molecular Biology*. 2006;35(6):689–696. Available from: <http://www.atsjournals.org/doi/abs/10.1165/rcmb.2006-01030C>.
42. Hardin BJ, Campbell KS, Smith JD, Arbogast S, Smith J, Moylan JS, et al. TNF-alpha acts via TNFR1 and muscle-derived oxidants to depress myofibrillar force in murine skeletal muscle. *Journal of Applied Physiology*. 2008;104(3):694–699. Available from: <http://jap.physiology.org/content/104/3/694>.
43. Supinski GS, Callahan LA. Caspase activation contributes to endotoxin-induced diaphragm weakness. *Journal of Applied Physiology*. 2006;100(6):1770–1777.

44. Jejuri SS, Henkelman EA, Cederna PS, Marcelo CL, Urbanchek MG, Jr WMK. Aging increases the susceptibility of skeletal muscle derived satellite cells to apoptosis. *Experimental Gerontology*. 2006;41(9):828 – 836.