Impacts of predators, habitat, recruitment, and disease on soft-shell clams *Mya arenaria* and stout razor clams *Tagelus plebeius* in Chesapeake Bay

AUTHORS AND ADDRESSES: Cassandra N. Glaspie^{a,*†}, Rochelle D. Seitz^a, Matthew B. Ogburn^b, Christopher F. Dungan^c, Anson H. Hines^b

^aVirginia Institute of Marine Science, College of William & Mary, P.O. Box 1346, Gloucester Point, Virginia 23062, U.S.A.

^bSmithsonian Environmental Research Center, 647 Contees Wharf Rd., Edgewater, Maryland 21037, U.S.A.

^cMaryland Department of Natural Resources, Cooperative Oxford Laboratory, 904 S. Morris St., Oxford, Maryland 21654, U.S.A

*Corresponding author contact information: glaspiec@oregonstate.edu; phone: 541-737-6105

†Present address: Oregon State University, Department of Fisheries and Wildlife, 104 Nash Hall, Corvallis, OR 97731, USA

RUNNING PAGE HEAD: Factors influencing clam decline

KEY WORDS: bivalve, seagrass, oyster, temperature, climate, *Perkinsus chesapeaki*

ABSTRACT

Soft-shell clams, *Mya arenaria*, and razor clams, *Tagelus plebeius*, in Chesapeake Bay have declined since the 1970s, with severe declines since the 1990s. These declines are likely caused by multiple factors including warming, predation, habitat loss, recruitment limitation, disease, and harvesting. A bivalve survey in Chesapeake Bay examined influential factors on bivalve populations, focusing on predation (crabs, fish, and cownose rays), habitat (mud, sand, gravel, shell, or seagrass), environment (temperature, salinity, and dissolved oxygen), recruitment, and disease. *M. arenaria* and *T. plebeius* were found more often in habitats with complex physical structures (seagrass, shell) than any other habitat. Pulses in bivalve density associated with recruitment were attenuated through the summer and fall when predators are most active, indicating that predators likely influence temporal dynamics in these species. Presence of *Mya arenaria*, which is near the southern extent of its range in Chesapeake Bay, was negatively correlated with water temperature. Recruitment of *M. arenaria* in Rhode River, MD, declined between 1980 and 2016. Infection by the parasitic protist *Perkinsus* sp. was associated with stressful environmental conditions, bivalve size, and environmental preferences of *Perkinsus* sp, but was not associated with bivalve densities. It is likely that habitat loss, low recruitment, and predators are major factors keeping *T. plebeius* and *M. arenaria* at low densities in Chesapeake Bay. Persistence at low densities may be facilitated by habitat complexity (presence of physical structures), whereas further reductions in habitats such as seagrass and shell hash could result in local extinction of these important bivalve species.

INTRODUCTION

The soft-shell clam *Mya arenaria* and the stout razor clam *Tagelus plebeius* are both large, deep-burrowing bivalves that are harvested in the Chesapeake Bay for human consumption and for bait (Dungan et al. 2002, Homer et al. 2011). *Mya arenaria*, in particular, supports a large commercial fishery in the U.S., and it accounted for 14% of domestic commercial bivalve dollar value in 2015 (NMFS 2016). In Chesapeake Bay, *M. arenaria* has supported a commercial hydraulic dredge fishery in Virginia and Maryland starting in the early 1950s with the invention of the hydraulic dredge. Commercial clammers also harvest *T. plebeius* for eel and crab bait (Dungan et al. 2002, Homer et al. 2011).

Historically, *M. arenaria* and *T. plebeius* served important roles as biomass dominants that contributed substantially to the food web and water quality of Chesapeake Bay (Abraham & Dillon 1986, Eggleston et al. 1992, Seitz et al. 2001). *Mya arenaria* and *T. plebeius* are key prey for numerous commercially and recreationally important fish and crab species (Eggleston et al. 1992, Seitz et al. 2005, Fisher 2010). These large-bodied filter-feeding clams likely played a large role in filtration of the water column when they were abundant; *M. arenaria* in the Baltic Sea can filter the entire water column in less than a day (Forster & Zettler 2004), and filtration by *M. arenaria* in the Gulf of Maine increased water clarity in some lagoons where their densities were high (Thiet et al. 2014). Non-oyster bivalves have recently gained attention for water filtration as an ecosystem service in Chesapeake Bay (Gedan et al. 2014), but the value of filtration by thin-shelled clams has not been assessed.

In Chesapeake Bay, *M. arenaria* has been in decline since the early 1970s, with more

pronounced declines since the 1990s, and this species now exists in Chesapeake Bay at record low levels (Fig. 1). Declines during and after 1972 are attributed to impacts from Tropical Storm Agnes, a 100-year storm that drastically reduced salinities and increased sedimentation throughout Chesapeake Bay (Hyer & Ruzecki 1976, Schubel 1976, Schubel et al. 1976), which resulted in a mass mortality event for *M. arenaria* (Cory & Redding 1976). Due to this storm, which followed declines in abundance of *M. arenaria* in the late 1960s, the commercial hydraulic dredge fishery for *M. arenaria* essentially ended in Virginia waters around 1968, and has failed to recover since then (Haven, 1970). More recent (post-1990) dramatic declines in abundance of *M. arenaria* have resulted in variable, low harvests in Maryland waters (Homer et al. 2011). Since 1980, commercial clammers have gradually switched to harvest of *T. plebeius* for eel and crab pot bait (Dungan et al. 2002, Homer et al. 2011). Like *M. arenaria*, *T. plebeius* populations have experienced declines in recent years, which were first documented in 2003 and resulted in the loss of 70-80% of the population in Maryland (Fig. 1; Homer et al. 2011). There are no historic landings records or long-term time series of *T. plebeius* abundance, so the history of decline and potential mechanisms for decline in this species are largely unknown.

Multiple factors likely control densities of *M. arenaria* and *T. plebeius* in the Chesapeake Bay and other systems, including predation (Seitz et al. 2001, Beal 2006), overfishing (Brousseau 2005), low recruitment (Beukema & Dekker 2005, Bowen & Hunt 2009), rising temperatures (Najjar et al. 2000), habitat loss (Glaspie 2017), and disease mortalities (Dungan et al. 2002). Some of these factors, in particular disease and overharvesting, have been blamed for the inability of *M. arenaria* to recover from Tropical Storm Agnes and the recent declines in *M. arenaria* and *T. plebeius* abundances (Fisher et al. 2011, Homer et al. 2011). Overharvesting is unlikely to be responsible for current trends in Chesapeake Bay clam densities because, as

previously mentioned, fishing pressure is not consistent throughout the Bay. The role of the remaining factors in determining local or regional population dynamics of *M. arenaria* and *T. plebeius* has yet to be determined.

There is some evidence that habitat preferences, including availability of refuge habitat that allows clams to avoid predation, may influence population dynamics of *M. arenaria* and *T. plebeius*. Habitat type has been shown to influence the distribution and abundances of *M. arenaria*. Growth rates can be impacted by sediment type, with higher growth rates observed in sand than mud-gravel-shell mixtures (Newell 1982). Habitats with more structure, such as gravel, shell hash, or seagrass, can also provide refuge from predators (Sponaugle & Lawton 1990, Skilleter 1994, Seitz et al. 2001). The availability of some refuge habitats, such as seagrass and oyster shell, is decreasing in the Chesapeake Bay (Rothschild et al. 1994, Lefcheck et al. 2017). The severe and persistent declines of Chesapeake Bay seagrass in the recent past have been attributed mostly to anthropogenic nutrient and sediment pollution, which decrease water clarity (Kemp et al. 2004). The dominant seagrass species in Chesapeake Bay, eelgrass *Zostera marina*, is also sensitive to warming, and the interaction between warming and poor water quality has resulted in recent declines of shallow beds (Lefcheck et al. 2017). In the past several years, seagrass die-offs induced by extreme high temperatures in Chesapeake Bay have resulted in the prediction that *Z. marina* may disappear from the Bay entirely (Moore & Jarvis 2008; though seagrass may be increasing in the coastal bays, see Orth & McGlathery 2012). Loss of oyster reefs in Chesapeake Bay due to overfishing, habitat destruction, and disease has resulted in the ecological extinction of oysters in the Bay (Rothschild et al. 1994, Wilberg et al. 2011). As oysters have been lost, so has a major source of shell material in the Bay. Oyster shell is a limited resource, and shell half-life is estimated to be as little as 3-10 years in Delaware Bay

(Mann & Powell 2007). Loss of this structural refuge may have implications for populations of deep-burrowing clams such as *M. arenaria* and *T. plebeius*.

The recent findings of high prevalences of the parasitic protist *Perkinsus chesapeaki* in *M. arenaria* and *T. plebeius* suggest that this could be a cause of the population declines of both bivalve species (Dungan et al. 2002, Reece et al. 2008). *Perkinsus chesapeaki* infections have reached epizootic levels in *M. arenaria* (Farley et al. 1991, Dungan et al. 2002, Reece et al. 2008). However, prevalence is not necessarily equal to pathogenicity, as seen in some disease-resistant oyster stocks in the Chesapeake Bay (Encomio et al. 2005). The cancer disseminated neoplasia also causes mortality of *M. arenaria*, but this disease is not as prevalent as *P. chesapeaki*, and is not reported to affect *T. plebeius* (Dungan et al. 2002); thus, the disease aspect of this study focuses only on the impact of *P. chesapeaki* on *M. arenaria* and *T. plebeius*.

Extreme temperature, dissolved oxygen, or salinity levels may be stressful to *M. arenaria* and *T. plebeius*, resulting in mortality or metabolic constraints that make these conditions not suitable for growth. *Mya arenaria* is near the southern extent of its geographic range in Virginia, and is not tolerant of temperatures above 28°C, a temperature that is frequently exceeded in Chesapeake Bay during summer months (Moore & Jarvis 2008). In contrast, *T. plebeius* is a warm-water species that is distributed into South America (Abraham et al. 2010), and high water temperatures (above 28°C) may not be stressful for this species. Benthic species in general are intolerant of dissolved oxygen levels below 1.5 mg L⁻¹ (Rosenberg et al. 1991), and *M. arenaria* will decrease burial depth and extend their siphons into the water column under severe hypoxia (< 1.5 mg L⁻¹) (Taylor & Eggleston 2000). Thus, low dissolved oxygen is likely to be stressful for both *M. arenaria* and *T. plebeius*. Finally, *T. plebeius* are rarely found at salinities under 5 ppt (Holland et al. 1987), and *M. arenaria* are intolerant of salinities below 4 ppt (Abraham &

Dillon 1986).

Settlement and post-settlement processes are also important in determining distribution of organisms in soft-sediment communities (Olafsson et al. 1994). Larval behavior, hydrodynamics, and intense predation of postlarval bivalves control patterns of *M. arenaria* recruitment in other systems (Beukema & Dekker 2005, Bowen & Hunt 2009). Recruitment of *M. arenaria* remains high in several tributaries of Chesapeake Bay (Lovall et al. 2017) though these individuals rarely survive to adulthood, except in habitats with sufficient structure to allow protection from predators (Seitz et al. 2005). Locally, recruitment of *M. arenaria* is not necessarily correlated with abundance of adult clams (Bowen & Hunt 2009), but low densities of *M. arenaria* and *T. plebeius* throughout the Chesapeake Bay may limit recruitment and contribute to the loss of these bivalve species, as seen in bay scallops *Argopecten irradians irradians* in Long Island Sound (Tettelbach et al. 2015). In addition, larval and post-larval individuals of *M. arenaria* are more susceptible to extreme environmental conditions such as high temperatures and low salinities (Abraham & Dillon 1986), and population dynamics for this species may be controlled by this sensitive life stage.

No single factor can likely be exclusively attributed to the declines of these two bivalve species; thus, multiple factors were considered to gain useful insight into the problem. In this study, we used field surveys to concurrently examine predation, habitat, recruitment, and disease prevalence/intensity to disentangle the relative effects of these factors on survival and persistence of *M. arenaria* and *T. plebeius*. We hypothesized that: (1) densities of *M. arenaria* and *T. plebeius* are positively associated with presence of complex habitat such as seagrass and shell, and negatively correlated with predator densities; (2) densities of *M. arenaria* are negatively correlated with temperature; and (3) intensity and prevalence of infection by presumed *Perkinsus*

chesapeaki (hereafter, *Perkinsus* sp.) increase under stressful environmental conditions (i.e., positively correlated with temperature and river discharge, negatively correlated with salinity and dissolved oxygen).

MATERIALS AND METHODS

Study system

The Chesapeake Bay provides an ideal location to study the effects of predators, habitat type, recruitment, disease, and physical factors on the distribution and abundance of *M. arenaria* and *T. plebeius*. The Bay offers a range of environmental conditions including temperature, salinity, dissolved oxygen, and availability of complex habitat. Lower Chesapeake Bay (the Virginia portion of Chesapeake Bay) is mostly polyhaline (except in the upper reaches of the tributaries). Upper Chesapeake Bay (the Maryland portion of the Bay) is mostly mesohaline. Bottom habitat type in the upper Bay is characterized by oyster shell hash, soft muds, fine sands, and gravel/pebbles (Smith et al. 2003). Bottom type in the lower Bay is similar to that of the upper Bay (Wright et al. 1987), except shallow shoals in the lower Bay south of the Potomac River often have mixed beds of eelgrass *Z. marina* and widgeongrass *Ruppia maritima* (Orth et al. 2010). The entire Chesapeake Bay experiences seasonal hypoxia in deep channel water, which occasionally wells up onto the shoals with the tides and wind events (Sanford 1990, Kemp et al. 2005). Summer temperatures in the upper Bay are likely still hospitable for *M. arenaria*, while the lower Bay experiences periods when they are above the tolerance limit for the species.

In Chesapeake Bay, the dominant predators of *M. arenaria* and *T. plebeius* include the

blue crab *Callinectes sapidus*, horseshoe crabs *Limulus polyphemus* (Botton 1984, Lee 2010), and demersal fishes (de Goeij et al. 2001, Seitz et al. 2001) including cownose rays *Rhinoptera bonasus* (Fisher 2010). Crabs forage for clams from the sediment and consume the entire clam (Beal 2006), whereas demersal fishes nip clam siphons, causing clams to reduce their burial depth and exposing them to increased predation by probing predators (de Goeij et al. 2001). High predation rates on infauna are also associated with seasonal migratory behavior of cownose rays (Blaylock 1993), which are able to consume bivalves that would otherwise avoid predation by burrowing, armor, and/or size refuges (Fisher 2010).

Survey design

Clams were collected from three subestuaries of lower Chesapeake Bay (Lynnhaven River, York River, Mobjack Bay), and three subestuaries of upper Chesapeake Bay (Western Shore, Eastern Bay, Chester River) in fall 2011; spring/summer/fall 2012; and either spring/summer 2013 (for lower Bay) or summer/fall 2013 (for upper Bay; see Table 1 for sampling dates). Each sampling season, four to nine sites within each subestuary were sampled; sites were specific tributaries or shorelines in the subestuary that were chosen to represent the range of available benthic habitat types (mud, sand, gravel, shell, or seagrass) available in the subestuary (Table 1).

At each sampling site in the lower Bay, three replicate samples were collected in each season from shallow water of 1.5-2 m depth mean high water using a suction sampling device that collects samples of 0.11 m² area and 40 cm depth. Replicate sample locations in each site were selected by throwing the suction cylinder from the boat, and samples were sieved through

3-mm mesh. All samples were assigned a habitat category (mud, sand, gravel, shell, or seagrass) based on observations made in the field and the lab during sample processing. In the upper Bay, due to the low density of clams and a lack of seagrass in clam habitat, samples were collected using a commercial hydraulic escalator dredge targeting areas fished for *M. arenaria*. This method of collection was not used in the lower Bay to prevent unnecessary destruction of existing seagrass beds, and because clams were consistently captured using less destructive methods such as suction sampling. Habitats were categorized by the presence/absence of oyster shell hash retained on the escalator dredge, and by diver surveys conducted in 2012. All *M. arenaria* and *T. plebeius* were counted, and clam densities at each site were determined by sample area (in lower Bay) or dredge distance (in upper Bay). All clams were measured to the nearest 1 mm for shell length. Bottom (e.g. < 1 m from the sediment surface) dissolved oxygen, salinity, and temperature were also recorded at each site using a YSI (Model 85, Yellow Springs Instruments).

Between spring 2012 and summer 2013, blue crab abundance was quantified at each lower Bay site within a few days of bivalve suction sampling using six replicate 20-m tows of a modified commercial crab scrape (usually used for harvesting soft-shell peeler crabs in seagrass in lower Chesapeake Bay; 6 mm mesh, 1 m width; Seitz et al., 2008) because it is effective sampling mobile fauna in seagrass. In the upper Bay, seagrass was not a concern and two replicate 4.9 m-wide otter trawl tows were conducted at each site in the spring, summer, and fall of 2012 for seven minutes each (approx.~ 500 m). Gear efficiencies of both methods are similar: 24-34% for capture of juvenile crabs with the crab scrape (Ralph 2014) and ~22% for capture of most predators, including fish and crabs, with the otter trawl (Homer et al. 1980). Any fish caught in tows were counted and released. A fish species list and number caught can be found in

Supplemental Table 1. Approximately 10% of bivalve sampling events were paired with trawls that occurred approximately a month before or after bivalve sampling, due to logistical constraints. In this case, an average of crab or fish abundance before and after the bivalve sampling was used in analyses. At each bivalve sampling site and for each season, the number of ray pits within 1 m to either side of a 50-m transect were counted, and are treated as a proxy of cownose ray density (Hines et al. 1997). Ray pits were about 0.3 m in diameter and 10 cm deep, and could be easily seen in good visibility, or detected by sweeping the sediment with hands in poor visibility. Horseshoe crabs were not quantified because they were not abundant enough in any of our samples. Fish and crab abundances per tow and ray pit counts were converted to density per 100 m² for analyses. Due to logistical constraints, certain data, such as environmental data and predator abundance, were missing from the survey dataset, especially for fall 2011, so models that included these missing variables were fit to data from spring 2012 through fall 2013. Environmental data from the CBIBS monitoring buoy at 38.963N 76.448W were used for Maryland western shore samples from summer and fall 2013.

We conducted a gear comparison of bivalve sampling methods by using suction samples collected in the same areas as hydraulic dredging in spring 2012. Site-average densities from suction samples were regressed against site-average densities from hydraulic dredging for both *M. arenaria* and *T. plebeius*. *Mya arenaria* densities calculated using suction sampling were 3.58 times higher than those calculated from hydraulic dredge samples ($R^2 = 0.95$; Supplementary Fig. 1a); thus, suction sample densities for *M. arenaria* were reduced by a factor of 3.58 to allow both data sets to be analyzed together. Densities calculated for *T. plebeius* using both methods were in close agreement (slope = 1.10, $R^2 = 0.90$; Supplementary Fig. 1b), so the raw data sets were combined for analyses. Suction samples appeared to collect more small and large *Mya*

arenaria (mean 42.33 mm, SD 13.73 mm; Supplementary Fig. 2a) than dredge sampling (mean 40.22 mm, SD 5.42 mm; Supplementary Fig. 2b), indicating that the discrepancy in densities was not an artifact of hydraulic dredge mesh size. Biomass of both *M. arenaria* and *T. plebeius* from the lower Bay was determined as ash free dry weight of bivalves dried in a drying oven for 24 hours, and ashed in a muffle furnace at 550 °C for five hours; bivalves in the upper Bay were not processed for biomass.

The long-term trend of *M. arenaria* recruitment in Maryland was evaluated using data from surveys conducted by SERC in the Rhode River, Maryland. Benthic core samples were collected approximately quarterly (typically March/April, June, October, December) at two sandy subtidal sites from 1981-2016. These sites were located at 38.885797N, 76.541790W and 38.868377N, 76.518276W. Cores were 10.2 cm in diameter and 35 cm long. Seven core samples were taken per site on each sample date and animals retained on a 500 µm sieve were preserved in 10% buffered formalin and stained with rose Bengal. All *M. arenaria* were measured for shell length and individuals <10 mm shell length were considered recruits. The total number of recruits in each year was calculated by site and then averaged across the two sites to provide an index of annual recruitment.

Disease

Whenever possible, 30 clams from each upper Bay site were held for 24-96 h in flow-through systems of ambient Rhode River or Tred Avon River waters before they were examined for *Perkinsus* sp. infections. Clams were dissected to secure labial palp tissues that were inoculated into tubes containing 3 mL of Ray's fluid thioglycollate medium (RFTM, Ray 1966),

and then incubated in the dark at 28 °C for 4-7 d. After incubation in RFTM, individual clam tissues were macerated in pools of 25% Lugol's iodine before microscopic examinations at 40x magnification. *Perkinsus* sp. hypnospores were quantified in the most heavily infected palps of individual clams, and infections were assigned ordinal intensity ranks of 0-5.

Perkinsus sp. prevalence and intensity data for 2011-2013 were analyzed together with similar data collected 2000-2009. Only data from clams collected at the same geographic locations by both investigations were used for the time series. For samples collected during our 2011-2013 survey, infection intensity categories included the rank of 0.5 for infections of very low intensities (Ray 1954). In accordance with suggested method for cross-calibrating such data (Dungan & Bushek 2015), infection intensity data for 2011-2013 were cross calibrated to those of 2000-2009 by pooling infection intensities of the lowest ranks (0.5, 1.0) (Dungan et al. 2002, Reece et al. 2008, Homer et al. 2011). Prevalences were estimated as the proportion of infected clams.

Clams tested for *Perkinsus* sp. infections were predominantly collected from upper Bay sites where hydraulic dredging was used to reliably collect adequate numbers of adult clams. Although hydraulic escalator dredges were not used in the lower Bay, 30 juvenile *M. arenaria* were collected by hand from Indian Field Creek, York River, VA, in April 2013, held in flow-through tanks for several months until they were 20-30 mm in shell length, and were assayed for *Perkinsus* sp. infections in July 2013.

Statistical analyses

Both *M. arenaria* and *T. plebeius* exhibited many instances of zero catch, so two models

were used to analyze the data: presence/absence was modeled with a binomial generalized additive model (GAM, logit link), and non-zero densities were modeled with a Gaussian GAM (identity link) on log-transformed data. GAMs for *T. plebeius* included the following as parametric predictors: temperature ($^{\circ}\text{C}$), salinity, dissolved oxygen (mg L^{-1}), ray pit density (m^{-2}), crab density (m^{-2}), fish density (m^{-2}), and habitat (5 categories: gravel, mud, sand, shell, and seagrass). *Mya arenaria* were rarer and data contained many more zeroes than *T. plebeius*; to allow for model convergence, avoid over-smoothing, and ensure homoscedasticity of residuals, habitat was reduced to two categories (simple, including mud, sand, and gravel; and complex, including seagrass and shell) in both presence/absence and non-zero density models, and the presence/absence model was run only on spring/summer data for 2012. *Perkinsus* sp. infection intensity rank-score was modeled as a log-transformed continuous variable using a Gaussian GAM (identity link). Disease GAMs for *M. arenaria* and *T. plebeius* included the following predictors: temperature ($^{\circ}\text{C}$), salinity, dissolved oxygen (mg L^{-1}), bivalve length (mm), and bivalve density (m^{-2}). To account for trends in data across space, a two-dimensional spline (trend-surface) was fit on latitude and longitude for all GAMs (Cressie 1993). To account for trends in time, sampling day was fit in each model using a spline with a kernel smoothing function (Kohn et al. 2002). The number of knots (k) in each model was chosen automatically using the Generalized Cross-validation (GCV) optimizer in the R package ‘mgcv’ (Wood 2017), except for the presence/absence model for *M. arenaria*, which was fit with a spline of order $k = 2$ because only two seasons were modeled and the GCV optimizer produced a model with skewed residuals.

All variables were examined for multicollinearity with scatter plots and Pearson correlation coefficients before inclusion in the model. There was no evidence of

multicollinearity; all variable combinations had Pearson correlation coefficients ≤ 0.8 (Berry & Feldman 1985; Supplementary Figure 3). For all GAMs, appropriateness of the smoothing parameter was assessed using the k-index method and the smoothing parameter chosen was deemed adequate if the simulated p-value was > 0.05 (Wood 2017). Homoscedasticity of residuals was assessed visually using residual and quantile-quantile plots. All models had appropriate smoothing and met the assumption of homogeneity of variance.

Time series (infection intensity, infection prevalence, and *M. arenaria* recruitment) were analyzed using automatic time series forecasting (ARIMA models; Hyndman & Khandakar 2008), and disease models included spring/summer temperature (April-August) obtained from the Chesapeake Bay Program Water Quality Database (<http://www.chesapeakebay.net/data>) for all tidal mainstem Chesapeake Bay stations, and annually averaged Susquehanna River discharge obtained from the USGS water quality monitoring station at Harrisburg, PA (National Water Information System at <https://waterdata.usgs.gov/usa/nwis>). No autocorrelation or non-stationarity was detected in ARIMA models of *Perkinsus* sp. infection intensities or prevalences ([p,d,q] = [0,0,0]), so analysis proceeded using linear models. For multiple comparisons, significant difference was determined using non-parametric bootstrap hypothesis testing with 10,000 simulations and $\alpha = 0.05$ (Efron & Tibshirani 1993), and Cohen's d was calculated as a measure of effect size for all two-group comparisons. All analyses were completed in R (R Core Team 2017), and code and data files have been provided as supplementary material.

RESULTS

Survey

Temperature over the course of the survey (fall 2011 – summer 2013) ranged 12.6 °C – 33.3 °C. Temperatures exceeding 28 °C and 30 °C were observed for 23.7% and 6.5% of all samples collected, respectively. Salinity ranged from 5.3 psu in the upper York River site to 25.7 psu in the Lynnhaven Bay sites. The minimum dissolved oxygen level observed was 3.3 mg L⁻¹ in Mobjack Bay, though the site means ranged 6.50 mg L⁻¹ - 9.35 mg L⁻¹ (Table 1).

Densities of *M. arenaria* were 2.75 times higher at sites in the lower Bay than at sites in the upper Bay ($p = 0.02$, $d = 0.01$, Fig. 2a). The maximum density observed was 53.33 m⁻² *M. arenaria* near the mouth of the York River, VA, in spring 2012. Out of all samples collected, 54.73% of upper Bay and 89.82% of lower Bay samples did not contain *M. arenaria*. Despite repeated sampling, samples from four sites in Virginia contained no *M. arenaria* (two in Mobjack Bay and two in Lynnhaven). Throughout all samples, densities of *Tagelus plebeius* were 13.07 times higher than *M. arenaria* densities ($p < 0.001$, $d = 0.02$, Fig. 2b). Densities of *Tagelus plebeius* were 8.37 times higher at sites in the lower Bay than at sites in the upper Bay ($p < 0.001$, $d = 0.04$). The maximum density observed for *T. plebeius* was 218.18 m⁻² in the lower Bay in Lynnhaven (Linkhorn Bay) in summer 2013. *Tagelus plebeius* were found at every lower Bay site, but were not found at two sites in the Eastern Bay of Maryland throughout the study period. Trends in biomass of both *M. arenaria* and *T. plebeius* in the lower Bay closely matched trends in densities (Supplementary Figure 4).

Average shell length of *M. arenaria* was 1.43 times greater at sites in the upper Bay than at sites in the lower Bay ($p < 0.001$, $d = 0.07$; Fig. 3a). Similarly, *T. plebeius* at sites in the upper Bay were 2.07 times larger than those at sites in the lower Bay ($p < 0.001$, $d = 0.04$; Fig. 3b).

Only 5.56% of *M. arenaria* collected in the lower Bay were greater than 50 mm in shell length, while 29.73% of upper Bay *M. arenaria* were 50 mm or larger.

The highest densities of *Mya arenaria* each year occurred in the spring, with declining density through the fall (Fig. 4a). *Tagelus plebeius* densities did not show consistent seasonal trends over the study period. In the lower Bay, *T. plebeius* densities were the lowest in spring 2012, increased through the summer and fall, and then remained high (generally 15-20 m⁻²) and stable through 2013 (Fig. 4b). In contrast, in the upper Bay the clearest seasonal trend involved a pulse of *T. plebeius* in spring 2012 (~5 m⁻²) that disappeared through the summer and fall (Fig. 4a).

Density of *M. arenaria* was higher in complex habitat (seagrass and shell) than in less complex habitat (mud, sand, and gravel) (Fig. 5a). The odds of finding *M. arenaria* in seagrass or shell were 392.04 times greater than in less-complex habitats such as mud, sand, and gravel. When *M. arenaria* were present, densities in seagrass or shell were 2.15 times greater than in less-complex habitats such as mud, sand, and gravel. The odds of finding *M. arenaria* decreased as temperature increased, and when they were present, *M. arenaria* density was negatively correlated with dissolved oxygen (Table 2). There was no significant relationship between *M. arenaria* density and predator density (Table 2).

Tagelus plebeius density in seagrass was higher than in less-complex habitats (shell, gravel, sand, and mud) (Fig. 5b). The odds of finding *T. plebeius* in seagrass were 11.65 times greater than in mud ($z = 1.48$, $p = 0.10$), 3.95 times greater than in sand ($z = 0.81$, $p = 0.09$), and 11.28 times greater than in shell ($z = 0.69$, $p = 0.0005$). When *T. plebeius* were present, densities in sand were 4.63 times greater than in gravel, 5.39 times greater than in seagrass, and 4.74 times greater than in shell. High densities in sand were only observed in the spring (Fig. 5b). High

densities of *T. plebeius* were also observed in shell, though this was not significant in the models (Table 2). When they were present, *T. plebeius* density increased with salinity (Table 2). There was no significant relationship between *T. plebeius* density and temperature, dissolved oxygen, or predator density (Table 2).

Mya arenaria recruitment in the Rhode River declined steadily between 1980 and 2016. Recruitment was characterized by an Autoregressive Integrated Moving Average (ARIMA[2, 1, 1]), with a negative trend (drift) of -0.03 million clams $\text{ha}^{-1} \text{y}^{-1}$. The survey captured no recruits post-2005 (Fig. 6).

Disease

Mean infection intensity for *Mya arenaria* was light at upper Bay sites, with a mean of 0.48 on a 5-point scale (Fig. 7a). Infection intensity in *M. arenaria* increased by a factor of 0.55 for each degree increase in temperature, and increased by a factor of 0.07 for each mm in shell length (back-transformed from Table 2). *Mya arenaria* infection intensity was positively correlated with temperature, salinity, and dissolved oxygen (Table 2). The mean infection intensity for *T. plebeius* was light to moderate and was 1.57 times greater than that for *M. arenaria* ($p < 0.001$, $d = 0.03$; Fig. 7b). Mean infection intensity for *T. plebeius* decreased by a factor of 0.16 with each unit increase in salinity, and increased by a factor of 0.21 with each unit increase in temperature (back-transformed from Table 2). Infection intensity increased with *T. plebeius* size, but did not increase with bivalve density for either *M. arenaria* or *T. plebeius* (Table 2).

Perkinsus sp. infection prevalence was low for *M. arenaria*, averaging 29.21% over the

course of the survey (2011-2013). *Perkinsus* sp. infection prevalence during the survey (2011-2013) was 2.17 times higher in *T. plebeius* (Fig. 7c) than in *M. arenaria* (Fig. 7d; $p < 0.001$, $d = 0.06$). Mean infection prevalence for both bivalve species increased with temperature and dissolved oxygen, and prevalence also increased with salinity for *M. arenaria* (Table 2). Infection prevalence increased with bivalve size for *M. arenaria*, but not for *T. plebeius* (Table 2). Infection prevalence and intensity were not related to *M. arenaria* or *T. plebeius* densities (Table 2). In the lower Bay, where 30 juvenile *M. arenaria* were assayed, hypnospores were present in two individuals at very low infection intensities (ordinal rank 0.05).

Prevalence of infection was moderate to high between 2000 and 2013, with mean infection rates of 68.46% for *M. arenaria* and 81.86% for *T. plebeius* (Fig. 8a). Across all years in the time series, average prevalence of infection was similar between species ($d = 0.25$, $p = 0.16$), and average infection intensities were light and similar between *M. arenaria* and *T. plebeius* ($d = 0.04$, $p = 0.18$). Infection prevalence and intensity were negatively correlated with annual Susquehanna River discharge for *M. arenaria* (prevalence: coef. = -1.25×10^{-5} , $t_9 = -3.40$, $p = 0.01$; intensity: coef. = -2.28×10^{-5} , $t_9 = -2.31$, $p = 0.05$) but not for *T. plebeius* (prevalence: coef. = -6.99×10^{-6} , $t_8 = -1.60$, $p = 0.15$; intensity: coef. = -1.17×10^{-5} , $t_8 = -1.28$, $p = 0.24$; Fig. 8c). Infection prevalence and intensity were not correlated with spring/summer temperature for *M. arenaria* (prevalence: coef. = 2.96×10^{-2} , $t_9 = -0.56$, $p = 0.59$; intensity: coef. = 6.23×10^{-2} , $t_9 = 0.44$, $p = 0.67$) or for *T. plebeius* (prevalence: coef. = -2.51×10^{-2} , $t_8 = -0.40$, $p = 0.70$; intensity: coef. = -9.95×10^{-2} , $t_8 = -0.76$, $p = 0.47$).

DISCUSSION

As we hypothesized, the distributions of thin-shelled commercial species *Mya arenaria* and *Tagelus plebeius* were positively associated with complex habitat. Densities of both *M. arenaria* and *T. plebeius* were associated with habitats with a high degree of complexity (seagrass and shell) as compared to some less-complex habitats (mud, sand, and gravel). Complex habitats may be more favorable for these species because they increase rates of larval settlement by baffling water currents (Heiss et al. 2010), provide increased food resources for both suspension and facultative deposit-feeding species (Peterson et al. 1984), and provide refuge from predators (Orth et al. 1984). However, this also implies that habitat loss may be an important factor in the decline of *M. arenaria* and *T. plebeius*, as both seagrass and oyster reef habitats are declining in Chesapeake Bay (Orth et al. 1984, 2006, Rothschild et al. 1994, Beck et al. 2011, Lefcheck et al. 2017).

In agreement with our hypotheses, densities of *M. arenaria* were negatively associated with increasing temperature. *Mya arenaria* is a cold-water species that is distributed from the sub-arctic regions to North Carolina along Atlantic coasts of North America (Abraham & Dillon 1986, Maximovich & Guerassimova 2003). Typically *M. arenaria* survives well in temperatures from 2 to 28 °C (Cohen 2005), with mortality usually occurring above 30 °C (Kennedy & Mihursky 1971). It is expected that with global climate change, Chesapeake Bay may become inhospitable for this species. The upper tolerance range for *M. arenaria* is frequently surpassed in the summer in Chesapeake Bay, especially in shallow waters where this sampling effort took place. High summer temperatures in particular are likely a factor in the decline of *M. arenaria* in lower Chesapeake Bay.

Bivalve presence/absence and density were not negatively associated with ray, blue crab, or fish densities for either species, in contrast to our hypothesis, and horseshoe crabs were not

abundant enough to determine their relative contribution to bivalve mortality. However, the snapshot in time provided by our predator and bivalve sampling may not reflect the legacy of predator-prey interactions experienced by local clam populations (Thrush et al. 1997, Seitz & Lipcius 2001, Seitz et al. 2016). Our seasonal time series for *M. arenaria* abundance in particular seem to indicate that predation is an important factor in determining density. In both the upper and lower Bay, densities of *M. arenaria* decline throughout the summer months when predation peaks (Hines et al. 1990). Observed temporal trends also correspond with *M. arenaria* reproductive behavior in Chesapeake Bay, where the fall spawn is more successful than the spring spawn, due to decreased predation pressure in the winter months (Blundon & Kennedy 1982a, Baker & Mann 1991). Individuals spawned in the fall are able to settle and grow throughout the winter, when risk of predation is minimal, and this new generation manifests as a springtime spike in density. However, even the clams that survive until the spring are almost completely consumed by predators each year, essentially resulting in an “annual crop” rather than a stable population with a sustainable age distribution. The declines in *M. arenaria* in summer may be associated with increasing temperature as well as predation, and we have insufficient evidence from our snapshot in time to distinguish between effects of the two. However, this effect persisted in both the upper and lower Bay, and under different temperature regimes, indicating that predation is likely a major driver of observed trends in *M. arenaria* density.

Despite occupying a similar niche in Chesapeake Bay, *M. arenaria* and *T. plebeius* exhibited different seasonal trends in densities. *Tagelus plebeius* spawns in the spring (Holland & Dean 1977, da Silva et al. 2015), and while a spring high-density event was noted in the upper Bay in 2012, the lower Bay had relatively steady densities that did not show a clear recruitment

signal or summer crash. *Tagelus plebeius* in the lower Bay likely do not exhibit the same seasonal crashes in abundance observed for *M. arenaria* because a robust adult *T. plebeius* population remains in lower Chesapeake Bay throughout the seasons. This population allows for high densities of many different size classes to exist at any given time. However, in the upper Bay, where densities are much lower, seasonal spring recruitment was evident in time series, indicating very different population dynamics in these two regions of Chesapeake Bay. *Tagelus plebeius* in the upper Bay seem to exhibit annual population fluctuations more similar to those of *M. arenaria*. This may be because of density-dependent mortality sources specific to the upper Bay, including fishing pressure and disease, which work to keep upper Bay *T. plebeius* at low densities that are sensitive to annual recruitment and predation.

Predation and low fecundity likely drive patterns in recruitment of *M. arenaria* in Chesapeake Bay, which is characterized by very low recruitment with some pockets of locally high recruitment, though studies examining recruitment in more locations would be necessary to identify factors contributing to low recruitment. Although there are some cases of locally high recruitment of *M. arenaria* in Virginia (Lovall et al. 2017), recruitment in the Rhode River, Maryland, declined drastically since the 1980s and now is essentially zero. Predation has been identified as the most important limiting factor in recruitment of *M. arenaria* in other regions (Beal et al. 2001, Hunt & Mullineaux 2002, Beal 2006), and likely plays an important role in Chesapeake Bay as well. In addition, low adult densities and small body sizes may limit recruitment, as fecundity increases exponentially with clam size (Brousseau 1978, Brousseau & Baglivo 1988).

In agreement with our hypothesis, *Perkinsus* sp. infection intensity and prevalence in both species was positively associated with high temperatures, conditions that were considered

stressful for *M. arenaria* in particular. Impacts of dissolved oxygen on *Perkinsus* sp. infection were less clear, though dissolved oxygen was generally within the tolerable range for *M. arenaria* and *T. plebeius* ($> 1.5 \text{ mg L}^{-1}$; Rosenberg et al. 1991, Taylor & Eggleston 2000), and may not have been extreme enough to induce a stress response during the course of the survey. Disease intensity and prevalence in *M. arenaria* was positively correlated with salinity, and long-term intensities and prevalences of *Perkinsus* sp. infections in *M. arenaria* were negatively associated with Susquehanna River discharge, a proxy of salinity. Salinity preference of *Perkinsus* sp. may explain the long-term correlation between infection intensities/prevalences and river discharge/salinity. *Perkinsus* sp. survives and proliferates best at moderate to high salinities, depending on the species. At 28°C , *Perkinsus chesapeaki* survival and proliferation is optimal at moderate salinities 15-25‰ (La Peyre et al. 2006) or 14 ‰ (McLaughlin et al. 2000). In oysters, infection intensity and prevalence of a similar protist, *Perkinsus marinus*, were both greater in higher salinity waters (Burrenson & Calvo 1996). Thus, it may be expected that *Perkinsus* sp. infection in *M. arenaria* would be greatest in high-salinity (low flow) years and locations.

It is possible that disease dynamics do not drive trends in *M. arenaria* and *T. plebeius* density in space and time, as intensities of infections were generally low for both species. In addition, infection intensity was associated with clam size (proxy of age), but not with clam densities. Previous studies have also failed to find an association between *P. chesapeaki* infection and mortality (Bushek et al. 2008). However, this study was unable to disentangle the impact of disease on mortality (which would manifest as a negative relationship between bivalve density and disease) from density-dependent disease dynamics (which would manifest as a positive relationship between bivalve density and disease). To our knowledge, this study

documents the most complete record of *Perkinsus* sp. infection in *M. arenaria* and *T. plebeius* to date, but impacts of disease mortalities on depleted clam populations of Chesapeake Bay remain to be carefully evaluated by rigorous experimental investigations of disease pathogenesis and outcomes.

Several major storms occurred during the study period, including Hurricane Sandy, which made landfall in New Jersey on October 29, 2012 and caused major flooding throughout Chesapeake Bay (Kunz et al. 2013), and Hurricane Lee in 2011, which was a major inflow event (Hirsch 2012). In contrast to Tropical Storm Agnes, these storms produced no noticeable mortality in our survey of *M. arenaria* and *T. plebeius* in the Chesapeake Bay. Although some of these storms caused extensive loss of human life and damage to infrastructure (Smith & Katz 2013), benthic ecosystems may be largely resistant to such disturbances. However, extreme storms have caused mass mortality of bivalves (Lohead et al. 2012, Freeman et al. 2013), and it is possible that these storms did not produce environmental conditions severe enough to produce a response in the bivalves surveyed as part of this study.

The lack of information on *T. plebeius* long-term abundances is an impediment to understanding the decline of this species and the consequences of this decline. Without landings records or fisheries-independent indices of *T. plebeius* abundance, it is difficult to assess the health of the population. In addition, basic population biology information for *T. plebeius* populations in the coastal U.S. is often lacking. Investigations of some populations of *T. plebeius* in Argentina (Lomovasky et al. 2016) and Brazil (da Silva et al. 2015) have established valuable baselines for the species. The ecological and commercial importance of this species, as well as its role harboring *Perkinsus* sp., warrant future studies on growth, reproduction, tolerance to environmental conditions, disease-related mortality, and distribution within U.S. waters.

Conclusions

It is likely that habitat loss, low recruitment, and predators are major factors keeping *Mya arenaria* and *Tagelus plebeius* at low densities in Chesapeake Bay, though continued harvest, which was not addressed in this study, may play a role as well. We found no evidence suggesting densities of *M. arenaria* or *T. plebeius* are negatively impacted by *Perkinsus* sp. infection prevalence or intensity, though the direct effect of *Perkinsus* sp. infection on bivalve mortality remains to be addressed. Extremely low densities of *M. arenaria*, decimated after tropical storm Agnes in 1972 (Cory & Redding 1976, Haven et al. 1976), and a susceptibility of this thin-shelled species to predation by blue crabs, likely fuel a feedback loop that leads to high per-capita rates of predation, which works to keep *M. arenaria* populations at low levels. In recent years, similar dynamics may have been at work in the upper Chesapeake Bay, where populations of *T. plebeius* have reached low densities through fishing pressure or possibly disease. Since *M. arenaria* and *T. plebeius* are preferred prey items for major predators such as *Callinectes sapidus* and *Rhinoptera bonasus* (Blundon and Kennedy, 1982b; Fisher, 2010), it is unlikely that predator switching will provide much relief. However, both species appear able to take advantage of refuge provided by complex habitats. Habitats such as seagrass and shell hash allow both bivalve species to persist at a low-density refuge, which may be stable. Further work should focus on predator-prey interactions, elucidating the existence and stability of low-density refuge habitats, the effect of population declines on ecosystem services such as filtration, and the likelihood that thin-shelled bivalve species may persist, recover, and sustain commercial fisheries in Chesapeake Bay.

ACKNOWLEDGMENTS

We gratefully acknowledge the assistance given by the students and staff of the Community Ecology and Marine Conservation Biology labs at the Virginia Institute of Marine Science, many Maryland Department of Natural Resources staff members who collected and analyzed clams for abundances and diseases during 2000-2009, and the Fish and Invertebrate Ecology Lab at the Smithsonian Environmental Research Center. This material is based upon work supported by the National Oceanic and Atmospheric Administration grant numbers NA11NMF4570218 and NA07NMF4570326. C. Glaspie was supported by an Environmental Protection Agency (EPA) Science to Achieve Results (STAR) Fellowship, grant number FP91767501; and the National Science Foundation Administration GK-12 program, grant number DGE-0840804. M. Ogburn was supported by a Postdoctoral Fellowship awarded by the Smithsonian Environmental Research Center. This work is based on a PhD dissertation of C. Glaspie, supervised by R. Seitz. This paper is contribution number XXXX from the Virginia Institute of Marine Science, College of William & Mary.

LITERATURE CITED

- Abraham BJ, Dillon PL (1986) Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Mid-Atlantic)- softshell clam. U.S. Fish and Wildlife Service Biological Report 82(11.68)
- Abrahao JR, Cardoso RS, Yokoyama LQ, Amaral ACZ (2010) Population biology and secondary production of the stout razor clam *Tagelus plebeius* (Bivalvia, Solecurtidae) on a sandflat in southeastern Brazil. *Zoologia* 27:54–64
- Baker PK, Mann R (1991) Soft shell clam. In: Funderburk S, Mihursky JA, Jordan SJ, Riley D (eds) Habitat Requirements for Chesapeake Bay Living Resources. U.S. Fish and Wildlife Service, Annapolis, MD
- Beal BF (2006) Relative importance of predation and intraspecific competition in regulating growth and survival of juveniles of the soft-shell clam, *Mya arenaria* L., at several spatial scales. *J Exp Mar Bio Ecol* 336:1–17
- Beal BF, Parker MR, Vencile KW (2001) Seasonal effects of intraspecific density and predator exclusion along a shore-level gradient on survival and growth of juveniles of the soft-shell clam, *Mya arenaria* L., in Maine, USA. *J Exp Mar Bio Ecol* 264:133–169
- Beck MW, Brumbaugh RD, Airoidi L, Carranza A, Coen LD, Crawford C, Defeo O, Edgar GJ, Hancock B, Kay MC, Lenihan HS, Luckenbach MW, Toropova CL, Zhang G, Guo X (2011) Oyster reefs at risk and recommendations for conservation, restoration, and management. *Bioscience* 61:107–116
- Berry WD, Feldman S (1985) Multiple Regression in Practice (Quantitative Applications in the Social Sciences). Sage, London

- Beukema JJ, Dekker R (2005) Decline of recruitment success in cockles and other bivalves in the Wadden Sea: Possible role of climate change, predation on postlarvae and fisheries. *Mar Ecol Prog Ser* 287:149–167
- Blaylock RA (1993) Distribution and abundance of the cownose ray, *Rhinoptera bonasus*, in lower Chesapeake Bay. *Estuaries* 16:255–263
- Blundon JA, Kennedy VS (1982a) Refuges for infaunal bivalves from blue crab, *Callinectes sapidus* (Rathbun), predation in Chesapeake Bay. *J Exp Mar Bio Ecol* 65:67–81
- Blundon JA, Kennedy VS (1982b) Mechanical and behavioral aspects of blue crab, *Callinectes sapidus* (Rathbun), predation on Chesapeake Bay bivalves. *J Exp Mar Bio Ecol* 65:47–65
- Botton ML (1984) The importance of predation by horseshoe crabs, *Limulus polyphemus*, to an intertidal sand flat community. *J Mar Res* 42:139–161
- Bowen JE, Hunt HL (2009) Settlement and recruitment patterns of the soft-shell clam, *Mya arenaria*, on the Northern Shore of the Bay of Fundy, Canada. *Estuaries and Coasts* 32:758–772
- Brousseau DJ (1978) Population dynamics of the soft-shell clam *Mya arenaria*. *Mar Biol* 50:63–71
- Brousseau DJ (2005) Effects of natural mortality and harvesting on inshore bivalve population trends. In: Buchsbaum R, Pederson J, Robinson WE (eds) *The Decline of Fisheries Resources in New England: Evaluating the Impact of Overfishing, Contamination, and Habitat Degradation*. MIT Sea Grant, Cambridge, MA
- Brousseau DJ, Baglivo JA (1988) Life tables for two field populations of soft-shell clam, *Mya arenaria*, (Mollusca: Pelecypoda) from Long Island Sound. *Fish Bull* 86:567–580
- Burreson EM, Calvo LMR (1996) Epizootiology of *Perkinsus marinus* disease of oysters in

644 Chesapeake Bay, with emphasis on data since 1985. J Shellfish Res 15:17–34

645 Bushek D, Landau B, Scarpa E (2008) *Perkinsus chesapeaki* in stout razor clams *Tagelus*

646 *plebeius* from Delaware Bay. Dis Aquat Organ 78:243–247

647 Clogg CC, Petkova E, Haritou A (1995) Statistical methods for comparing regression

648 coefficients between models. Am J Sociol 100:1261–1293

649 Cohen AN (2005) Guide to the Exotic Species of San Francisco Bay. San Francisco Estuary

650 Institute, Oakland, CA

651 Cory RL, Redding MJ (1976) Mortalities caused by Tropical Storm Agnes to clams and oysters

652 in the Rhode River area of Chesapeake Bay. In: Anderson AM (ed) The Effects of Tropical

653 Storm Agnes on the Chesapeake Bay Estuarine System. The Johns Hopkins University

654 Press, Baltimore, MD, p 478–487

655 Cressie NAC (1993) Statistics for Spatial Data. Wiley, New York, NY

656 da Silva CF, Corte GN, Yokoyama LQ, Abrahão JR, Amaral ACZ (2015) Growth, mortality, and

657 reproduction of *Tagelus plebeius* (Bivalvia: Solecurtidae) in Southeast Brazil. Helgol Mar

658 Res 69:1–12

659 Dungan CF, Bushek D (2015) Development and applications of Ray’s fluid thioglycollate media

660 for detection and manipulation of *Perkinsus* spp. pathogens of marine molluscs. J Invertebr

661 Pathol 131:68–82

662 Dungan CF, Hamilton RM, Hudson KL, McCollough CB, Reece KS (2002) Two epizootic

663 diseases in Chesapeake Bay commercial clams, *Mya arenaria* and *Tagelus plebeius*. Dis

664 Aquat Organ 50:67–78

665 Efron B, Tibshirani RJ (1993) An Introduction to the Bootstrap. Chapman & Hall, New York,

666 NY

- 667 Eggleston DB, Lipcius RN, Hines AH (1992) Density-dependent predation by blue crabs upon
 668 infaunal clam species with contrasting distribution and abundance patterns. *Mar Ecol Prog*
 669 *Ser* 85:55–68
- 670 Encomio VG, Stickler SM, Allen SK, Chu F-L (2005) Performance of “natural dermo-resistant”
 671 oyster stocks — survival, disease, growth, condition and energy reserves. *J Shellfish Res*
 672 24:143–155
- 673 Farley CA, Plutschak DL, Scott RF (1991) Epizootiology and distribution of transmissible
 674 sarcoma in Maryland softshell clams, *Mya arenaria*, 1984-1988. *Environ Health Perspect*
 675 90:35–41
- 676 Fisher RA (2010) Life history, trophic ecology, and prey handling by cownose ray, *Rhinoptera*
 677 *bonasus*, from Chesapeake Bay. Gloucester Point, VA
- 678 Fisher RA, Call GC, Grubs RD (2011) Cownose ray (*Rhinoptera bonasus*) predation relative to
 679 bivalve ontogeny. *J Shellfish Res* 30:187–196
- 680 Forster S, Zettler ML (2004) The capacity of the filter-feeding bivalve *Mya arenaria* L. to affect
 681 water transport in sandy beds. *Mar Biol* 144:1183–1189
- 682 Freeman RL, Dattilo BF, Morse A, Blair M, Felton S, Pojeta J (2013) The “Curse of
 683 Rafinesquina:” Negative taphonomic feedback exerted by strophomenid shells on storm-
 684 buried lingulids in the Cincinnati Series. *Palaios* 28:359–372
- 685 Gedan KB, Kellogg L, Breitburg DL (2014) Accounting for multiple foundation species in
 686 oyster reef restoration benefits. *Restor Ecol* 22:517–524
- 687 Glaspie CN (2017) *Mya arenaria* and *Mercenaria mercenaria* predator-prey dynamics. *Knowl*
 688 *Netw Biocomplexity*
- 689 Goeij P de, Luttikhuisen PC, Meer J van der, Piersma T (2001) Facilitation on an intertidal

mudflat: The effect of siphon nipping by flatfish on burying depth of the bivalve *Macoma*
balthica. *Oecologia* 126:500–506

Haven DS (1970) A study of the hard and soft clam resources of Virginia: Annual contract report
for the period 1 July 1969 through 30 June 1970. Gloucester Point, VA

Haven DS, Hargis WJ, Loesch JG, Whitcomb JP (1976) The effect of Tropical Storm Agnes on
oysters, hard clams, soft clams, and oyster drills in Virginia. In: Anderson AM (ed) The
Effects of Tropical Storm Agnes on the Chesapeake Bay Estuarine System. The Johns
Hopkins University Press, Baltimore, MD, p 488–508

Heiss WM, Smith AM, Probert PK (2010) Influence of the small intertidal seagrass *Zostera*
novazelandica on linear water flow and sediment texture. *New Zeal J Mar Freshw Res*
34:689–694

Hines A, Haddon A, Wiechert L (1990) Guild structure and foraging impact of blue crabs and
epibenthic fish in a sub-estuary of Chesapeake Bay. *Mar Ecol Prog Ser* 67:105–126

Hines AH, Whitlatch RB, Thrush SF, Hewitt JE, Cummings VJ, Dayton PK, Legendre P (1997)
Nonlinear foraging response of a large marine predator to benthic prey: Eagle ray pits and
bivalves in a New Zealand sandflat. *J Exp Mar Bio Ecol* 216:191–210

Hirsch RM (2012) Flux of nitrogen, phosphorus, and suspended sediment from the Susquehanna
River Basin to the Chesapeake Bay during Tropical Storm Lee, September 2011, as an
indicator of the effects of reservoir sedimentation on water quality.

Holland AF, Dean JM (1977) The biology of the stout razor clam *Tagelus plebeius*: II. Some
aspects of the population dynamics. *Chesap Sci* 18:188–196

Holland AF, Shaughnessy AT, Hiegel MH (1987) Long-term variation in mesohaline
Chesapeake Bay macrobenthos spatial and temporal patterns. *Estuaries* 10:227–245

713 Homer ML, Dungan CF, Tarnowski ML (2011) Assessment of Chesapeake Bay commercial
714 softshell clams *Mya arenaria* and *Tagelus plebeius*. Report to NOAA Chesapeake Bay
715 Fisheries Science Program, Award NA07NMF4570326

716 Homer M, Mihursky JA, Jones P (1980) Quantitative approaches towards characterizing
717 estuarine fish populations and communities. Am Fish Soc Potomac Chapter Proc Fourth
718 Annu Meet:39–99

719 Hunt HL, Mullineaux LS (2002) The roles of predation and postlarval transport in recruitment of
720 the soft shell clam (*Mya arenaria*). Limnol Oceanogr 47:151–164

721 Hyer P V., Ruzecki EP (1976) Changes in salinity structure of the James, York and
722 Rappahannock estuaries resulting from the effects of Tropical Storm Agnes. In: Ruzecki EP
723 (ed) The Effects of Tropical Storm Agnes on the Chesapeake Bay Estuarine System. The
724 Johns Hopkins University Press, Baltimore, MD, p 66–80

725 Hyndman RJ, Khandakar Y (2008) Automatic time series forecasting: The forecast package for
726 R. J Stat Softw 26

727 Kemp WM, Batleson R, Bergstrom P, Carter V, Gallegos CL, Hunley W, Karrh L, Koch EW,
728 Landwehr JM, Moore KA, Murray L, Naylor M, Rybicki NB, Court Stevenson J, Wilcox
729 DJ (2004) Habitat requirements for submerged aquatic vegetation in Chesapeake Bay:
730 Water quality, light regime, and physical-chemical factors. Estuaries 27:363–377

731 Kemp WM, Boynton WR, Adolf JE, Boesch DF, Boicourt WC, Brush G, Cornwell JC, Fisher
732 TR, Glibert PM, Hagy JD, Harding LW, Houde ED, Kimmel DG, Miller WD, Newell RIE,
733 Roman MR, Smith EM, Stevenson JC (2005) Eutrophication of Chesapeake Bay: Historical
734 trends and ecological interactions. Mar Ecol Prog Ser 303:1–29

735 Kennedy VS, Mihursky JA (1971) Upper temperature tolerances of some estuarine bivalves.

736 Chesap Sci 12:193–204

737 Kohn R, Schimek MG, Smith M (2002) Spline and kernel regression for dependent data. In:

738 Schimek MG (ed) Smoothing and Regression: Approaches, Computation and Application.

739 John Wiley & Sons, Inc.

740 Kunz M, Mühr B, Kunz-Plapp T, Daniell JE, Khazai B, Wenzel F, Vannieuwenhuyse M, Comes

741 T, Elmer F, Schröter K, Fohringer J, Münzberg T, Lucas C, Zschau J (2013) Investigation

742 of superstorm Sandy 2012 in a multi-disciplinary approach. Nat Hazards Earth Syst Sci

743 13:2579–2598

744 Lee WJ (2010) Intensive use of an intertidal mudflat by foraging adult American horseshoe crabs

745 *Limulus polyphemus* in the Great Bay estuary, New Hampshire. Curr Zool 56:611–617

746 Lefcheck JS, Wilcox DJ, Murphy RR, Marion SR, Orth RJ (2017) Multiple stressors threaten the

747 imperiled coastal foundation species eelgrass (*Zostera marina*) in Chesapeake Bay, USA.

748 Glob Chang Biol:1–11

749 Lohead J, Gillespie D, Hand C (2012) Storm-Induced anastrophic burial of the Pacific Geoduck

750 (*Panopea generosa*) on the West Coast of Vancouver Island. J Shellfish Res 31:959–967

751 Lomovasky BJ, Brey T, Klügel A, Iribarne O (2016) Distribution pattern, density and growth of

752 the stout razor clam *Tagelus plebeius* in a South-west Atlantic estuarine system. J Mar Biol

753 Assoc United Kingdom

754 Lovall CD, Seitz RD, Knick KE (2017) Direct and indirect impacts of shoreline development on

755 shallow-water benthic communities in a depauperate estuarine system. Bull Mar Sci

756 93:715–741

757 Mann R, Powell EN (2007) Why oyster restoration goals in the Chesapeake Bay are not and

758 probably cannot be achieved. J Shellfish Res 26:905–917

- Maximovich N V., Guerassimova A V. (2003) Life history characteristics of the clam *Mya arenaria* in the White Sea. Helgol Mar Res 57:91–99
- McLaughlin SM, Elsayed EE, Faisal M (2000) Analysis of extracellular proteins of two *Perkinsus* spp. isolated from the softshell clam *Mya arenaria* in vitro. Comp Biochem Physiol Part B Biochem Mol Biol 126:587–598
- Moore KA, Jarvis JC (2008) Eelgrass diebacks in the lower Chesapeake Bay: Implications for long-term persistence. J Coast Res 55:135–147
- Najjar RG, Walker HA, Anderson PJ, Barron EJ, Bord RJ, Gibson JR, Kennedy VS, Knight CG, Megonigal JP, O'Connor RE, Polsky CD, Psuty NP, Richards BA, Sorenson LG, Steele EM, Swanson RS (2000) The potential impacts of climate change on the mid-Atlantic coastal region. Clim Res 14:219–233
- National Marine Fisheries Service (2016) Fisheries of the United States 2015.
- Newell CL (1982) The effects of sediment type on growth rate and shell allometry in the soft shelled clam *Mya arenaria* L. J Exp Mar Bio Ecol 65:285–295
- Olafsson EB, Peterson CH, Ambrose WG (1994) Does recruitment limitation structure populations and communities of macro-invertebrates in marine soft sediments: The relative significance of pre- and post-settlement processes. Oceanogr Mar Biol Annu Rev 32:65–109
- Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, Hughes AR, Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M, Williams SL (2006) A global crisis for seagrass ecosystems. Bioscience 56:987
- Orth RJ, Heck, Jr. KL, Montfrans J van (1984) Faunal communities in seagrass beds: A review of the influence of plant structure and prey characteristics on predator-prey relationships.

782 Estuaries 7:339–350

783 Orth RJ, Marion SR, Moore KA, Wilcox DJ (2010) Eelgrass (*Zostera marina* L.) in the

784 Chesapeake Bay Region of Mid-Atlantic Coast of the USA: Challenges in conservation and

785 restoration. *Estuaries and Coasts* 33:139–150

786 Orth RJ, McGlathery KJ (2012) Eelgrass recovery in the coastal bays of the Virginia Coast

787 Reserve, USA. *Mar Ecol Prog Ser* 448:173–176

788 Peterson CH, Summerson HC, Duncan PB (1984) The influence of seagrass cover on population

789 structure and individual growth rate of a suspension-feeding bivalve, *Mercenaria*

790 *mercenaria*. *J Mar Res* 42:123–138

791 Peyre M La, Casas S, Peyre J La (2006) Salinity effects on viability, metabolic activity and

792 proliferation of three *Perkinsus* species. *Dis Aquat Organ* 71:59–74

793 R Core Team (2017) R: A language and environment for statistical computing.

794 Ralph GM (2014) Quantification of nursery habitats for blue crabs in Chesapeake Bay. Virginia

795 Institute of Marine Science, College of William and Mary

796 Ray SM (1954) Biological studies of *Dermocystidium marinum*, a fungus parasite of oysters. Rice

797 University

798 Ray SM (1966) A review of the culture method for detecting *Dermocystidium marinum*, with

799 suggested modifications and precautions. *Proc Natl Shellfish Assoc* 54:55–69

800 Reece KS, Dungan CF, Burreson EM (2008) Molecular epizootiology of *Perkinsus marinus* and

801 *P. chesapeaki* infections among wild oysters and clams in Chesapeake Bay, USA. *Dis*

802 *Aquat Organ* 82:237–248

803 Rosenberg R, Hellman B, Johansson B (1991) Hypoxia tolerance of marine benthic fauna. *Mar*

804 *Ecol Prog Ser* 79:127–131

805 Rothschild BJ, Ault JS, Goulletquer P, Héral M (1994) Decline of the Chesapeake Bay oyster
806 population: A century of habitat destruction and overfishing. *Mar Ecol Prog Ser* 111:29–39

807 Sanford LP (1990) Covariability of dissolved oxygen with physical processes in the summertime
808 Chesapeake Bay. *J Mar Res* 48:567–590

809 Schubel JR (1976) Effects of Agnes on the suspended sediment on the Chesapeake Bay and
810 contiguous shelf waters. In: Schubel JR (ed) *The Effects of Tropical Storm Agnes on the*
811 *Chesapeake Bay Estuarine System*. The Johns Hopkins University Press, Baltimore, MD, p
812 179–200

813 Schubel JR, Carter HH, Cronin WB (1976) Effects of Agnes on the distribution of salinity along
814 the main axis of the Bay and in contiguous shelf waters. In: Ruzicki EP (ed) *The Effects of*
815 *Tropical Storm Agnes on the Chesapeake Bay Estuarine System*. The Johns Hopkins
816 University Press, Baltimore, MD, p 33–65

817 Seitz R, Lipcius R (2001) Variation in top-down and bottom-up control of marine bivalves at
818 differing spatial scales. *ICES J Mar Sci* 58:689–699

819 Seitz R, Lipcius R, Hines A (2017) Consumer versus resource control and the importance of
820 habitat heterogeneity for marine bivalves. *Oikos* 126:121–135

821 Seitz RD, Lipcius RN, Hines AH, Eggleston DB (2001) Density-dependent predation, habitat
822 variation, and the persistence of marine bivalve prey. *Ecology* 82:2435–2451

823 Seitz RD, Lipcius RN, Knick KE, Seebo MS, Long WC, Brylawski BJ, Smith A (2008) Stock
824 enhancement and carrying capacity of blue crab nursery habitats in Chesapeake Bay. *Rev*
825 *Fish Sci* 16:329–337

826 Seitz RD, Lipcius RN, Seebo MS (2005) Food availability and growth of the blue crab in
827 seagrass and unvegetated nurseries of Chesapeake Bay. *J Exp Mar Bio Ecol* 319:57–68

828 Skilleter G (1994) Refuges from predation and the persistence of estuarine clam populations.
829 Mar Ecol Prog Ser 109:29–42

830 Smith AB, Katz RW (2013) US billion-dollar weather and climate disasters: Data sources,
831 trends, accuracy and biases. Nat Hazards 67:387–410

832 Smith GF, Roach EB, Bruce DG (2003) The location, composition, and origin of oyster bars in
833 mesohaline Chesapeake Bay. Estuar Coast Shelf Sci 56:391–409

834 Sponaugle S, Lawton P (1990) Portunid crab predation on juvenile hard clams: Effects of
835 substrate type and prey density. Mar Ecol Prog Ser 67:43–53

836 Taylor DL, Eggleston DB (2000) Effects of hypoxia on an estuarine predator-prey interaction:
837 Foraging behavior and mutual interference in the blue crab *Callinectes sapidus* and the
838 infaunal clam prey *Mya arenaria*. Mar Ecol Prog Ser 196:221–237

839 Tettelbach ST, Peterson BJ, Carroll JM, Furman BT, Hughes SWT, Havelin J, Europe JR, Bonal
840 DM, Weinstock AJ, Smith CF (2015) Aspiring to an altered stable state: Rebuilding of bay
841 scallop populations and fisheries following intensive restoration. Mar Ecol Prog Ser
842 529:121–136

843 Thiet RK, Smith SM, Rubino V, Clark R, Lee K, Oset J (2014) Soft shell clams (*Mya arenaria*)
844 contribute to macroalgal blooms in a partially-restored Gulf of Maine back-barrier lagoon.
845 Ecol Restor 32:46–58

846 Thrush SF, Cummings VJ, Dayton PK, Ford R, Grant J, Hewitt JE, Hines AH, Lawrie SM,
847 Pridmore RD, Legendre P, McArdle BH, Schneider DC, Turner SJ, Whitlatch RB,
848 Wilkinson MR (1997) Matching the outcome of small-scale density manipulation
849 experiments with larger scale patterns an example of bivalve adult/juvenile interactions. J
850 Exp Mar Bio Ecol 216:153–169

Wilberg MJ, Livings ME, Barkman JS, Morris BT, Robinson JM (2011) Overfishing, disease, habitat loss, and potential extirpation of oysters in upper Chesapeake Bay. *Mar Ecol Prog Ser* 436:131–144

Wood SN (2017) *Generalized Additive Models: An Introduction with R*, 2nd Editio. Chapman & Hall

Wright LD, Prior DB, Hobbs CH, Byrne RJ, Boon JD, Schaffner LC, Green MO (1987) Spatial variability of bottom types in the lower Chesapeake Bay and adjoining estuaries and inner shelf. *Estuar Coast Shelf Sci* 24:765–784

TABLES

Table 1. Locations and dates of suction sampling in the lower Chesapeake Bay and hydraulic dredge sampling in the Upper Chesapeake Bay. Sites (latitude = Lat. and longitude = Lon.) were each sampled on multiple dates, listed here for each subestuary. Means are shown for temperature (Temp., °C), salinity (Sal.) and dissolved oxygen (DO, mg L⁻¹).

Region	Subestuary	Lat.	Lon.	Temp.	Sal.	DO	Dates
Lower Bay	Lynnhaven River	36.88367	-76.01586	24.25	22.25	7.27	11/9/11,
		36.89569	-76.08216	22.35	22.87	7.77	4/30/12,
		36.89569	-76.10185	22.29	23.21	7.67	7/12/12,
		36.89872	-76.04659	23.55	23.18	7.82	10/4/12,
	Mobjack Bay	37.29100	-76.38210	21.20	19.06	8.99	5/1/13,
		37.31217	-76.40636	21.62	18.82	8.53	7/23/13
		37.34371	-76.41560	21.76	18.84	8.63	11/3/11,
		37.37065	-76.36165	21.54	19.36	8.53	4/20/12,
	York River	37.26266	-76.39778	22.87	20.33	8.21	7/30/12,
		37.28126	-76.58297	22.41	17.40	7.46	10/17/12,
		37.41419	-76.67194	23.17	12.94	8.41	4/30/13,
		37.43980	-76.70376	22.78	12.13	7.08	7/15/13
Upper Bay	Chester River	38.93225	-76.22830	23.8	12.36	9.35	11/2/11,
		38.99975	-76.20897	22.20	11.80	8.60	4/19/12,
		39.04367	-76.18493	22.71	11.71	7.70	7/10/12,
		39.08914	-76.15221	22.35	10.80	6.80	9/13/12,
	Eastern Bay	38.83910	-76.24934	22.97	13.33	8.43	4/24/13,
		38.87986	-76.33017	22.66	13.34	8.14	7/10/13
		38.90133	-76.25754	22.22	13.23	8.28	10/25/11,
		38.91160	-76.25772	23.63	12.94	7.70	4/20/12,
		38.94111	-76.25729	23.65	12.54	7.75	7/26/12,
	Western Shore	38.86741	-76.51770	22.49	11.73	7.06	10/23/12,
		38.86819	-76.50999	21.68	11.21	7.01	6/11/13,
		38.88224	-76.48127	22.04	12.33	6.96	9/18/13
		38.89855	-76.43542	26.80	13.20	8.60	10/18/11,
		38.90902	-76.45731	21.71	11.84	7.11	5/31/12,
		38.93742	-76.43924	24.01	11.76	6.50	7/31/12,
		38.98781	-76.41432	22.78	12.95	7.94	6/18/13,
		38.9808	-76.44611	22.95	13.05	7.77	9/20/13
		39.01121	-76.39606	22.82	12.65	7.58	

898 Table 2. Generalized additive model (GAM) coefficients and standard deviations (in parentheses) for presence/absence (P/A), density,
899 *Perkinsus* sp. infection intensity (average ordinal intensity rank of infected individuals, scale 0-5) and prevalence (Dis. Prevalence) of
900 infection (proportion of individuals infected) models for *Mya arenaria* and *Tagelus plebeius*. Transformations (either log or logit) are
901 indicated next to the variable name and coefficients are not back transformed. Parametric predictor variables were temperature (temp.,
902 °C), salinity (Sal.), dissolved oxygen (DO, mg L⁻¹), ray pit density (rays m⁻²), crab density (crabs m⁻²), fish density (fish m⁻²), and
903 habitat (gravel [GR], mud [MU], sand [SA], shell [SH], and seagrass [included in the intercept, int.]). Disease GAMs also included
904 bivalve length (mm), and bivalve density (m⁻²). Coefficients significant at $\alpha = 0.05$ are bolded. N = number of observations used to fit
905 each model.

	<i>Mya arenaria</i>				<i>Tagelus plebeius</i>			
	logit(P/A)	log(Density)	log(Intensity)	Dis. Prevalence	logit(P/A)	log(Density)	log(Intensity)	Dis. Prevalence
<i>Int.</i>	147.01 (62.41)	6.11 (7.09)	-21.90 (2.09)	-23.01 (2.10)	12.14 (5.45)	-4.56 (4.22)	-4.23 (1.73)	-3.18 (1.70)
<i>Temp.</i>	-4.79 (2.34)	-0.27 (0.29)	0.44 (0.12)	0.60 (0.07)	-0.28 (0.16)	0.01 (0.11)	0.19 (0.07)	0.14 (0.07)
<i>Sal.</i>	-1.27 (0.75)	0.17 (0.17)	0.27 (0.13)	0.48 (0.04)	-0.29 (0.16)	0.31 (0.14)	-0.18 (0.06)	0.06 (0.05)
<i>DO</i>	-0.55 (0.78)	-0.32 (0.16)	0.54 (0.06)	0.19 (0.03)	0.22 (0.17)	-0.04 (0.1)	-0.03 (0.03)	0.10 (0.03)
<i>Rays</i>	-7.94 (15.51)	0.09 (1.76)	NA	NA	1.45 (2.89)	0.63 (1.25)	NA	NA
<i>Crabs</i>	0.34 (0.77)	-0.04 (0.37)	NA	NA	-0.09 (0.18)	-0.01 (0.15)	NA	NA
<i>Fish</i>	-0.37 (0.65)	0.10 (0.17)	NA	NA	-0.12 (0.15)	-0.15 (0.12)	NA	NA
<i>Habitat</i>	SG & SH v.	SG & SH v.	NA	NA	GR: -0.46 (1.32)	GR: 0.15 (0.97)	NA	NA
	MU, SA & GR	MU, SA & GR			MU: -2.46 (1.48)	MU: 1.27 (0.96)	NA	NA
	-5.97 (2.82)	-0.77 (0.38)			SA: -1.37 (0.81)	SA: 1.68 (0.65)	NA	NA
					SH: -2.42 (0.69)	SH: 0.13 (0.59)	NA	NA
<i>Bivalve length</i>	NA	NA	0.06 (0.01)	2.89x10⁻² (4.01x10⁻³)	NA	NA	0.04 (0.01)	-0.02 (0.01)
<i>Bivalve density</i>	NA	NA	8.13x10 ⁻⁵ (2.26x10 ⁻³)	4.11x10 ⁻⁴ (1.62x10 ⁻³)	NA	NA	1.78x10 ⁻³ (3.52x10 ⁻³)	1.04x10 ⁻³ (2.92x10 ⁻³)
<i>N</i>	150	62	102	88	310	179	154	128

917 **FIGURES**

932 Figure 1. (a) *Mya arenaria* abundance ($\pm$ S.E.) and fishery landings (solid line) for the period
933 1951-2015, with inset showing amplification of abundance and landings for 2000-2015.
934 Abundance data (shaded bars) are from Maryland DNR (2001-2008) and SERC (2011-2013)
935 fishery-independent escalator dredge sampling. Landings are for the Maryland and Virginia
936 portions of Chesapeake Bay combined (NMFS Commercial Landings Database). Vertical dashed
937 line represents Tropical Storm Agnes (1972). Data source: NMFS Commercial Landings
938 Database. (b) *Tagelus plebeius* abundance ($\pm$ S.E.) from Maryland DNR (2001-2008) and SERC
939 (2011-2013) fishery-independent escalator dredge sampling. Note differences in y-axis ranges.

Figure 2. Mean density (m^{-2}) of (a) *Mya arenaria* and (b) *Tagelus plebeius* captured in suction and hydraulic dredge samples between fall 2011 and summer 2013. Point size is a linear function of mean density at each site: $\text{Radius} = (0.25 \times \text{Density}) + 1$

Figure 3. Average shell length in lower and upper Chesapeake Bay for (a) *Mya arenaria* and (b) *Tagelus plebeius*, with boxes from the first to third quartile, a horizontal line at the median, and a triangle at the mean. Whiskers extend from the lowest data point that is still within 1.5 inter-quartile range (IQR) of the lower quartile, to the highest data point still within 1.5 IQR of the upper quartile, and data outside this range (outliers) are shown as circle symbols.

Figure 4. Mean densities and standard errors for (a) *Mya arenaria* and (b) *Tagelus plebeius* from fall 2011 through fall 2013 for the lower Bay (black) and upper Bay (white). NA = not available (not sampled). For the lower Bay, N = 36 for all seasons. For the upper Bay, N = 62, 51, 67, 59, NA, 43, 67 for Fall 2001 to Fall 2013, respectively.

Figure 5. Trends in density for (a) *Mya arenaria* and (b) *Tagelus plebeius* in different habitats. Error bars denote standard error. For *M. arenaria*, N = 314 and 251 for simple and complex habitat, respectively. For *T. plebeius*, N = 57, 32, 182, 187, 64 for mud, sand, gravel, shell, and seagrass, respectively.

Figure 6. Mean annual *Mya arenaria* recruitment from fishery-independent core sampling in the Rhode River, Maryland (1980-2016). Recruitment data include only clams ≤ 10 mm shell length. Shown are means of two sampling sites (bars), standard error (error bars) and fit from ARIMA model (solid black line). Years where sampling occurred but no *M. arenaria* were captured are denoted by a “0”; no clams were captured in samples after 2005.

Figure 7. Average *Perkinsus* sp. (a) infection intensity (average ordinal intensity rank of individuals 0-5) for *Mya arenaria*, (b) infection intensity for *Tagelus plebeius*, (c) prevalence of infection (proportion of individuals infected) for *M. arenaria*, and (d) prevalence of infection for *T. plebeius* for the upper Chesapeake Bay from hydraulic dredge samples collected between fall 2011 and fall 2013. Point size is a linear function of mean disease intensity ($\text{Radius} = [2.0 \times \text{Intensity}] + 0.5$) or mean disease prevalence ($\text{Radius} = [2.5 \times \text{Prevalence}] + 0.5$) at each site.

Figure 8. Time series of *Perkinsus* sp. (a) mean annual infection intensity (average ordinal intensity rank) and (b) mean annual prevalence (proportion of individuals infected) for *Mya arenaria* (black) and *Tagelus plebeius* (white) from 2000-2013. Error bars indicate standard error. Also shown is (c) average annual Susquehanna River discharge as a proxy for salinity (from the USGS National Water Information System). NA = not available (not sampled). See supplementary data files for sample sizes.

1001 **SUPPLEMENTARY TABLES**

1002 Supplemental Table 1. Fish species list and total number caught at sites in the lower and upper
 1003 Chesapeake Bay. In the lower Bay, a total of 432 20-m tows were conducted with a modified
 1004 crab scrap. In the upper Bay, a total of 96 500-m tows were conducted with an otter trawl.
 1005

Common Name	Scientific name	Number caught	
		Lower Bay	Upper Bay
White catfish	<i>Ameiurus catus</i>	0	1
Bay Anchovy	<i>Anchoa mitchelli</i>	many	many
American eel	<i>Anguilla rostrata</i>	3	0
Silver perch	<i>Bairdiella chrysoura</i>	518	0
Spadefish	<i>Chaetodipterus faber</i>	10	0
Spotted seatrout	<i>Cynoscion nebulosus</i>	5	0
Weakfish	<i>Cynoscion regalis</i>	3	0
Gizzard shad	<i>Dorosoma cepedianum</i>	0	2
Mummichog	<i>Fundulus heteroclitus</i>	3	0
Skilletfish	<i>Gobiesox strumosus</i>	35	0
Channel catfish	<i>Ictalurus punctatus</i>	0	1
Pinfish	<i>Lagodon rhomboides</i>	71	0
Spot	<i>Leiostomus xanthurus</i>	380	213
Rainwater killifish	<i>Lucania parva</i>	12	0
Atlantic croaker	<i>Micropogonias undulatus</i>	381	0
White perch	<i>Morone americana</i>	0	2095
Striped bass	<i>Morone saxatilis</i>	19	84
Oyster toadfish	<i>Opsanus tau</i>	1	0
Summer flounder	<i>Paralichthys dentatus</i>	77	0
Harvestfish	<i>Peprilus paru</i>	1	0
Yellow perch	<i>Perca flavescens</i>	0	2
Norther puffer	<i>Sphoeroides maculatus</i>	6	0
Tonguefish	<i>Symphurus plagiusa</i>	12	0
Hogchoker	<i>Trinectes maculatus</i>	183	0
Spotted hake	<i>Urophycis regia</i>	5	0

1006

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Relationship between density of (a) *Mya arenaria* and (b) *Tagelus plebeius* in suction samples and hydraulic dredge samples, including equations from simple linear regressions. The inclusion of the high-density data point in regressions did not significantly impact slopes for *M. arenaria* ($z = 1.39$, $p = 0.17$) or *T. plebeius* ($z = 1.36$, $p = 0.18$), as determined by a two-sided hypothesis test of z values calculated from model coefficients and standard errors (Clogg et al. 1995). High-density data points for both species do not come from the same location, and these densities were within the range of expected values for both species; thus, these data points were not omitted from analyses.

1019
1020 Supplementary Figure 2. Size (length) density histograms for *Mya arenaria* captured in (a)
1021 suction samples and (b) hydraulic dredge samples.

1022

Supplementary Figure 3. Pearson correlation coefficients (top panels) and scatter plots with splines (bottom plots) for all environmental and predator variables. Color denotes direction (negative = red, positive = blue) and magnitude (shading) of correlation coefficients.

1028
 1029 Supplemental Figure 4. Lower Chesapeake Bay biomass (ash-free dry weight) of (a) *Mya*
 1030 *arenaria* and (b) *Tagelus plebeius* captured in suction samples between fall 2011 and summer
 1031 2013. Point size is a linear function of mean biomass at each site: $\text{Radius} = (4.0 \times \text{Density}) + 1$.
 1032