

1 **Title**

2 A novel environmental DNA (eDNA) sampling method for aye-ayes from their feeding

3 traces

4

5 **Author list**

6 Megan L. Aylward¹, Alexis P. Sullivan², George H. Perry^{2,3}, Steig E. Johnson¹, Edward E.

7 Louis Jr⁴

8 ¹ Department of Anthropology and Archaeology, University of Calgary, Calgary, Alberta, T2N 1N4, Canada

9 ² Department of Biology, Pennsylvania State University, State College, PA, 16802, USA

10 ³ Department of Anthropology, Pennsylvania State University, State College, PA, 16802, USA,

11 ⁴ Grewcock Center for Conservation and Research, Omaha's Henry Doorly Zoo and Aquarium, Omaha, NE,

12 USA

13 **Correspondence address**

14 Megan Aylward

15 Department of Anthropology and Archaeology, University of Calgary

16 2500 University Drive

17 Calgary, Alberta T2N 1N4

18 mlylwar@ucalgary.ca

19

20

21

22

23

24

25

26

27 **Abstract**

28 Non-invasive sampling is an important development in population genetic monitoring of
29 wild animals. Particularly, the collection of environmental DNA (eDNA) which can be
30 collected without needing to encounter the target animal, facilitates the genetic analysis
31 of cryptic and threatened species. One method that has been applied to these types of
32 sample is target capture and enrichment which overcomes the issue of high proportions
33 of exogenous (non-host) DNA from these lower quality samples. We tested whether
34 target capture of mitochondrial DNA from sampled feeding traces of wild aye-eyes would
35 yield mitochondrial DNA sequences for population genetic monitoring. We sampled
36 gnawed wood from feeding traces where aye-eyes excavate wood-boring insect larvae
37 from trees. We designed RNA probes complementary to the aye-aye's mitochondrial
38 genome and used these to isolate aye-aye DNA from other non-target DNA in these
39 samples. We successfully retrieved six near-complete mitochondrial genomes from two
40 sites within the aye-aye's geographic range that had not been sampled previously. This
41 method can likely be applied to alternative foraged remains to sample species other than
42 aye-eyes. Our method demonstrates the application to next-generation molecular
43 techniques to species of conservation concern.

44 **Key words:** Aye-aye, Endangered, Environmental DNA, Lemur, Madagascar, Mitogenome,
45 Non-invasive sampling, Target capture

46

47 **Introduction**

48 Genetic sampling of wild populations can help us to address questions of demography,
49 individual relatedness, population structure, and other important aspects of biodiversity
50 that cannot be answered by behavioural monitoring alone (Allendorf, Hohenlohe, &

51 Luikart, 2010). However, the practicality of sampling wild animals can be limited by
52 ethical implications due to the implicit risks of sedating animals to collect blood or tissue
53 samples, especially for arboreal species, and for rare or elusive species (including
54 nocturnal animals), by infrequent encounter rates. (Kohn & Wayne, 1997; Goossens,
55 Chikhi, Utami, de Ruiter, & Bruford, 2000). Even when it is possible and safe to collect
56 invasive samples from wild individuals, the number of samples that can be obtained with
57 this approach may be fewer than analytically desirable (Sikes & Gannon, 2011). Under
58 these conditions, non-invasive sampling is a valuable tool in advancing our
59 understanding of wild populations.

60

61 Over the last two decades, advances in molecular biology have facilitated the use of non-
62 invasive sampling for genetic analyses of wild populations (Morin, Wallis, Moore,
63 Chakraborty, & Woodruff, 1993; Joost et al., 2007). Furthermore, traces of DNA left in the
64 environment from biological material (eDNA) can be collected without even
65 encountering target individuals. Source materials for non-invasive or eDNA sampling that
66 have been used in wildlife monitoring and forensic studies include feathers, faeces, egg
67 shells, hair, saliva, urine, and snail trails (Valiere & Taberlet, 2000; Inoue, Inoue-
68 Murayama, Takenaka, & Nishida, 2007; Beja-Pereira, Oliveira, Alves, Schwartz, & Luikart,
69 2009; Sastre, Francino, Sanchez, & Ramirez, 2009). Non-invasive and eDNA sampling are
70 particularly valuable in research surrounding unhabituated populations, cryptic species,
71 and those at risk of extinction (Goossens et al., 2000; Cushman, McKelvey, Hayden, &
72 Schwartz, 2006; Kendall et al., 2009; Cleverger & Sawaya, 2010; Schubert et al., 2011;
73 Morin, Kelly, & Waits, 2016; Orkin, Yang, Yang, Yu, & Jiang, 2016).

74

75 One species that could especially benefit from the advances in non-invasive genetic
76 sampling is the aye-aye (*Daubentonia madagascariensis*), which is a rare and elusive
77 Malagasy primate of major conservation concern (Sterling & McCreless, 2007; Schwitzer
78 et al., 2013). The low population density, large home range size, cryptic nature, and
79 nocturnal activity of aye-ayes make them particularly difficult to locate, and precise
80 distributions and population densities are unclear (Sterling, 1994b). Challenges of
81 monitoring aye-ayes and obtaining reliable population dynamics data are reflected in the
82 volatility of aye-aye's conservation status designations over the last 70 years. In the
83 1950's, the aye-aye was thought to be extinct (Sterling, 1994b). After aye-ayes were
84 rediscovered in 1957, they were classified as Endangered; in 2008, their status was
85 changed to Near Threatened before being reassessed as Endangered in 2012
86 (Andriaholinirina et al., 2015). In addition to being listed as Endangered, aye-ayes are
87 currently considered one of the world's top 25 most endangered primates
88 (Randimbiharirina et al., 2017 in Schwitzer et al. 2016-2018). Few encounters, along
89 with the aye-aye's solitary social organization and long maternal investment suggest low
90 population densities. Low nuclear genomic diversity in aye-ayes reflects these
91 assumptions; genomic analyses estimates of heterozygosity of 0.051%, and genetic
92 diversity across synonymous sites of $\pi = 0.073$, are the lowest of any primate species
93 studied to date (Perry, Melsted, et al., 2012; Perry, Reeves, et al., 2012). Therefore, despite
94 the wide distribution of aye-ayes, there are likely few individuals, increasing the risk of
95 local and global extinction (Schwitzer et al., 2013; Gross, 2017).

96

97 The IUCN's lemur survival strategy recognises the need for biological monitoring of aye-
98 ayes to better assess population status and conserve genetic diversity within this lineage
99 (Schwitzer et al., 2013). In addition to their elusive behaviour making aye-ayes difficult

100 to find and monitor, genetic sampling of individuals once they have been located is
101 challenging; invasive collection of blood and tissue can only be achieved during
102 immobilization, which is risky and must be conducted by trained and experienced
103 personnel (Cunningham, Unwin, & Setchell, 2015). Therefore, to assess genetic diversity
104 and identify priority populations for conservation a new, reliable means of non-invasive
105 sampling in aye-eyes is required.

106

107 Reliable genotyping from non-invasively collected material holds much promise in
108 sampling threatened species. Environmental DNA samples that are degraded due to
109 exposure to both biotic and abiotic factors or contain high proportions of exogenous DNA
110 can now provide reliable genetic markers (Beja-Pereira et al., 2009; Carpenter et al.,
111 2013; Snyder-mackler et al., 2016). Techniques such as high-throughput sequencing
112 technologies reduce sequencing errors and improve genotyping accuracy by providing
113 greater depth of coverage across loci. One particularly promising approach is target
114 capture, which provides a means of isolating endogenous DNA from the high proportions
115 of exogenous DNA in non-invasive samples (Bi, Linderoth, Vanderpool, Good, & Nielsen,
116 2013; Hawkins et al., 2015; Kirillova et al., 2015; Kistler et al., 2015; Mohandesan et al.,
117 2017). Specifically, RNA probes which are complementary to particular regions or
118 markers in the genome of the target organism are specially designed and synthesized
119 (Gnirke et al., 2009). These probes hybridize to the endogenous DNA in the sample. After
120 hybridization, the biotin coating of the probes allows streptavidin-coated magnetic beads
121 to bind; the bound probes and hybridized endogenous DNA can then be isolated from the
122 exogenous DNA by using a magnet (Gnirke et al., 2009; Giolai et al., 2016). These
123 developments make monitoring and sampling of wild populations where individuals are
124 difficult to locate increasingly feasible.

125

126 For primates, the application of target capture from low quality samples has largely been
127 applied to ancient DNA studies, but Perry et al., (2010) and Snyder-Mackler et al., (2016)
128 have presented effective approaches for sampling endogenous DNA from faeces in extant
129 primate species. Recently, Chiou & Bergey , (2018) demonstrated a different method that
130 isolates endogenous DNA from primate faeces by targeting the higher CpG-methylation
131 density of vertebrate taxa relative to the exogenous bacterial and plant DNA in faeces.
132 Although faecal collection is a popular source of non-invasive sampling of primates (Oka
133 & Takenaka, 2001; Quéméré, Crouau-Roy, Rabarivola, Louis, & Chikhi, 2010; Chancellor
134 et al., 2011), it is unlikely to be a feasible method of sampling for aye-ayes. Factors such
135 as aye-aye's nocturnal behaviour, the height that they travel in the canopy, and their large
136 nightly travel distances mean that defecation is difficult to observe and locating and
137 collecting faecal material is problematic (Randimbiharirina et al., 2017); accordingly,
138 at two sites where aye-ayes have been monitored by Madagascar Biodiversity
139 Partnership since 2010, aye-aye faecal samples have only been collected through routine
140 immobilizations.. Therefore, to gain information on the genetics of aye-aye populations
141 to meet the IUCN aims, in this paper, we explore the possibility of sampling eDNA from
142 aye-ayes (Schwitzer et al., 2013).

143

144 One potential source of eDNA in aye-ayes is from their distinct feeding traces left on trees.
145 These traces are associated with their adaptations for extracting the larvae of wood-
146 boring insects from tree trunks and branches, after identifying suitable foraging locations
147 via a process of sniffing, lightly tapping, and listening (Sterling, 1994a; Erickson, 1995).
148 Aye-ayes gnaw into selected areas of trees with their elongated and continuously growing
149 incisors and extract larvae using their thin flexible third digit (Sterling, 1994a). During

150 this foraging process, the buccal cavity of the aye-aye comes into contact with the wood
151 (Sterling, 1994a; Erickson, 1995; Sterling & McCreless, 2007; arkive.org, 2017). Thus,
152 trace amounts of aye-aye biological material in the form of epithelial cells of the buccal
153 mucosa or from saliva may be deposited and accessible as an eDNA source.

154

155 We investigated the application of target capture and enrichment to obtain aye-aye DNA
156 from aye-aye feeding traces. We aimed to determine whether this method is a feasible
157 alternative to invasive sampling. If eDNA samples provide a means of remotely sampling
158 wild aye-aye populations, we predict that (1) target enrichment is an effective method of
159 obtaining aye-aye DNA from the exogenous DNA in feeding traces, and (2) we will be able
160 to obtain full mitochondrial genomes (hereafter 'mitogenomes') for population genomic
161 analysis.

162

163 **Methods**

164 **Sampling sites and techniques**

165 Initial training for researchers and local technicians on accurate identification of aye-aye
166 feeding traces and the sample collection method was conducted through the Kianjavato
167 Ahmanson Field Station (KAFS) located in the Kianjavato commune of the Vatovavy-
168 Fitovinany region in southeast Madagascar during June 2014. KAFS is a long-term field
169 site of the Madagascar Biodiversity Partnership (MBP). Individual aye-ayes present in the
170 region have been fitted with ATS (Applied Telemetry Systems®) VHS radio-collars for
171 behavioural monitoring, which afforded the opportunity to view known feeding sites in
172 an area of bamboo to become familiar with density and frequency of feeding traces, their
173 characteristics and to practice the collection technique (see below).

174

175 To then test the sampling method on unmonitored populations, we selected two study
176 sites near to the limits of the aye-aye's geographic distribution, where aye-ayes have been
177 sighted but not sampled: Manombo Special Reserve in the Atsimo-Atsinanana region in
178 southeastern Madagascar, operated by Madagascar National Parks and at that time
179 supported by Durrell Wildlife Conservation Trust, and the Tsingy de Beanka, Ambinda in
180 the Melaky region in west-central Madagascar and supported by Biodiversity
181 Conservation Madagascar (Figure 1). Between July and December 2014, MA surveyed the
182 forest at each site with the assistance of three local technicians. Prior to sampling, local
183 technicians cleared undergrowth from existing trail systems in the forest to use as
184 transects. During daylight hours, these transects were walked at a slow pace of
185 approximately 1 km/hr to look for feeding traces and searched areas on either side of the
186 transect at 200 m intervals. Traces were observed in both live and dead plants where
187 characteristic holes gnawed by aye-aye were identified. These were confirmed by the
188 distinct shape of feeding traces and typically presence of teeth marks (Figure 2). The
189 research permit (N^o 162/14/MEF/SG/DGF/DCB.SAP/SCB) for sample collection was
190 obtained from the Ministere de l'environnement, des eaux et forets et du tourisme, in
191 Madagascar.

192

193 To collect samples, we selected areas around the edge of the trace where the buccal cavity
194 and teeth of the animal were likely to have come into contact with the tree (Figure 2). MA
195 took 3-6 samples of tree material per trace; each sample was approximately 2 g⁻¹ of wood
196 shavings. MA collected samples using a scalpel to cut wood from the site directly into a
197 1.5 mL Eppendorf[®] tube containing 500 µL RNAlater. To reduce the chance of
198 contamination, MA wore latex gloves and used a new, sterile scalpel blade for each
199 sample. At Manombo, we walked 9.5 km of transects over a 6 km² area. At Beanka, we

200 surveyed 10.8 km of transects covering a total 9 km² area. MA sampled a total of 59 traces
201 across the two sites (Manombo: n=27; Beanka: n=32). We stored samples at ambient
202 temperature until they could be transferred to 4 °C for storage in Antananarivo,
203 Madagascar. MBP exported samples from Madagascar (Export Permit Number 233N-
204 EV07/MG16) and stored the samples at Omaha's Henry Doorly Zoo and Aquarium at 4 °C
205 for four months prior to processing.

206

207 **DNA extraction, library preparation, capture, and sequencing**

208 After samples were removed from storage, we extracted DNA from all samples. We
209 vortexed each sample for 30 seconds in PBS to wash aye-aye DNA from the surface of the
210 wood and then removed pieces of wood using sterile forceps. We added 500 µL of PBS
211 solution to the sample and centrifuged at 10,000 g to pellet the cellular DNA and then
212 carefully removed the supernatant so as not to disturb the precipitate. We performed all
213 DNA extractions on the precipitate using EZNA® blood DNA mini kit following the buccal
214 swab protocol with the following modifications: we lysed samples overnight, and we
215 eluted DNA in 50 µL elution buffer, incubated for 20 minutes and then re-eluted with the
216 same eluate. Preliminary tests to confirm the presence of aye-aye DNA were conducted
217 using a microsatellite marker for loci AYE33 of approximately 279 bp, with the following
218 primer pair: forward 5'-3' GTCTGCTACTCCTTAGGTGCTG and reverse 5'-
219 TGGCTCAGGGCAATACAAT-3'. We amplified the extracted DNA in a 15 µL reaction
220 containing 0.25 µL of each 10 uM primer, 7.5 µL buffer containing 1.5 mM MgCl₂ (1X) and
221 0.2 mM of each dNTP (1X), 0.12 µL KAPA3G plant DNA polymerase, 4.9 µL H₂O, and 2 µL
222 DNA sample. Cycling parameters began with an initial enzyme activation temperature of
223 95 °C for 3 min, followed by 32 cycles of: denaturation at 98°C for 20 s, primer annealing
224 at 58 °C for 15 s and elongation at 72 °C for 30 s, then one cycle of final elongation at 72

225 °C for 1 min. We visually confirmed the presence of the expected 279 bp PCR product by
226 gel electrophoresis with 5 µL of PCR product and using 2 µL of a 1 kb ladder.

227

228 Extracted DNA was transported to Pennsylvania State University, University Park, PA for
229 library preparation and DNA capture. We quantified the DNA concentration and fragment
230 size of ten samples using a bioanalyzer (Agilent Technologies, Santa Clara, CA) at the Huck
231 Institute of the Life Sciences (HILS), University Park, PA, genomics core facility. We
232 selected a subset of 22 samples that were estimated to have been between 14 h-6 months
233 old at the time of collection based on observations during surveys and discolouration of
234 wood. We prepared DNA sequencing libraries from each of these samples using the
235 Illumina TruSeq nano[®] kit. The TruSeq nano[®] library preparation protocol recommends
236 200 ng⁻¹ of input DNA. Due to the typically low quantities of DNA extracted from each
237 individual wood sample (<300 pg/µL), we combined DNA extracted from multiple
238 samples taken from the same trace into one sample per feeding trace to increase total
239 amounts of library preparation input DNA per sample to approximately 5-42 ng⁻¹ DNA
240 (based on bioanalyzer readings for a subset of samples). To account for the low input DNA
241 quantities, we adapted the protocol to use half quantities of all reagents and volumes. We
242 sheared each DNA sample to a mean size of 500 bp with the Covaris-M220[®] using the
243 following settings: 34 seconds, peak incident of 50, duty factor of 20% and 200 cycles per
244 burst. For library preparation wash steps, 180 µL ethanol was used and an additional
245 initial wash step was included to remove residual salts from eluting extracted DNA. We
246 used the Bioanalyzer (Agilent Technologies, Santa Clara, CA) to quantify the final library
247 and confirm a mean library size of 750 bp fragments, as expected for 550 bp libraries
248 once the sequencing adapters and unique barcodes are attached to the 550bp DNA
249 fragments.

250

251 As a preliminary test to check for the presence of aye-aye DNA in these libraries, we
252 shotgun sequenced ten DNA libraries. Libraries were included as part of a larger
253 multiplexed sequencing pool (Table A1) and sequenced using the Illumina NextSeq® 500
254 2x150 pair end chemistry at UCLA Clinical Microarray Core Facility. Samples were
255 demultiplexed at the sequencing facility.

256

257 RNA probes for capture of the aye-aye mitogenome were designed through MyBaits® at
258 Arbor Biosciences™. Baits were designed based on the aye-aye reference mitochondrial
259 DNA genome sequence (GenBank® accession NC_010299.1), bait length was 80 bp at 4x
260 tiling and these are available as a predesigned MyBaits® Mito panel. We conducted
261 captures as per MyBaits® manual v.3 and a single round of captures was conducted on 19
262 of the eDNA (foraging remains) samples. Post capture products were quantified using
263 Bioanalyzer (Agilent Technologies, Santa Clara, CA) at the genomics core facility at the
264 HILS and pooled for sequencing (Table A2). These capture products were sequenced as
265 one pool of 17 samples and two of the samples (BTS38 and BTS108) were included as
266 part of a different sequencing pool. The final mitochondrial capture pool of 17 Mito Bait
267 captured DNA libraries was sequenced on one lane of the NextSeq 500 mid-output 2x150
268 bp pair end chemistry UCLA Clinical Microarray Core Facility. Samples were
269 demultiplexed and barcodes removed at the sequencing facility.

270

271 **Data analysis**

272 All computational analyses were conducted on servers provided by WestGrid
273 (www.westgrid.ca) at Compute Canada (www.computecanada.ca). Raw reads were
274 submitted to the NCBI SRA project under the accession PRJNA434884. The following

275 processing and alignment pipeline was used for the shotgun and captured DNA
276 sequencing pools. We assessed sequence read quality using Fast-QC (Andrews, 2007) and
277 filtered reads based on minimum read length of 120bp and mean quality of 20 using
278 PRINSEQ-lite (Schmieder & Edwards, 2011). We removed any potential human
279 contaminant reads using the bbsplit package of bbmap (Bushnell, 2016); sequencing
280 reads were aligned to both the aye-aye and human genome and any reads which matched
281 better to the human genome than the aye-aye genome were removed from the dataset.
282 Paired reads were aligned to the aye-aye reference genome GCA_000241425.1 (Perry,
283 Reeves, et al., 2012), using the bwa-mem alignment algorithm (Li & Durbin, 2009; Li,
284 2013). We used the MarkDuplicates tool in PicardToolsv.1106 (Broad Institute, 2017) to
285 remove sequencing and PCR duplicates, and reads flagged as clipped were filtered from
286 the dataset. We generated consensus sequences for the eDNA samples with near-
287 complete mitogenomes using samtoolsv.1.3, bcftoolsv.1.3 and vcfUtilsv.1.3 (Li et al.,
288 2009; Danecek et al., 2011; Li, 2011). To confirm the base calls in these consensus
289 sequences, we compared the samtoolsv.1.3 generated sequences to those generated
290 using IGV (Robinson et al., 2012) to ensure the same nucleotide bases were found by each
291 software program.

292

293 We compared the mitogenome sequences from our eDNA samples with the GenBank®
294 aye-aye mitogenome data to confirm that they were novel (Perry et al., 2013; Kistler et
295 al., 2015). We called variants using GATK (McKenna et al., 2010), and used haplotype
296 caller to generate individual gVCFs and then the joint variant caller. To filter for biallelic
297 SNPs only, we used VCFtoolsv.1.12 and set maximum and minimum alleles to two.

298

299 We used PGDSpiderv2.3 to convert SNP variant calls from vcf to ped format (Lischer &
300 Excoffier, 2012). We used the --cluster and --matrix commands in Plinkv.1.7 (Purcell et
301 al., 2007) to calculate the pairwise proportion (identity by state, IBS) of shared alleles at
302 SNP loci among all complete mitogenome sequences. For the Plink analysis, we used the
303 complete mitochondrial genomes excluding the hyper-variable region given the potential
304 for homoplasmy in this region and the possibility for multiple mutations per site (Ballard
305 & Whitlock, 2004; Gonder, Mortensen, Reed, De Sousa, & Tishkoff, 2007).

306

307 **Results**

308 **Pre-capture processing**

309 After library preparation, shotgun sequencing of libraries from eDNA samples did not
310 yield sufficient mitochondrial DNA for mitogenome analysis. Of the 10 libraries that were
311 shotgun sequenced, seven yielded fewer than 10 reads aligning to the aye-aye
312 mitochondrial genome; one sample, with the highest number of mitochondrial reads, was
313 MSR58 with 189 unique reads (0.000274%) (Table 1).

314

315 **Target capture DNA sequencing**

316 Target captures of the mitogenome increased the proportion of on-target mitochondrial
317 DNA reads by several orders of magnitude compared to reads yielded from shotgun
318 sequencing without capture (Table 1). For samples for which it was possible to calculate
319 the fold enrichment (i.e., the samples that were both shotgun sequenced and sequenced
320 following DNA capture), the percent of unique on-target reads increased by over 100
321 times. For captured samples, the percent of on-target reads ranged from 0.009% to 27.5%
322 of the total reads, whereas the percent unique on-target reads ranged from 0.001% to
323 0.05% (Table 2).

324

325 We recovered near-complete mitogenomes for six of the 19 captured samples (31.6%).
326 The depth and breadth of coverage varied between samples (Table 1). For these six
327 mitogenomes, the mean depth of coverage was 16x (range 8-25x). For the 13 samples
328 with less than 90% coverage, the percent of the genome covered ranged from 10-71%
329 (mean 32.58%) with a mean depth of 1.7 (+/-0.3-2.8). Five of these 13 samples had a
330 breadth of coverage between 45-70% and were all identical at SNP loci to the samples
331 with greater breadth of coverage. We only included samples with coverage greater than
332 90% in downstream analysis.

333

334 **Mitogenome analysis**

335 Based on multiple sequence alignment of these seven near-complete mitogenomes, we
336 identified one unique mitogenome sequence from Manombo Special Reserve and two
337 unique mitogenomes from Tsingy de Beanka. Alignment with previously published
338 GenBank® sequences that the mitogenome sequences obtained from the eDNA samples
339 were unique to all previously published aye-aye mitogenomes. For the GenBank
340 sequences sample locations from northern Madagascar, we follow the population names
341 in Perry et al., (2013); North, East, and West (Figure 1). When we compared SNP loci
342 among samples, all mitogenomes clustered into distinct populations according to their
343 collection location (Figure 3). Pairwise comparison of genotypes at polymorphic loci
344 indicate that there are four distinct populations (Figure 3). The Tsingy de Beanka
345 population showed moderate differentiation from the other populations (Figure 3).
346 There was also a split between the East population and Manombo in the southeast, while
347 the North population shared the fewest alleles with all other populations.

348

349 **Discussion**

350 We tested whether sampling of aye-aye feeding traces is a feasible method of non-
351 invasive sampling of this species. Our first prediction, that target capture would provide
352 a means of sampling aye-aye DNA from these feeding traces, was supported by the
353 increased amount of endogenous DNA obtained from target capture compared to shotgun
354 sequencing. The number of on-target reads from the shotgun sequencing was not at
355 sufficient coverage across the mitogenome to identify polymorphic sites and genotype
356 individuals. The high level of enrichment from captured libraries compared to shotgun
357 sequencing indicates the efficacy of a target capture approach (Gnirke et al., 2009). Our
358 second prediction, that mitogenomes could be obtained from these samples and used for
359 population genomic analysis, was also supported. We were able to obtain near complete
360 mitogenomes for 31.6% of the processed samples. Previous studies that have sampled
361 eDNA from foraged material have typically only retrieved partial D-loop fragments rather
362 than whole mitogenomes (Nichols, Konigsson, Danell, & Spong, 2012; Caniglia, Fabbri,
363 Mastrogiuseppe, & Randi, 2013; Wheat, Allen, Miller, Wilmers, & Levi, 2016). While
364 partial fragments can be useful in metabarcoding studies for species detection, whole
365 mitogenomes provide greater phylogenetic resolution and reduce the likelihood of
366 including nuclear copies of mitochondrial DNA(numts) (Johnsen, Kearns, Omland, &
367 Anmarkrud, 2017).

368

369 **Future directions for methodological improvements for sampling aye-aye eDNA**

370 The method developed here is a valuable tool to extend sampling of unmonitored aye-aye
371 populations. Yet, improved efficacy of the method to increase amount of aye-aye DNA
372 obtained and therefore reduce the cost per sample could make it a more attractive
373 application for conservation monitoring (Ekblom & Galindo, 2010). Low proportions of

374 on-target, unique sequence reads recovered are common when working with
375 environmental samples (Ávila-Arcos et al., 2015; Nielsen et al., 2017) and adequate
376 sampling effort is required to account for this issue. A mode of three different samples
377 was taken per feeding trace, and it would be feasible to double this number. Therefore,
378 we recommend more intensive sampling of each feeding trace to allow for multiple
379 library preparations per trace, whilst maintaining sufficient amounts of input DNA for
380 TruSeq nano® library preparation. Multiple library preparations for each trace prior to
381 capture may increase the number of unique DNA fragments within the samples and
382 reduce PCR bias. The proportion of unique reads aligning to the target region was
383 relatively low (0.4-12 %). To better assess the number of PCR cycles to run post-capture
384 to achieve sufficient concentrations for sequencing, qPCR quantification may be valuable
385 to ensure reduced PCR duplication, therefore improved sequencing efficacy and
386 ultimately increased cost effectiveness (Enk, Rouillard, & Poinar, 2013).

387

388 **Application towards addressing the IUCN's objectives for aye-ayes**

389 We demonstrated a novel method that can be applied across the species range for
390 population genomic monitoring. This method helps to meet objective seven of the IUCN's
391 lemur survival plan set out by Schwitzer et al., (2013:page32) which is "Fill knowledge
392 gaps in population ecology and biodiversity of lemurs, and increase training of Malagasy
393 scientists". Our method provides a means of genomic biodiversity monitoring using
394 cutting edge molecular techniques. Application of this method allowed us to successfully
395 recover six aye-aye mitogenomes from sites in the south-east and west of Madagascar
396 where aye-ayes have not been sampled previously. The wide distribution of aye-ayes
397 across Madagascar means sampling across environments is key in effective population
398 monitoring at local and national scales (Schwitzer et al., 2013). The different habitat types

399 that aye-eyes are distributed in range from rainforests in southeast Madagascar, a habitat
400 with relatively humidity and rainfall, to the dry deciduous forest in the west of
401 Madagascar, with less dense canopy cover and likely greater exposure to direct UV
402 radiation (Rakoto-Joseph, Garde, David, Adelard, & Randriamanantany, 2009;
403 Andriamisedra, Aylward, Johnson, Louis, & Raharivololona, 2015). Our recovery of
404 mitogenomes from distinct environments indicates these abiotic factors do not
405 necessarily preclude the ability to sample eDNA from these areas and this method can
406 likely be applied to sample in habitat across the species' geographic distribution (Sterling
407 & McCreless, 2007). The mobility of this sampling technique along with little training
408 required for sample collection provides a means of surveying large areas of aye-aye
409 habitat in a relatively short time frame. The ease of sampling and ability to survey large
410 areas provide the opportunity for sampling multiple individuals across a population and
411 accessing mitogenome diversity within forest fragments.

412

413 Our preliminary analysis revealed high differentiation between the North population and
414 all other sampled locations, consistent with the whole genome analysis of Perry et al.,
415 (2013) which showed the North population to be the most genetically distinct of three
416 regional populations. Given that our relatively sparse sampling strategy revealed two
417 genetically distinct populations, we encourage the application of this method and the use
418 of the predesigned aye-aye Mito panel from MyBaits® to sample additional areas of aye-
419 aye habitat.

420

421 **Applications to other species**

422 The aye-aye is arguably one of the most elusive lemur species in Madagascar (Schwitzer
423 et al., 2013), and other endangered species that are difficult to locate could also benefit

424 from eDNA sampling. The Critically Endangered *Prolemur simus* leave traces on bamboo
425 plants which are similar to those of sympatric bamboo lemurs that feed on the same plant
426 species (Tan, 1999; Ravaloharimanitra et al., 2011). Application of the method presented
427 here could confirm the presence of these threatened species which are typically sparsely
428 distributed and difficult to locate in the wild (Wright et al., 2008; Ravaloharimanitra et
429 al., 2011; Frasier et al., 2015). Similarly, Critically Endangered *Eulemur cinereiceps* could
430 be sampled via traces on gnawed fronds of *Cecropia peltata* trees (Ralainasolo,
431 Ratsimbazafy, & Stevens, 2006). Although the method developed here is novel, sampling
432 of saliva from foraged material has been used previously as a means of non-invasive
433 sampling; gorillas and golden monkeys discard plant material (Smiley et al., 2010),
434 ungulates leave saliva traces on foraged twigs (Nichols et al., 2012), and large carnivores
435 leave saliva on prey (Blejwas, Williams, Shin, Dale, & Jaeger, 2006; Sundqvist, Ellegren, &
436 Vila, 2008; Glen et al., 2010; Caniglia et al., 2013; Wheat et al., 2016). This method of target
437 capture could be applied to these sources to sample mitogenome eDNA from a range of
438 species.

439

440 **NGS molecular techniques and conservation**

441 This method provides an example of the utility of next-generation molecular techniques
442 for non-invasive samples towards sampling wild populations in a species that requires
443 conservation attention. The application of target capture and enrichment has been used
444 previously in non-human primates to sample faecal and ancient DNA (Perry et al., 2010;
445 Kistler et al., 2015; Snyder-mackler et al., 2016). We demonstrated the application of
446 similar molecular techniques to sample from an Endangered lemur species, revealing
447 broad-scale population structure across the aye-aye's geographic range. One area of
448 discussion that arises from reviews of the application of next-generation molecular

449 techniques to conservation is often the necessity of these novel techniques over more
450 conventional approaches, given the increased cost and amount of data generated
451 (Allendorf et al., 2010; McMahon, Teeling, & Höglund, 2014; Shafer et al., 2015). Here we
452 present both a unique challenge and solution to sampling a low-density, elusive, and
453 Endangered species. These methodological developments are valuable tools that have
454 enabled us to sample and monitor a cryptic species that otherwise has limited genomic
455 sampling potential. As the field of conservation genomics expands methods such as the
456 one presented here can be applied to achieve direct conservation action.

457

458

459 **Author contributions**

460 EL, GP, MA and SJ conceived idea and designed methodology; MA collected samples; MA
461 and AS conducted laboratory work. All authors contributed critically to drafts and gave
462 final approval for publication.

463

464

465 **Data accessibility**

466 Raw sequence reads are available from GenBank SRA accession SRP133213

Figures and tables




Figure 1. A map of the island of Madagascar showing sample locations targeted in this study. The orange circles highlight the two locations surveyed for aye-aye feeding traces and eDNA samples were collected. The blue circles indicate the locations that were sampled in Perry et al. 2013, from which the mitogenome sequences from Kistler et al. 2015 were sampled.



Figure 2. Aye-aye feeding traces. These images indicate the typical aye-aye feeding traces showing distinct shapes and teeth marks. These traces centre on a hole where the aye-aye has used its specialized extractive foraging strategy to access wood boring insect larvae. We sampled pieces of wood around the edge of the trace where the buccal cavity of the aye-aye comes into contact with the tree.




Figure 3. Identity by state matrix on the left, which shows the proportion of pairwise shared alleles at each SNP site for mitogenomes excluding the d-loop region. We compare eDNA samples from feeding traces with previously published mitogenome sequences (Kistler et al., 2015). Warmer colours indicate greater proportions of shared alleles at these loci.

Table 1. The fold increase in percent of unique on-target reads aligning to the aye-aye mitochondrial genome target MitoBait captures compared to shotgun sequencing of the DNA libraries. Data show for the ten libraries that were shotgun sequenced prior to capture.

Sample ID	Number of on-target reads	Unique on-target reads, shotgun sequencing (%)	Unique on-target reads, from captured DNA libraries (%)	Fold enrichment
MSR01	0	0.000000	0.0027	-
MSR44	8	0.000038	0.0538	1416
MSR46	0	0.000000	0.0019	-
MSR50	31	0.000063	0.0248	394
MSR58	189	0.000274	0.0408	149
BTS04	0	0.000000	0.0051	-
BTS38	6	0.000103	0.0267	259
BTS60	0	0.000000	0.0005	-
BTS108	24	0.000124	0.0139	112
BTS112	0	0.000000	0.0056	-

Table 2. Sequencing summary statistics showing number of reads and depth and breadth of coverages for MitoBait captures. Grey shading indicates samples with over 90% breadth of coverage that were used for comparisons with published mitogenomes.

Sample site	Sample ID	Reads on-target (%)	Unique reads on-target (%)	Mean read depth (SD)	Mean % coverage
Tsingy de Beanka	BTS04	0.06	0.0015	1.56 (0.9)	59.3
Tsingy de Beanka	BTS27	0.38	0.0063	13.0 (5.3)	99.4
Tsingy de Beanka	BTS38	0.22	0.027	23.3 (5.9)	98.0
Tsingy de Beanka	BTS60	0.02	0.0005	1.8 (2.2)	45.6
Tsingy de Beanka	BTS68	4.3	0.0088	8.5 (6.0)	98.9
Tsingy de Beanka	BTS102	0.26	0.0013	1.8 (1.4)	27.3
Tsingy de Beanka	BTS105	0.5	0.0005	1.7 (1.2)	22.1
Tsingy de Beanka	BTS108	0.4	0.014	15.1 (5.6)	99.5
Tsingy de Beanka	BTS112	0.46	0.0056	1.3 (0.6)	10.9
Manombo Special	MSR01	0.05	0.0027	2.0 (1.4)	65.9
Manombo Special	MSR23	0.04	0.039	1.2 (0.4)	12.3
Manombo Special	MSR37	0.16	0.0002	1.1 (0.3)	11.1
Manombo Special	MSR39	27.54	0.0024	5.0 (11.83)	45.5
Manombo Special	MSR44	22.25	0.05	1.5 (1.2)	21.9
Manombo Special	MSR46	4.11	0.0019	1.3 (0.5)	10.8
Manombo Special	MSR50	12.19	0.025	2.5 (2.07)	70.9
Manombo Special	MSR56	0.024	0.0001	1.3 (0.6)	15.0
Manombo Special	MSR58	2.71	0.04	12.4 (7.2)	98.5
Manombo Special	MSR62	2.20	0.015	25.1 (8.8)	99.8

468 **References**
469

- 470 Allendorf, F. W., Hohenlohe, P. A., & Luikart, G. (2010). Genomics and the future of
471 conservation genetics. *Nature Reviews Genetics*, *11*(10), 697–709.
472 doi:10.1038/nrg2844
- 473 Andriaholinirina, N., Baden, A., Blanco, M., Chikhi, L., Cooke, A., Davies, N., ... Zaramody.
474 (2015). *Daubentonia madagascariensis*. *The IUCN Red List of Threatened Species*,
475 8235. doi:10.2305/IUCN.UK.2014-
- 476 Andriamisedra, R., Aylward, M., Johnson, S. E., Louis, E. E., & Raharivololona, B. M.
477 (2015). Détermination de quelques aspects de l'écologie de *Daubentonia*
478 *madagascariensis* (Gmelin, 1788) dans deux forêts... *Lemur News*, *19*.
479 arhive.org. (2017). Aye-aye feeding on grubs using elongated finger. BBC Motion
480 Gallery.
- 481 Ávila-Arcos, M. C., Sandoval-Velasco, M., Schroeder, H., Carpenter, M. L., Malaspinas, A. S.,
482 Wales, N., ... Gilbert, M. T. P. (2015). Comparative performance of two whole-
483 genome capture methodologies on ancient DNA Illumina libraries. *Methods in*
484 *Ecology and Evolution*, *6*(6), 725–734. doi:10.1111/2041-210X.12353
- 485 Ballard, J. W. O., & Whitlock, M. C. (2004). The incomplete natural history of
486 mitochondria. *Molecular Ecology*, *13*(4), 729–744. doi:10.1046/j.1365-
487 294X.2003.02063.x
- 488 Beja-Pereira, A., Oliveira, R., Alves, P. C., Schwartz, M. K., & Luikart, G. (2009). Advancing
489 ecological understandings through technological transformations in noninvasive
490 genetics. *Molecular Ecology Resources*, *9*(5), 1279–301. doi:10.1111/j.1755-
491 0998.2009.02699.x
- 492 Bi, K. E., Linderoth, T., Vanderpool, D. A. N., Good, J. M., & Nielsen, R. (2013). Unlocking
493 the vault : next-generation museum population genomics. *Molecular Ecology*, *22*,
494 6018–6032. doi:10.1111/mec.12516
- 495 Blejwas, K. M., Williams, C. L., Shin, G. T., Dale, R., & Jaeger, M. M. (2006). Salivary DNA
496 Evidence Convicts Breeding Male Coyotes of Killing Sheep. *Journal of Wildlife*
497 *Management*, *70*(4), 1087–1093.
- 498 Bushnell, B. (2016). BBMap short read aligner. *University of California, Berkeley,*
499 *California*. URL [Http://Sourceforge. Net/Projects/Bbmap](http://Sourceforge.Net/Projects/Bbmap).
- 500 Caniglia, R., Fabbri, E., Mastrogiuseppe, L., & Randi, E. (2013). Who is who ?
501 Identification of livestock predators using forensic genetic approaches. *Forensic*
502 *Science International : Genetics*, *7*, 397–404.
- 503 Carpenter, M. L., Buenrostro, J. D., Valdiosera, C., Schroeder, H., Allentoft, M. E., Sikora,
504 M., ... Gilbert, M. T. P. (2013). Pulling out the 1 %: Whole-Genome Capture for the
505 Targeted Enrichment of Ancient DNA Sequencing Libraries. *The American Journal*
506 *of Human Genetics*, *93*, 852–864. doi:10.1016/j.ajhg.2013.10.002
- 507 Chancellor, R. L., Satkoski, J., George, D., Olupot, W., Lichti, N., Smith, D. G., & Waser, P. M.
508 (2011). Do Dispersing Monkeys Follow Kin? Evidence from Gray-cheeked
509 Mangabeys (*Lophocebus albigena*). *International Journal of Primatology*, *32*(2),
510 474–490. doi:10.1007/s10764-010-9483-6
- 511 Chiou, K. L., & Bergey, C. M. (2018). Methylation-based enrichment facilitates low-cost,
512 noninvasive genomic scale sequencing of populations from feces. *Scientific*
513 *Reports*, *8*(1). doi:10.1038/s41598-018-20427-9

- 514 Clevenger, A. P., & Sawaya, M. A. (2010). Piloting a Non-Invasive Genetic Sampling
515 Method for Evaluating Population-Level Benefits of Wildlife Crossing Structures.
516 *Ecology and Society*, 15(1).
- 517 Cunningham, E. P., Unwin, S., & Setchell, J. M. (2015). Darting Primates in the Field: A
518 Review of Reporting Trends and a Survey of Practices and Their Effect on the
519 Primates Involved. *International Journal of Primatology*, 36(5), 911–932.
520 doi:10.1007/s10764-015-9862-0
- 521 Cushman, S. a, McKelvey, K. S., Hayden, J., & Schwartz, M. K. (2006). Gene flow in
522 complex landscapes: testing multiple hypotheses with causal modeling. *The*
523 *American Naturalist*, 168(4), 486–99. doi:10.1086/506976
- 524 Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., Depristo, M. A., ... Vcf, T.
525 (2011). The variant call format and VCFtools. *Bioinformatics (Oxford, England)*,
526 27(15), 2156–2158. doi:10.1093/bioinformatics/btr330
- 527 Ekblom, R., & Galindo, J. (2010). Applications of next generation sequencing in
528 molecular ecology of non-model organisms. *Heredity*, 107(1), 1–15.
529 doi:10.1038/hdy.2010.152
- 530 Enk, J., Rouillard, J., & Poinar, H. (2013). Quantitative PCR as a predictor of aligned
531 ancient DNA. *Biotechniques*. doi:10.2144/000114114
- 532 Erickson, C. J. (1995). Feeding Sites for Extractive Foraging by the Aye-Aye, *Dauben-*
533 *tonia madagascariensis*, 240, 235–240.
- 534 Frasier, C. L., Rakotonirina, J.-N., Razanajatovo, L. G., Nasolonjanahary, T. S.,
535 Rasolonileniraka, Mamiaritiana, S. B., ... Louis, E. E. (2015). Expanding
536 Knowledge on Life History Traits and Infant Development in the Greater Bamboo
537 Lemur (*Prolemur simus*): Contributions from Kianjavato, Madagascar. *Primate*
538 *Conservation*, 29(1), 75–86. doi:10.1896/052.029.0110
- 539 Giolai, M., Paajanen, P., Verweij, W., Percival-Alwyn, L., Baker, D., Witek, K., ... Clark, M. D.
540 (2016). Targeted capture and sequencing of gene-sized DNA molecules.
541 *BioTechniques*, 61(6), 315–322. doi:10.2144/000114484
- 542 Glen, A. S., Berry, O., Sutherland, D. ., Garretson, S., Robinson, T., & de Tores, P. J. (2010).
543 Forensic DNA confirms intraguild killing of a chuditch (*Dasyurus geoffroii*) by a
544 feral cat (*Felis catus*). *Conservation Genetics*, 11(September 2014), 1099–1101.
545 doi:10.1007/s10592-009-9888-y
- 546 Gnirke, A., Melnikov, A., Maguire, J., Rogov, P., LeProust, E. M., Brockman, W., ...
547 Nusbaum, C. (2009). Solution hybrid selection with ultra-long oligonucleotides
548 for massively parallel targeted sequencing. *Nature Biotechnology*, 27(2), 182–
549 189. doi:10.1038/nbt.1523
- 550 Gonder, M. K., Mortensen, H. M., Reed, F. A., De Sousa, A., & Tishkoff, S. A. (2007). Whole-
551 mtDNA genome sequence analysis of ancient african lineages. *Molecular Biology*
552 *and Evolution*, 24(3), 757–768. doi:10.1093/molbev/msl209
- 553 Goossens, B., Chikhi, L., Utami, S. S., de Ruiter, J., & Bruford, M. W. (2000). A multi-
554 samples, multi-extracts approach for microsatellite analysis of faecal samples in
555 an arboreal ape. *Conservation Genetics*, 1(2), 157–162.
556 doi:10.1023/A:1026535006318
- 557 Gross, M. (2017). Primates in peril. *Current Biology*, 27(12), R573–R576.
558 doi:10.1016/j.cub.2017.06.002
- 559 Hawkins, M. T. R., Hofman, C. A., Callicrate, T., Mcdonough, M. M., Tsuchiya, M. T. N., Ecer,
560 E. L. I., & Errez, E. G. (2015). In-solution hybridization for mammalian
561 mitogenome enrichment : pros , cons and challenges associated with

- 562 multiplexing degraded DNA. *Molecular Ecology Resources*. doi:10.1111/1755-
563 0998.12448
- 564 Inoue, E., Inoue-Murayama, M., Takenaka, O., & Nishida, T. (2007). Wild chimpanzee
565 infant urine and saliva sampled noninvasively usable for DNA analyses. *Primates*,
566 (48), 156–159. doi:10.1007/s10329-006-0017-y
- 567 Johnsen, A., Kearns, A. M., Omland, K. E., & Anmarkrud, J. A. (2017). Sequencing of the
568 complete mitochondrial genome of the common raven *Corvus corax* (Aves:
569 Corvidae) confirms mitogenome-wide deep lineages and a paraphyletic
570 relationship with the Chihuahuan raven *C. cryptoleucus*. *PLOS ONE*, 12(10),
571 e0187316. doi:10.1371/journal.pone.0187316
- 572 Joost, S., Bonin, a, Bruford, M. W., Després, L., Conord, C., Erhardt, G., & Taberlet, P.
573 (2007). A spatial analysis method (SAM) to detect candidate loci for selection:
574 towards a landscape genomics approach to adaptation. *Molecular Ecology*,
575 16(18), 3955–69. doi:10.1111/j.1365-294X.2007.03442.x
- 576 Kendall, K. C., Stetz, J. B., Boulanger, J., MacLeod, A. M. Y. C., Paetkau, D., & White, G. C.
577 (2009). Demography and Genetic Structure of a Recovering Grizzly Bear
578 Population. *The Journal of Wildlife Management*, 73(1), 3–16. doi:10.2193/2008-
579 330
- 580 Kirillova, I. V., Zanina, O. G., Chernova, O. F., Lapteva, E. G., Trofimova, S. S., Lebedev, V. S.,
581 ... Shapiro, B. (2015). An ancient bison from the mouth of the Rauchua River
582 (Chukotka, Russia). *Quaternary Research*, 84(2), 232–245.
583 doi:10.1016/j.yqres.2015.06.003
- 584 Kistler, L., Ratan, A., Godfrey, L. R., Crowley, B. E., Hughes, C. E., Lei, R., ... Perry, G. H.
585 (2015). Comparative and population mitogenomic analyses of Madagascar 's
586 extinct, giant 'subfossil' lemurs. *Journal of Human Evolution*, 79, 45–54.
587 doi:10.1016/j.jhevol.2014.06.016
- 588 Kohn, M. H., & Wayne, R. K. (1997). Facts from faeces revisited. *TREE*, 12, 223–227.
- 589 Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association
590 mapping and population genetical parameter estimation from sequencing data.
591 *Bioinformatics*, 27(21), 2987–2993. doi:10.1093/bioinformatics/btr509
- 592 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with
593 BWA-MEM.
- 594 Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
595 Wheeler transform. *Bioinformatics*, 25(14), 1754–1760.
596 doi:10.1093/bioinformatics/btp324
- 597 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... Durbin, R. (2009).
598 The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16),
599 2078–2079. doi:10.1093/bioinformatics/btp352
- 600 Lischer, H. E. L., & Excoffier, L. (2012). PGDSpider: an automated data conversion tool
601 for connecting population genetics and genomics programs. *Bioinformatics*,
602 28(2), 298–298. doi:10.1093/bioinformatics/btr642
- 603 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., ... DePristo,
604 M. A. (2010). The genome analysis toolkit: A MapReduce framework for
605 analyzing next-generation DNA sequencing data. *Genome Research*, 20(9), 1297–
606 1303. doi:10.1101/gr.107524.110
- 607 McMahan, B. J., Teeling, E. C., & Höglund, J. (2014). How and why should we implement
608 genomics into conservation? *Evolutionary Applications*, 7(9), 999–1007.
609 doi:10.1111/eva.12193

- 610 Mohandesan, E., Speller, C. F., Peters, J., Uerpmann, H., Uerpmann, M., Cupere, B. E. A. D.
611 E., ... Burger, P. A. (2017). Combined hybridization capture and shotgun
612 sequencing for ancient DNA analysis of extinct wild and domestic dromedary
613 camel. *Molecular Ecology Resources*, 17, 300–313. doi:10.1111/1755-0998.12551
- 614 Morin, D. J., Kelly, M. J., & Waits, L. P. (2016). Monitoring coyote population dynamics
615 with fecal DNA and spatial capture–recapture. *The Journal of Wildlife*
616 *Management*, 80(5), 824–836. doi:10.1002/jwmg.21080
- 617 Morin, P. A., Wallis, J., Moore, J. J., Chakraborty, R., & Woodruff, D. S. (1993). Non-
618 invasive sampling and DNA amplification for paternity exclusion, community
619 structure, and phylogeography in wild chimpanzees. *Primates*, 34(3), 347–356.
620 doi:10.1007/BF02382630
- 621 Nichols, R. V., Königsson, H., Danell, K., & Spong, H. (2012). Browsed twig environmental
622 DNA : diagnostic PCR to identify ungulate species. *Molecular Ecology Resources*,
623 12, 983–989. doi:10.1111/j.1755-0998.2012.03172.x
- 624 Nielsen, E. E., Morgan, J. A. T., Maher, S. L., Edson, J., Gauthier, M., Pepperell, J., ...
625 Ovenden, J. R. (2017). Extracting DNA from ‘jaws’: high yield and quality from
626 archived tiger shark (*Galeocerdo cuvier*) skeletal material. *Molecular Ecology*
627 *Resources*, 17(3), 431–442. doi:10.1111/1755-0998.12580
- 628 Oka, T., & Takenaka, O. (2001). Wild gibbons’ parentage tested by non-invasive DNA
629 sampling and PCR-amplified polymorphic microsatellites. *Primates*, 42(1), 67–
630 73. doi:10.1007/BF02640690
- 631 Orkin, J. D., Yang, Y., Yang, C., Yu, D. W., & Jiang, X. (2016). Cost-effective scat-detection
632 dogs: unleashing a powerful new tool for international mammalian conservation
633 biology. *Scientific Reports*, 6(1), 34758–34758. doi:10.1038/srep34758
- 634 Perry, G. H., Louis, E. E., Ratan, A., Bedoya-Reina, O. C., Burhans, R. C., Lei, R., ... Miller, W.
635 (2013). Aye-aye population genomic analyses highlight an important center of
636 endemism in northern Madagascar. *Proceedings of the National Academy of*
637 *Sciences of the United States of America*, 110(15), 5823–8.
638 doi:10.1073/pnas.1211990110
- 639 Perry, G. H., Marioni, J. C., Melsted, P., & Gilad, Y. (2010). Genomic-scale capture and
640 sequencing of endogenous DNA from feces. *Molecular Ecology*, 19(24), 5332–44.
641 doi:10.1111/j.1365-294X.2010.04888.x
- 642 Perry, G. H., Melsted, P., Marioni, J. C., Wang, Y., Bainer, R., Pickrell, J. K., ... Gilad, Y.
643 (2012). Comparative RNA sequencing reveals substantial genetic variation in
644 endangered primates. *Genome Research*, 22(4), 602–10.
645 doi:10.1101/gr.130468.111
- 646 Perry, G. H., Reeves, D., Melsted, P., Ratan, A., Miller, W., Michelini, K., ... Gilad, Y. (2012).
647 A genome sequence resource for the aye-aye (*Daubentonia madagascariensis*), a
648 nocturnal lemur from Madagascar. *Genome Biology and Evolution*, 4(2), 126–35.
649 doi:10.1093/gbe/evr132
- 650 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., ... Sham,
651 P. C. (2007). REPORT PLINK : A Tool Set for Whole-Genome Association and
652 Population-Based Linkage Analyses. *The American Journal of Human Genetics*,
653 81(September), 559–575. doi:10.1086/519795
- 654 Quéméré, E., Crouau-Roy, B., Rabarivola, C., Louis, E. E., & Chikhi, L. (2010). Landscape
655 genetics of an endangered lemur (*Propithecus tattersalli*) within its entire
656 fragmented range. *Molecular Ecology*, 19(8), 1606–21. doi:10.1111/j.1365-
657 294X.2010.04581.x

- 658 Rakoto-Joseph, O., Garde, F., David, M., Adelard, L., & Randriamanantany, Z. A. (2009).
659 Development of climatic zones and passive solar design in Madagascar. *Energy*
660 *Conversion and Management*, 50(4), 1004–1010.
661 doi:10.1016/j.enconman.2008.12.011
- 662 Ralainasolo, F. B., Ratsimbazafy, J. H., & Stevens, N. J. (2006). Behavior and diet of the
663 Critically Endangered *Eulemur cinereiceps* in Manombo forest , southeast
664 Madagascar. *Madagascar Conservation and Development*, 3(1).
- 665 Randimbiharirinina, D. R., Raharivololona, B. M., Hawkins, M. T. R., Frasier, C. L.,
666 Culligan, R., Sefczek, T., ... Louis, E. E. (2017). Behavior and Ecology of male aye-
667 ayes (*Daubentonia madagascariensis*) in the Kianjavato Classified Forest ,
668 southeastern Madagascar. *Folia Primatologica, Manuscript*.
- 669 Randimbiharirinina, D., Sefczek, T., Raharivololona, B. M., Andriamalala, Y.,
670 Ratsimbazafy, Jonah, & Louis, E. E. (2017). Aye-Aye *Daubentonia*
671 *madagascariensis* (Gmelin, 1788) Madagascar (2016). In C. Schwitzer, R. A.
672 Mittermeier, Rylands, Anthony B., F. chiozza, E. Williamson, E. Macfie, ... A. Cotton
673 (Eds.), *Primates in Peril The world's 25 most endangered primates 2016-2018*.
- 674 Ravaloharimanitra, M., Ratolojanahary, T., Rafalimandimby, J., Rajaonson, A.,
675 Rakotonirina, L., Rasolofoharivelo, T., ... King, T. (2011). Gathering Local
676 Knowledge in Madagascar Results in a Major Increase in the Known Range and
677 Number of Sites for Critically Endangered Greater Bamboo Lemurs (*Prolemur*
678 *simus*). *International Journal of Primatology*, 32(3), 776–792.
679 doi:10.1007/s10764-011-9500-4
- 680 Robinson, J. T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &
681 Mesirov, J. P. (2012). Integrative Genomics Viewer. *Nature Biotechnology*, 29(1),
682 24–26. doi:10.1038/nbt.1754.Integrative
- 683 Sastre, N., Francino, O., Sanchez, A., & Ramirez, O. (2009). Sex identification of wolf
684 (*Canis lupus*) using non-invasive samples, (July 2015). doi:10.1007/s10592-008-
685 9565-6
- 686 Schmieder, R., & Edwards, R. (2011). Quality control and preprocessing of metagenomic
687 datasets. *Bioinformatics (Oxford, England)*, 27(6), 863–864.
688 doi:10.1093/bioinformatics/btr026
- 689 Schubert, G., Stoneking, C. J., Arandjelovic, M., Boesch, C., Eckhardt, N., Hohmann, G., ...
690 Vigilant, L. (2011). Male-mediated gene flow in patrilocal primates. *PloS One*,
691 6(7), e21514–e21514. doi:10.1371/journal.pone.0021514
- 692 Schwitzer, C., Mittermeier, R. a., Davies, N., Johnson, S. E., Ratsimbazafy, J. H.,
693 Razafindramanana, J., ... Nash, S. (2013). *Lemurs of Madagascar: a strategy for*
694 *their conservation 2013-2016* (No. 9781934151624).
- 695 Shafer, A., Wolf, J., Alves, P., Bergström, L., .M, B., Brannstrom, I., ... Zielinski, P. (2015).
696 Genomics and the challenging translation into conservation practice. *Trends in*
697 *Ecology & Evolution*, 30(2), 78–87. doi:10.1016/j.tree.2014.11.009
- 698 Sikes, R. S., & Gannon, W. L. (2011). Guidelines of the American Society of Mammalogists
699 for the use of wild mammals in research. *Journal of Mammalogy*, 92(1), 235–253.
700 doi:10.1644/10-MAMM-F-355.1
- 701 Smiley, T., Spelman, L., Lukasik-, M., Mukherjee, J., Ph, D. V. M. D., Kaufman, G., ...
702 Cranfield, M. (2010). Noninvasive Saliva Collection Techniques for Free-Ranging
703 Mountain Gorillas and Captive Eastern Gorillas E. *Journal of Zoo and Wildlife*
704 *Medicine*, 41(2), 201–209.
- 705 Snyder-mackler, N., Majoros, W. H., Yuan, M. L., Shaver, A. O., Gordon, J. B., Kopp, G. H., ...
706 Alberts, S. C. (2016). Ef fi cient Genome-Wide Sequencing and Noninvasively

- 707 Collected Samples. *Genetics*, 203(June), 699–714.
708 doi:10.1534/genetics.116.187492
- 709 Sterling, E. J. (1994a). Aye-aye: Specialists on Structurally Defended Resources. *Folia*
710 *Primatologica*, 62, 142–154.
- 711 Sterling, E. J. (1994b). Taxonomy and Distribution of *Daubentonia madagascariensis*: A
712 Historical Perspective. *Folia Primatologica*, 62, 8–13.
- 713 Sterling, E. J., & McCreless, E. E. (2007). Adaptations in the Aye-aye : A Review. In *Lemurs*
714 (pp. 159–184).
- 715 Sundqvist, A.-K., Ellegren, H., & Vila, C. (2008). Wolf or dog ? Genetic identification of
716 predators from saliva collected around bite wounds on prey. *Conservation Ge*, 9,
717 1275–1279. doi:10.1007/s10592-007-9454-4
- 718 Tan, C. L. (1999). Group composition, home range size, and diet of three sympatric
719 bamboo lemur species (genus *Haplemur*) in Ranomafana National Park,
720 Madagascar. *International Journal of Primatology*, 20(4), 547–566.
- 721 Valiere, N., & Taberlet, P. (2000). Urine collected in the field as a source of DNA for
722 species and individual identification, 2150–2153.
- 723 Wheat, R. E., Allen, J. M., Miller, S. D. L., Wilmers, C. C., & Levi, T. (2016). Environmental
724 DNA from Residual Saliva for Efficient Noninvasive Genetic Monitoring of Brown
725 Bears (*Ursus arctos*). *PloS One*, 1–18. doi:10.5281/zenodo.58748
- 726 Wright, P. C., Johnson, S. E., Irwin, M. T., Jacobs, R., Schlichting, P., Lehman, S., ... Tan, C.
727 (2008). The Crisis of the Critically Endangered Greater Bamboo Lemur (*Prolemur*
728 *simus*), 2008(23), 5–17.
729

730

Supporting information

Table A1. Sequencing pool parameters for shotgun sequencing of eDNA libraries.

Sample	Conc. ng/ul	Vol. in pool	Total ng	% pool by amount
MSR01	0.26	2	0.528	6.81
MSR44	0.17	5.88	0.999	6.58
MSR46	0.46	2.17	0.999	6.58
MSR50	0.39	2.56	0.998	6.58
MSR58	0.96	1.04	0.994	6.58
BTS04	0.53	1.89	1	6.6
BTS38	0.03	1	0.03	0.2
BTS60	0.70	1.43	1	6.59
BTS108	0.20	1	0.2	1.32
BTS112	0.34	2.94	0.999	6.58

Table A2. Sequencing pool parameters for MitoBait captures from eDNA samples.

Sample	Conc. ng/ul	Vol. in pool	ng total	% pool by amount
BTS04	0.36	11	3.96	6.7
BTS27	6.8	3	4.08	6.9
BTS60	0.515	7.8	4.02	6.8
BTS68	0.633	6.3	3.99	6.75
BTS102	4.4	0.8	3.52	5.96
BTS105	1.56	2.6	4.06	6.87
BTS112	0.23	16.7	2.87	4.86
MSR01	0.43	9.4	4	6.77
MSR23	0.546	7.3	3.99	6.75
MSR37	0.389	10.3	4.01	6.79
MSR39	0.848	4.7	3.9	6.75
MSR44	0.003	10	0.03	0.05
MSR46	0.003	10	0.03	0.05
MSR50	0.0	10	0	0.00
MSR56	0.932	4.3	4.01	6.79
MSR58	0.209	10	2.09	3.54
MSR62	2.66	1.5	3.99	6.75
BTS108	0.42	6	2.5	N/A
BTS38	2.2	1.8	3.96	N/A