Title: Ratiometric assays of autophagic flux in zebrafish for analysis of familial Alzheimer's disease-like mutations

Authors: Haowei Jiang^{1,*}, Morgan Newman¹, Dhanushika Ratnayake^{1,2}, Michael Lardelli¹

Affiliations: ¹University of Adelaide, School of Biological Sciences, Alzheimer's Disease Genetics Laboratory, North Terrace, Adelaide, SA 5005, AUSTRALIA

²Current Address: Monash University, Australian Regenerative Medicine Institute, Wellington Road, Clayton 3800, AUSTRALIA

***Corresponding Author:** Haowei Jiang, University of Adelaide, School of Biological Sciences, Alzheimer's Disease Genetics Laboratory, North Terrace, Adelaide, SA 5005, AUSTRALIA. Email: haowei.jiang@adelaide.edu.au

Key words: autophagy, familial Alzheimer's disease, PRESENILIN genes, zebrafish

Abbreviations: A β , amyloid- β ; AD, Alzheimer's Disease; APP, Amyloid beta A4 protein; Atg, autophagy-related gene; AV, autophagic vacuole; CQ, chloroquine; GFP, green fluorescent protein; LC3, microtubule-associated protein 1 light chain 3; PE, phosphatidylethanolamin; polyQ, poly-glutamine; PSEN, presenilin; Rapa, rapamycin

Financial Disclosure Statement: This research was supported by grants from the National

23 Health and Medical Research Council of Australia, GNT1061006 and GNT1126422, and by
24 funds from the School of Biological Sciences of the University of Adelaide. HJ is supported
25 by an Adelaide Scholarship International from the University of Adelaide.

26

27 **Conflict of Interest Statement:** The authors declare no conflict of interest.

28

29 **Abstract**

30 Protein aggregates such as those formed in neurodegenerative diseases can be degraded via
 31 autophagy. To assess changes in autophagic flux in zebrafish models of familial Alzheimer's
 32 disease (fAD) mutations, we first developed a transgene, polyQ80-GFP-v2A-GFP, expressing
 33 equimolar amounts of aggregating polyQ80-GFP and a free GFP internal control in zebrafish
 34 embryos and larvae. This assay detects changes in autophagic flux by comparing the relative
 35 strength of polyQ80-GFP and free GFP moiety signals on western immunoblots probed with
 36 an antibody detecting GFP. However, the assay's application is limited by the toxicity of
 37 polyQ80-GFP, and because aggregation of this protein may, itself, induce autophagy. To
 38 overcome these issues, we subsequently developed a similar ratiometric assay where
 39 expression of a GFP-Lc3a-GFP transgene generates initially equimolar amounts of GFP-Lc3a
 40 (directed to autophagic degradation) and a free GFP internal control. The sensitivity of this
 41 latter assay is reduced by a cellular protease activity that separates Lc3a from GFP-Lc3a, thus
 42 contributing to the apparent free GFP signal and somewhat masking decreases in autophagic
 43 flux. Nevertheless, the assay demonstrates significantly decreased autophagic flux in
 44 zebrafish lacking *presenilin2* gene activity supporting that the Presenilin2 protein, like human
 45 PRESENILIN1, plays a role(s) in autophagy. Zebrafish heterozygous for a typical fAD-like,
 46 reading-frame-preserving mutation in *psen1* show decreased autophagic flux consistent with
 47 observations in mammalian systems. Unexpectedly, a zebrafish model of the only confirmed
 48 reading-frame-truncating fAD mutation in a human *PRESENILIN* gene, the K115Efs
 49 mutation of human *PSEN2*, shows possibly increased autophagic flux in young zebrafish
 50 (larvae).

51

52 **Introduction**

53

54 Autophagic/lysosomal dysfunction is thought to be involved in the neurodegenerative process
 55 of Alzheimer's disease (AD) (1). The endosomal-lysosomal system is a prominent site for the
 56 processing of the AMYLOID BETA A4 PRECURSOR PROTEIN (APP) to form the
 57 aggregating peptide amyloid β (A β) (2), which is the major component of the amyloid plaques
 58 observed in AD brains. Autophagic vacuoles (AVs) are thought to be the major reservoirs of
 59 intracellular A β , and accumulation of immature AVs has been detected in AD brains ,
 60 suggesting that the maturation of AVs to lysosomes may be impaired (3, 4). Both full-length
 61 APP and β -secretase-cleaved APP are found in such AVs, which are also highly enriched in
 62 PRESENILIN (PSEN) proteins (5) (components of the γ -secretase complexes that cleave
 63 APP to form A β) suggesting that a link may exist between A β production and cell survival
 64 pathways through activated autophagy in AD. The A β accumulation is thought to be induced
 65 by the combination of increased autophagy induction and defective clearance of
 66 A β -generating AVs (1).

67

68 Dominant mutations in the *PRESENILIN* (*PSEN*) genes cause the majority of familial, early
 69 onset AD (fAD, (6)). We have previously argued that a body of evidence supports that the
 70 AD-relevant effect of these mutations may be to alter the activity of PSEN holoproteins
 71 rather than the endoproteolysed forms that are active in the γ -secretase complexes that cleave
 72 APP (7). Indeed, in 2010 Lee et al. showed that changes in PSEN1 holoprotein function,

73 rather than γ -secretase activity, appear to affect lysosomal function in *PSEN1* fAD mutant
 74 human fibroblasts (8). However, the effects of *PSEN* fAD mutations on autophagy are
 75 currently debated (9) and it has not yet been shown that the related protein, PSEN2, plays a
 76 role in autophagic flux. Thus, the roles of the multifunctional PSEN proteins in cellular
 77 function and AD require further investigation.

78

79 Non-mammalian animal models including the zebrafish can facilitate AD research through
 80 development of rapid, novel assays for cellular processes such as autophagy. An excellent
 81 example is the fluorescent protein-based assay published by Kaizuka et al in 2016 that allows
 82 qualitative visualization of differences in autophagic flux within cells (10). However, it is
 83 also desirable to make simple, quantitative assessments of relative differences in autophagic
 84 flux in whole tissues/animals e.g. when these are subjected to different drug treatments or
 85 have different genotypes. In this paper we describe the development of two,
 86 internally-controlled assays using zebrafish embryos/larvae that allow rapid quantitative
 87 comparison of relative autophagic flux using western immunoblotting. We describe
 88 exploitation of the superior of these two assays to demonstrate that activity of the zebrafish
 89 *PSEN2*-orthologous gene, *psen2*, is required for efficient autophagy. We also investigate
 90 changes in autophagic flux in two novel zebrafish models of fAD-like mutations in the
 91 human *PSEN* genes. While a typical, reading frame-preserving fAD-like mutation in
 92 zebrafish *psen1* significantly decreases autophagic flux, a model of the only known reading
 93 frame-truncating *PSEN* fAD mutation does not and may possibly increase autophagic flux in
 94 young fish.

95

96 **Methods and Materials**

97

98 **Zebrafish husbandry and animal ethics**

99 All the wild-type and mutant zebrafish were maintained in a recirculated water system. All
100 work with zebrafish was conducted under the auspices of the Animal Ethics Committee of the
101 University of Adelaide.

102

103 **Construction of the polyQ80-GFP-v2A-GFP transgene**

104 A DNA sequence coding for polyQ80-GFP-v2A-GFP (Fig 2A and S1 File A) was synthesized
105 by Biomatik Corp. and subsequently ligated into the pT2AL200R150G (Tol2
106 transposon-based) (S1 File D) gene transfer vector for expression from the
107 ubiquitously-transcribed elongation factor 1 alpha promoter (EF1 α -p) (11). The
108 polyQ80-GFP-v2A-GFP transgene codes for two proteins, an 80-residue polyglutamine
109 repeat sequence (polyQ80) fused to the N-terminal of GFP protein as well as a free GFP
110 protein. A viral 2A peptide (v2A) sequence between the two GFP sequences allows their
111 synthesis as separate entities by a “ribosomal-skip” mechanism (12). Thus, translation of
112 polyQ80-GFP-v2A-GFP mRNA gives 1:1 stoichiometric synthesis of polyQ80-GFP and free
113 GFP. PolyQ80-GFP subsequently aggregates and should be degraded by autophagy, while the
114 free GFP remains soluble in the cytosol to act as an internal control.

115

116 **Fig 2. Design of three GFP-based constructs for assay of autophagic flux.**

(A) Transgene PolyQ80-GFP-v2A-GFP.

(B) Transgene GFP-Lc3a-GFP.

(C) Transgene GFP-Lc3a.

Construction of the GFP-Lc3a-GFP transgene

A sequence coding for GFP-Lc3a-GFP (Fig 2B and S1 File B) was synthesized by Biomatik and ligated into the pT2AL200R150G (13) gene transfer vector (S1 File D). This construct encodes a fusion protein where GFP is linked to the N-terminal of zebrafish Lc3a which, at its C-terminal, is linked to an additional GFP. When this transgene is expressed in cells, the most C-terminal glycine residue of Lc3a is cleaved by an endogenous ATG4 family protease, producing equimolar amounts of GFP-Lc3a and free GFP. GFP-Lc3a is conjugated to PE and localizes to autophagosomes (10). The GFP-Lc3a molecules attached to the autophagosomal inner membrane are subsequently degraded after fusion with lysosomes, while those on the outer membrane are deconjugated by Atg4 proteins and recycled back to the cytosol. The free GFP exists in the cytoplasm and functions as an internal control. Relative autophagic activity is measured as changes in the ratio of GFP-Lc3a / free GFP via western immunoblotting.

Construction of the GFP-Lc3a transgene

The GFP-Lc3a transgene (Fig 2C and S1 File C) in the pT2AL200R150G vector was derived from GFP-Lc3a-GFP by PCR amplification using 5'-phosphorylated primers to exclude the downstream GFP coding sequence followed by ligation to recircularise the plasmid. The

sequences of the PCR primers used were 5'-TAGATCGATGATGATCCAGACATGA-3' and 5'-GCAGCCGAAGGTCTCCT-3'.

Zebrafish embryos

Three mutations were involved in our research, a putatively null mutation in zebrafish *psen2*, *psen2*^{S4Ter}, a model of a typical reading-frame-preserving fAD-type mutation in *psen1*, *psen1*^{Q96K97del}, and a model of human *PSEN2*^{K115Efs} (14), *psen1*^{K97Gfs}. Descriptions of the generation of these mutations were listed in S3 File.

Five different genotypes of zebrafish embryos, Tübingen (TU) wildtype (+/+), putatively *psen2* null heterozygous (*psen2*^{S4Ter}/+), putatively *psen2* null homozygous (*psen2*^{S4Ter}/*psen2*^{S4Ter}), *psen1*^{Q96K97del} heterozygous (*psen1*^{Q96K97del}/+), and *psen1*^{K97Gfs} heterozygous (*psen1*^{K97Gfs}/+), were spawned either by mass mating or pair-mating of individuals.

Microinjection of zebrafish embryos

For the polyQ80-GFP-v2A-GFP assay, zebrafish zygotes were injected with ~5-10 nL of a solution containing 25ng/μL of the polyQ80-GFP-v2A-GFP transgene with 25ng/μL of Tol2 transposase mRNA (11). For the GFP-Lc3a-GFP and GFP-Lc3a assays, zebrafish zygotes were injected with ~5-10 nL of a solution of 50ng/μL of the GFP-Lc3a-GFP or GFP-Lc3a transgenes respectively with 25ng/μL of the Tol2 transposase mRNA. The injected embryos were incubated at 28.5°C in E3 medium (15). At ~24 hours post fertilization (hpf), embryos

showing widely distributed GFP expression (as visualized by fluorescence microscopy) were selected for subsequent analysis (Fig 1).

Fig 1. Experimental flow chart.

Rapamycin and Chloroquine treatments

For the PolyQ80-GFP-v2A-GFP assay, GFP-expressing zebrafish embryos from microinjection of polyQ80-GFP-v2A-GFP were separated randomly into three groups at ~30 hpf. Two groups were treated with either 1μM rapamycin (Rapa) (SIGMA, R8781) or 50mM chloroquine (CQ) (SIGMA, C6628) from 30 hpf until 48 hpf, with one group remaining untreated as a control. The embryos were chilled and then lysed for western immunoblotting at ~48 hpf.

For the GFP-Lc3a-GFP assay (or GFP-Lc3a expression analysis) at 48 hpf in +/+ embryos, GFP-expressing zebrafish embryos from microinjection were separated randomly into four groups at ~24 hpf. Three of these groups were treated with rapamycin (1μM) or chloroquine (50μM or 50mM) from 30 hpf until 48 hpf, and the remaining group remained untreated as a control. The embryos were chilled and then lysed immediately for western immunoblotting at ~48 hpf.

For the GFP-Lc3a-GFP assay (or GFP-Lc3a expression analysis) at 96 hpf in +/+ larvae, GFP-expressing zebrafish embryos from microinjection were selected at ~24 hpf and

randomly separated into several groups for treatment with rapamycin (1 μ M) or chloroquine (various concentrations) from 78 hpf until 96 hpf. Larvae were then chilled and lysed immediately for western immunoblotting at ~96 hpf.

Western immunoblot analyses

48 and 52 hpf-old embryos were firstly dechorionated and deyolked and then placed in sample buffer (2% sodium dodecyl sulphate (SDS), 5% β -mercaptoethanol, 25% v/v glycerol, 0.0625 M Tris-HCl [pH 6.8], and bromophenol blue) (16), heated immediately to 95°C for 10 min and then sonicated (Diagenode Bioruptor UCD-200) in an ice-water bath at high power mode for 10 min before loading onto polyacrylamide gels for electrophoresis (see below). The 96 hpf-old larvae were not deyolked before lysis in sample buffer.

Samples were loaded onto NuPAGE™ 4-12% Bis-Tris Protein Gels (Invitrogen, NP0323BOX), and the separated proteins were subsequently transferred to nitrocellulose membrane (BIO-RAD, 1620115) using the Mini Gel Tank and Blot Module Set (Life technologies, NW2000). The nitrocellulose membranes were subsequently blocked with blocking reagent (Roche, 11921681001) and then probed with the primary antibody, polyclonal anti-GFP goat (ROCKLAND™, 600-101-215), followed by secondary antibody, horseradish peroxidase (HRP) conjugated anti-goat antibody (ROCKLAND™, 605-703-125). Finally, bound antibody was detected by chemiluminescence using SuperSignal™ West Pico PLUS Chemiluminescent Substrate (ThermoFisher, 34580). The ChemiDoc™ MP Imaging System (Bio-Rad) was used to image all the western immunoblots. The intensity of each band

from western immunoblots was measured by Image Lab™ Software (Bio-Rad). All these intensity data are presented in S2 File.

Statistical tests

F-tests were first applied between different groups of data. If the p value of the F-test was >0.05 , reflecting no significant difference between the variances of two groups, then a two-tailed t-test assuming equal variances was applied to the two groups. If the p value of the F-test was <0.05 , reflecting a significant difference between the variances of two groups, then a two-tailed t-test assuming unequal variances was applied to the two groups.

Results

PolyQ80-based autophagy assay in zebrafish

The viral 2A (v2A) system for simultaneous, stoichiometric expression of two peptides from single mRNA transcripts was first applied in zebrafish by Provost et al. (12). The presence of the v2A linker in a coding sequence causes ribosomes to “skip” production of a peptide bond without terminating translation. We sought to exploit this to adapt Ju et al.’s polyQ80-based assay of autophagic flux (described in 2009) to zebrafish. Ju et al expressed fusions to luciferase of aggregating polyQ80 and non-aggregating polyQ21 in separate thigh muscles of mice (17). As a more direct measure of protein concentration we sought to measure by western immunoblotting the relative amounts of a polyQ80-GFP fusion and free GFP when these are translated in a 1:1 ratio from a single mRNA (Figures 1 and 2A). The

polyQ80-GFP-v2A-GFP transgene expressing these proteins is carried in a Tol2 transposon vector and introduced into fertilized zebrafish eggs by microinjection together with transposase mRNA (see Materials and Methods, Fig 2A) (Note: We initially developed a polyQ80-GFP-v2A-mCherry-based system but this required immunoblotting against GFP followed by stripping and reprobing the blot to detect mCherry, and the expression ratios thus produced were then only comparable between the samples on individual blots – data not shown.) To allow stable propagation of the polyQ80-GFP-v2A-GFP transgene in *Escherichia coli* (i.e. to avoid recombination between the directly repeated GFP coding sequences) we introduced numerous silent mutations into the degenerate codon positions in the downstream, free GFP coding sequence (see Supplementary Data File 1).

Injection of the polyQ80-GFP-v2A-GFP transgene into fertilized zebrafish eggs results in its widespread insertion into the chromosomes of embryonic cells. Expression of GFP is detectable for weeks afterwards although the transgene does not appear to transmit through the germline (data not shown). Western immunoblotting of embryos (up to 48 hours post fertilization, hpf) or larvae (after 48 hpf) from these injected eggs identified three expected bands of protein: a small fraction of full-length polyQ80-GFP-v2A-GFP, and greater amounts of separated polyQ80-GFP and free GFP proteins (Fig 3A).

Fig 3. polyQ80-GFP-v2A-GFP assay of autophagy.

(A) Western immunoblots from polyQ80-GFP-v2A-GFP-injected *+/+* embryos at 48 hpf after treatment with rapamycin (Rapa) or chloroquine (CQ).

(B) Ratios (polyQ80-GFP / free GFP) in polyQ80-GFP-v2A-GFP-injected $+/+$ embryos at 48 hpf after treatment with Rapa or CQ. Means with standard deviations (SDs) are shown.

(C) Western immunoblots from polyQ80-GFP-v2A-GFP-injected $+/+$ embryos at 24 hpf, 48 hpf, 72 hpf, and 96 hpf.

(D) Ratios (polyQ80-GFP / free GFP) in polyQ80-GFP-v2A-GFP-injected $+/+$ embryos at 24 hpf, 48 hpf, 72 hpf, and 96 hpf. Means with SDs are indicated.

An advantage of zebrafish over mammalian models is the easy treatment of their living embryos with drugs by placement of these in the embryos' aqueous support medium for direct absorption. (The absorption of less soluble drugs can be facilitated by co-exposure to 1% DMSO, e.g. (18)). To test whether polyQ80-GFP was being degraded by autophagy selectively relative to free GFP we sought to enhance autophagy induction or block autophagic flux using rapamycin, (19) or chloroquine (20). Both drugs have previously been used successfully to modulate autophagy in zebrafish (19, 21).

In zebrafish embryos, autophagy is up-regulated at the pharyngula stage (24~48 hpf), and the earliest time point at which the phosphatidylethanolamin (PE)-Lc3-II conjugate (critical for autophagy induction) has been detected is 32 hpf (22). Therefore, in our research, injected embryos were exposed to 1 μ M rapamycin or 50mM chloroquine from 30 hpf before lysis at 48 hpf for analysis. The ratios of polyQ80-GFP to free GFP observed from both drug treatments compared to non-treatment controls are presented in Fig 3B. The ratio (poly80Q-GFP / free GFP) was significantly reduced through rapamycin treatment

($p=0.0090$), indicating that autophagy was induced. The ratio (poly80Q-GFP / free GFP) was apparently increased through chloroquine treatment ($p=0.0708$), consistent with inhibition of autophagy. Since the changes in the ratios of poly80Q-GFP / free GFP are consistent with the changes in autophagy expected from these different drug treatments, the polyQ80-GFP-v2A-GFP assay appears able to measure autophagic flux in zebrafish.

The polyQ80-GFP / free GFP ratio observed from the polyQ80-GFP-v2A-GFP transgene possibly decreases between 24 hpf and 96 hpf as indicated by a statistically non-significant trend (Fig 3C and 3D). Also, we noticed that this assay itself is somewhat toxic to zebrafish embryos with approximately half of the injected embryos showing abnormal development at 24 hpf (data not shown). This may be due to the previously observed toxicity of polyglutamine proteins in zebrafish (23, 24). Furthermore, in 2007, Schiffer et al. reported that polyQ102-GFP could aggregate to form large SDS-insoluble inclusions, while free GFP was observed to be produced by removal of polyQ moieties from polyQ-GFP fusion proteins (23). All these phenomena might introduce unanticipated variability into observed polyQ80 / free GFP ratios (although this can be overcome somewhat by the extensive experimental replication that is facilitated by use of the zebrafish model system). For this reason we sought a less-toxic, aggregation-independent, but still internally-controlled alternative assay system.

GFP-LC3-GFP probe to quantify autophagic flux by western immunoblotting

In 2016, Kaizuka et al. (10) described constructs to visualize autophagic flux in zebrafish embryos. Their GFP-LC3-RFP fusion protein is cleaved into separate GFP-LC3 and RFP

proteins by embryos' endogenous ATG4 activity. The GFP-LC3 then associates with autophagosomes while the RFP acts as an internal control.

We wished to adapt the assay from Kaizuka et al. to allow quantification of autophagic flux by western immunoblotting while avoiding the difficulties posed by use of polyQ80-GFP. However, we knew from previous unpublished work with GFP-v2A-RFP fusion proteins that use of an RFP internal control for normalization of GFP expression is problematic since results are only comparable between samples on the same immunoblot. This difficulty could be overcome if GFP served both as autophagy target and as the internal control. Therefore we applied the tandem GFP principle of the polyQ80-GFP-v2A-GFP construct to create GFP-LC3-GFP. This is cleaved by ATG4 within cells to give, initially, equimolar amounts of GFP-LC3 and free GFP proteins (Fig 2B). When assayed, these proteins have discernibly different molecular masses identifiable on western immunoblots probed with a single anti-GFP primary antibody (Fig 2B). The mean GFP-Lc3a / free GFP ratio observed in wild type (+/+) embryos at 48 hpf is slightly higher than at 96 hpf (1.22 vs 1.13 in Fig 5A and 5B respectively). However, similar to the polyQ80-GFP-v2A-GFP assay, this difference is not statistically significant ($p=0.42$).

To test whether changes in the GFP-Lc3a / free GFP ratio reflect changes in autophagic flux, the GFP-Lc3a-GFP construct was expressed in embryos subjected to treatment with rapamycin or chloroquine. Treatment with rapamycin produced a significant decrease in the GFP-Lc3a / free GFP ratio ($p=0.001$, Fig 5A), reflecting the expected increase in autophagy.

Unexpectedly, treatment with chloroquine, both at high (50mM) and low (50μM) concentrations resulted in significant decreases in the GFP-Lc3a / free GFP ratio (Fig 5A). This apparent increase in autophagy conflicted with chloroquine's activity as an autophagy inhibitor, which implied that an unanticipated factor could be distorting the observed ratio during the chloroquine treatments.

Fig 5. Ratios (GFP-Lc3a / free GFP) from the GFP-Lc3a-GFP assay of autophagy.

(A) Embryos at 48 hpf. *+/+* embryos were treated with rapamycin, Rapa, or various concentrations of chloroquine, CQ. *psen2^{S4Ter}* mutant embryos were not treated with these drugs.

(B) *+/+* embryos at 96 hpf after treatment with various concentrations of chloroquine.

(C) Zebrafish embryos of various genotypes at 96 hpf. Only *+/+* embryos were treated with rapamycin or chloroquine as indicated.

p-values are from two-tailed t-tests assuming either equal or unequal variances as appropriate.

After investigating the scientific literature we discovered a report from Ni et al. in 2011 describing that both chloroquine and rapamycin treatments of HeLa cells can cause increased lysosomal pH and cleavage of GFP-LC3 to release free GFP (similar to the removal of polyQ from polyQ-GFP as observed by Schiffer et al (23)). If this form of cleavage also existed in zebrafish it might provide an additional source of free GFP that would affect observed GFP-Lc3a / GFP ratios in embryos expressing the GFP-Lc3a-GFP assay transgene. To test

this possibility we constructed a simple fusion of GFP to Lc3a by deleting the downstream GFP coding sequences from the GFP-Lc3a-GFP transgene to create GFP-Lc3a (Fig 2C). The GFP-Lc3a transgene was then expressed in zebrafish embryos in the same way as for GFP-Lc3a-GFP and the rapamycin and chloroquine treatments were then repeated. At 48 hpf, free GFP (i.e. not fused to Lc3a) can clearly be observed on western immunoblots (Fig 6A), confirming that a protease activity that separates GFP from Lc3a exists in zebrafish. Interestingly, the GFP-Lc3a protein from this transgene is apparently observable as both GFP-Lc3a-I and GFP-Lc3a-II forms whereas we have only observed a single form of GFP-Lc3a protein when this is produced from the GFP-Lc3a-GFP transgene. A similar observation was made by Kaizuka et al. (10) for expression of GFP-LC3 from their GFP-LC3-RFP-LC3ΔG construct when this was expressed from a transgene (as our protein fusions are) rather than from injected mRNA. The free-GFP derived from our GFP-Lc3a transgene is certainly non-negligible relative to the observed GFP-Lc3a-I (e.g. in untreated, wild type embryos, see Fig 6A) at 48 hpf. This implies that free GFP from cleavage of GFP-Lc3a may significantly affect GFP-Lc3a / GFP ratios from GFP-Lc3a-GFP-injected embryos at this developmental time point. When we observed cleavage of GFP-Lc3a in 96 hpf larvae, far less free GFP was seen relative to GFP-Lc3a-I. Therefore, the ratio of GFP-Lc3a to GFP may be less distorted by free GFP from cleavage of GFP-Lc3a at 96 hpf.

Fig 6. Western immunoblots from GFP-Lc3a-injected embryos and larvae.

(A) 48 hpf and (B) 96 hpf respectively. +/- embryos and larvae were also treated with rapamycin (Rapa) or chloroquine (CQ) as indicated.

358

359

360 When we repeated the chloroquine treatments on GFP-Lc3a-GFP transgene-injected
361 zebrafish embryos with lysis at 96hpf we then observed a slightly increased GFP-Lc3a / free
362 GFP ratio at the lowest chloroquine dosage (50 μ M) although without apparent statistical
363 significance ($p=0.6232$, Fig 5B). At higher concentrations of chloroquine (12.5 mM and
364 above) the GFP-Lc3a / free GFP ratio was decreased or was very variable (Fig 5B). Ni et al.
365 reported that saturating doses of chloroquine in HeLa cells are able to block GFP-LC3
366 cleavage completely (25) and this may also be occurring in some zebrafish embryos treated
367 with 50 mM chloroquine. A large proportion of larvae (~80%) were dead at 96 hpf after the
368 50mM chloroquine treatment indicating high lethality of this chloroquine concentration.

369

370 We conclude that exploitation of the GFP-Lc3a-GFP assay is best performed at 96hpf or later
371 to minimize the influence of generation of free GFP from cleavage of GFP-Lc3a. Generation
372 of free GFP from cleavage of GFP-Lc3a reduces the sensitivity of the GFP-Lc3a-GFP
373 construct for detecting decreases in autophagic flux.

374

375 **Autophagic flux in zebrafish *pсен* mutants expressing GFP-Lc3a-GFP**

376 Our tests of the GFP-Lc3a-GFP assay indicated that, when used at 96 hpf, it can reveal
377 changes in autophagic flux without the toxicity and other problems encountered with the
378 aggregation-based polyQ80-GFP-v2A-GFP assay. Therefore, we exploited the
379 GFP-Lc3a-GFP assay to examine autophagic flux changes in a number of zebrafish

presenilin mutants that we have generated in our laboratory (S3 File). These are: a typical fAD-like mutation in *psen1* lacking two codons but preserving the open reading frame, *psen1*^{Q96K97del}; a putatively null mutation of the *psen2* gene, *psen2*^{S4Ter}; and a zebrafish model of the unique, open reading frame-truncating PSEN fAD mutation *PSEN2*^{K115Efs} (14). We have previously suggested that this latter mutation causes fAD by inappropriate mimicry of expression of an hypoxia-induced, naturally-occurring truncated isoform of PSEN2 denoted PS2V (26). In zebrafish, the PS2V-equivalent isoform is expressed from the *PSEN1*-orthologous gene, *psen1*. Therefore, we have modeled this mutation in zebrafish by generation of the *psen1* allele, *K97Gfs*.

The effects of the fAD-like mutations were tested in the heterozygous state corresponding to the dominant inheritance pattern shown by such mutations in humans. The effects of the putatively null mutation of *psen2* were tested in both homozygous and heterozygous embryos to examine, respectively, whether *psen2* activity has any effect on autophagy and whether this effect is haploinsufficient. Fertilized eggs bearing *presenilin* mutations were injected with the GFP-Lc3a-GFP transgene and then examined by western immunoblotting at 96 hpf (Fig 5C). Both heterozygosity for the fAD-like *psen1*^{Q96K97del} mutation and loss of *psen2* activity were observed to decrease autophagic flux significantly ($p=0.0268$ and $p=0.0086$ respectively) as expected from previous analysis of fAD mutations in human *PSEN1* (8) and confirming that *psen2* also functions in autophagy. Heterozygosity for the putatively null mutation in *psen2* also appeared to reduce the GFP-Lc3a / free GFP ratio indicating decreased autophagic flux although this did not reach statistical significance ($p=0.063$). Unexpectedly, heterozygosity

for the mutation *psen1*^{K97Gfs} that models the unique ORF-truncating *K115E* fAD mutation of human *PSEN2* appeared possibly to increase autophagic flux ($p=0.0565$) reflecting complexity in the relationship between fAD mutations and this crucial cellular process.

Discussion

To monitor autophagic flux in zebrafish embryos, we first designed the polyQ80-GFP-v2A-GFP assay that is based on the assumption that polyQ80-GFP aggregates and is a target for autophagy while free GFP is primarily degraded via the proteasome. Since this assay provides 1:1 stoichiometric co-expression of polyQ80-GFP and free GFP protein within the same cells *in vivo* it should reflect changes in autophagic flux by changes in the ratio of polyQ80-GFP / free GFP as observed by western immunoblotting. Since both the polyQ80-GFP and the free GFP protein are detected simultaneously by the same primary antibody, polyQ80-GFP / free GFP ratios can be compared on separate immunoblots which greatly facilitates experimental replication and statistical analysis. While rapamycin and chloroquine treatments of embryos injected with the polyQ80-GFP-v2A-GFP transgene produced ratio changes indicative of the expected changes in autophagic flux, the aggregation of polyQ80 may, itself, induce autophagy and this would limit the assay's sensitivity. This issue is overcome by replacing the polyQ80-GFP protein with a non-aggregating GFP-Lc3a fusion protein, based on the approach taken by Kaizuka et al. (10). Like the polyQ80-GFP-v2A-GFP construct, the GFP-Lc3a-GFP construct produces 1:1 stoichiometric co-expression of two proteins in the same cells *in vivo* (GFP-Lc3a and free GFP proteins).

424

425 The GFP-Lc3a-GFP assay assumes that GFP-Lc3a protein is a target of autophagy while the
 426 free GFP protein remains in the cytosol as an internal control. However, the GFP-Lc3a-GFP
 427 transgene should not affect autophagic flux itself nor TORC1 activity as the related
 428 GFP-LC3-RFP-LC3ΔG construct of Kaizuka et al. did not appear to do so (10). Nevertheless,
 429 the GFP-Lc3a-GFP assay has its own particular limitations. Cleavage of LC3 away from GFP
 430 was reported in HeLa cells when lysosomal pH was increased by chloroquine or rapamycin
 431 treatments (25). Consistent with this, we also saw production of free GFP by cleavage of
 432 GFP-Lc3a in zebrafish embryos and larvae (Fig 6A and 5B). Since the free GFP produced by
 433 GFP-Lc3a cleavage provided an additional source of free GFP, this can affect the measured
 434 ratio of GFP-Lc3a / free GFP and this reduces the sensitivity of the assay to detect decreases
 435 in autophagic flux. We found that this effect was minimized at 96 hpf compared to 48 hpf.
 436 (Note that the observations by Schiffer et al. imply that polyQ80-GFP will also form free
 437 GFP by removal of polyQ80 in zebrafish embryos (23).)

438

439 It is possible that development of a GFP-Lc3a-RFP-GFP transgene-based autophagy assay
 440 might overcome the insensitivity caused by generation of free GFP from GFP-Lc3a since an
 441 RFP-GFP fusion would be easy to identify separately from any free GFP in western
 442 immunoblotting. The accumulation of free GFP versus GFP-Lc3a in such an assay would be
 443 additionally informative regarding autophagic flux versus lysosomal accumulation.

444

Using the GFP-Lc3a-GFP assay, we were able to compare the levels of autophagic flux in zebrafish mutant and wild type larvae. In both *psen2*^{S4Ter/+} and *psen2*^{S4Ter/psen2}^{S4Ter} larvae, we detected decreased autophagic flux (Fig 5C), demonstrating that *psen2*, like *psen1*, plays a role in regulating cells' autophagic activity. This has not previously been shown. However, we should note that, in terms of Psen2 protein's role in γ -secretase activity, our previous work has shown that zebrafish Psen2 plays a greater role in Notch signaling than its corresponding mammalian orthologue (27) and we cannot exclude that zebrafish Psen2 is more similar to Psen1 in its role in autophagy than mammalian PSEN2 is similar to PSEN1. In heterozygous larvae of the "typical" (reading frame-preserving) fAD mutation model *psen1*^{Q96K97del}, we also observed decreased autophagic flux (Fig 5C). Interestingly, our zebrafish model of the human *PSEN2*^{K115Efs} fAD mutation, *psen1*^{K97Gfs}, did not show decreased autophagic flux. Instead, heterozygosity for this mutation appeared possibly to increase autophagic flux in the GFP-Lc3a-GFP assay.

The *K115Efs* mutation of human *PSEN2* is unique among the human *PRESENILIN* fAD mutations in that it truncates the open reading frame of the gene. Nevertheless, our unpublished analyses of aged adult fish modelling this mutation show similar alterations in expression of the genes responding to hypoxia to those seen in aged fish carrying the more "typical" fAD-like mutation *Q96K97del* (Newman et al. *manuscript in preparation*). Therefore, it will be interesting to examine autophagic flux in such aged fish to see whether it subsequently becomes inhibited (as is typical in Alzheimer's disease (4)). To this end, we are

currently determining whether the GFP-Lc3a-GFP transgene can be transmitted through the zebrafish germline for widespread expression in adult tissues.

The possibly increased autophagic flux observed due to heterozygosity for the reading frame-truncating *psen1*^{K97Gfs} mutation argues against the idea that the human *PSEN2*^{K115Efs} mutation, or the zebrafish *psen1*^{K97Gfs} model of it, represent hypomorphic alleles as such alleles would be expected to decrease autophagic flux (e.g. see (8) and the putatively *psen2* null allele analysis of this paper). In fact, our results suggest that this truncated form of PRESENILIN, and the naturally occurring PS2V isoform that it supposedly mimics, may interact with PRESENILIN holoproteins to increase their function in autophagy. Induction of autophagy is a pro-survival response to cellular stresses such as hypoxia (28) and we have previously shown using zebrafish that PS2V acts to decrease cell death under such conditions (26). The observation of increased autophagic flux in zebrafish larvae expressing *psen1*^{K97Gfs} is consistent with this.

In summary, we tested two GFP-based assays intended to detect changes in autophagic flux in zebrafish embryos and larvae by western immunoblotting: polyQ80-GFP-v2A-GFP and GFP-Lc3a-GFP. Both transgenes provide 1:1 stoichiometric co-expression of an autophagy target protein and of free GFP as an internal control. The two assays both appeared able to reflect changes in autophagic flux although each assay displayed particular limitations. Using the GFP-Lc3a-GFP assay, we found that lack of *psen2* activity in zebrafish reduces autophagic flux as does heterozygosity for a typically reading-frame-preserving fAD-like

mutation of *psen1* (*Q96K97del*).

References

1. Boland B, Kumar A, Lee S, Platt FM, Wegiel J, Yu WH, et al. Autophagy induction and autophagosome clearance in neurons: relationship to autophagic pathology in Alzheimer's disease. *J Neurosci*. 2008;28(27):6926-37.
2. Orr ME, Oddo S. Autophagic/lysosomal dysfunction in Alzheimer's disease. *Alzheimers Res Ther*. 2013;5(5).
3. Yu WH, Cuervo AM, Kumar A, Peterhoff CM, Schmidt SD, Lee JH, et al. Macroautophagy--a novel Beta-amyloid peptide-generating pathway activated in Alzheimer's disease. *J Cell Biol*. 2005;171(1):87-98.
4. Nixon RA, Wegiel J, Kumar A, Yu WH, Peterhoff C, Cataldo A, et al. Extensive Involvement of Autophagy in Alzheimer Disease: An Immuno-Electron Microscopy Study. *Journal of Neuropathology & Experimental Neurology*. 2005;64(2):113-22.
5. Area-Gomez E, de Groof AJ, Boldogh I, Bird TD, Gibson GE, Koehler CM, et al. Presenilins are enriched in endoplasmic reticulum membranes associated with mitochondria. *Am J Pathol*. 2009;175(5):1810-6.
6. Pottier C, Hannequin D, Coutant S, Rovelet-Lecrux A, Wallon D, Rousseau S, et al. High frequency of potentially pathogenic SORL1 mutations in autosomal dominant early-onset Alzheimer disease. *Mol Psychiatry*. 2012;17(9):875-9.
7. Jayne T, Newman M, Verdile G, Sutherland G, Munch G, Musgrave J, et al. Evidence For and Against a Pathogenic Role of Reduced gamma-Secretase Activity in Familial Alzheimer's Disease. *J Alzheimers Dis*. 2016;52(3):781-99.
8. Lee JH, Yu WH, Kumar A, Lee S, Mohan PS, Peterhoff CM, et al. Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted by Alzheimer-related PS1 mutations. *Cell*. 2010;141(7):1146-58.
9. Zhang X, Garbett K, Veeraghavalu K, Wilburn B, Gilmore R, Mirnics K, et al. A role for presenilins in autophagy revisited: normal acidification of lysosomes in cells lacking PSEN1 and PSEN2. *J Neurosci*. 2012;32(25):8633-48.
10. Kaizuka T, Morishita H, Hama Y, Tsukamoto S, Matsui T, Toyota Y, et al. An Autophagic Flux Probe that Releases an Internal Control. *Mol Cell*. 2016;64(4):835-49.
11. Clark KJ, Urban MD, Skuster KJ, Ekker SC. Transgenic zebrafish using transposable elements. *Methods Cell Biol*. 2011;104:137-49.
12. Provost E, Rhee J, Leach SD. Viral 2A peptides allow expression of multiple proteins from a single ORF in transgenic zebrafish embryos. *Genesis*. 2007;45(10):625-9.
13. Kawakami K. Tol2: a versatile gene transfer vector in vertebrates. *Genome Biol*. 2007;8(1).
14. Jayadev S, Leverenz JB, Steinbart E, Stahl J, Klunk W, Yu CE, et al. Alzheimer's disease phenotypes and genotypes associated with mutations in presenilin 2. *Brain*. 2010;133(Pt 4):1143-54.
15. E3 medium (for zebrafish embryos) [database on the Internet]2011 [cited October 1, 2011]. Available from: <http://cshprotocols.cshlp.org/content/2011/10/pdb.rec66449.short>.
16. Ganesan S, Moussavi Nik SH, Newman M, Lardelli M. Identification and expression analysis of the zebrafish orthologues of the mammalian MAP1LC3 gene family. *Exp Cell Res*. 2014;328(1):228-37.
17. Ju J-S, Miller SE, Jackson E, Cadwell K, Piwnicka-Worms D, Weihl CC. Quantitation of selective autophagic protein aggregate degradation in vitro and in vivo using luciferase reporters. *Autophagy*. 2009

- 2009/05/16;5(4):511-9.
18. Grunwald DJ, Eisen JS. Headwaters of the zebrafish [mdash] emergence of a new model vertebrate. *Nat Rev Genet.* [10.1038/nrg892]. 2002;3(9):717-24.
19. Fleming A, Rubinsztein DC. Zebrafish as a model to understand autophagy and its role in neurological disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease.* 2011;1812(4):520-6.
20. Yoon YH, Cho KS, Hwang JJ, Lee SJ, Choi JA, Koh JY. Induction of lysosomal dilatation, arrested autophagy, and cell death by chloroquine in cultured ARPE-19 cells. *Invest Ophthalmol Vis Sci.* 2010;51(11):6030-7.
21. Cui J, Sim TH-F, Gong Z, Shen H-M. Generation of transgenic zebrafish with liver-specific expression of EGFP-Lc3: A new in vivo model for investigation of liver autophagy. *Biochemical and Biophysical Research Communications.* 2012 2012/06/01/;422(2):268-73.
22. He C, Bartholomew CR, Zhou W, Klionsky DJ. Assaying autophagic activity in transgenic GFP-Lc3 and GFP-Gabarap zebrafish embryos. *Autophagy.* 2009;5(4):520-6.
23. Schiffer NW, Broadley SA, Hirschberger T, Tavan P, Kretzschmar HA, Giese A, et al. Identification of anti-prion compounds as efficient inhibitors of polyglutamine protein aggregation in a zebrafish model. *J Biol Chem.* 2007;282(12):9195-203.
24. van Bebber F, Paquet D, Hruscha A, Schmid B, Haass C. Methylene blue fails to inhibit Tau and polyglutamine protein dependent toxicity in zebrafish. *Neurobiol Dis.* 2010;39(3):265-71.
25. Ni HM, Bockus A, Wozniak AL, Jones K, Weinman S, Yin XM, et al. Dissecting the dynamic turnover of GFP-LC3 in the autolysosome. *Autophagy.* 2011;7(2):188-204.
26. Moussavi Nik SH, Newman M, Wilson L, Ebrahimie E, Wells S, Musgrave I, et al. Alzheimer's disease-related peptide PS2V plays ancient, conserved roles in suppression of the unfolded protein response under hypoxia and stimulation of gamma-secretase activity. *Hum Mol Genet.* 2015;26.
27. Selkoe D, Kopan R. Notch and Presenilin: regulated intramembrane proteolysis links development and degeneration. *Annu Rev Neurosci.* 2003;26:565-97.
28. Mazure NM, Pouyssegur J. Hypoxia-induced autophagy: cell death or cell survival? *Current Opinion in Cell Biology.* 2010 2010/04/01/;22(2):177-80.

Supporting information

S1 File. Sequence design and sub-cloning for the GFP-based constructs.

(A) Sequence design for the polyQ80-GFP-v2A-GFP construct

(B) Sequence design for the GFP-Lc3a-GFP construct

(C) Sequence design for the GFP-Lc3a construct

(D) Sub-cloning of the GFP-based constructs into the Tol2 vector

S2 File. Intensity ratios of western immunoblots

566 Table 1. Intensity ratios of western immunoblots for Figure 3B

567 Table 2. Intensity ratios of western immunoblots for Figure 3D

568 Table 3. Intensity ratios of western immunoblots for Figure 5A

569 Table 4. Intensity ratios of western immunoblots for Figure 5B

570 Table 5. Intensity ratios of western immunoblots for Figure 5C

571

572 **S3 File. Mutagenesis and breeding of the mutant fish**

573 (A) Mutagenesis and breeding of the *psen2*^{S4Ter} zebrafish

574 (B) Mutagenesis and breeding of the *psen1*^{Q96K97del} and *psen1*^{K97Gfs} zebrafish

575

Mass mating or pair-mating

Microinjection

Select embryos with widespread GFP expression

DRUG TREATMENT ETC.

Larva (after 48 hpf)

Lysis

Experimental replicates etc. are comparable across multiple western immunoblots

...

Embryo (up to 48 hpf)

Remove yolk

Lysis

Experimental replicates etc. are comparable across multiple western immunoblots

...

A

B

bioRxiv preprint doi: <https://doi.org/10.1101/272351>; this version posted February 26, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

C

bioRxiv preprint doi: <https://doi.org/10.1101/272351>; this version posted February 26, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

C
96 hpf

A

B