Variability and compensation of cardiomyocyte ionic conductances at the population level

Colin Rees^{1†}, Jun-Hai Yang^{2†}, Marc Santolini¹, Aldons J. Lusis^{2,3}, James N. Weiss², Alain Karma^{1*}

*For correspondence:
a.karma@neu.edu (AK)

†These authors contributed equally to this work

¹Physics Department and Center for Interdisciplinary Research on Complex Systems, Northeastern University, Boston MA 02115; ²Departments of Medicine (Cardiology) and Physiology, Cardiovascular Research Laboratory, David Geffen School of Medicine, University of California, Los Angeles, CA 90095; ³Departments of Microbiology and Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, CA 90095

Abstract Conductances of ion channels and transporters controlling cardiac excitation may vary in a population of subjects with different cardiac gene expression patterns. However, the amount of variability and its origin are not quantitatively known. We propose a new computational method to predict this variability that consists of finding combinations of conductances generating a normal intracellular Ca^{2+} transient without any constraint on the action potential. Furthermore, we validate experimentally its predictions using the Hybrid Mouse Diversity Panel, a model system of genetically diverse mouse strains that allows us to quantify inter-subject versus intra-subject variability. The method predicts that conductances of inward Ca^{2+} and outward K^{+} currents compensate each other to generate a normal Ca^{2+} transient in good quantitative agreement with current measurements in ventricular myocytes from hearts of different isogenic strains. Our results suggest that a feedback mechanism sensing the aggregate Ca^{2+} transient of the heart suffices to regulate ionic conductances.

Introduction

Following the landmark publication of the Hodgkin-Huxley model of nerve-cell action potential over six decades ago *Hodgkin and Huxley (1952)*, electrophysiological models of increasing complexity have been developed to describe the cardiac action potential (AP) and its interaction with the intracellular calcium (Ca^{2+}) signal, which links electrical signaling to mechanical contraction in cardiomyocytes *Bers (2001)*. As illustrated in Fig. 1 for a mouse ventricular myocyte *Bondarenko et al. (2004)*, those models typically involve a large set of interacting cellular components that includes various voltage-gated membrane ion channels and transporters, the $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger, and Ca^{2+} handling proteins such as the ryanodine receptor (RyR) Ca^{2+} release channels, which open in response to Ca^{2+} entry into the cell via L-type Ca^{2+} channels, and the sarcoplasmic reticulum (SR) Ca^{2+} ATPase (SERCA), which uptakes Ca^{2+} back into the SR. Ca^{2+} release and uptake from the SR causes a transient rise in cytosolic Ca^{2+} concentration, the calcium transient (CaT), which activates myocyte contraction. Those cellular-scale models have been traditionally constructed by piecing together separate mathematical models describing molecular-scale components in the same species (guinea pig *Luo and Rudy (1991)*, mouse *Bondarenko et al. (2004)*, rabbit *Shannon et al. (2004)*;

Figure 1. Schematic representation of the sarcolemmal currents and intracellular Ca^{2+} cycling proteins of the mouse ventricular myocyte model.

41 *Mahajan et al. (2008)*, dog *Fox et al. (2002)*, etc), and by sometimes mixing models in different
 42 species. Experimental measurements of voltage-current relationships and other properties used to
 43 develop those models are typically averaged over several cells from one or a few hearts. While such
 44 prototypical “population-averaged” (and even “species-averaged”) models have proven useful to
 45 investigate basic mechanisms of cardiac arrhythmias on cellular to tissue scales *Krogh-Madsen and*
 46 *Christini (2012)*; *Karma (2013)*; *Qu et al. (2014)*, they fall short of predicting how different individuals
 47 in a genetically diverse population respond to perturbations (such as physiological stressors, ion
 48 channel mutations, drug or gene therapies, etc) affecting one or several cellular components.

49 In a neuroscience context, the limitation of population-averaged models has been highlighted
 50 by pioneering theoretical and experimental studies by Abbot, Marder, and co-workers demon-
 51 strating that ion channel conductances can exhibit a high degree of activity-dependent plasticity
 52 as well as variability between individuals *LeMasson et al. (1993)*; *Siegel et al. (1994)*; *Liu et al.*
 53 *(1998)*; *Golowasch et al. (1999)*; *Prinz et al. (2004)*; *Schulz et al. (2006)*; *Marder and Goaillard (2006)*;
 54 *Grashow et al. (2009)*; *Marder (2011)*; *O’Leary et al. (2013)*. Theoretical studies along this line first
 55 originated from an attempt to explain why neurons can maintain fixed electrical activity patterns
 56 despite a high rate of ion channel turnover. This question was addressed by treating ion channel
 57 conductances as dynamical variables in models of neuronal activity and by using the intracellular
 58 Ca^{2+} concentration as an activity-dependent feedback mechanism regulating their values *LeMas-*
 59 *son et al. (1993)*; *Siegel et al. (1994)*; *Liu et al. (1998)*, a mechanism supported by experiments
 60 *Golowasch et al. (1999)*. Those model neurons displayed remarkable properties such as the ability
 61 to modify their conductances to maintain a given behavior when perturbed or to develop different
 62 properties in response to different patterns of presynaptic activity. Subsequently, a different type of
 63 computational study in which model parameters (conductances and synaptic strengths of a circuit
 64 model of the crustacean somato-gastric ganglion) were varied randomly, demonstrated that a
 65 similar bursting activity could be obtained with multiple parameter sets, i.e. multiple “good enough
 66 solutions” (GES) *Prinz et al. (2004)*. This prediction agreed qualitatively with an experimental study
 67 demonstrating that neurons of the same circuits obtained from different crabs could have markedly
 68 different ion channel densities, corresponding to different gene expression, but yet different circuits
 69 could generate similar bursting activities *Schulz et al. (2006)*.

70 A later experimental study further showed that different circuits, while operating similarly under
 71 controlled conditions, could respond differently to perturbations such as serotonin addition, which
 72 increases the bursting frequency at a population level, but lowers it in some individuals *Grashow*
 73 *et al. (2009)*. Those findings shed light on why pharmacological treatments may work in some
 74 individuals but not others. In addition, they suggest that the existence of different good enough
 75 solutions provides an evolutionary advantage for the survival of a genetically diverse population by

allowing different individuals to better adapt to different environmental challenges so as to survive and restore the population.

Those studies and related ones in the broader context of systems biology *Daniels et al. (2008)*; *Gutenkunst et al. (2007)*; *Transtrum et al. (2015)* and cardiac electrophysiology *Sarkar and Sobie (2009, 2010)*; *Weiss et al. (2012)*; *Sarkar et al. (2012)*; *Britton et al. (2013)*; *Groenendaal et al. (2015)*; *Muszkiewicz et al. (2016)*; *Krogh-Madsen et al. (2016)* have produced a paradigm shift from population-averaged models, with unique fine-tuned parameter sets, to populations of models characterized by multiple parameter sets. In this new paradigm, each set representing a different individual in a population can produce a similar behavior under controlled conditions, but a starkly different response to perturbations for some individuals. This paradigm shift, however, creates new theoretical and experimental challenges.

On the theoretical side, an open question is how to search for GES representing different individuals in a population. The results of this search, which has been carried out using various methods (e.g. random search *Prinz et al. (2004)*, multivariate regression analysis *Sarkar and Sobie (2009, 2010)*, or genetic algorithms *Groenendaal et al. (2015)*), depend critically on what outputs are selected to constrain model parameters. To date, parameter searches in a cardiac context have been “AP centric”, focusing primarily on features of the membrane voltage (V_m) signal. Sarkar and Sobie showed that very different combinations of ion conductances can produce almost identical cardiac AP waveforms, albeit different CaT amplitudes, and that adding additional constraints on the V_m and Ca^{2+} signals can further constrain model parameters *Sarkar and Sobie (2009, 2010)*. Groenendaal et al. *Groenendaal et al. (2015)* constrained model parameters using V_m traces with variable AP waveforms recorded from guinea pig cardiomyocytes under randomly timed electrical stimuli, as opposed to a unique AP waveform recorded during periodic pacing. This search yielded parameter sets that are potentially better suited to describe more complex aperiodic forms of V_m dynamics relevant for arrhythmias. Britton et al. *Britton et al. (2013)* observed experimentally a significant variability in AP waveform in rabbit Purkinje fibers and searched for model parameter combinations that reproduce this waveform variability. They then used those parameter sets to predict different effects of pharmacological blockade of cardiac HERG (I_{Kr} current) potassium channel in different subjects *Britton et al. (2013)*.

All those GES searches have relied in part or entirely on using measured AP features *Sarkar and Sobie (2009, 2010)*; *Sarkar et al. (2012)*; *Britton et al. (2013)*; *Groenendaal et al. (2015)*; *Muszkiewicz et al. (2016)*; *Krogh-Madsen et al. (2016)* to constrain model parameters. However, given that there is no known voltage sensing mechanisms regulating ion channel expression, it is unclear if natural biological variability can be predicted based on AP features. Here, we adopt a different “ Ca^{2+} centric” view *Weiss et al. (2012)*, which postulates as in a neuroscience context *Golowasch et al. (1999)*; *LeMasson et al. (1993)*; *Siegel et al. (1994)*; *Liu et al. (1998)*; *O’Leary et al. (2013)* that model parameters are predominantly constrained by feedback sensing of Ca^{2+} , and potentially other ions (e.g. Na^+) affecting ion channel regulation. This assumption is motivated by the fact that the CaT is directly linked to mechanical contraction and hence to arterial blood pressure in the organism, which is subject to strong feedback regulation by baroreceptors in the autonomic nervous system. Hence, multiple parameter combinations producing a normal CaT could potentially represent different GES in a genetically diverse population. In addition, unlike voltage, intracellular concentrations of Ca^{2+} and Na^+ ions ($[Ca]_i$ and $[Na]_i$, respectively) have a known interactive role in transcriptional regulation of cardiac ion channel proteins and their function *Rosati and McKinnon (2004)*. For example, the Ca^{2+} /calcineurin/NFAT pathway regulates L-type Ca^{2+} channel (LCC) expression *Qi et al. (2008)* and Na^+ modulates cAMP-dependent regulation of ion channels in the heart *Harvey et al. (1991)* including phosphorylation of LCCs via cAMP-dependent protein kinase *Balke and Wier (1992)*. To test this hypothesis, we perform a GES search in which parameters of a mouse ventricular myocyte model are only constrained by CaT features and $[Na]_i$. This search yields GES with different conductances of the L-type Ca^{2+} current ($I_{Ca,L}$) and K^+ currents ($I_{to,f}$ and I_{Kur}) and reveals that conductances are strongly correlated due to compensatory effects of those currents on

the CaT.

On the experimental side, a major challenge is to test whether different GES produced by any given search method are representative of different individuals in a genetically diverse population. Performing this test generally requires distinguishing quantitatively the variability of conductances and electrophysiological phenotype observed in cells extracted from the same heart (intra-heart cell-to-cell variability) from the variability of the same quantities between different subjects (inter-subject variability). Making this distinction is made extremely difficult by the fact that AP features and conductances vary significantly between cells extracted from same region of the heart *Banyasz et al. (2011)*; *Groenendaal et al. (2015)* and that regional (e.g. ventricular base-to-apex and epicardium to endocardium) variations of ion channel expression are also present. The existence of large intra-heart cell-to-cell variability, and the practical limitation that only a finite number of cells can typically be extracted from a single heart for current measurements, raises the question of whether it is feasible to distinguish electrophysiological parameters between genetically distinct individuals.

To cope with this challenge, we use here the Hybrid Mouse Diversity Panel (HMDP) that is a collection of approximately 100 well-characterized inbred strains of mice that can be used to analyze the genetic and environmental factors underlying complex traits. Because inbred strains are isogenic and renewable, we are able to use multiple hearts from the same strain to obtain enough statistics to differentiate quantitatively between intra-heart and inter-subject variability in conductances of key currents ($I_{Ca,L}$, $I_{to,f}$ and I_{Kur}) affecting the AP and CaT of mouse ventricular myocytes from different strains. The results show that, despite large cell-to-cell variability, some strains have clearly distinguishable mean conductances (i.e. conductances averaged over all cells for the same strain). Mean conductances can vary by up to two-and-a-half fold between strains. The results further show that, remarkably, variations of mean conductances for individual strains follow the same correlation ($I_{Ca,L}$ current is large or small when the sum of $I_{to,f}$ and I_{Kur} currents is large or small, respectively) predicted by our computational Ca^{2+} centric GES search. The central hypothesis that parameters are constrained predominantly by features of the CaT (as a surrogate for arterial blood pressure) is further validated experimentally by showing that strains with very different conductances have similar contractile activity. In addition, we use tissue scale simulations to show that compensation remains effective at an organ scale despite large cell-to-cell variability within an individual heart. Finally, we use our findings to interpret the results of recent studies of cardiac hypertrophy and heart failure induced by a stressor in the HMDP *Ghazalpour et al. (2012)*; *Rau et al. (2014, 2017)*; *Santolini et al. (2018)*.

Results

Effects of individual conductances on the calcium transient

We first used a mouse ventricular myocyte model (see Methods) to investigate the effects of changing a single electrophysiological parameter on the CaT. The CaT was characterized by its amplitude, defined as the difference $\Delta[Ca]_i$ between the diastolic and peak value of the cytosolic Ca^{2+} concentration $[Ca]_i$, and the time-averaged value of $[Ca]_i$ over one pacing period, denoted by $\langle[Ca]_i\rangle$. The CaT amplitude $\Delta[Ca]_i$ is a major determinant of the contractile force while $\langle[Ca]_i\rangle$ provides an average measure of the cytosolic Ca^{2+} concentration in the cell. Both quantities will be used as Ca^{2+} sensors for our multi-parameter search of GES and examining individual parameter effects will be useful later to interpret the results of that search. We vary the conductances of sarcolemmal currents and transporters depicted in Fig. 2A and the expression levels of Ca^{2+} handling proteins that include the ryanodine receptor (RyR) Ca^{2+} release channels and the sarcoplasmic reticulum (SR) Ca^{2+} ATPase SERCA, which pumps Ca^{2+} from the cytosol into the SR. For each parameter value, we pace the myocyte at a 4 Hz frequency for many beats until a steady-state is reached where the CaT profile used to calculate $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$ and the intracellular sodium concentration $[Na]_i$ no longer vary from beat to beat.

Fig. 2 shows the effects of individual parameter changes on the steady-state CaT amplitude

Figure 2. Effects of individual conductances on the Ca²⁺ transient (CaT). A) CaT amplitude defined as the difference $\Delta[Ca]_i$ between the peak and diastolic values of the cytosolic Ca²⁺ concentration $[Ca]_i$ versus G/G_{ref} where G is the individual conductance value and G_{ref} some fixed reference value. B) Time-averaged $[Ca]_i$ over one pacing period ($\langle[Ca]_i\rangle$) versus G/G_{ref} . Illustration of the effect of varying $I_{Ca,L}$ conductance (C) and $I_{to,f}$ conductance (D) on AP and CaT profiles, where 50%, 100%, and 150% correspond to $G/G_{ref}=0.5$, 1.0, and 1.5, respectively. E) Effect of varying RyR conductance on SR Ca²⁺ concentration $[Ca]_{SR}$ and CaT.

(Fig. 2A) and average $[Ca]_i$ (Fig. 2B). Both quantities are plotted as a function of conductance fold change G/G_{ref} where G_{ref} is a reference value producing a normal CaT. Increasing the conductance of the inward L-type Ca²⁺ current $I_{Ca,L}$ is seen to strongly increase both $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$ but has a weak effect on the AP waveform (Fig. 2C). The effects on the CaT stem from the fact that $I_{Ca,L}$ is the main trigger of Ca²⁺-induced Ca²⁺ release (CICR), which transfers a large amount of Ca²⁺ from the SR to the cytosol. In contrast, increasing the conductances of K⁺ currents that are dominant in the mouse such as $I_{to,f}$ (fast inactivating component of the transient outward current) and I_{Kur} causes both $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$ to decrease. Increasing either of those K⁺ currents speeds up repolarization (as illustrated for $I_{to,f}$ in Fig. 2D) and hence inactivation of $I_{Ca,L}$, thereby reducing the magnitude of CICR. Increasing the conductance of the sodium-calcium exchanger current I_{NaCa} also causes both $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$ to decrease by enhancing the forward mode of this current that extrudes Ca²⁺ from the cytosol. Changing RyR expression from its reference value is seen to leave $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$ almost unchanged, even though it strongly affects the SR Ca²⁺ concentration $[Ca]_{SR}$ (Fig. 2E). This behavior reflects the well-known effect that making RyR channels more leaky (e.g. by addition of caffeine that increases RyR activity or, similarly here, by increasing the magnitude of the Ca²⁺ release flux through $RyRs$) yields a transient increase in CaT amplitude, but no change in the steady-state CaT amplitude after $[Ca]_{SR}$ adjusts to a lower steady-state level *Bers (2001)*. This effect is illustrated by time traces of $[Ca]_{SR}$ and $[Ca]_i$ in steady-state for different RyR expression levels in Fig. 2E. Finally, changing the expression level of $SERCA$ has opposite effects on $\Delta[Ca]_i$ and $\langle[Ca]_i\rangle$. Increasing $SERCA$ magnitude increases SR Ca²⁺ load, thereby increasing the amount of SR Ca²⁺ release and CaT amplitude, but at the same time depletes Ca²⁺ from the cytosol.

Computationally determined good enough solutions

Next, we performed a computational search for combinations of parameters that yield a normal electrophysiological output as defined by the steady-state CaT amplitude $\Delta[Ca]_i$, time averaged cytosolic Ca²⁺ concentration $\langle[Ca]_i\rangle$, and intracellular sodium concentration $[Na]_i$ at a 4 Hz pacing

Figure 3. Computationally determined good enough solutions (GES). A) Examples of GES representing combinations of 6 conductances that produce a normal CaT and intracellular Na^+ concentration. Each color represents a different GES and the corresponding AP and CaT profiles are shown in B) and C), respectively. D) Histograms of individual normalized conductances G/G_{ref} for a collection of 7,263 GES showing that some conductances are highly variable while others are highly constrained. E) Three-dimensional (3D) plot revealing a 3-way compensation between conductances of $I_{Ca,L}$, $I_{to,f}$ and I_{Kur} . Each GES is represented by a red dot. All GES lie close to a 2D surface in this 3D plot. Pairwise projections (grey shadows) do not show evidence of 2-way compensation between pairs of conductances. F) Alternate representation of 3-way compensation obtained by plotting $I_{Ca,L}$ versus the sum of $I_{to,f}$ and I_{Kur} . Peak values of those currents after a voltage step from -50 to 0 mV are used to make this plot that can be readily compared to experiment.

Figure 3-Figure supplement 1. Histograms of individual ion channel conductances in 8320 GESs found by a GES search constrained only by Ca^{2+} transient amplitude and average, but not constrained by intracellular sodium concentration.

Figure 3-Figure supplement 2. Correlation between $I_{Ca,L}$ and the sum of $I_{to,f}$ and I_{Kur} is weaker but still significant when intracellular sodium concentration is not constrained.

frequency. A GES search that uses the diastolic and peak $[Ca]_i$ values as Ca^{2+} sensors, instead of $\Delta[Ca]_i$ (the difference between the peak and diastolic $[Ca]_i$ values) and time averaged $[Ca]_i$, gives nearly identical results. So our Ca^{2+} sensors can be straightforwardly interpreted physiologically as requirements of normal diastolic and systolic contractile function necessary for a normal arterial blood pressure at the organism scale. A “good enough solution” (GES) was defined as a combination of electrophysiological parameters that produces output values of those three quantities that are close enough to normal target values, which are defined as the values $\Delta[Ca]_i^*$, $\langle [Ca]_i \rangle^*$, and $[Na]_i^*$ corresponding to the reference set of parameters (G_{ref} values) of the ventricular myocyte model. The search was conducted by defining a cost function

$$E(\mathbf{p}) = \sqrt{\sum_{n=1}^N \left(\frac{S_n(\mathbf{p}) - S_n^*}{S_n^*} \right)^2} \leq \epsilon, \quad (1)$$

which is an aggregate measure of the deviation of output sensors $S_n(\mathbf{p})$ from their desired target values S_n^* . Here, $N = 3$ with $S_1 = \Delta[Ca]_i$, $S_2 = \langle [Ca]_i \rangle$, and $S_3 = [Na]_i$, and ϵ is a small tolerance that we choose to be 5%. E is a function of model parameters $\mathbf{p} = (p_1, p_2, \dots)$ chosen to consist of the conductances of $I_{Ca,L}$, $I_{to,f}$, I_{Kur} , and I_{NaCa} as well as RyR and SERCA expression levels. Effects of individual changes of those parameters on CaT properties measured by S_1 and S_2 are shown in Figs. 2A,B. Conductances of other sarcolemmal currents that were found to have a negligible effect on the CaT were kept constant. The search for GES was conducted by first generating a large population of $\sim 10,000$ randomly chosen candidate models, with each model represented by a single parameter set $\mathbf{p}$. A candidate model was generated by randomly assigning each parameter ($p_1, p_2, \dots$) a value comprised between 0 and 300% of its reference value G_{ref} . We then utilized a multivariate minimization algorithm (see Methods for details) that evolves $\mathbf{p}$ until the GES optimization constraint defined by Eq. 1 is satisfied. This method typically yields a large number of GES (7263 of the $\sim 10,000$ trials yield a GES with 6 parameters and 3 sensors described above, with 2737 either not converging or not producing a physiological output) and is computationally more efficient than a random search without optimization that yields very few GES.

Results of the GES search are shown in Fig. 3. Fig. 3A shows the parameters of six representative GES and their corresponding AP waveforms (Fig. 3B) and CaT profiles (Fig. 3C). The CaT profiles are all very close to each other, which holds for all GES, while the AP waveforms exhibit larger variations owing to the fact that the GES search does not involve any voltage sensing. Fig. 3D shows histograms of parameters for all GES. Conductances of sarcolemmal currents tend to be highly variable except for I_{NaCa} , which turns out to be constrained by the intracellular sodium concentration sensor ($S_3 = [Na]_i$). This is revealed by a GES search with only Ca^{2+} sensing (S_1 and S_2) that yields a broader histogram for the I_{NaCa} conductance (Figure 3–Figure Supplement 1). The histogram of RyR expression level is very broad. This is consistent with the fact that this parameter was found to have a very weak effect on the CaT (Figs. 2A,B) due to the compensatory adjustment of SR Ca^{2+} load (Fig. 2E). In contrast, the histogram of SERCA is very narrow. This feature, which persists even if Na^+ sensing is removed (Figure 3–Figure Supplement 1), is predominantly due to Ca^{2+} sensing. It stems from the fact that changing SERCA expression level has opposite effects on the CaT amplitude (Fig. 2A) and average $[Ca]_i$ (Fig. 2B), increasing one while decreasing the other or vice-versa. Therefore, those opposite effects cannot be compensated by changes of conductance of sarcolemmal currents that simultaneously increase or decrease both Ca^{2+} sensors, or by changes of RyR expression level that has a negligible effect on the CaT due to SR load adjustment. However, conductances of inward and outward currents that change both Ca^{2+} sensors in opposite directions can in principle compensate each other. This compensation is revealed by representing each GES as a point in a 3D plot (Fig. 3E) whose axes are the conductances of $I_{Ca,L}$, $I_{to,f}$ and I_{Kur} . This plot shows that all GES lie close to a 2D surface in this 3D conductance space due to a 3-way compensation between the effects of $I_{Ca,L}$, $I_{to,f}$, and I_{Kur} on the CaT. GES lie inside a smeared 2D surface (i.e. a 2D surface of finite thickness) in the 3D conductance space of Fig. 3E. This feature stems from the fact that the GES parameter space considered here is in principle 6-dimensional (4 sarcolemmal

Figure 4. Good enough solutions in the Hybrid Mouse Diversity Panel (HMDP). A) Central result of this paper showing quantitative agreement between theoretically predicted and experimentally measured compensation of inward Ca^{2+} and outward K^{+} currents. Equivalent plot of Fig. 3F showing the sum of $I_{to,f}$ and I_{Kur} versus $I_{Ca,L}$ for 9 different mouse strains using peak values of those currents (proportional to conductances) after a voltage step from -50 to 0 mV. Mean current values (green filled squares) are shown together with standard errors of the mean (thin bars) for each strain. Computationally determined GES are superimposed and shown as faded red points using all 3 sensors (CaT amplitude, average $[\text{Ca}]_i$, and diastolic $[\text{Na}]_i$) and faded blue points for 2 sensors (CaT amplitude and average $[\text{Ca}]_i$). Lines represent linear regression fits using the method of Chi-squared minimization with errors in both coordinates including (solid line, $p = 0.0144$) and excluding (dashed line, $p = 0.0007$) the outlier strain BXA12/Pgn marked by a red box. The small p values of those fit validate the computationally predicted 3-way compensation of Ca^{2+} and K^{+} currents. The 3 strains selected for the organ scale study (C57BL/6J, CXB1/ByJ, and BXA25/Pgn) with low, medium, and high $I_{Ca,L}$ conductance, respectively, are highlighted by blue circles. B) Cell shortening in different HMDP strains where thick and thin bars correspond to standard error of the mean and standard deviation, respectively. A standard ANOVA test shows no significant differences in cell shortening between strains ($p = 0.4136$) supporting the hypothesis that different combinations of conductances produce a similar CaT and contractile activity.

Figure 4-Figure supplement 1. Patch clamp measurements of mean $I_{Ca,L}$, $I_{to,f}$, and I_{Kur} functional current density averaged over multiple cells for 9 HMDP mouse strains.

current conductances and 2 Ca^{2+} protein expression levels). However, SERCA expression and I_{NaCa} conductance are constrained by Ca^{2+} and Na^{+} sensing, respectively, and RyR expression has a negligible effect on both Ca^{2+} and Na^{+} sensors, thereby reducing the relevant parameter space to the 3 conductance axes of Fig. 3C. The subspace of GES that minimizes the cost function E must therefore lie on the 2D surface $E = 0$. This surface is smeared because I_{NaCa} is only constrained by Na^{+} sensing within a finite range and the GES search only minimizes E within a finite tolerance ($E \leq \epsilon$ instead of $E = 0$).

To facilitate the comparison with experiments presented in the next subsection, it is useful to represent the 3-way compensation between $I_{\text{Ca,L}}$, $I_{\text{to,f}}$, and I_{Kur} conductances by plotting the sum of the peak currents of $I_{\text{to,f}}$ and I_{Kur} versus the peak current of $I_{\text{Ca,L}}$ with all three currents measured under voltage-clamp with a step from -50 to 0 mV. Those peak currents are proportional to conductances up to proportionality factors fixed by intra- and extra-cellular ionic concentrations and voltage. In this peak-current representation, the smeared 2D surface of GES of Fig. 3E takes on the simpler form of a thick nearly straight line (Fig. 3F).

Good enough solutions in the HMDP

In order to test the computational modeling predictions, and at the same time differentiate intra-heart cell-to-cell from inter-subject variability, we performed electrophysiological and contractile measurements on ventricular myocytes obtained from mouse hearts of 9 different strains from the HMDP listed in the Methods, each strain assumed to represent a different good enough solution. Peak values of $I_{\text{Ca,L}}$, $I_{\text{to,f}}$, and I_{Kur} were measured under voltage-clamp with a step from -50 to 0 mV following established protocols (see Methods). The K^{+} currents were measured in the same cell and the Ca^{2+} currents in different cells. Contraction was analyzed by measuring myocyte shortening during several paced beats in separate cells for 6 strains that include 5 of the strains in which conductances were measured. In order to collect enough statistics to distinguish cell-to-cell from inter-strain variability, several hearts of each isogenic strain were used. The number of cells that could be obtained from one heart for L-type Ca^{2+} current, several K^{+} currents, or contraction analysis varied from 1 to 7 so that several hearts of each strain were needed to obtain enough independent measurements to statistically distinguish intra-heart cell-to-cell from inter-strain variability (see data in Methods).

The results of current and contraction measurements are shown in Fig. 4. In Fig. 4A, we plot the sum of the peak currents of $I_{\text{to,f}}$ and I_{Kur} versus the peak current of $I_{\text{Ca,L}}$ together with the standard errors of the mean (SEM) of those quantities. Bar plots showing mean current values together with both SEM and standard deviation (SD) characterizing cell-to-cell variability are given in the Methods. We also superimpose on this plot the computationally predicted GES of Fig. 3F. Different HMDP strains, each representing a GES, are seen to function with different combinations of Ca^{2+} and K^{+} currents that compensate each other in a non-trivial 3-way fashion that closely follows the GES computationally determined with a 3-sensor search in which both the CaT and Na^{+} concentration are constrained (faded red points in Fig. 4A). The sum of $I_{\text{to,f}}$ and I_{Kur} follows a linear correlation with $I_{\text{Ca,L}}$ ($p=0.0007$) using 8 out of 9 strains and the correlation remains statistically significant ($p=0.0144$) if the outlier strain (BXA12/PgnJ) is included. Interestingly, this outlier strain still falls within the larger range of computationally predicted GES using a 2-sensor search without Na^{+} sensing (faded blue points in Fig. 4A). To distinguish cell-to-cell from inter-strain variability, we performed a one-way ANOVA F-test on the $I_{\text{Ca,L}}$ measurements. The result shows that $I_{\text{Ca,L}}$ measurements for all strains do not originate from the same distribution (p -value $p=0.000024$). Furthermore, we performed a student T-test using raw data of $I_{\text{Ca,L}}$ measurements for all pairs of strains. The results yield very small statistically significant p -values for pairs of strains with sufficiently different average current values (e.g. BXA25/PgnJ, CXB1/ByJ, and C57BL/6J in Fig. 4A). Those results are consistent with the fact that mean currents differ much more than their standard error for those strains, as can be seen by visual inspection of means and SEM values corresponding to thin bars on both axes of Fig. 4A. We conclude that inter-strain variability of ion channel conductances can be distinguished from

cell-to-cell variability of those same quantities for a significant number of the strains investigated. While the Ca^{2+} and K^{+} currents were measured for the 9 strains reported in Fig. 4A, the Ca^{2+} current was measured in 7 additional strains (total of 16 strains). Those additional measurements reported in the Methods confirm that some strains can have markedly different $I_{\text{Ca,L}}$ conductances.

Unlike ion channel conductances, CaT properties were assumed not to vary in the computationally-enabled GES search, which rests on the hypothesis that Ca^{2+} sensing provides a feedback mechanism that regulates ion channel gene and protein expression. The results in Fig. 4B, which use contraction as a surrogate for CaT amplitude, support this hypothesis by showing that mean values of cell shortening do not vary substantially across strains. This is confirmed by performing a standard ANOVA statistical test, which shows that cell shortening measurements for the six strains reported in Fig. 4B do not originate from different distributions within statistical uncertainty (p-value $p=0.4136$).

Compensation at the organ scale

Current measurements discussed in the previous section show that mean conductances of Ca^{2+} and K^{+} currents vary between strains in a compensatory way so as to produce a normal CaT. They also show that conductances vary significantly from cell to cell around their mean values. This is illustrated in Fig. 5A for 3 mouse strains that have statistically distinguishable mean $I_{\text{Ca,L}}$ conductances (low, medium, and high) as measured by standard errors (thick bars), but exhibit large cell-to-cell variability as measured by the standard deviations (thin bars) of the distributions of conductance measurements in individual cells. This raises the question of whether compensation remains operative at the organ scale in the presence of large cell-to-cell variability. There are two interlinked aspects to this question. The first relates to the cellular-level dynamical coupling between membrane voltage and intracellular Ca^{2+} dynamics that is inherently nonlinear *Krogh-Madsen and Christini (2012); Karma (2013); Qu et al. (2014)*. Even when cells are uncoupled, this nonlinearity could potentially cause the mean CaT amplitude in an ensemble of cells with highly variable conductances to differ from the CaT amplitude computed in a single cell with conductances set to the mean values of the ensemble, as traditionally done in cardiac modeling. The second aspect relates to the additional effect of gap-junctional coupling between cells. This effect is well-known to smooth out cell-to-cell variation of AP waveforms on a mm scale that is much larger than the individual myocyte length. However, whether this smoothing translates into increased organ-scale uniformity of CaT amplitude and contractility is unclear.

To address those two aspects, we constructed tissue scale computational models for 3 mouse strains with statistically distinguishable average conductances (as illustrated for $I_{\text{Ca,L}}$ in Fig. 5A). Tissues of each strain consisted of 56×56 electrically coupled cells (see Methods for details). Simulations were carried out with and without electrical coupling to assess the effect of the latter. The conductances of $I_{\text{Ca,L}}$, $I_{\text{to,f}}$, and I_{Kur} were assumed to vary randomly from cell to cell. Their values were drawn randomly from Gaussian distributions with average values and standard deviations that match experimental current measurements in each strain. All other parameters were kept fixed to reference values. The resulting cell-to-cell variation of conductances is shown in Fig. 5B) using the same peak-current representation of Figs. 3F and 4A. In this representation, each point represents a different cell, and clouds of points of the same color represent all cells in a tissue of the same strain. Furthermore, the center of each cloud falls on the thick line corresponding to the computationally determined GES surface where compensation is operative at the single cell level.

The results of simulations with populations of uncoupled and coupled cells with randomly varying conductances are shown in Figs. 5C to 5G. Fig. 5C shows that AP waveforms are highly variable when cells are uncoupled, reflecting the variability in conductances. Fig. 5D shows that AP waveforms becomes uniform when cells are coupled, as expected, even though interstrain variability is still significant. Fig. 5E compares histograms of CaT amplitude and AP duration (APD) when cells are uncoupled and coupled. Consistent with the AP waveforms of Figs. 5C and 5D, APD histograms in Fig. 5E show that junctional coupling strongly reduces APD variability, as expected.

Figure 5. Organ scale compensation. A) Mean $I_{Ca,L}$ conductance in 3 different HMDP strains where thick and thin bars denote standard error and standard deviation, respectively. B) Sets of conductances generated to be representative of individual cells within ventricular tissue of the 3 strains by assigning normally distributed random values to the $I_{Ca,L}$, $I_{to,f}$ and I_{Kur} conductances using experimentally determined means and standard deviations. The blue, green, and red points correspond to the 3 HMDP strains with low (C57BL/6J), medium (CXB1/ByJ), and high (BXA25/PgnJ) $I_{Ca,L}$ conductance, respectively, and the grey points are the results of the 3-sensor GES search (same as Fig. 3F). C) Variable AP waveforms in uncoupled myocytes with conductances randomly chosen from the distribution shown in B for C57BL/6J and D) AP waveforms for coupled myocytes in tissue for C57BL/6J and the two other strains. AP waveforms of uncoupled cells vary significantly from cell to cell as observed experimentally (Fig. *Figure 5-Figure Supplement 1*) but are uniform in electrotonically coupled cells, as expected. E) Histograms of Ca^{2+} transient (CaT) amplitude (ΔCa) and action potential duration (APD) for C57BL/6J in electrotonically uncoupled and coupled cells. Importantly, in coupled cells, the more uniform APD translates into a much more uniform CaT amplitude, reflecting the strong effect of the cell's APD on its CaT amplitude. F) Distribution of CaT amplitudes within electrotonically coupled cells in tissue scale simulations using the parameter distributions from B. The 3 strains have the same mean CaT amplitude averaged over all cells marked by a thick vertical gray line, thereby demonstrating that compensation of Ca^{2+} and K^+ currents remains operative at a tissue scale. G) Distribution of CaT amplitudes obtained by varying only $I_{Ca,L}$ conductance and with $I_{to,f}$ and I_{Kur} conductances fixed to their reference values. Lack of compensation between Ca^{2+} and K^+ currents in this case yields different mean CaT amplitude.

Figure 5-Figure supplement 1. Action potential recordings from isolated myocytes for mouse strain C57BL/6J paced at 4 Hz under current clamp.

Figure 6. Correlation between L-type Ca^{2+} current conductance and cardiac hypertrophic response to a stressor for different HMDP strains. The Pearson correlation is $r=0.86$ ($p=3e-4$).

CaT amplitude histograms in turn reveal that, in coupled cells, the more uniform APD translates into a more uniform CaT amplitude (i.e. a narrower ΔCa histogram), reflecting the strong effect of the cell's APD on its CaT amplitude. At the organ scale, this ensures that cells in tissue contract with similar force (proportionate to the CaT amplitude), so that force generation is homogeneous throughout the tissue.

Fig. 5F and 5G show that compensation remain operative at the organ scale. Fig. 5F shows that, even though the strains have different average conductances (Fig. 5B), they produce CaT amplitude histograms with approximately the same mean and width. In contrast, if the same simulation is repeated by fixing the conductance of K^+ currents to the value of the strain with the medium value of $I_{Ca,L}$ conductance (CXB1/ByJ), the different $I_{Ca,L}$ conductances are not compensated by different $I_{to,f}$ and I_{Kur} conductances, yielding CaT amplitude distributions with shifted peaks and hence different aggregate contraction (Fig. 5G).

In summary, our results show that compensation remains operative at the organ scale because CaT amplitude histograms have similar means with and without electrical coupling (Fig. 5E). This implies that cell-to-cell variability of conductances and hence APD causes variability of CaT amplitude without significantly affecting its mean, so that $I_{Ca,L}$ and potassium currents can compensate each other even though conductances exhibit large cell-to-cell variations from their mean values. Gap junctional coupling has the additional important effect of reducing CaT amplitude variability, thereby promoting tight organ-level behavior despite high cell-to-cell variability.

Cardiac hypertrophic response to a stressor

From a functional standpoint, the most relevant implication of the present study is that different GES may exhibit markedly different responses to perturbations, as previously demonstrated in a neuroscience context *Grashow et al. (2009)*. To examine this possibility, we reviewed data from separate studies of isoproterenol (ISO)-induced cardiac hypertrophy and heart failure in approximately 100 HMDP strains that include most of the strains used in the present study. In those studies, heart mass was measured in those strains before (m_{pre}) and three weeks after (m_{post}) implantation of a pump continuously delivering isoproterenol (Table 3). Fig. 6 reveals the existence of a statistically very significant correlation between baseline $I_{Ca,L}$ conductance and the hypertrophic response (m_{post}/m_{pre}). Although many factors contribute to the hypertrophic response in the HMDP *Rau et al. (2017)*; *Santolini et al. (2018)*, intracellular Ca^{2+} overload activating the Ca^{2+} -calcein-NFAT

signaling pathway has been shown to play a major role **Bers (2008)**. Since $I_{Ca,L}$ is the major pathway of Ca^{2+} entry into the cytoplasm, it is intriguing to speculate that strains with a larger baseline $I_{Ca,L}$ conductance under baseline conditions have a more robust increase in $I_{Ca,L}$ that is not adequately compensated by repolarizing K^+ currents, making those strains more susceptible to Ca^{2+} overload when $I_{Ca,L}$ is enhanced during sustained β -adrenergic stimulation by isoproterenol. Hypothetically, this may result in a stronger cardiac hypertrophic response. To make this case convincingly, however, would require demonstrating that Ca^{2+} overload is chronically worsened in strains with a high baseline $I_{Ca,L}$ conductance and ruling out other strain-dependent hypertrophy-promoting pathways that are not Ca^{2+} -dependent, which is beyond the scope of the present work.

Compensation and gene expression

In a neuroscience context, ionic conductances of neurons from the stomatogastric ganglion of different crabs were previously found by **Schulz et al. (2006)** to be correlated with gene expression, as shown by independent measurements in the same subjects of functional densities of different ion channels, used to determine conductances, and mRNA levels of genes encoding for pore forming subunits of those channels. In the present study, we did not perform independent measurements of gene expression in the same ventricular myocytes used to measure ionic conductances. However, to examine the possible relationship between compensation of conductances and gene expression, we reviewed the gene expression data from the aforementioned studies of ISO-induced cardiac hypertrophy and heart failure in approximately 100 HMDP strains that include most of the strains used in the present study. Gene expression was measured both in control (pre-ISO) and after injection of ISO for 21 days in 8-10 week old female mice (post-ISO). Details of heart biopsies conducted pre- and post-ISO and microarray data analysis are given in the Methods section of **Santolini et al. (2018)**.

No statistically significant pairwise correlation (Pearson correlation coefficient $r < 0.25$ and p -value $p > 0.05$) were found between expression levels of the genes *Cacna1c*, *Kcnd2*, and *Kcna5*, encoding for the pore forming subunit of the *Cav1.2*, *Kv4.2*, and *Kv1.5* channels associated with $I_{Ca,L}$, $I_{to,f}$, and I_{Kur} , respectively. However, a statistically very significant correlation was found between expression levels of *Cacna1c* and *Kcnp2* that encodes the KChIP2 accessory β subunits directly interacting with *Kv4.2* (see Fig. 7 and caption for r and p values). This correlation is present both in control (pre-ISO), which is the condition relevant to our conductance measurements in selected strains (Fig. 4), and post-ISO. Since increased KChIP2 level is known to increase the functional current density of $I_{to,f}$ **Kuo et al. (2001)**; **Jin et al. (2010)**, the strong positive correlation between *Cacna1c* (*Cav1.2*) and *Kcnp2* (KChIP2) expression levels may partially contribute to the positive correlation between $I_{Ca,L}$ and $I_{to,f}+I_{Kur}$ functional current densities (Fig. 4).

Discussion

In the present study, we have proposed a new methodology for searching for combinations of electrophysiological parameters representing different individuals in a genetically diverse population. While previous studies have used primarily AP features to constrain parameters **Sarkar and Sobie (2009, 2010)**; **Weiss et al. (2012)**; **Sarkar et al. (2012)**; **Britton et al. (2013)**; **Groenendaal et al. (2015)**; **Muszkiewicz et al. (2016)**; **Krogh-Madsen et al. (2016)**, we have chosen to constrain parameters using the Ca^{2+} transient that plays a key role to regulate ion channel expression and activity. We have also examined the effect of additionally constraining the intracellular Na^+ concentration that is also known to modulate ion channel activity. Regulatory mechanisms traverse different levels of biological organization from transcriptional regulation to post-transcriptional and post-translational modification to ion channel trafficking and phosphorylation. Those mechanisms are presently not known in sufficient detail to be modeled quantitatively. However, there is sufficient experimental evidence of feedback sensing of cellular activity via Ca^{2+} and Na^+ concentrations to make a search that constrains model parameters based on those signals plausible. The Ca^{2+} transient determines the contractile force underlying arterial blood pressure generation regulated by baroreceptor

Figure 7. Compensation and gene expression. Plot showing the existence of a statistically very significant correlation (Pearson correlation coefficient $r = 0.47$ and p-value $p = 8.1 \cdot 10^{-13}$) between the expression level of Kcnp2, encoding the KChIP2 accessory β subunits that interact with Kv4.2 channels ($I_{to,f}$) and of Cacna1c, a gene encoding the α_{1C} subunit of the Cav1.2 L-type calcium channels ($I_{Ca,L}$) across 206 mice. Cardiac gene expression was measured in 106 control (Pre-ISO) strains and 21 days after injection of isoproterenol (post-ISO) in 100 HMDP strains (a smaller number due to higher mortality of certain strains). Note that the significant correlation holds when considering separately pre-ISO (blue points, $r = 0.59$, $p = 2 \cdot 10^{-11}$) and post-ISO (red points, $r = 0.42$, $p = 1.5 \cdot 10^{-5}$) data. Lines show best fits of a linear model for pre-ISO (blue), post-ISO (red), and pre- and post-ISO combined (black). Expression data is taken from *Santolini et al. (2018)* and is averaged over all microarray probes for each gene.

feedback via the autonomic nervous system. Hence, fixing the diastolic and peak $[Ca]_i$ values is a physiologically meaningful choice to search for parameter combinations that produce a normal diastolic and systolic function, which we have adopted here.

A remarkable and nontrivial finding of the present computational study is that this choice suffices to produce a physiological AP waveform whose duration matches the experimentally observed variation in isolated myocytes even though the voltage signal is not used to constrain model parameters. This is further highlighted by the fact that, even though the GES search is performed using target values of Ca^{2+} sensors for a 4 Hz pacing frequency, different GES exhibit similar force-frequency relationships over a broad range of pacing frequencies from 0.25 to 4 Hz. Force-frequency curves (with CaT amplitude used as force equivalent) for parameters corresponding to three different strains with low, medium and high L-type Ca^{2+} current conductance all exhibit a negative staircase (i.e. increased CaT amplitude with decreasing frequency < 1 Hz) characteristic of mouse electrophysiology *Bers (2001)*. This property stems from the knock on effect of voltage on the L-type Ca^{2+} current and SR Ca^{2+} release via CICR. As a result of this effect, constraining the CaT indirectly constrains the relative magnitudes of depolarizing and repolarizing currents affecting the AP, i.e. the same CaT amplitude can be obtained with combinations of Ca^{2+} and K^+ currents that are both large or both small, thereby compensating each other, but not with combinations in which the Ca^{2+} current is large and the sum of K^+ currents is small or vice-versa. It remains that the AP waveform and duration are only partially constrained by the CaT and are thus more variable than in a GES search that uses AP features such as duration, plateau voltage, etc, to constrain parameter sets *Sarkar and Sobie (2009, 2010)*.

While the additional constraint to keep the intracellular Na^+ concentration $[Na]_i$ within a normal physiological range is not necessary to produce a physiological AP waveform, it constrains more tightly the conductance of the Na^+-Ca^{2+} exchanger current (compare I_{NaCa} histograms of Fig. 3D and *Figure 3–Figure Supplement 1*). This leads in turn to a tighter 3-way compensation between the L-type Ca^{2+} current and two dominant repolarizing K^+ currents in the mouse, which is less pronounced in a GES search in which $[Na]_i$ is not constrained (*Figure 3–Figure Supplement 2*) and $I_{Ca,L}$ can be compensated by I_{NaCa} in addition to those K^+ currents (4-way compensation).

The present GES search can identify possible combinations of electrophysiological parameters underlying a normal cardiac electrophysiological phenotype. However, it cannot by itself predict which GES among those found are actually represented, even approximately, in a genetically diverse population. Previous electrophysiological studies have indicated that ion channel/transporter conductances and AP waveform can vary from cell to cell and between different subjects. However, distinguishing intra-heart cell-to-cell variability from inter-subject variability has so far proven difficult since different subjects have different genetic backgrounds. Therefore, electrophysiological experiments that use cells isolated from the heart of a subject with a specific genetic background cannot be assumed to be repeatable in other subjects.

In the present study, we have leveraged the fact that different mice in the same HMDP inbred strain are isogenic to distinguish for the first time intra-heart cell-to-cell variability from inter-subject (inter-strain in the HMPD context) variability. This has allowed us to use several hearts for each strain and perform current measurements in enough cells to statistically distinguish mean conductances of several currents in different strains. The results (Fig. 4A, Table 1 and *Figure 4–Figure Supplement 1*) clearly show that conductances differ between strains. Statistical testing shows that $I_{Ca,L}$ conductance measurements for different strains are very unlikely to belong to the same distribution (as indicated by the very small p-value) and mean conductances can vary by as much as two-and-a-half fold between pairs of strains (e.g. C57BL/6J and BXA25/PgnJ), far in excess of the typical standard error of the mean. Furthermore, guided by the predictions of our computational GES search, we have also measured conductances of two dominant repolarizing currents in the mouse, $I_{to,f}$ and I_{Kur} , to test for the existence of compensation between those currents and $I_{Ca,L}$. The results (Fig. 4A) show that for 8 out of the 9 strains in which all three currents were measured, the currents accurately compensate each other as predicted by the GES

search in which both the CaT and $[\text{Na}]_i$ are constrained. Compensation is evidenced by the linear regression fit of $I_{\text{to},f} + I_{\text{Kur}}$ versus $I_{\text{Ca,L}}$. One outlier strain (BXA12/PgnJ) deviates from this fit but still falls within the larger ensemble of computationally predicted GES without the $[\text{Na}]_i$ constraint. Importantly, cells isolated from mouse strains with very different $I_{\text{Ca,L}}$ conductance have statistically indistinguishable contractile function (Fig. 4B). This suggests that compensation between $I_{\text{Ca,L}}$ and K^+ currents in different HMDP strains is present to maintain Ca^{2+} homeostasis, as assumed in the computational GES search. As a whole, the results clearly support the hypothesis that Ca^{2+} concentration play a major role in feedback sensing of cellular activity and regulation of ion channel expression. While more strains would need to be studied to more accurately determine the role of the Na^+ concentration, measurements reported in Fig. 4A suggest that it plays at least an auxiliary role in further constraining conductances beyond Ca^{2+} sensing.

Finally, the correlation between $I_{\text{Ca,L}}$ conductance and cardiac hypertrophic response of HMDP strains to sustained β -adrenergic stimulation (Fig. 6) also highlights the importance of considering the inherent variability of electrophysiological parameters in a genetically diverse population to interpret the variability of phenotypic response in the setting of complex diseases such as heart failure. It also suggests that pharmacological treatments such as anti-arrhythmic L-type calcium channel blockers, or pathologies such as hyperkalemia (elevated potassium level), will have different effects in different subjects. For example, in the setting of the HMDP, an L-type calcium channel blocker or hyperkalemia would be expected to have a stronger effect on the calcium transient and action potential of mice strains that function under normal conditions with larger $I_{\text{Ca,L}}$ and potassium current conductances. Taking into account this population-variability seems ultimately needed to develop personalized therapies for cardiac arrhythmias and heart failure.

Methods and Materials

Overview of the HMDP

The hybrid mouse diversity panel (HMDP) consists of a population of over 100 inbred mouse strains selected for usage in systematic genetic analyses of complex traits *Ghazalpour et al. (2012)*. The main goals in selecting the strains were to (i) increase resolution of genetic mapping, (ii) have a renewable resource that is available to all investigators world-wide, and (iii) provide a shared data repository¹ that would allow the integration of data across multiple scales, including genomic, transcriptomic, metabolomic, proteomic, and clinical phenotypes.

Electrophysiological and contraction measurements

Cell Isolation

Ventricular myocytes were enzymatically isolated from the hearts of adult female mice (8-12 wk old) using a procedure previously developed and utilized to isolate rabbit cardiomyocytes by Yang et al. *Yang et al. (2008)*. Briefly, hearts were removed from mice anesthetized with intravenous pentobarbital and perfused retrogradely at 37°C in Langendorff fashion with nominally Ca^{2+} -free Tyrode's buffer containing 1.2 mg/ml collagenase type II (catalog number 4176; Worthington) and 0.12 mg/ml protease type XIV (catalog number P5147; Sigma) for 10-17 min. After washing out the enzyme solution, the ventricles were cut from the atria and aorta and transferred to a separate glass dish containing Tyrode's solution. Cells were isolated by gentle mechanical dissociation, stored at room temperature, and used within 5 h. This procedure typically yielded 30-50% of rod-shaped and Ca^{2+} -tolerant myocytes.

Patch Clamping

Isolated ventricular myocytes were patch clamped in the whole cell ruptured patch configuration using borosilicate glass pipettes (1–3-megaohm tip resistance). Myocytes were superfused at $34\text{--}36^\circ\text{C}$ with Tyrode's solution modified accordingly. Currents were measured under voltage clamp

¹<https://systems.genetics.ucla.edu/about/hmdp2>

conditions, using an Axopatch 200B amplifier with a Digidata 1440A interface (Axon Instruments, Union City, CA). Data were acquired and analyzed using pClamp (Axon instruments) and Origin (Origin).

L-type calcium current measurements

For characterization of I_{Ca} properties, the pipette solution, designed to eliminate K^+ and Cl^- currents, contained (in mM) 100 CsMeS, 30 CsCl, 5 MgATP, 5 phosphocreatine di(tris), 5 N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES), 5 NaCl and 0.1 1,2-Bis(2-Aminophenoxy)ethane-N,N',N'-tetraacetic acid (BAPTA) (pH adjusted with HEPES to 7.1-7.2). The superfusate, designed to eliminate K^+ currents, contained (in mM), 136 NaCl, 5.4 CsCl, 1 MgCl₂, 0.33 NaH₂PO₄, 10 glucose, 5 HEPES, and 1.2 CaCl₂ (pH adjusted with Trizma base to 7.4). Voltage-clamp protocols to assess activation were as previously described by Delbridge et al. *Delbridge et al. (1997)*. In all experiments, peak I_{Ca} current was recorded after a voltage step from -50 to 0 mV.

Potassium current measurements

For characterization of K^+ current properties, the pipette solution, designed to eliminate Ca^{2+} currents, contained (in mM) 130 KCl, 5 MgATP, 5 phosphocreatine di(tris), 5 HEPES, 5 NaCl, and 10 BAPTA (pH adjusted with HEPES to 7.1-7.2). The superfusate, designed to eliminate Ca^{2+} currents, contained (in mM) 136 NaCl, 5.4 KCl, 1 MgCl₂, 0.33 NaH₂PO₄, 10 glucose, 5 HEPES, and 0.2 CdCl₂ (pH adjusted with Trizma base to 7.4). To access the activation of K^+ current components: I_{Kur} , $I_{to,f}$ and I_{Kss} , we adopted a voltage protocol, similar to the one reported by *Zhou et al. (1998)*, in combination with the usage of 4-aminopyridine (4-AP) that has the following pharmacological properties: (1) I_{Kur} is markedly blocked by 4-AP at submillimolar concentration (e.g., 0.1 mM); (2) a higher concentration (i.e., > 1 mM) blocks $I_{to,f}$ effectively; and (3) I_{Kss} is 4-AP resistant. Using this procedure, peak I_{Kur} , $I_{to,f}$ and I_{Kss} currents could be deduced from three current measurements without 4-AP and with 0.1 mM and 2 mM 4-AP after a voltage step from -50 to 0 mV.

Results of patch clamp measurements

Results of patch clamp measurements for all strains are summarized in Table 1 and *Figure 4-Figure Supplement 1*. Both the L-type Ca^{2+} current and two K^+ currents ($I_{to,f}$ and I_{Kur}) were measured in 9 strains and the L-type Ca^{2+} current alone was measured in 16 strains.

Contraction analysis

The length of ventricular myocytes was measured and analyzed using the method described by *Sdek et al. (2011)*. Briefly, myocytes were imaged during pacing using a high-speed charge-coupled device-based camera (128 × 128 pixels; Cascade 128+; Photometrics) at 290 frames per second. The acquired video image data were then processed using Imaging Workbench software (version 6.0; INDEC BioSystems). Myocyte length was measured and analyzed using ImageJ software. Cell shortening was calculated from the ratio of peak systolic length to resting diastolic length averaged over 10 contractions evoked by the stimulus train. Results of cell shortening measurements for six strains are summarized in Table 2.

Heart extraction and mass measurement for cardiac hypertrophic response

At sacrifice, hearts were excised, drained of excess blood and weighed. Each chamber of the heart (LV with inter-ventricular septum, RV free wall, RA and LA) was isolated and subsequently weighed. Cardiac hypertrophy was calculated as the increase in total heart weight after isoproterenol (ISO) treatment compared to control animals (see Table 3). As described previously *Wang et al. (2016)*, the ISO treatment consisted of 30 mg per kg body weight per day of Isoproterenol (ISO) administered for 21 days in 8-10 week old female mice using ALZET osmotic mini-pumps, which were surgically implanted intraperitoneally. The average number of control hearts per strain was 2.75. The average number of treated hearts per strain was 3.5.

Table 1. Patch clamp measurements of $I_{Ca,L}$, $I_{to,f}$, and I_{Kur} functional current density. Mean current density averaged over n cells isolated from multiple hearts for each strain is given together with the standard error.

Strain	$I_{Ca,L}$ (pA/pF)	n	I_{Kur} (pA/pF)	n	$I_{to,f}$ (pA/pF)	n	I_{Kss} (pA/pF)	n
A/J	13.71947 ± 1.23085	19	11.61804 ± 2.79089	13	13.03735 ± 1.64818	14	5.23445 ± 0.48377	13
BALB/cByJ	10.72278 ± 1.3951	9	11.496 ± 2.03274	10	7.46938 ± 0.82615	9	8.197 ± 0.62438	10
BTBR T+tf/J	14.75667 ± 1.14159	6	16.42364 ± 2.78295	11	9.271 ± 1.43985	10	6.093 ± 0.39553	9
BXA12/PgnJ	11.01333 ± 0.98995	9	15.66429 ± 1.70548	8	11.89875 ± 2.18027	8	8.34556 ± 1.30065	9
BXA25/PgnJ	17.57 ± 4.81376	5	15.8049 ± 1.74509	7	11.62704 ± 1.59613	7	7.11986 ± 1.31554	6
BXH6/TyJ	7.32625 ± 0.59327	16	5.954 ± 0.43731	10	8.73172 ± 0.67617	11	7.98545 ± 0.87545	11
C57BL/6J	7.3925 ± 0.93181	10	7.82727 ± 1.45134	11	7.594 ± 0.92097	10	9.16273 ± 1.62148	11
CXB1/ByJ	11.93909 ± 0.81022	11	11.69556 ± 1.42095	9	8.73883 ± 1.11087	10	6.089 ± 0.5724	10
CXB11/HiA/J	7.63625 ± 0.89344	8	8.90316 ± 2.49755	6	8.42537 ± 1.24635	7	6.8481 ± 1.3638	7
AXB8/PgnJ	7.355 ± 1.09612	6	-	-	-	-	-	-
BXA14/PgnJ	11.28091 ± 1.00796	15	-	-	-	-	-	-
BXA4/PgnJ	11.77615 ± 1.19476	13	-	-	-	-	-	-
BXD34/TyJ	11.12556 ± 1.13741	9	-	-	-	-	-	-
CBA/J	7.956 ± 0.81269	10	-	-	-	-	-	-
CXB7/ByJ	9.05386 ± 0.6087	16	-	-	-	-	-	-
SJL/J	11.2745 ± 0.99708	20	-	-	-	-	-	-

Table 2. Cell Shortening at 1 Hz pacing

Strain	Shortening ($\Delta L/L$)	n
BXA12/PgnJ	0.124 ± 0.02521	5
BXA14/PgnJ	0.13346 ± 0.01223	14
BTBR T+tf/J	0.11029 ± 0.01015	7
BALB/cByJ	0.1348 ± 0.01525	5
C57BL/6J	0.13533 ± 0.00914	6
BXA25/PgnJ	0.1156 ± 0.01456	5

Mathematical Model of Mouse Ventricular Myocytes

We have developed a novel mathematical model of mouse ventricular myocytes that combines elements of previously published ventricular myocyte models *Shiferaw et al. (2003)*; *Shannon et al. (2004)*; *Bondarenko et al. (2004)*; *Mahajan et al. (2008)*. For this purpose, we kept the mathematical formulation of intracellular calcium cycling of the Mahajan model developed by *Shiferaw et al. (2003)* and replaced several sarcolemmal currents by those formulated by *Bondarenko et al. (2004)* for mouse ventricular myocytes. This model allowed us to explore efficiently the space of good enough solutions that we can compare to experimentally measured variability found in the hybrid mouse diversity panel (HMDP). The *Bondarenko et al. (2004)* mouse model has sarcolemmal currents fitted to detailed experimental measurements of sarcolemmal currents in mouse ventricular myocytes. The *Mahajan et al. (2008)* model is a rabbit model that integrates a Markov model of $I_{Ca,L}$ together with the *Shannon et al. (2004)* formulation of other sarcolemmal currents and the *Shiferaw et al. (2003)* model of calcium cycling and SR calcium release. The *Shiferaw et al. (2003)* model represents the release of calcium from the SR as a sum of individual spark events, which reproduces important observed instabilities such as Ca^{2+} transient alternans. Even though the *Shiferaw et al. (2003)* model of calcium cycling model was developed for rabbit myocytes, it can in principle describe Ca^{2+} cycling in other species including mouse. Therefore, we constructed a mouse model starting with the *Mahajan et al. (2008)* model, which incorporates the *Shiferaw et al. (2003)* model of calcium cycling, and using the formulation of *Shannon et al. (2004)* for $I_{Ca,L}$ and

Table 3. Heart mass before and 3 weeks after Isoproterenol (ISO) injection.

Strain	Heart mass pre-ISO, m_{pre} (g)	Heart mass post-ISO, m_{post} (g)
A/J	0.088666667	0.133
AXB8/PgnJ	0.087	0.0992
BALB/cByJ	0.10105	0.1389
BTBR T+tf/J	0.14162	0.223
BXA-12/PgnJ	0.064	NA
BXA-14/PgnJ	0.0975	0.1248
BXA-4/PgnJ	0.1031	0.14675
BXD-34/TyJ	0.1215	NA
BXH-6/TyJ	0.0845	0.0878
C57BL/6J	0.096716667	0.1222
CBA/J	0.095333333	0.13
CXB-11/HiA/J	0.1135	0.1255
CXB-7/ByJ	0.109	0.1475
SJL/J	0.087	0.123333333

Bondarenko et al. (2004) for other sarcolemmal currents fitted to mouse data. We did not use the *Bondarenko et al. (2004)* detailed Markov formulations of the gating of I_{Na} and $I_{Ca,L}$ channels with fast transition rates that were computationally prohibitive for the number of simulations we performed in this study. However, we checked that our combined model reproduces well the mouse electrophysiological phenotype of the *Bondarenko et al. (2004)* model while being computationally efficient and incorporating a realistic description of Ca^{2+} cycling. We verified that the *Shiferaw et al. (2003)* model of Ca^{2+} cycling integrated with the *Shannon et al. (2004)* model of $I_{Ca,L}$ produced a normal bell-shaped SR release as a function of step-voltage. We also verified that the force-frequency relationship produced by the model has the correct negative staircase observed experimentally. The equations for the model are described below. The reference values of the 6 parameters varied in this study are given in Table 4. Table 5 lists all other parameters used that were kept fixed.

Table 4. Reference values of ionic current parameters varied in the GES search

Parameter	Definition	Reference Value	Reference Source
g_{Ca}	Ca^{2+} current flux	333.32 mmol/(Cm C)	Measured
v_{up}	Peak uptake rate	1.17 μ M/ms	Chosen*
g_{NaCa}	Peak NaCa rate	36.6 μ M/s	<i>Bondarenko et al. (2004)</i>
g_{RyR}	Release current strength	12.9 sparks cm^2 /mA	<i>Mahajan et al. (2008)</i>
$g_{to,f}$	$I_{to,f}$ peak conductance	0.16 A/F	Measured
g_{Kur}	I_{Kur} peak conductance	0.144 A/F	Measured

* v_{up} was chosen such that the reference Ca^{2+} transient amplitude was normal.

Table 5. Mouse ventricular myocyte model parameters.

Parameter	Definition	Value
Physical constants and ionic concentrations		
C_m	Cell capacitance	$3.1 \times 10^{-4} \mu\text{F}$
V_i	Cell volume	$2.58 \times 10^{-5} \mu\text{l}$
V_s	Submembrane volume	$0.02 V_i$
F	Faraday Constant	96.485 C/mmol
R	Universal gas constant	$8.314 \text{ J mol}^{-1} \text{ K}^{-1}$
T	Temperature	298 K
$[\text{Na}]_o$	External Na^+ concentration	140 mM
$[\text{K}]_i$	Internal K^+ concentration	143.5 mM
$[\text{K}]_o$	External K^+ concentration	5.4 mM
$[\text{Ca}^{2+}]_o$	External Ca^{2+} concentration	1.8 mM
Cytosolic buffering parameters		
B_T	Troponin C concentration	70 $\mu\text{mol/l}$ cyt
k_{on}^T	on rate for Troponin C binding	$0.0327 (\mu\text{M ms})^{-1}$
k_{off}^T	off rate for Troponin C binding	$0.0196 (\text{ms})^{-1}$
B_{SR}	SR binding site concentration	47 $\mu\text{mol/l}$ cyt
K_{SR}	SR binding site disassociation constant	0.6 μM
B_{Cd}	Calmodulin binding site concentration	24 $\mu\text{mol/l}$ cyt
K_{Cd}	Calmodulin binding site disassociation constant	7 μM
B_{mem}	Membrane binding site concentration	15 $\mu\text{mol/l}$ cyt
K_{mem}	Membrane binding site disassociation constant	0.3 μM
B_{sar}	Sarcolemma binding site concentration	42 $\mu\text{mol/l}$ cyt
K_{sar}	Sarcolemma binding site disassociation constant	13 μM

SR release parameters

τ_r	Spark lifetime	10 ms
τ_a	NSR-JSR diffusion time	20 ms
u	Release slope	4 ms ⁻¹
c_{sr}	Release slope threshold	90 μ M / I cytosol
τ_d	$c_p - c_s$ diffusion time	0.50 ms*
τ_s	$c_s - c_i$ diffusion time	0.75 ms

Exchanger, uptake, and SR leak parameters

c_{up}	Uptake threshold	0.5 μ M
k_{sat}	NaCa saturation threshold	0.1
ξ	NaCa energy barrier position	0.35
K_{m,Na_i}	ion mobility constant	21 mM
K_{m,Na_o}	ion mobility constant	87.5 mM
K_{m,Ca_o}	ion mobility constant	1380 μ M
g_l	Leak current conductance	1.74×10^{-5} ms ⁻¹

Ionic current parameters

g_{Na}	Na ⁺ current conductance	13 mS/ μ F
$g_{Na,b}$	Na ⁺ background current conductance	0.0026 mS/ μ F
$g_{Ca,b}$	Ca ²⁺ background current conductance	0.000367 mS/ μ F
$\bar{g}_{Ca}$	Strength of local LCC calcium flux	9000 mM/(cm C)
g_{K1}	I_{K1} conductance	0.2938 mS/ μ F
g_{NaK}	I_{NaK} conductance	1.716 mS/ μ F
g_{Kss}	I_{Kss} conductance	0.025 mS/ μ F
$g_{to,s}$	$I_{to,s}$ conductance	0 mS/ μ F
$J_{PMCA,max}$	Maximal J_{PMCA} flux	1 pA/pF
K_{PMCA}	Saturation constant for Ca ²⁺ current	0.5 μ M
P_{Ca}	Constant	0.00054 cm/s
$P_{o,max}$	Constant	0.083

*We have reduced this value from the original value of *Mahajan et al. (2008)* so that the Ca²⁺ transient increases when SERCA uptake rate is increased. See Section for details.

Equations for Ca^{2+} cycling

We use the model for Ca^{2+} cycling developed by *Shiferaw et al. (2003)* and subsequently implemented in *Mahajan et al. (2008)*. The equations for Ca^{2+} cycling are

$$\frac{dc_s}{dt} = \beta_s \left[\frac{V_i}{V_s} (J_{rel} - J_d + J_{Ca} + J_{NaCa}) - J_{trpn}^s \right], \quad (2)$$

$$\frac{dc_i}{dt} = \beta_i \left[J_d - J_{up} + J_{leak} - J_{trpn}^i - J_{PMCA} \right], \quad (3)$$

$$\frac{dc_j}{dt} = -J_{rel} + J_{up} - J_{leak}, \quad (4)$$

$$\frac{dc'_j}{dt} = \frac{c_j - c'_j}{\tau_a}, \quad (5)$$

$$J_d = \frac{c_s - c_i}{\tau_s}, \quad (6)$$

$$\frac{dc_p}{dt} = \tilde{J}_{SR} + \tilde{J}_{Ca} - \frac{(c_p - c_s)}{\tau_d}, \quad (7)$$

where the SR leak flux and RyR release flux are given by

$$J_{leak} = g_l (12.4 c_j - c_i), \quad (8)$$

$$\frac{dJ_{rel}}{dt} = N'_s(t) c_j \frac{Q(c'_j)}{c_{sr}} - \frac{J_{rel}}{T}, \quad (9)$$

$$T = \frac{\tau_r}{1 - \tau_r \left(\frac{dc_j}{dt} / c_j \right)} \quad (10)$$

$$Q(c'_j) = \begin{cases} 0 & 0 < c'_j < 50, \\ c'_j - 50 & 50 \leq c'_j \leq c_{sr}, \\ uc'_j + (1 - u)c_{sr} - 50 & c'_j > c_{sr}, \end{cases} \quad (11)$$

$$N'_s = -g_{RyR}(V) P_o i_{Ca}, \quad (12)$$

$$g_{RyR}(V) = g_{RyR} \frac{e^{-0.05(V+30)}}{1 + e^{-0.05(V+30)}}, \quad (13)$$

$$\tilde{J}_{SR} = -g_{SR}(V) Q(c'_j) P_o i_{Ca}, \quad (14)$$

$$g_{SR}(V) = 50 g_{RyR}(V) \quad (15)$$

Intracellular Ca^{2+} buffering

Similarly to *Mahajan et al. (2008)*. All buffering parameters are experimentally based and summarized in *Shannon et al. (2004)*. Buffering to SR, calmodulin, membrane, and sarcolemma binding sites are modeled using the instantaneous buffering approximation given by

$$\beta_i = \left(1 + \frac{B_{SR} K_{SR}}{(c_i + K_{SR})^2} + \frac{B_{cd} K_{cd}}{(c_i + K_{cd})^2} + \frac{B_{mem} K_{mem}}{(c_i + K_{mem})^2} + \frac{B_{sar} K_{sar}}{(c_i + K_{sar})^2} \right)^{-1}, \quad (16)$$

$$\beta_s = \left(1 + \frac{B_{SR} K_{SR}}{(c_s + K_{SR})^2} + \frac{B_{cd} K_{cd}}{(c_s + K_{cd})^2} + \frac{B_{mem} K_{mem}}{(c_s + K_{mem})^2} + \frac{B_{sar} K_{sar}}{(c_s + K_{sar})^2} \right)^{-1} \quad (17)$$

Buffering to Troponin C is given by

$$\frac{d[CaT]_i}{dt} = J_{trpn}^i = k_{on}^T c_i (B_T - [CaT]_i) - k_{off}^T [CaT]_i, \quad (18)$$

$$\frac{d[CaT]_s}{dt} = J_{trpn}^s = k_{on}^T c_s (B_T - [CaT]_s) - k_{off}^T [CaT]_s \quad (19)$$

605 The SERCA uptake pump
606 Similarly to *Shiferaw et al. (2003)*

$$J_{up} = \frac{v_{up} c_i^2}{c_i^2 + c_{up}^2} \quad (20)$$

607 Na^+ dynamics
608 Intracellular Na^+ dynamics are given by

$$\frac{[Na^+]_i}{dt} = 100 \frac{C_m}{F v_i} (I_{Na} + 3I_{NaCa} + 3I_{NaK} + I_{Na,b}) \quad (21)$$

609 In order to reduce computation time, we have sped up the rate at which the system reaches
610 steady-state by increasing $d[Na^+]_i/dt$ by a factor of 100. This will make sodium converge to steady-
611 state on a time-scale fast enough to perform the number of simulations necessary for this study.
612 Once the cell reaches steady-state, $d[Na^+]_i/dt$ is zero, so this modification will not affect the sodium
613 dynamics at steady-state. By doing this, we can save up to 90% of calculation before the system
614 reaches steady-state.

615 Ionic currents

The rate of change of the membrane voltage V is described by the equation

$$\frac{dV}{dt} = -(I_{stim} + I_{Ca,L} + I_{PMCA} + I_{NaCa} + I_{Na} + I_{to,f} + I_{to,s} + I_{Kur} + I_{Kss} + I_{K1} + I_{NaK} + I_{Ca,b} + I_{Na,b}) \quad (22)$$

$$I_{Ca,L} = \frac{-2Fv_i}{C_m} J_{Ca} \quad (23)$$

$$I_{NaCa} = \frac{Fv_i}{C_m} J_{NaCa} \quad (24)$$

$$I_{PMCA} = \frac{-2Fv_i}{C_m} J_{PMCA} \quad (25)$$

616 where I_{stim} is the external stimulus current driving the cell.

617 The L-type Ca current ($I_{Ca,L}$)

Similarly to the *Shannon et al. (2004)* model,

$$J_{Ca} = g_{Ca} P_o i_{Ca} \quad (26)$$

$$\tilde{J}_{Ca} = -\tilde{g}_{Ca} P_o i_{Ca} \quad (27)$$

$$i_{Ca} = \frac{4P_{Ca} V F^2}{RT} \frac{c_s e^{2VF/RT} - 0.341[Ca^{2+}]_o}{e^{2VF/RT} - 1} \quad (28)$$

$$P_o = P_{o,max} \times d \times f \times f_{Ca} \quad (29)$$

$$\frac{df_{Ca}}{dt} = 0.12(1 - f_{Ca}) - \frac{1.4025}{1 + (30/c_p)^4} f_{Ca} \quad (30)$$

$$d_{\infty} = \frac{1}{1 + e^{-(V+4.6)/6.3}} \quad (31)$$

$$\tau_d = d_{\infty} \frac{1 - e^{-(V+4.6)/6.3}}{0.035(V + 4.6)} \quad (32)$$

$$f_{\infty} = 1 - \frac{1}{1 + e^{-(V+22.8)/6.1}} \quad (33)$$

$$\tau_f = \frac{1}{0.02 - 0.007e^{-(0.0337(V+10.5))^2}} \quad (34)$$

$$\frac{dd}{dt} = \frac{d_{\infty} - d}{\tau_d} \quad (35)$$

$$\frac{df}{dt} = \frac{f_{\infty} - f}{\tau_f} \quad (36)$$

618 Calcium background leak ($I_{Ca,b}$)

$$I_{Ca,b} = g_{Ca,b}(V - E_{Ca}) \quad (37)$$

$$E_{Ca} = \frac{RT}{2F} \log \left(\frac{[Ca^{2+}]_o}{c_i} \right) \quad (38)$$

619 The Sarcolemmal Ca^{2+} ATPase (I_{PMCA})

620 The sarcolemmal Ca^{2+} pump (I_{PMCA}) provides another mechanism, in addition to the exchanger
621 (I_{NaCa}), for the extrusion of Ca^{2+} ions out of the cell. This pump is not included in *Mahajan et al.*
622 (2008). We added this current using the formula used by *Bondarenko et al. (2004)*.

$$J_{PMCA} = J_{PMCA,max} \frac{c_i^2}{K_{PMCA}^2 + c_i^2} \quad (39)$$

623 The Na^+ - Ca^{2+} exchange flux ($NaCa$)

Similarly to the *Bondarenko et al. (2004)* model,

$$J_{NaCa} = g_{NaCa} \frac{e^{\xi VF/RT} [Na^+]_i^3 [Ca^{2+}]_o - e^{(\xi-1)VF/RT} [Na^+]_o^3 c_i}{(1 + k_{sat} e^{(\xi-1)VF/RT}) (K_{m,Na}^3 + [Na^+]_o^3) (K_{m,Ca} + [Ca^{2+}]_o)} \quad (40)$$

624 The fast sodium current (I_{Na})

Similarly to the *Shannon et al. (2004)* model,

$$I_{Na} = g_{Na} m^3 h j (V - E_{Na}) \quad (41)$$

$$\frac{dh}{dt} = \alpha_h (1 - h) - \beta_h h \quad (42)$$

$$\frac{dj}{dt} = \alpha_j (1 - j) - \beta_j j \quad (43)$$

$$\frac{dm}{dt} = \alpha_m (1 - m) - \beta_m m \quad (44)$$

$$\alpha_m = 0.32 \frac{V + 47.13}{1 - e^{-0.1(V+47.13)}} \quad (45)$$

$$\beta_m = 0.08 e^{-V/11} \quad (46)$$

For $V \geq -40$ mV,

$$\alpha_h = 0 \quad (47)$$

$$\alpha_j = 0 \quad (48)$$

$$\beta_h = \frac{1}{0.13(1 + e^{(V+10.66)/-11.1})} \quad (49)$$

$$\beta_j = 0.3 \frac{e^{-2.535 \times 10^{-7} V}}{1 + e^{-0.1(V+32)}} \quad (50)$$

For $V \leq -40$ mV,

$$\alpha_h = 0.135 e^{(V+80)/-6.8} \quad (51)$$

$$\beta_h = 3.56 e^{0.079 V} + 3.1 \times 10^5 e^{0.35 V} \quad (52)$$

$$\alpha_j = \frac{(-1.2714 \times 10^5 e^{0.2444 V} - 3.474 \times 10^{-5} e^{-0.04391 V}) \times (V + 37.78)}{1 + e^{0.311(V+79.23)}} \quad (53)$$

$$\beta_j = \frac{0.1212 e^{-0.01052 V}}{1 + e^{-0.1378(V+40.14)}} \quad (54)$$

625 Sodium background leak ($I_{Na,b}$)

$$I_{Na,b} = g_{Na,b}(V - E_{Na}) \quad (55)$$

$$E_{Na} = \frac{RT}{F} \log \left(\frac{[Na^+]_o}{[Na^+]_i} \right) \quad (56)$$

626 Inward rectifier K⁺ current (I_{K1})

Similarly to the *Bondarenko et al. (2004)* model,

$$I_{K1} = g_{K1} \frac{[K^+]_o}{[K^+]_o + 0.21} \left(\frac{V - E_K}{1 + e^{0.0896(V - E_K)}} \right) \quad (57)$$

$$E_K = \frac{RT}{F} \log \left(\frac{[K^+]_o}{[K^+]_i} \right) \quad (58)$$

627 The fast component of the transient outward K⁺ current ($I_{to,f}$)

This current is modified from the formulation of *Bondarenko et al. (2004)* as:

$$I_{to,f} = g_{to,f} a_{to,f}^3 i_{to,f} (V - E_K) \quad (59)$$

$$\frac{da_{to,f}}{dt} = \frac{a_{\infty} - a_{to,f}}{\tau_a} \quad (60)$$

$$\frac{di_{to,f}}{dt} = \frac{i_{\infty} - i_{to,f}}{\tau_i} \quad (61)$$

$$\alpha_a = 0.18264e^{0.03577(V+45)} \quad (62)$$

$$\beta_a = 0.3956e^{0.06237(V+45)} \quad (63)$$

$$\alpha_i = \frac{0.00152e^{-(V+13.5)/7.0}}{0.067083e^{-(V+33.5)/7.0} + 1} \quad (64)$$

$$\beta_i = \frac{0.0095e^{(V+33.5)/7.0}}{0.051335e^{(V+33.5)/7.0} + 1} \quad (65)$$

$$\tau_a = \frac{1}{\alpha_a + \beta_a} \quad (66)$$

$$\tau_i = \frac{1}{\alpha_i + \beta_i} \quad (67)$$

$$a_{\infty} = \frac{\alpha_a}{\alpha_a + \beta_a} \quad (68)$$

$$i_{\infty} = (1 - r) \frac{\alpha_i}{\alpha_i + \beta_i} + r \quad (69)$$

628 where $r = 0.37$ accounts for the presence of a persistent outward potassium current in patch clamp
629 measurements of $I_{to,f}$. We have increased the rates of the inactivation gate (α_i and β_i) from the
630 original formulation to match experimental measurements of $I_{to,f}$ inactivation rate under voltage
631 clamp.

632 The slow component of the transient outward K⁺ current ($I_{to,s}$)

Similarly to the *Bondarenko et al. (2004)* model,

$$I_{to,s} = g_{to,s} a_{to,s} i_{to,s} (V - E_K) \quad (70)$$

$$\frac{da_{to,s}}{dt} = \frac{a_{ss} - a_{to,s}}{\tau_{ta,s}} \quad (71)$$

$$\frac{di_{to,s}}{dt} = \frac{i_{ss} - i_{to,s}}{\tau_{ti,s}} \quad (72)$$

$$a_{ss} = \frac{1}{1 + e^{-(V+22.5)/7.7}} \quad (73)$$

$$i_{ss} = \frac{1}{1 + e^{(V+45.2)/5.7}} \quad (74)$$

$$\tau_{ta,s} = 0.493e^{-0.0629V} + 2.058 \quad (75)$$

$$\tau_{ti,s} = 270.0 + \frac{1050}{1 + e^{(V+45.2)/5.7}} \quad (76)$$

633 The ultra-rapidly activating component of the delayed rectifier K^+ current ($I_{Kur}/I_{K,slow}$)
Similarly to the **Bondarenko et al. (2004)** model,

$$I_{Kur} = g_{Kur} a_{ur} i_{ur} (V - E_K) \quad (77)$$

$$\frac{da_{ur}}{dt} = \frac{a_{ss} - a_{ur}}{\tau_{aur}} \quad (78)$$

$$\frac{di_{ur}}{dt} = \frac{i_{ss} - i_{ur}}{\tau_{iur}} \quad (79)$$

$$\tau_{aur} = 0.493e^{-0.0629V} + 2.058 \quad (80)$$

$$\tau_{iur} = 1200 - \frac{170}{1 + e^{(V+45.2)/5.7}} \quad (81)$$

634 We have reduced the timescale of inactivation (τ_{iur}) from the original formulation to match experi-
635 mental measurements of I_{Kur} inactivation rate under voltage clamp.

636 The Non-inactivating steady-state K^+ current (I_{Kss})
Similarly to the **Bondarenko et al. (2004)** model,

$$I_{Kss} = g_{Kss} a_{Kss} (V - E_K) \quad (82)$$

$$\frac{da_{Kss}}{dt} = \frac{a_{ss} - a_{Kss}}{\tau_{Kss}} \quad (83)$$

$$\tau_{Kss} = 39.3e^{-0.0862V} + 13.17 \quad (84)$$

637 The Na^+ - K^+ pump current (I_{NaK})

Similarly to the **Bondarenko et al. (2004)** model,

$$I_{NaK} = g_{NaK} f_{NaK} \frac{1}{1 + (K_{m,NaK}/[Na^+]_i)^3} \frac{[K^+]_o}{[K^+]_o + K_{m,Ko}} \quad (85)$$

$$f_{NaK} = \frac{1}{1 + 0.1245e^{-0.1VF/RT} + 0.01548767\sigma e^{-VF/RT}} \quad (86)$$

$$\sigma = \frac{1}{7}(e^{[Na^+]_o/67300} - 1) \quad (87)$$

638 Effect of c_s - c_i diffusion rate on how the Ca^{2+} transient depends on SERCA

639 The effect of modifications of the SERCA pump on the steady-state Ca^{2+} transient is shown in Fig.
640 1. While increasing the SERCA's peak uptake current has the effect of sequestering Ca^{2+} back into
641 the SR which would reduce the Ca^{2+} transient, the dominant effect is to increase the Ca^{2+} transient
642 due to higher SR Ca^{2+} load at steady-state. This is consistent with experiments showing restoration
643 of Ca^{2+} transient amplitude when SERCA is upregulated(**del Monte et al., 1999; Del Monte et al.,**
644 **2002).**

645 Initial simulations using the original Ca^{2+} cycling parameters of **Mahajan et al. (2008)** found the
646 SR Ca^{2+} load to decrease as the uptake rate was increased, as more Ca^{2+} was extruded from the
647 submembrane region of the myocyte via NCX before it had an opportunity to diffuse into the c_i
648 compartment to be re-uptaken in into the SR. In order to rectify this, we increased the diffusion
649 rate between the c_s and c_i compartments. We found that when this rate is faster (smaller τ_s) the
650 Ca^{2+} increases with increasing SERCA uptake rate. For this reason we use a value of $\tau_s = 0.75$ ms for
651 all simulations in this study.

652 Computational reproduction of patch clamp experiments

653 In order to compare the GEs found by the computational search to the phenotype variability found
654 in the HMDP, we iterate the model with voltage held constant, reproducing the experimental patch
655 clamp procedure described in the Methods section of the main text. Each model is simulated for 1
656 s with V_m held at -50 mV in order to reach steady-state. V_m is then raised to 0 mV, and peak values
657 $I_{Ca,L}$, $I_{to,f}$ and I_{Kur} are recorded.

Tissue scale modeling

Tissue scale modeling is performed using a 56×56 array of myocytes, each with individual values of ionic conductances. Electrotonic coupling is simulated by introducing a diffusive into the V_m evolution equation,

$$\frac{\partial V_m}{\partial t} = -\frac{1}{C_m}(I_{\text{stim}} + I_{\text{ion}}) + D\nabla^2 V_m. \quad (88)$$

The applied stimulus current occurs at a pacing rate of 4 Hz and is applied to each myocyte simultaneously. The diffusive term is applied isotropically, with diffusive co-efficient, $D = 1 \text{ cm}^2/\text{s}$. In the discretized diffusion equation,

$$V_{i,j}^{t+1} = V_{i,j}^t + dt \left[-\frac{1}{C_m}(I_{\text{stim}} + I_{\text{ion}}) + \frac{D}{\Delta x^2} \left(\sum_{i \pm 1, j \pm 1} V_{i,j}^t - 4V_{i,j}^t \right) \right], \quad (89)$$

We use a lattice size of $\Delta x = 225 \mu\text{m}$, such that the 56×56 lattice represents a $1.25 \text{ cm} \times 1.25 \text{ cm}$ tissue.

GES search

In this study we consider variation in 6 important ionic currents: L-type Ca current ($I_{Ca,L}$), the SR ATPase SERCA, $\text{Na}^+ - \text{Ca}^{2+}$ exchange (NaCa), ryanodine receptor (RyR), the transient outward K^+ current ($I_{to,f}$), and the ultra-rapidly-activating K^+ current (I_{Kur}). The strength of these ionic currents is determined by their conductance g_i . Any given set of parameters ($\mathbf{p} = \{p_1, p_2, \dots, p_n\}$) corresponds to a different candidate myocyte model and produces a different phenotype, which we characterize by quantifiable measurements of its steady-state behaviour (sensors) i.e. steady state calcium transient amplitude, action potential duration and sarcoplasmic reticulum (SR) Ca^{2+} concentration. When stimulating a model with a given period, these parameters (once the simulation has reached steady state) produce a phenotype which we can compare to the phenotype produced by the standard parameters of our model ($\mathbf{p}_{\text{ref}}$).

We define a cost function E that quantifies how much each model's phenotype differs from our reference phenotype as

$$E(\mathbf{p})^2 = \sum_{n=1}^N \left(\frac{S_n(\mathbf{p}) - S_n(\mathbf{p}_{\text{ref}})}{S_n(\mathbf{p}_{\text{ref}})} \right)^2, \quad (90)$$

where the $S_n(\mathbf{p})$ s are sensors characterizing the electrophysiological phenotype of the model's output. $E(\mathbf{p}_{\text{ref}})$ is zero by definition. The three sensors used in this study are listed in table 6.

Table 6. Simulation outputs corresponding to physiological sensors

Abbreviation	Description	Reference Value
$\langle [\text{Ca}]_i \rangle$	Average cytosolic Ca^{2+} over one beat	$0.24 \mu\text{M}$
$\Delta [\text{Ca}]_i$	Ca^{2+} transient amplitude	$0.5 \mu\text{M}$
$[\text{Na}]_i$	Diastolic Na^+	14 mM

For any given set of conductances, a simulation is performed that outputs a value for each sensor. All values are calculated after the simulation is paced with a pacing cycle length PCL = 250 ms for 12.5 s when the system has reached steady state.

We define a *good enough solution* (GES) as a set of conductances with a phenotype such that the value of cost function $E(\mathbf{p})$ is less than a threshold $\epsilon = 0.05$. None of the cost function sensors S_n are based on the membrane potential, and therefore a GES does not necessarily have an action potential shape close to the reference action potential shape. A GES is required to achieve steady-state. Therefore parameters that produce parameter sets that do not reach a steady state during pacing at constant cycle length, such as those which exhibit calcium transient alternans, are not considered GESs. We additionally reject any set of parameters for which the output steady-state

SR Ca^{2+} load is above a threshold $[\text{Ca}^{2+}]_{\text{SR}} > 130 \mu\text{M}_{\text{Cyt}}$, which we consider to be unphysiologically overloaded.

Completing an exhaustive search of the parameter space becomes increasingly computationally intensive as the number of parameters grows. An exhaustive search of an M dimensional space, considering K possible values for each conductance requires K^M evaluations of the cost function. Additionally, as the number of sensors (N) that we use to calculate the cost function increases, the fraction of models tested that are good enough solutions (Φ) will decrease. We calculated $E(\mathbf{p})$ for 10^7 random parameter sets, $\mathbf{p}$, and found only 11 which were GESs ($E(\mathbf{p}) < 0.05$). It is therefore apparent that an exhaustive search is not efficient for finding GES in high dimensional parameter space and for this reason we use a minimization scheme to find GESs. We start by randomly assigning values to each parameter such that $0 < p_i < 3p_{i,\text{ref}}$ for each i in $\mathbf{p}$, and then minimize them with respect to the cost function, $E(\mathbf{p})$, using the Nelder-Mead simplex algorithm **Nelder and Mead (1964)** (also known as the Amoeba algorithm) until $E(\mathbf{p}) < \epsilon$. Running this procedure 10,000 times yields 7263 GESs, with the remaining trials being rejected either because the minimization algorithm does not converge, the SR load constraint is not satisfied, or the system is found to not be in steady-state (determined by comparing $\Delta[\text{Ca}]_i$ of the 50th and 51st beat).

The Nelder-Mead Simplex Algorithm

The Nelder-Mead algorithm (**Nelder and Mead, 1964**) maintains a non-degenerate simplex at each iteration, a geometric figure in n dimensions of nonzero volume that is the convex hull of $n + 1$ vertices, $\vec{x}_0, \vec{x}_1, \dots, \vec{x}_n$, and their respective function values. Suppose we start from the vector $\vec{x}_0$, the simplex can be initialized as $\vec{x}_i = \vec{x}_0 + \delta \vec{e}_i$, where $\vec{e}_i$ is a unit vector, and where δ is our guess of the problem's characteristic length scale. In each iteration, new points are computed, along with their function values, to form a new simplex. The algorithm terminates when the function values at the vertices of the simplex satisfy a predetermined condition. One iteration of the Amoeba algorithm consists of the following steps (the standard values for the coefficients are: $\alpha = 1, \beta = 2, \gamma = 0.5, \sigma = 0.5$):

1. Order: order and re-label the $n + 1$ vertices as $\vec{x}_0, \vec{x}_1, \dots, \vec{x}_n$, such that $F(\vec{x}_0) \leq F(\vec{x}_1) \leq \dots \leq F(\vec{x}_n)$. Since we want to minimize $F(\vec{x}_0)$, we refer to $\vec{x}_0$ as the best point, to $\vec{x}_n$ as the worst point, and to $\vec{x}_{n-1}$ as the next worst point. Let $\vec{x}_c$ refer to the centroid of the n points in the vertex.
$$\vec{x}_c = \sum_{i=0}^{n-1} \vec{x}_i / n.$$
2. Reflect: compute the reflected point, $\vec{x}_r = \vec{x}_c + \alpha(\vec{x}_c - \vec{x}_n)$. Evaluate $F(\vec{x}_r)$. If $F(\vec{x}_0) \leq F(\vec{x}_r) < F(\vec{x}_{n-1})$, then obtain a new simplex by replacing the worst point $\vec{x}_n$ with the reflected point $\vec{x}_r$ and go to step 1.
3. Expand: if $F(\vec{x}_r) < F(\vec{x}_0)$, compute the expanded point, $\vec{x}_e = \vec{x}_c + \beta(\vec{x}_r - \vec{x}_c)$. If $F(\vec{x}_e) < F(\vec{x}_r)$, then obtain a new simplex by replacing the worst point $\vec{x}_n$ with the expanded point $\vec{x}_e$ and go to step 1; otherwise then obtain a new simplex by replacing the worst point $\vec{x}_n$ with the reflected point $\vec{x}_r$ and go to step 1.
4. Contract: At this step, where it is certain that $F(\vec{x}_r) > F(\vec{x}_{n-1})$, compute the contracted point $\vec{x}_{con} = \vec{x}_c + \gamma(\vec{x}_n - \vec{x}_c)$. If $F(\vec{x}_{con}) \leq F(\vec{x}_n)$, obtain a new simplex by replacing the worst point $\vec{x}_n$ with the expansion point $\vec{x}_{con}$ and go to step 1.
5. Shrink: replace all vertices except the best $\vec{x}_0$ with $\vec{x}_i = \vec{x}_0 + \sigma(\vec{x}_i - \vec{x}_0)$ and go to step 1.

Two sensor search only constraining the Ca^{2+} transient

The results of the GES search described in the main text were reproduced using only 2 sensors, constraining $\langle [\text{Ca}]_i \rangle$ and $\Delta[\text{Ca}]_i$ but not constraining $[\text{Na}]_i$. This results in a broader histogram for the I_{NaCa} conductance (**Figure 3-Figure Supplement 1** compared to Fig. 3).

Acknowledgments

This research was supported by NIH/NHLBI grant 5R01HL114437 and by the Laubisch and Kawata Endowments. The authors thank Yibin Wang for valuable input into this study. A.K. acknowledges stimulating discussions with Eve Marder in the early stage of this work.

References

- Balke CW**, Wier WG. Modulation of L-type calcium channels by sodium ions. *Proceedings of the National Academy of Sciences*. 1992; 89(10):4417–4421.
- Banyasz T**, Horvath B, Jian Z, Izu LT, Chen-Izu Y. Sequential dissection of multiple ionic currents in single cardiac myocytes under action potential-clamp. *Journal of molecular and cellular cardiology*. 2011; 50(3):578–581.
- Bers D**. Excitation-contraction coupling and cardiac contractile force, vol. 237. Springer Science & Business Media; 2001.
- Bers DM**. Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol*. 2008; 70:23–49.
- Bondarenko VE**, Szigeti GP, Bett GC, Kim SJ, Rasmusson RL. Computer model of action potential of mouse ventricular myocytes. *American Journal of Physiology-Heart and Circulatory Physiology*. 2004; 287(3):H1378–H1403.
- Britton OJ**, Bueno-Orovio A, Van Ammel K, Lu HR, Towart R, Gallacher DJ, Rodriguez B. Experimentally calibrated population of models predicts and explains intersubject variability in cardiac cellular electrophysiology. *Proceedings of the National Academy of Sciences*. 2013; 110(23):E2098–E2105.
- Daniels BC**, Chen YJ, Sethna JP, Gutenkunst RN, Myers CR. Sloppiness, robustness, and evolvability in systems biology. *Current opinion in biotechnology*. 2008; 19(4):389–395.
- Del Monte F**, Harding SE, Dec GW, Gwathmey JK, Hajjar RJ. Targeting phospholamban by gene transfer in human heart failure. *Circulation*. 2002 feb; 105(8):904–907. doi: [10.1161/hc0802.105564](https://doi.org/10.1161/hc0802.105564).
- Delbridge L**, Satoh H, Yuan W, Bassani J, Qi M, Ginsburg KS, Samarel AM, Bers DM. Cardiac myocyte volume, Ca²⁺ fluxes, and sarcoplasmic reticulum loading in pressure-overload hypertrophy. *American Journal of Physiology-Heart and Circulatory Physiology*. 1997; 272(5):H2425–H2435.
- Fox JJ**, McHarg JL, Gilmour RF. Ionic mechanism of electrical alternans. *American Journal of Physiology-Heart and Circulatory Physiology*. 2002; 282(2):H516–H530.
- Ghazalpour A**, Rau CD, Farber CR, Bennett BJ, Orozco LD, Van Nas A, Pan C, Allayee H, Beaven SW, Civelek M, et al. Hybrid mouse diversity panel: a panel of inbred mouse strains suitable for analysis of complex genetic traits. *Mammalian genome*. 2012; 23(9-10):680–692.
- Golowasch J**, Abbott L, Marder E. Activity-dependent regulation of potassium currents in an identified neuron of the stomatogastric ganglion of the crab *Cancer borealis*. *Journal of Neuroscience*. 1999; 19:RC33–1.
- Grashow R**, Brookings T, Marder E. Reliable neuromodulation from circuits with variable underlying structure. *Proceedings of the National Academy of Sciences*. 2009; 106(28):11742–11746.
- Groenendaal W**, Ortega FA, Kherlopian AR, Zygmunt AC, Krogh-Madsen T, Christini DJ. Cell-specific cardiac electrophysiology models. *PLoS Comput Biol*. 2015; 11(4):e1004242.
- Gutenkunst RN**, Waterfall JJ, Casey FP, Brown KS, Myers CR, Sethna JP. Universally sloppy parameter sensitivities in systems biology models. *PLoS computational biology*. 2007; 3(10):e189.
- Harvey R**, Jurevicius J, Hume J. Intracellular Na⁺ modulates the cAMP-dependent regulation of ion channels in the heart. *Proceedings of the National Academy of Sciences*. 1991; 88(16):6946–6950.
- Hodgkin AL**, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. *The Journal of physiology*. 1952; 117(4):500.
- Jin H**, Hadri L, Palomeque J, Morel C, Karakikes I, Kaprielian R, Hajjar R, Lebeche D. KCHIP2 attenuates cardiac hypertrophy through regulation of Ito and intracellular calcium signaling. *Journal of molecular and cellular cardiology*. 2010; 48(6):1169–1179.
- Karma A**. Physics of cardiac arrhythmogenesis. *Annu Rev Condens Matter Phys*. 2013; 4(1):313–337.

- 780 **Krogh-Madsen T**, Christini DJ. Nonlinear dynamics in cardiology. Annual review of biomedical engineering.
781 2012; 14:179–203.
- 782 **Krogh-Madsen T**, Sobie EA, Christini DJ. Improving cardiomyocyte model fidelity and utility via dynamic
783 electrophysiology protocols and optimization algorithms. The Journal of physiology. 2016; .
- 784 **Kuo HC**, Cheng CF, Clark RB, Lin JJC, Lin JLC, Hoshijima M, Nguyen-Tran VT, Gu Y, Ikeda Y, Chu PH, et al. A defect
785 in the Kv channel-interacting protein 2 (KCHIP2) gene leads to a complete loss of Ito and confers susceptibility
786 to ventricular tachycardia. Cell. 2001; 107(6):801–813.
- 787 **LeMasson G**, Marder E, Abbott L. Activity-dependent regulation of conductances in model neurons. SCIENCE-
788 NEW YORK THEN WASHINGTON-. 1993; 259:1915–1915.
- 789 **Liu Z**, Golowasch J, Marder E, Abbott L. A model neuron with activity-dependent conductances regulated by
790 multiple calcium sensors. Journal of Neuroscience. 1998; 18(7):2309–2320.
- 791 **Luo Ch**, Rudy Y. A model of the ventricular cardiac action potential. Depolarization, repolarization, and their
792 interaction. Circulation research. 1991; 68(6):1501–1526.
- 793 **Mahajan A**, Shiferaw Y, Sato D, Baher A, Olcese R, Xie LH, Yang MJ, Chen PS, Restrepo JG, Karma A, et al. A rabbit
794 ventricular action potential model replicating cardiac dynamics at rapid heart rates. Biophysical journal. 2008;
795 94(2):392–410.
- 796 **Marder E**. Variability, compensation, and modulation in neurons and circuits. Proceedings of the National
797 Academy of Sciences. 2011; 108(Supplement 3):15542–15548.
- 798 **Marder E**, Goaillard JM. Variability, compensation and homeostasis in neuron and network function. Nature
799 Reviews Neuroscience. 2006; 7(7):563–574.
- 800 **del Monte F**, Harding SE, Schmidt U, Matsui T, Kang ZB, Dec GW, Gwathmey JK, Rosenzweig A, Hajjar RJ.
801 Restoration of Contractile Function in Isolated Cardiomyocytes From Failing Human Hearts by Gene Transfer
802 of SERCA2a. Circulation. 1999 dec; 100(23):2308–2311. doi: [10.1161/01.CIR.100.23.2308](https://doi.org/10.1161/01.CIR.100.23.2308).
- 803 **Muszkiewicz A**, Britton OJ, Gemmell P, Passini E, Sánchez C, Zhou X, Carusi A, Quinn TA, Burrage K, Bueno-Orovio
804 A, et al. Variability in cardiac electrophysiology: using experimentally-calibrated populations of models to
805 move beyond the single virtual physiological human paradigm. Progress in biophysics and molecular biology.
806 2016; 120(1):115–127.
- 807 **Nelder BJA**, Mead R. A simplex method for function minimization. The Computer Journal. 1964; 7(4):308–313.
808 doi: [10.1093/comjnl/7.4.308](https://doi.org/10.1093/comjnl/7.4.308).
- 809 **O’Leary T**, Williams AH, Caplan JS, Marder E. Correlations in ion channel expression emerge from homeostatic
810 tuning rules. Proceedings of the National Academy of Sciences. 2013; 110(28):E2645–E2654.
- 811 **Prinz AA**, Bucher D, Marder E. Similar network activity from disparate circuit parameters. Nature neuroscience.
812 2004; 7(12):1345–1352.
- 813 **Qi XY**, Yeh YH, Xiao L, Burstein B, Maguy A, Chartier D, Villeneuve LR, Brundel BJ, Dobrev D, Nattel S. Cellular
814 signaling underlying atrial tachycardia remodeling of L-type calcium current. Circulation research. 2008;
815 103(8):845–854.
- 816 **Qu Z**, Hu G, Garfinkel A, Weiss JN. Nonlinear and stochastic dynamics in the heart. Physics reports. 2014;
817 543(2):61–162.
- 818 **Rau CD**, Romay MC, Tuteryan M, Wang JJC, Santolini M, Ren S, Karma A, Weiss JN, Wang Y, Lusis AJ. Systems genet-
819 ics approach identifies gene pathways and Adamts2 as drivers of isoproterenol-induced cardiac hypertrophy
820 and cardiomyopathy in mice. Cell systems. 2017; 4(1):121–128.
- 821 **Rau CD**, Wang J, Avetisyan R, Romay M, Martin L, Ren S, Wang Y, Lusis AJ. Mapping genetic contributions to
822 cardiac pathology induced by Beta-adrenergic stimulation in mice. Circulation: Cardiovascular Genetics. 2014;
823 p. CIRCGENETICS–113.
- 824 **Rosati B**, McKinnon D. Regulation of ion channel expression. Circulation Research. 2004; 94(7):874–883.
- 825 **Santolini M**, Romay MC, Yukhtman CL, Rau CD, Ren S, Saucerman JJ, Wang JJ, Weiss JN, Wang Y, Lusis AJ, et al. A
826 personalized, multiomics approach identifies genes involved in cardiac hypertrophy and heart failure. npj
827 Systems Biology and Applications. 2018; 4(1):12.

828 **Sarkar A**, Sobie EA. Regression Analysis for Constraining Free Parameters in Electrophysiological Models of
829 Ventricular Cells. *Biophysical Journal*. 2009; 96(3):667a.

830 **Sarkar AX**, Christini DJ, Sobie EA. Exploiting mathematical models to illuminate electrophysiological variability
831 between individuals. *The Journal of physiology*. 2012; 590(11):2555–2567.

832 **Sarkar AX**, Sobie EA. Regression analysis for constraining free parameters in electrophysiological models of
833 cardiac cells. *PLoS Comput Biol*. 2010; 6(9):e1000914.

834 **Schulz DJ**, Goaillard JM, Marder E. Variable channel expression in identified single and electrically coupled
835 neurons in different animals. *Nature neuroscience*. 2006; 9(3):356–362.

836 **Sdek P**, Zhao P, Wang Y, Huang Cj, Ko CY, Butler PC, Weiss JN, MacLellan WR. Rb and p130 control cell cycle
837 gene silencing to maintain the postmitotic phenotype in cardiac myocytes. *The Journal of cell biology*. 2011;
838 194(3):407–423.

839 **Shannon TR**, Wang F, Puglisi J, Weber C, Bers DM. A mathematical treatment of integrated Ca dynamics within
840 the ventricular myocyte. *Biophysical journal*. 2004; 87(5):3351–3371.

841 **Shiferaw Y**, Watanabe M, Garfinkel A, Weiss J, Karma A. Model of intracellular calcium cycling in ventricular
842 myocytes. *Biophysical journal*. 2003; 85(6):3666–3686.

843 **Siegel M**, Marder E, Abbott L. Activity-dependent current distributions in model neurons. *Proceedings of the*
844 *National Academy of Sciences*. 1994; 91(24):11308–11312.

845 **Transtrum MK**, Machta BB, Brown KS, Daniels BC, Myers CR, Sethna JP. Perspective: Sloppiness and emergent
846 theories in physics, biology, and beyond. *The Journal of chemical physics*. 2015; 143(1):07B201_1.

847 **Wang JJC**, Rau C, Avetisyan R, Ren S, Romay MC, Stolin G, Gong KW, Wang Y, Lusis AJ. Genetic dissection of cardiac
848 remodeling in an isoproterenol-induced heart failure mouse model. *PLoS genetics*. 2016; 12(7):e1006038.

849 **Weiss JN**, Karma A, MacLellan WR, Deng M, Rau CD, Rees CM, Wang J, Wisniewski N, Eskin E, Horvath S, et al.
850 “Good enough solutions” and the genetics of complex diseases. *Circulation research*. 2012; 111(4):493–504.

851 **Yang JH**, Yang L, Qu Z, Weiss JN. Glycolytic oscillations in isolated rabbit ventricular myocytes. *Journal of*
852 *Biological Chemistry*. 2008; 283(52):36321–36327.

853 **Zhou J**, Jeron A, London B, Han X, Koren G. Characterization of a slowly inactivating outward current in adult
854 mouse ventricular myocytes. *Circulation Research*. 1998; 83(8):806–814.

Figure 3-Figure supplement 1. Histograms of individual ion channel conductances in 8320 GESs found by a GES search constrained only by Ca^{2+} transient amplitude and average, but not constrained by $[\text{Na}]_i$.

Figure 3-Figure supplement 2. Correlation between $I_{\text{Ca,L}}$ and the sum of $I_{\text{to,f}}$ and I_{Kur} is not as strong when $[\text{Na}]_i$ is not constrained.

Figure 4-Figure supplement 1. Patch clamp measurements of mean $I_{\text{Ca,L}}$ (A), $I_{\text{to,f}}$ (B), and I_{Kur} (C) functional current density averaged over multiple cells for 9 HMDP mouse strains with standard errors (thick bars) and standard deviations (thin bars).

Figure 5-Figure supplement 1. Action potential recordings from isolated myocytes for mouse strain C57BL/6J paced at 4 Hz under current clamp. The recordings illustrate the typical degree of cell-to-cell variability of AP morphology observed in all strains.