Individual repeatability and heritability of divorce in a wild population

Ryan R. Germain^{1,2†}, Matthew E. Wolak^{1,3†}, and Jane M. Reid¹

¹School of Biological Sciences, University of Aberdeen, Aberdeen, UK

Current addresses:

²Lab of Ornithology, Cornell University, Ithaca, NY, USA

³Department of Biological Sciences, Auburn University, Auburn, AL, USA

[†]Equal author contributions

Authors for correspondence:

Ryan Germain

email: ryan.r.germain@gmail.com

Matthew Wolak

email: matthew.wolak@auburn.edu

Subject Areas:

behaviour, evolution, ecology

Keywords: indirect genetic effects, mating system evolution, social monogamy, quantitative genetics

ABSTRACT

Understanding micro-evolutionary responses of mating systems to contemporary selection requires estimating sex-specific additive genetic variances and cross-sex genetic covariances in key reproductive strategy traits. One key trait comprises the occurrence of divorce versus mate-fidelity across sequential reproductive attempts. If divorce represents an evolving behavioural strategy it must have non-zero individual repeatability and heritability, but quantitative estimates from wild populations are scarce. We used 39 years of individual breeding records and pedigree data from free-living song sparrows (*Melospiza melodia*) to quantify sex-specific permanent individual and additive genetic variances, and hence estimate repeatability and heritability, in liability for divorce. We estimated moderate repeatability in females, but little repeatability in males. Estimates of additive genetic variance were small in both sexes, and the cross-sex genetic covariance was close to zero. However, the total heritability was likely non-zero but small, indicating low potential for micro-evolutionary response to selection. Rapid micro-evolution of divorce rate therefore appears unlikely, even if there were substantial fitness benefits of divorce and resulting selection.

INTRODUCTION

Pair-bond resilience, resulting from mate-fidelity (i.e. maintaining pair-bonds over multiple breeding attempts) versus divorce (i.e. dissolving pair-bonds to re-pair with a new mate) [1,2], is a key feature of animal mating systems that results from sexual selection and can fundamentally influence the distribution of offspring across parents and the resulting population-wide distribution of fitness. Divorce occurs in numerous socially-monogamous taxa and may increase individual fitness by counter-acting constraints on initial mate choice, and hence be adaptive [2,3]. However, if there is to be ongoing adaptive micro-evolution of divorce rate in response to selection [2,4–6] then individual liability for divorce must have non-zero heritability, and hence vary consistently among individuals (i.e. have non-zero individual repeatability).

In general, repeatable expression of mating and reproductive traits implies that selection may act consistently on individuals over their lifetimes, and defines the maximum possible heritability [7,8]. Decomposing total phenotypic variance into repeatable individual variance (V_I), additive genetic variance (V_A) and environmental variance allows estimation of heritability and, in principle, indicates the potential for micro-evolutionary responses to contemporary selection [9]. However, divorce represents an interesting class of jointly-expressed (i.e. ‘emergent’) traits that result from female-male interactions, and hence can be simultaneously influenced by genetic effects of both sexes. Further, correlated expression of sex-specific genetic effects, leading to non-zero cross-sex genetic covariance ($COV_{A\text{♀♂}}$), can generate evolutionary sexual conflict [10] and further alter the potential for evolutionary responses to selection on either or both sexes that could result from sex-specific fitness consequences of key mating decisions (e.g. [11]). Understanding the evolutionary dynamics of divorce, and the implications for other co-evolving reproductive behaviours contributing to mating systems, therefore requires

explicit estimation of sex-specific V_A ($V_{A\text{♀}}$ and $V_{A\text{♂}}$), and $\text{COV}_{A\text{♀♂}}$, in divorce in populations experiencing un-manipulated natural and sexual selection environments.

Advances in quantitative genetic methods mean that $V_{A\text{♀}}$, $V_{A\text{♂}}$, and $\text{COV}_{A\text{♀♂}}$ underlying emergent traits can be estimated given complex relatedness structures arising in wild populations (e.g. [9]). Since divorce versus mate-fidelity represent alternative outcomes of pairing decisions across consecutive breeding attempts, divorce is appropriately modelled as a ‘threshold trait’, where breeding pairs’ underlying continuous joint liabilities for divorce translate into expression at some threshold (e.g. [12]). Such models also permit estimation of ‘total heritability’ of divorce, resulting from the combination of $V_{A\text{♀}}$, $V_{A\text{♂}}$, and $\text{COV}_{A\text{♀♂}}$ [13], which represents the overall potential for micro-evolutionary responses to selection on the population-wide distribution of liabilities. Such analyses require phenotypic observations of divorce versus mate-fidelity, conditional on survival between consecutive breeding attempts, from diverse relatives [14]. We use 39 years of comprehensive observations of free-living song sparrows (*Melospiza melodia*) to estimate (i) population-level divorce rate, (ii) $V_{A\text{♀}}$, $V_{A\text{♂}}$ and $\text{COV}_{A\text{♀♂}}$ in divorce, and (iii) individual repeatability and the sex-specific and total heritability, thereby assessing the potential for ongoing evolution of divorce as a key reproductive strategy.

MATERIAL AND METHODS

A resident population of individually colour-ringed song sparrows on Mandarte Island, Canada has been intensively studied since 1975. Each year, all individuals alive in late April (typical start of breeding [15]) are recorded in a comprehensive census to determine over-winter survival and pairing status (re-sighting probability >0.99 [16]), and all breeding attempts are closely

monitored (ESMS1). Females and males form socially persistent pairings that cooperate to rear offspring (1–4 broods per year), but can form new pairings within and between breeding seasons following divorce or mate-death.

To identify cases of divorce versus mate-fidelity, we extracted each female’s lifetime sequence of breeding events (≥ 1 egg laid) where re-pairing could have occurred (i.e. she initiated a subsequent breeding event) and categorized these events as divorce, mate-fidelity or mate-death according to the pair-bond’s fate and whether her current mate was still alive during her subsequent event (ESMS1). Instances of mate-death were identified from daily field observations and April censuses, and excluded from our dataset.

We fitted two generalized linear mixed models to decompose total variance in liability for divorce. Model 1 estimated variances attributable to permanent effects of individual females ($V_{I\text{♀}}$), males ($V_{I\text{♂}}$) and of unique female-male social pairings (V_S), and the year when a focal breeding event occurred (V_Y). Model 2 additionally estimated $V_{A\text{♀}}$, $V_{A\text{♂}}$, and $\text{COV}_{A\text{♀♂}}$ in liability for divorce, using comprehensive pedigree data to quantify relatedness (i.e. an ‘animal model’ [9]; ESMS2). $\text{COV}_{A\text{♀♂}}$ is the covariance between additive genetic effects of alleles expressed in all females versus all males, not the genetic covariance between a female and her socially-paired mate (e.g. [11,15]; ESMS3). Our current aim was to partition the total naturally-occurring variation in liability for divorce and thereby appropriately estimate heritability, not to explain variation in the occurrence of divorce. Fixed effects were consequently restricted to a two-level factor that defined whether an observation of divorce versus mate-fidelity spanned breeding events separated by the non-breeding season (‘between-season’) versus consecutive events within the same season (‘within-season’, ESMS1) and separate regressions on female and male

individual coefficients of inbreeding (f) (Model 2 only), thereby estimating inbreeding depression in liability for divorce and minimising any associated bias in estimated V_A [17]. Sex-specific repeatabilities in liability for divorce were estimated from Model 1 as:

$$Repeatability_{\varphi} = \frac{V_{I\varphi}}{V_{P_Model1}} \text{ and } Repeatability_{\sigma} = \frac{V_{I\sigma}}{V_{P_Model1}}$$

Sex-specific heritabilities (h^2) were estimated from Model 2 as:

$$h^2_{\varphi} = \frac{V_{A\varphi}}{V_{P_Model2}} \text{ and } h^2_{\sigma} = \frac{V_{A\sigma}}{V_{P_Model2}}$$

The 'total heritability' (T^2) was calculated from Model 2 as:

$$T^2 = \frac{V_{A\varphi} + V_{A\sigma} + 2(COV_{A\varphi\sigma})}{V_{P_Model2}}$$

The total variance in liability (V_{P_Model1} , V_{P_Model2}) was calculated separately for each model (ESMS3). Models were fitted using Bayesian inference in R, using relatively uninformative priors (ESMS5). Posterior distributions of repeatabilities, heritabilities, and T^2 were calculated from marginal posterior distributions of underlying variance components. We report liability-scale posterior modes, means and 95% credible intervals (95%CI) from 5000 posterior samples, and present prior distributions alongside posteriors to facilitate inference. Estimates and conclusions were robust to alternative model specifications (ESMS5).

RESULTS

There were 1,419 breeding events where divorce could have occurred, involving 566 unique social pairings among 358 females and 341 males. Divorce occurred on 166 (11.7%) occasions (details in ESMS1).

Model 1 revealed that the largest component of variance in liability for divorce was $V_{I\text{♀}}$, while $V_{I\text{♂}}$ was comparatively small (table 1, ESMS4). Because V_S and V_Y were also small, female repeatability for divorce was moderate (~16%; table 1). The lower 95%CI limit converged towards zero, but 98% of posterior samples exceeded 0.01, departing from the prior distribution (figure 1a), and indicating that female repeatability is most likely greater than zero. In contrast, male repeatability was smaller (table 1); only 76% of posterior samples exceeded 0.01 (figure 1b). Divorce was less likely to occur within a breeding season than between seasons (table 1).

Model 2 showed that $V_{A\text{♀}}$ and $V_{A\text{♂}}$ in liability for divorce were both small, and $\text{COV}_{A\text{♀♂}}$ was estimated as close to zero (table 1, ESMS4). Sex-specific heritabilities were therefore small for both females and males. However, posterior distributions departed from the priors, with higher density at higher values (e.g. at the posterior means, figure 2a,b) implying that heritabilities, and underlying $V_{A\text{♀}}$ and $V_{A\text{♂}}$, exceed zero. Indeed, $V_{A\text{♀}}$, $V_{A\text{♂}}$, and $\text{COV}_{A\text{♀♂}}$ combined to generate a small but likely non-zero total heritability (T^2) for divorce (table 1); 92% of posterior samples exceeded 0.01 (ESMS5). This again deviates from the prior distribution (figure 2c), and from the posterior distribution that would have resulted given zero $V_{A\text{♀}}$, $V_{A\text{♂}}$ and hence T^2 (ESMS5). Liability for divorce tended to increase with increasing f , especially in females, but the 95% CIs overlapped zero (table 1).

146

147 **DISCUSSION**

148 The ~12% divorce rate observed in song sparrows is relatively low compared to other temperate-
 149 breeding passerine birds (~20–50%; [14,18,19]). However, there was evidence of moderate $V_{I\text{♀}}$
 150 and hence female repeatability, but lower $V_{I\text{♂}}$ and male repeatability, in liability for divorce
 151 (figure 1a,b, ESMS4). These estimates imply that sex-specific h^2 is not *a priori* zero. However in
 152 practice, h^2 was estimated to be small in both sexes.

153 Most previous quantitative genetic analyses of divorce come from human twin-studies,
 154 and show relatively high heritabilities with divorce defined as a sex-specific trait (e.g. 0.3–0.6
 155 [20,21]). However, such estimates may be inflated by shared environmental and cultural effects
 156 [9], and often only consider whether individuals ever divorced over their lifetime. Our focus on
 157 sequential breeding events, considering among-individual variances across repeat observations,
 158 allows estimation of individual repeatability as well as pair and year variances, which encompass
 159 variances stemming from ecological and/or social environmental effects that could influence the
 160 occurrence of divorce. The only previous quantitative genetic analysis of divorce in a wild (non-
 161 human) population also estimated low female heritability in savannah sparrows (*Passerculus*
 162 *sandwichensis*) [14]. Because [14]’s estimate of male repeatability for divorce was not
 163 distinguishable from zero, heritability estimates were restricted to females and did not consider
 164 potential contributions of $V_{A\text{♂}}$ or $\text{COV}_{A\text{♀♂}}$. Our results suggest that both song sparrow sexes do
 165 contribute to the total additive genetic variance, and hence to the total heritability for divorce
 166 (T^2). Indeed small, potentially undetectable, effects in each sex can combine to generate
 167 detectable total T^2 [13]. Thus the overall apparent potential for micro-evolutionary responses to

selection on divorce is greater when considering the interactive effects of the sexes jointly (i.e. considering divorce as an emergent trait) than when considering either sex alone.

Many studies have investigated the potential costs and benefits of divorce in wild populations, particularly in socially-monogamous birds (reviews: [2,3,5]). Divorce is generally considered to be adaptive, and increases an individual's subsequent breeding success and hence fitness under certain conditions [2,3]. However, ongoing micro-evolutionary responses to such contemporary selection require V_A . Our results indicate that such V_A , and consequent potential for evolutionary response to selection, while probably non-zero, is smaller than is often implicitly assumed [2,4–6]. Further, relatively low divorce rates, such as observed in song sparrows, will intrinsically limit the intensity of selection [12]. Overall, rapid and marked micro-evolutionary changes in the frequency of divorce appear unlikely, even if divorce were beneficial for one or both members of a breeding pair. Consequently, there is also limited potential for genetic covariance between liability for divorce and other key reproductive strategy traits, such as female extra-pair reproduction (EPR), negating the suggestion that divorce and EPR both represent manifestations of an underlying 'weak-pair' syndrome [22] and limiting the potential for indirect selection.

Ethics

All field procedures approved by the University of British Columbia Animal Care Committee ethical review (certificates UBCACC A12-0229 and A14-0336).

189 **Data accessibility**

190 Data and code are available as electronic supporting information.

191

192 **Authors' contributions**

193 RRG and MEW designed the study and conducted analyses with input from JMR. All authors
194 contributed to writing, agree to be held accountable for all aspects of the work, and approve the
195 final version of the manuscript.

196

197 **Competing interests**

198 We declare no competing interests.

199

200 **Funding**

201 All authors were supported by a European Research Council grant to JMR. Fieldwork was
202 supported by the Natural Sciences and Engineering Research Council of Canada and the
203 University of British Columbia.

204

205 **Acknowledgments**

We thank the Tsawout and Tseycum First Nation bands for access to Mandarte, Peter Arcese, Lukas Keller, Pirmin Nietlisbach, and the University of Aberdeen Maxwell High Performance Computing cluster.

References

1. Black JM. 1996 *Partnerships in Birds - The Study of Monogamy*. Oxford: Oxford University Press.
2. Culina A, Radersma R, Sheldon BC. 2015 Trading up: the fitness consequences of divorce in monogamous birds. *Biol. Rev.* **90**, 1015–1034. (doi:10.1111/brev.12143)
3. Dubois F, Cézilly F. 2002 Breeding success and mate retention in birds: a meta-analysis. *Behav. Ecol. Sociobiol.* **52**, 357–364. (doi:10.1007/s00265-002-0521-z)
4. Diamond JM. 1987 A darwinian theory of divorce. *Nature* **329**, 765–766.
5. Choudhury S. 1995 Divorce in birds: a review of the hypotheses. *Anim. Behav.* **50**, 413–429. (doi:10.1006/anbe.1995.0256)
6. McNamara JM, Forslund P. 1996 Divorce rates in birds: predictions from an optimization model. *Am. Nat.* **147**, 609–640. (doi:10.1086/285869)
7. Boake CRB. 1989 Repeatability: Its role in evolutionary studies of mating behavior. *Evol. Ecol.* **3**, 173–182. (doi:10.1007/BF02270919)
8. Dingemanse NJ, Dochtermann NA. 2013 Quantifying individual variation in behaviour: mixed-effect modelling approaches. *J. Anim. Ecol.* **82**, 39–54. (doi:10.1111/1365-2656.12013)
9. Kruuk LEB. 2004 Estimating genetic parameters in natural populations using the ‘animal

model'. *Philos. Trans. R. Soc. London B* **359**, 873–890. (doi:10.1098/rstb.2003.1437)

10. Bonduriansky R, Chenoweth SF. 2009 Intralocus sexual conflict. *Trends Ecol. Evol.* **24**, 280–288. (doi:10.1016/J.TREE.2008.12.005)

11. Wolak ME, Reid JM. 2016 Is pairing with a relative heritable? Estimating female and male genetic contributions to the degree of biparental inbreeding in song sparrows (*Melospiza melodia*). *Am. Nat.* **187**, 736–752. (doi:10.1086/686198)

12. Roff DA. 1996 The evolution of threshold traits in animals. *Q. Rev. Biol.* **71**, 3–35. (doi:10.1086/419266)

13. Bijma P, Muir WM, Van Arendonk JAM. 2007 Multilevel selection 1: quantitative genetics of inheritance and response to selection. *Genetics* **175**, 277–288. (doi:10.1534/genetics.106.062711)

14. Wheelwright NT, Teplitsky C. 2017 Divorce in an island bird population: causes, consequences, and lack of inheritance. *Am. Nat.* **190**, 557–569. (doi:10.1086/693387)

15. Germain RR, Wolak ME, Arcese P, Losdat S, Reid JM. 2016 Direct and indirect genetic and fine-scale location effects on breeding date in song sparrows. *J. Anim. Ecol.* **85**, 1613–1624. (doi:10.1111/1365-2656.12575)

16. Wilson S, Norris DR, Wilson AG, Arcese P. 2007 Breeding experience and population density affect the ability of a songbird to respond to future climate variation. *Proc. R. Soc. London B* **274**, 2539–2545. (doi:10.1098/rspb.2007.0643)

17. Reid JM, Keller LF. 2010 Correlated inbreeding among relatives: occurrence, magnitude, and implications. *Evolution* **64**, 973–985. (doi:10.1111/j.1558-5646.2009.00865.x)

18. Lindén M. 1991 Divorce in great tits -- chance or choice? An experimental approach. *Am. Nat.* **138**, 1039–1048. (doi:10.1086/285267)

- 251 19. Ramsay SM, Otter KA, Mennill DJ, Ratcliffe LM, Boag PT. 2000 Divorce and extrapair
252 mating in female black-capped chickadees (*Parus atricapillus*): separate strategies with a
253 common target. *Behav. Ecol. Sociobiol.* **49**, 18–23. (doi:10.1007/s002650000270)
- 254 20. Jocklin V, McGue M, Lykken DT. 1996 Personality and divorce: a genetic analysis. *J.*
255 *Pers. Soc. Psychol.* **71**, 288–99. (doi:10.1037//0022-3514.71.2.288)
- 256 21. Jerskey BA, Panizzon MS, Jacobson KC, Neale MC, Grant MD, Schultz M, Eisen SA,
257 Tsuang MT, Lyons MJ. 2010 Marriage and divorce: a genetic perspective. *Pers. Individ.*
258 *Dif.* **49**, 473–478. (doi:10.1016/j.paid.2010.05.007)
- 259 22. Forstmeier W, Nakagawa S, Griffith SC, Kempenaers B. 2014 Female extra-pair mating:
260 adaptation or genetic constraint? *Trends Ecol. Evol.* **29**, 456–464.
261 (doi:10.1016/j.tree.2014.05.005)

262

Table 1: Marginal posterior modes, means and 95% CIs from models decomposing total variance in liability for divorce. V_I and V_A represent permanent individual and additive genetic variances for females (♀) and males (♂). $\text{COV}_{A\text{♀♂}}$ is the cross-sex genetic covariance. V_S and V_Y are the social pair and year variances, respectively. Posterior statistics for sex-specific repeatabilities and heritabilities (h^2), ‘total heritability’ (T^2), fixed effects of within-season versus between-season (intercept) and regressions on individual female or male coefficient of inbreeding (f) are also shown.

270

	Model 1		Model 2	
	mode, mean	95%CI	mode, mean	95%CI
variance components				
$V_{I\varnothing}$	0.24, 0.28	4×10^{-6} , 0.53	0.003, 0.22	6×10^{-7} , 0.48
$V_{I\sigma}$	0.001, 0.09	5×10^{-8} , 0.26	0.002, 0.08	4×10^{-8} , 0.25
V_S	0.001, 0.14	2×10^{-9} , 0.46	0.003, 0.15	4×10^{-8} , 0.48
V_Y	0.001, 0.07	4×10^{-7} , 0.19	0.001, 0.07	8×10^{-9} , 0.19
$V_{A\varnothing}$			0.001, 0.07	5×10^{-9} , 0.25
$V_{A\sigma}$			0.001, 0.08	4×10^{-8} , 0.21
$COV_{A\varnothing\sigma}$			-0.0002, -0.003	-0.08, 0.06
variance ratios				
repeatability $\varnothing$	0.16, 0.17	3×10^{-6} , 0.30		
repeatability σ	0.001, 0.05	3×10^{-8} , 0.15		
$h^2_{\varnothing}$			0.001, 0.04	2×10^{-9} , 0.14
h^2_{σ}			0.001, 0.04	3×10^{-8} , 0.12
T^2			0.02, 0.08	1×10^{-4} , 0.20
fixed effects				
intercept	-0.64, -0.66	-0.89, -0.44	-0.81, -0.90	-1.28, -0.59
within-season	-1.08, -1.10	-1.34, -0.85	-1.11, -1.12	-1.37, -0.87
$f_{\varnothing}$			1.92, 1.98	-0.66, 4.45
f_{σ}			0.59, 0.94	-1.72, 3.61

271

272 **Figure 1.** Marginal posterior samples (bars), density (solid black line), mean (red dotted line)
 273 and 95%CI limits (dashed lines) of sex-specific repeatabilities for divorce in (a) female and (b)
 274 male song sparrows. Blue lines illustrate prior distributions (ESMS5). Areas where the posterior
 275 density exceeds the prior density highlight parameter values that are likely supported by the data.

276

277 **Figure 2.** Marginal posterior distributions for (a) female and (b) male heritabilities (h^2), and (c)
 278 the total heritability (T^2) for divorce in song sparrows. See figure 1 for plot description.

(b)

(b)

(c)