Article: Beta diversity patterns derived from island biogeography theory

Muyang Lu^{*1}, David Vasseur¹, Walter Jetz^{1,2}

1. Department of Ecology and Evolutionary Biology, Yale University, 165 Prospect Street, New Haven, CT, 06520, USA.

2. Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Road, SL5 7PY Ascot, Berks, UK.

Running head: beta diversity of islands

Words in the abstract: 164

Words in main body: 3820

References: 85

Table: 1

Figure: 3

* Correspondence: Muyang Lu (email: muyang.lu@yale.edu)

16 **Abstract**

17 The Theory of Island Biogeography (TIB) has been successful in predicting alpha
 18 diversity patterns such as species-area relationships and species-abundance
 19 distributions. Although beta diversity (i.e. the dissimilarity of community composition)
 20 has long been recognized as an important element of the TIB and is crucial for
 21 understanding community assembly processes, it has never been formally
 22 incorporated into the theory. Here we derive theoretical predictions for the expected
 23 pairwise beta diversity values under a species-level neutral scenario where all species
 24 have equal colonization and extinction rates. We test these predictions for the avian
 25 community composition of 42 islands (and 93 species) in the Thousand Island Lake,
 26 China. We find that alpha diversity patterns alone do not distinguish a species-level
 27 neutral model from a non-neutral model. In contrast, beta diversity patterns clearly
 28 reject a species-level neutral model. We suggest that the presented theoretical
 29 integration beta diversity offers a powerful path for testing the presence of neutral
 30 processes in ecology and biogeography.

31

32 **Keywords:** neutral theory; null model; beta diversity; meta-community; incidence
 33 function; community assembly; regional species pool

INTRODUCTION

Island biogeography is undergoing a renaissance (Warren et al. 2015; Santos et al. 2016; Patiño et al. 2017), at the heart of which lies the equilibrium theory of island biogeography (TIB) formulated by MacArthur and Wilson 50 years ago (1967). The TIB is most celebrated for elegantly linking species richness with colonization and extinction rates, which in turn are influenced by island area and distance to the mainland. Equilibrium TIB makes two key predictions: 1) equilibrium species richness depends on colonization and extinction rates; 2) community composition of an island is undergoing constant turnover when at equilibrium. While early empirical studies have focused on testing the equilibrium and temporal turnover predictions (Diamond 1969; Simberloff 1969; Simberloff and Wilson 1969, 1970; Diamond and May 1977), recent interest in TIB is stimulated by the success of Hubbell's unified neutral theory (a descendant of the TIB; Hubbell 2001), the advance of food web theories (Holt 2009), the developments of phylogenetic analyses (Valente et al. 2015; Lim and Marshall 2017), and the accumulation of island datasets (Weigelt et al. 2013, 2016). Motivated by the desire for a general island biogeography theory, more and more complexities have been added to the original framework, including allometric scaling (Jacquet et al. 2017), trophic interactions (Gravel et al. 2011; Cazelles et al. 2016), speciation (Chen and He 2009; Rosindell and Phillimore 2011; Rosindell and Harmon 2013), habitat heterogeneity (Kadmon and Allouche 2007), and island ontogenies (Whittaker et al. 2008; Borregaard et al. 2016). While the complexity of

the theory has increased, examinations to date have focused on alpha diversity patterns such as species richness (MacArthur and Wilson 1967), species-abundance relationships (Rosindell and Phillimore 2011; Rosindell and Harmon 2013; Kessler and Shnerb 2015), functional diversity (Jacquet et al. 2017), and endemism (Chen and He 2009).

One particular omission in the theory and tests of the TIB to date is beta diversity, i.e. the compositional dissimilarities among communities, a widely used metric in biodiversity studies (Leprieur et al. 2011; Stegen et al. 2013; Segre et al. 2014; Si et al. 2016). The integration of beta diversity patterns into the TIB has the potential to unlock a range of uses in community assembly studies and the development of a unified meta-community theory (Leibold et al. 2004).

The concept of beta diversity was first introduced by Whittaker (1960, 1965) and was quickly linked to an array of important mechanisms in community assembly: biotic interaction (Graham and Fine 2008), environment filtering (Veech and Crist 2007; Buckley and Jetz 2008), dispersal limitation (Ojima and Jiang 2017; Wu et al. 2017), and historical contingency (Fukami and Nakajima 2011). However, despite the widespread recognition of the beta diversity concept, there is little consensus about the best way to measure it (Tuomisto 2010a, 2010b; Anderson et al. 2011). Depending on the purpose of a study, beta diversity could be measured as turnover (directional) or variation (non-directional) of compositional similarity (Nekola and

White 1999; Legendre and De Cáceres 2013); based on pairwise or multiple-sites comparisons (Hui and McGeoch 2014; Arita 2017; Latombe et al. 2017; Marion et al. 2017); decomposed in additive or multiplicative framework (with respect to alpha and gamma diversity; Jost 2007; Cabral *et al.* 2014); calculated by incidence (presence-absence) or abundance data (Chao et al. 2016). Even for the commonly used pairwise beta diversity, there is a heated debate about what is the best way to partition it into richness difference and replacement (or nestedness and turnover) components (Baselga 2010; Podani and Schmera 2011; Legendre 2014; Baselga and Leprieur 2015). It is generally agreed that different measures quantify different aspects of communities and that their use should be guided by the question asked (Anderson et al. 2011; Legendre 2014). Although beta-diversity measures have been increasingly synthesized under more general frameworks such as Hill's number and variance of community matrix (Chao et al. 2016), only a few of them have been examined analytically from process-based theories (Chave and Leigh 2002; Condit et al. 2002; Zillio et al. 2005; Connolly et al. 2017). Moreover, theoretical considerations of beta-diversity patterns to date have been abundance-based and hence limited to more restricted datasets, such as vegetation plots (Chave and Leigh 2002; Condit et al. 2002; Zillio et al. 2005). In summary, the link between beta diversity patterns and fundamental processes such as colonization and extinction in the TIB remain ill developed.

In this study, we set out to address this shortcoming and provide a theoretical and example empirical integration of beta diversity into TIB. Specifically, we present the first theoretical predictions for the expected pairwise beta diversity patterns for both Jaccard and Sørensen indices in the classic MacArthur and Wilson framework. This framework constitutes in essence a species-level neutral model (as opposed to the individual-level neutral theory; Hubbell 2009; Rosindell and Phillimore 2011). The species-level neutral model assumes that all species have the same colonization and extinction rates, while the individual-level neutral model assumes that all individuals have the same colonization and extinction rates, which allows species-level asymmetry to arise because of different abundance (Rosindell and Harmon 2013). We first derived the probability mass function and expected value of pairwise beta diversity in a general case where the equilibrium assumption is not required. We then examined our results under the classic MacArthur-Wilson equilibrium framework. Finally, we test the derived predictions for a dataset of 93 avian species occurring across 42 islands of the Thousand Island Lake Region in China (Appendix, Wang *et al.* 2010).

MATERIAL AND METHODS

To derive expected beta diversity, we need to consider the joint statistical distribution of multiple islands. We here focus on pairwise community comparisons, because average pairwise dissimilarity is shown to be the only unbiased and consistent

estimator when applied to empirical data (Marion et al. 2017). We also calculated the expected value of the partitioned components of pairwise beta dissimilarity for both Baselga and Podani families (Baselga 2010; Podani and Schmera 2011). To test our predictions, we applied an “Incidence Function” approach pioneered by Diamond (1975) and used by Connor and Simberloff (1978, 1979) during early debates of null models.

Probability mass function of pairwise beta diversity

All pairwise beta diversity indices share the same probability mass function to describe island community composition because they are calculated from four quantities: the total number of species of two islands N (regional species richness), the number of species shared by both islands k , the number of species unique to the first island i , and the number of species unique to the second island j (thus $N = k + i + j$). Then the number of species present on the first island is $i+k$, and the number of species present on the second island is $j+k$.

Denoting the size of the mainland species pool as M , the probability mass function of pairwise beta diversity, conditioning on N , k and i , can be derived as follows: let the occurrence probability of the ith species on the first island be p , and the jth species on the second island be q . Furthermore, assuming that the occurrence probabilities are properties of the island rather than of species, the probability of obtaining exactly i unique species and k shared species from a regional pool of size M is given by

$p^{k+i}(1-p)^{M-(k+i)}$. Similarly, the probability of obtaining j unique and k shared species on the second island is $q^{k+j}(1-q)^{M-(k+j)}$, which yields the probability of obtaining the regional community specified by i, j, k and M , as $p^{k+i}(1-p)^{M-(k+i)}q^{k+j}(1-q)^{M-(k+j)}$.

Assuming that species are neutral with respect to their occurrence probabilities within an island, we then count how many different combinations of island communities could be obtained with regional richness N , k shared species, and i unique species to the first island (the number of unique species to the second island is then fixed as $k-i$). There are exactly $\binom{M}{N}$ ways of choosing N out of M species to be present in either or both islands from the mainland species pool. Similarly, there are $\binom{N}{k}$ ways to choose k out of N species to be shared by both islands and for the unique $N-k$ species, there are $\binom{N-k}{i}$ ways of assigning i species to the first island. After simplification, the probability of pairwise beta diversity conditioning on the total number of species of both islands N , the number of species shared by both islands k , and the number of species unique to the first island i is:

$$\Pr(\text{regional community} | N, k, i) = \binom{M}{N} \binom{N}{k, i} p^{k+i} q^{k+j} (1-p)^{M-(k+i)} (1-q)^{M-(k+j)}$$

(1)

Because beta diversity measures (e.g. Jaccard and Sørensen indices) are usually undefined when $N = 0$, the expected pairwise beta diversity should be normalized by the term $1 - (1-p)^M(1-q)^M$ to exclude “double-absence” scenarios (Anderson et al. 2011). Only the expected Jaccard dissimilarity (Jaccard dissimilarity is defined as

160 $1 - \frac{k}{N}$) has a simple analytical form that is independent of the size of mainland
 161 species pool M (see Appendix S1 in Supporting Information for derivation; we do not
 162 show the analytical form of the expected Sørensen dissimilarity because the specific
 163 form depends on the size of mainland species pool):

$$164 \quad E(\text{Jaccard.dissimilarity}) = \frac{p + q - 2pq}{p + q - pq} \quad (2)$$

165
 166 It has a form that is similar to the expected pairwise Jaccard derived by Chase *et al.*
 167 (2011) but allows two islands to have different occurrence probabilities. p and q could
 168 also be interpreted as probabilities of the same island at different times, and in that
 169 case (2) becomes the expected temporal Jaccard dissimilarity. This quantity does not
 170 require any equilibrium assumptions of alpha diversity and hence could be seen as a
 171 generalized version of the apparent turnover derived by Diamond and May (1977)
 172 which is essentially a pairwise Sørensen dissimilarity index in equilibrium (Morrison
 173 2017).

174 We also calculated the expected pairwise beta diversity for Jaccard and Sørensen
 175 families as well as their partitioned components (for detailed formulas of the Baselga
 176 and Podani families see Baselga & Leprieur 2015) conditioning on both islands
 177 having species ($i+k > 0$ and $j+k > 0$) because empty islands are sometimes excluded
 178 from analysis (Wang et al. 2016). In Baselga's family, pairwise dissimilarity is
 179 partitioned into the nestedness component and the turnover component. The
 180 nestedness component measures the extent to which species in a smaller community

are a subset of species in a larger community, while the turnover component measures how much of the dissimilarity is caused by species replacement (Baselga 2010). The turnover component and the nestedness component in Baselga's family, respectively, correspond to the replacement component and the richness difference component in Podani's family (Legendre 2014).

Island biogeography theory

Following the stochastic version of the TIB (MacArthur and Wilson 1967), the occurrence probability of a species on an island is a function of colonization rate c and extinction rate e :

$$\frac{dp}{dt} = c(1-p) - ep \quad (3)$$

When occurrence probability of a species is at stochastic equilibrium (stationary distribution), the occurrence probability is:

$$p_{i,eq} = \frac{c}{c+e} \quad (4)$$

Let $\theta = \frac{e}{c}$, which is the relative extinction rates, equation (4) becomes:

$$p_{i,eq} = \frac{1}{1+\theta} \quad (5)$$

Substitute $p = q = \frac{1}{1+\theta}$ into equation (2), the expected Jaccard dissimilarity when two islands have equal colonization and extinction probabilities is:

$$E(\text{Jaccard.dissimilarity}) = \frac{2\theta}{1+2\theta} \quad (6)$$

To take into account asymmetry between islands (e.g. the effect of area, isolation or habitat types), let $p = \frac{1}{1+\theta}$ and $q = \frac{1}{1+z\theta}$, so that the relative extinction rate on the

second island is z times the relative extinction rate on the first island. Equation (2)

becomes:

$$E(\text{Jaccard.dissimilarity}) = \frac{(1+z)\theta}{1+(1+z)\theta} \quad (7)$$

Empirical tests

Data

We use a published dataset of avian community composition for 93 birds and 42 islands in The Thousand Island Lake, China to test our predictions (see Appendix in Wang *et al.* 2010). The region (29°22" – 29°50" N, 118°34"– 119°15" E) consists of an inundated lake with more than 1000 islands created by dam constructions in 1959. Because the islands were formed recently, there was no in situ speciation in this region. The relative small area of the region (573 km²) ensures that the islands are accessible to most of the bird species (Si et al. 2016). Bird occupancies from 2006 to 2009 on 42 islands were documented using line-transect method. Island variables measured in the dataset include area, distance to mainland, habitat diversity and plant species richness.

Incidence function and statistical analysis

Because extinction rates and colonization rates are difficult to measure directly, an alternative way is to use an “Incidence function” approach to estimate parameters from a snapshot of occupancy patterns (Diamond 1975; Hanski 2009). To test the

predictions of island biogeography theory, equation (4) is modeled as a function of isolation and area. Parameters are fitted by maximum likelihood with binomial distribution. Three neutral models and one non-neutral model are examined in this study:

Neutral model 1: Inverse ratio

Colonization rates are modeled as an inverse ratio function of isolation: $c = \frac{a}{I}$, and extinction rates are modeled as an inverse ratio of area: $e = \frac{b}{A}$. a and b are fitted parameters.

Neutral model 2: Exponential

Colonization rates are modeled as an exponential function of isolation: $c = \exp(-aI)$, and extinction rates are modeled as an exponential function of area: $e = \exp(-bA)$. a and b are fitted parameters. Substitute $c = \exp(-aI)$ and $e = \exp(-bA)$ into equation (4), the occurrence probability becomes

$$p = \frac{1}{1 + \exp(-bA + aI)}$$
. Thus the exponential neutral model is equivalent to a logistic

regression without intercepts, which could be directly fitted by ‘glm’ in R with binomial distribution and logit link.

Neutral model 3: GLM

This model adds habitat diversity and plant richness to the predictors in addition to area and isolation in Neutral Model 2. AIC is used to select the best model. Area and isolation are log-transformed to ensure better linearity.

Non-neutral model: Aggregate species-level GLM

To take into account species-level non-neutrality, species identity is included as a fixed effect categorical variable in the GLM, which allows each species to have a different intercept (baseline occurrence probability) but share the same response to island area, isolation, plant richness and habitat types. This is essentially a stacked species distribution modeling approach (Calabrese et al. 2014; Ko et al. 2016), which increases the number of parameters by the number of species minus one (92 parameters in this case). This procedure aims to capture the observed presence-absence variation among species but does not pin down the causes of species-level non-neutrality such as traits and abundance differences. More realistic assumptions of non-neutrality such as different responses to area and isolation could be made but are not the main focus of this paper. AUC and AIC values are calculated to compare the overall performance of the models. AUC is calculated with a Mann-Whitney U statistic using R package ‘PresenceAbsence’. AIC is calculated from the best-fitted likelihood function. ΔAIC are calculated by subtracting the AIC of the null model (GLM with only one intercept) from the AIC of the fitted model.

Predictions of alpha diversity and beta diversity

The predicted species richness of each island is given by the summed fitted occurrence probabilities of all species as used in conventional stacked species distribution modeling (Calabrese et al. 2014; Ko et al. 2016). While predicted pairwise Jaccard dissimilarity can be calculated analytically from equation (2), the

partitioned nestedness and turnover components can only be estimated by simulations (we only present the results of Baselga's family because its partitioned components are independent with each other when gamma diversity is fixed; Baselga & Leprieur 2015). We therefore estimate predicted pairwise beta diversity as the mean of 1000 community matrices simulated from the fitted occurrence probabilities. Observed values are regressed against predicted values using ordinary least square. R^2 is used as a measure of predictive power for alpha and beta diversity patterns. If the model predicts the observed patterns well, the fitted regression line should be close to the 1:1 ratio line when observed values are plotted against predicted values. All statistical analyses are performed in R 3.3.2.

RESULTS

Symmetric islands (same colonization and extinction rates for both islands)

The expected Jaccard dissimilarity conditional on both islands having non-zero species richness increases monotonically with relative extinction rate θ and the size (number of species) of the mainland species pool M . As M increases, the expected Jaccard dissimilarity converges to $\frac{2\theta}{1+2\theta}$ (Fig. 1a). When decomposed into turnover and nestedness (or replacement and richness difference) components, contrasting patterns are observed: while the turnover component and the replacement component increase monotonically with θ and M (Fig. 1b, e), the nestedness component and richness difference component are both unimodal functions of θ with maximum

values less than 0.3 (Fig. 1c, f). The maximum nestedness decreases as M increases (Fig. 1c), while the maximum richness difference changes little with the increase of M (Fig. 1f). The relative importance of turnover increases monotonically with θ and M (Fig. 1d). In a special case of 2 mainland species, the ratio of expected turnover and expected nestedness equals θ (Fig. 1d). The Sørensen family indices have similar quantitative behaviors as the Jaccard family indices (see Fig. S1 in Appendix).

Asymmetric islands (different colonization and extinction rates for two islands)

The expected Jaccard dissimilarity conditional on both islands having non-zero species richness converges to $\frac{(1+z)\theta}{1+(1+z)\theta}$ as M increases. But the deviation from $\frac{(1+z)\theta}{1+(1+z)\theta}$ also increases with islands asymmetry z (Fig. 2a, e, i). The turnover component increases monotonically with θ and M , and decreases with z (Fig. 2b, f, j). The nestedness component decreases with M when z is small (Fig. 2c), but this relationship with M is reversed when z gets larger (Fig. 2g, k). The ratio of expected turnover and expected nestedness also increases with M when z is small (Fig. 2d), and decreases with M when z is large (Fig. 2h, l).

Empirical test

We tested the above predictions for an inland lake island system using 93 bird species and 42 islands. We used an incidence function approach to fit the observed occupancy

patterns and calculated predicted alpha and beta diversity patterns from the fitted models. The non-neutral GLM has the lowest AIC and highest AUC in all models (Table 1). Among three neutral models, the neutral GLM model has the lowest AIC and the highest AUC. All models are better at predicting alpha diversity patterns than at predicting beta diversity patterns (Fig. 3). The neutral exponential model and GLM systematically underestimate the nestedness component, but overestimate the turnover component (Fig. 3f, j). In contrast, the neutral inverse ratio model systematically underestimates all observed pairwise beta diversity at the lower range of the predictions, yet it overestimates pairwise beta diversity at the higher range of the predictions (Fig. 3a-d). Both the neutral GLM and the non-neutral GLM successfully predict the observed alpha diversity pattern (Fig. 3i, m), but only the non-neutral GLM successfully predicts the observed beta diversity patterns (Fig. 3n-p).

DISCUSSION

Theoretical results

We derived a set of novel predictions of beta diversity patterns from the island biogeography theory. Under the classic MacArthur and Wilson framework, pairwise beta diversity patterns are determined by three factors: the size of the mainland species pool, extinction rates and colonization rates. Conditional on at least one island having species ($N > 0$), only the expected Jaccard dissimilarity is independent of the size of mainland species pool. When conditioning on both islands having non-zero

richness ($i+k > 0$ and $j+k > 0$), all indices are dependent on the size of mainland species pool. Ignoring empty islands has a huge influence on beta diversity patterns when the size of mainland species pool is small (e.g. less than 10 species), but the effect becomes negligible toward larger mainland species pool sizes, because of the rapidly decreasing probability of empty islands (Fig. 1a). This result stresses the need to include empty islands in empirical tests of the island biogeography theory (Wang et al. 2016).

Pairwise Jaccard dissimilarity increases with relative extinction rates (the ratio between extinction rates and colonization rates). This is because when extinction events become more frequent the chance of forming “checkerboard” patterns (Diamond 1975) grows, as shown by the increasing turnover component (Fig. 1b). The nestedness component first increases then decreases with relative extinction rates (Fig. 1c). The maximum nestedness is achieved when extinction rates are less than colonization rates (Fig. 1c). When relative extinction rates or the size of mainland species pool gets larger, the increase of Jaccard dissimilarity is mainly driven by the increasing turnover component (Fig. 1d). Indices in Podani’s family (the replacement and richness difference components) have similar qualitative behaviors, consistent with previous finding that the partitioned components in Baslega’s family are correlated with the partitioned components in Podani’s family (Legendre 2014; Baselga and Leprieur 2015). Island asymmetry, which takes into account the effect of

isolation and area, does not change the qualitative behaviors of the indices (Fig 2), but increases the level of nestedness, because species in a small (distant) island are more likely to be a subset of species in a large (near) island.

Neutral theory and null models

Our theoretical results are derived from a species-level neutral theory. We show that both the neutral GLM and the non-neutral GLM are successful in predicting species richness (alpha diversity) of birds in the Thousand Islands Lake (Fig. 3i, m). But only the non-neutral model predicts the observed beta diversity patterns. The patterns of partitioned components further reveal that the neutral GLM fails to predict the observed pairwise Jaccard dissimilarity because it overestimates turnover and underestimates nestedness (Fig. 3j-l). Our analysis does not include biological factors that may cause species-level non-neutrality, but the presented framework is flexible to include such information if available.

Null models have become one of the most important tools in ecology. Despite their increasing popularity, it has also been recognized that their assumptions (e.g. the widely used random shuffling approaches) can have serious limitations, especially in the presence of multiple processes (Gotelli and Ulrich 2012; Pigot and Etienne 2015; Miller et al. 2017; O'Dwyer et al. 2017). In terms of beta diversity, the entangled links among alpha, beta and gamma diversity are known to reduce the statistical power of randomization tests and generate ambiguity in their interpretations (Chase et

al. 2011; Kraft et al. 2011; Qian et al. 2012, 2013; Ulrich et al. 2017). Our results support the use of mechanistic null models such as those based on neutral theory (O'Dwyer et al. 2009; Rosindell et al. 2012), maximum entropy theory (Xiao et al. 2015, 2016; O'Dwyer et al. 2017), and incidence functions (Hanski et al. 1996; Helm et al. 2006; Hanski 2009) to improve upon random-shuffling null models in hypothesis testing. These mechanistic models could generate multiple diagnostic patterns and allows stronger test for ecological theories.

Future directions

Our results are derived from a mainland-island model where the colonization and extinction events on islands are independent of one another. More realistic modifications should be considered in order to predict biodiversity changes in a real landscape (Hanski 2009). For example, environmental heterogeneity and local dispersal are found to be important drivers of beta diversity in experimental studies (Grainger and Gilbert 2016; Gianuca et al. 2017; Ojima and Jiang 2017; Rodrigues and Diniz-Filho 2017). Moreover, in oceanic islands, high endemism arises from speciation events (Chen and He 2009; Gascuel et al. 2016; Steinbauer et al. 2016), which are likely to increase the level of species turnover. Future studies should also extend presence-based measures to abundance-based beta diversity measures which may provide additional insights into the dynamics of meta-community (Bashan et al. 2016; Kalyuzhny and Shnerb 2017). We believe that the integration of beta diversity

patterns into the Theory of Island Biogeography offers new opportunities to infer community assembly processes.

ACKNOWLEDGEMENTS

We thank Dr. James Rosindell for providing helpful comments for earlier drafts. We thank members in Jetz Lab and Vasseur Lab for feedbacks and discussions.

AUTHORSHIP

ML and DV conceived of the idea. The idea was further developed with DV and WJ's input. ML conducted the analyses and wrote the manuscript. All authors assisted with revisions.

REFERENCE

- Anderson, M. J., T. O. Crist, J. M. Chase, M. Vellend, B. D. Inouye, A. L. Freestone, N. J. Sanders, et al. 2011. Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology Letters* 14:19–28.
- Arita, H. T. 2017. Multisite and multispecies measures of overlap, co-occurrence, and co-diversity. *Ecography* 40:709–718.
- Baselga, A. 2010. Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography* 19:134–143.
- Baselga, A., and F. Leprieur. 2015. Comparing methods to separate components of beta diversity. *Methods in Ecology and Evolution* 6:1069–1079.
- Bashan, A., T. E. Gibson, J. Friedman, V. J. Carey, S. T. Weiss, E. L. Hohmann, and

412 Y.-Y. Liu. 2016. Universality of human microbial dynamics. *Nature* 534:259–
413 262.

414 Borregaard, M. K., T. J. Matthews, and R. J. Whittaker. 2016. The general dynamic
415 model: towards a unified theory of island biogeography? *Global Ecology and*
416 *Biogeography* 25:805–816.

417 Buckley, L. B., and W. Jetz. 2008. Linking global turnover of species and
418 environments. *Proceedings of the National Academy of Sciences* 105:17836–
419 17841.

420 Cabral, J. S., P. Weigelt, W. D. Kissling, and H. Kreft. 2014. Biogeographic, climatic
421 and spatial drivers differentially affect α -, β - and γ -diversities on oceanic
422 archipelagos. *Proceedings. Biological sciences / The Royal Society*
423 281:20133246.

424 Calabrese, J. M., G. Certain, C. Kraan, and C. F. Dormann. 2014. Stacking species
425 distribution models and adjusting bias by linking them to macroecological
426 models. *Global Ecology and Biogeography* 23:99–112.

427 Cazelles, K., N. Mouquet, D. Mouillot, and D. Gravel. 2016. On the integration of
428 biotic interaction and environmental constraints at the biogeographical scale.
429 *Ecography* 39:921–931.

430 Chao, A., C. H. Chiu, and D. Warton. 2016. Bridging the variance and diversity
431 decomposition approaches to beta diversity via similarity and differentiation
432 measures. *Methods in Ecology and Evolution* 7:919–928.

- 433 Chase, J. M., N. J. B. Kraft, K. G. Smith, M. Vellend, and B. D. Inouye. 2011. Using
434 null models to disentangle variation in community dissimilarity from variation in
435 α -diversity. *Ecosphere* 2:art24.
- 436 Chave, J., and E. G. Leigh. 2002. A spatially explicit neutral model of β -diversity in
437 tropical forests. *Theoretical Population Biology* 62:153–168.
- 438 Chen, X.-Y., and F. He. 2009. Speciation and endemism under the model of island
439 biogeography. *Ecology* 90:39–45.
- 440 Condit, R., N. Pitman, E. G. Leigh Jr., J. Chave, J. Terborgh, R. B. Foster, P. Nunez
441 V., et al. 2002. Beta-diversity in tropical forest trees. *Science* 295:666–669.
- 442 Connolly, S. R., T. P. Hughes, and D. R. Bellwood. 2017. A unified model explains
443 commonness and rarity on coral reefs. *Ecology Letters* 20:477–486.
- 444 Connor, E. F., and D. Simberloff. 1978. Species number and compositional similarity
445 of the Galapagos Flora and Avifauna. *Ecological Monographs* 48:219–248.
- 446 ———. 1979. The assembly of species communities: chance or competition?
447 *Ecology* 60:1132.
- 448 Diamond, J. M. 1969. Avifaunal equilibria and species turnover rates on the Channel
449 Islands of California. *Proceedings of the National Academy of Sciences* 64:57–
450 63.
- 451 ———. 1975. Assembly of species communities. Pages 342–444 in *Ecology and*
452 *Evolution of communities*. Harvard University Press.
- 453 Diamond, J. M., and R. M. May. 1977. Species turnover rates on islands: dependence

454 on census interval. *Science* 197:266–270.

455 Fukami, T., and M. Nakajima. 2011. Community assembly: alternative stable states or
456 alternative transient states? *Ecology Letters* 14:973–984.

457 Gascuel, F., F. Laroche, A. S. Bonnet-Lebrun, and A. S. L. Rodrigues. 2016. The
458 effects of archipelago spatial structure on island diversity and endemism:
459 predictions from a spatially-structured neutral model. *Evolution* 70:2657–2666.

460 Gianuca, A. T., S. A. J. Declerck, P. Lemmens, and L. De Meester. 2017. Effects of
461 dispersal and environmental heterogeneity on the replacement and nestedness
462 components of β -diversity. *Ecology* 98:525–533.

463 Gotelli, N. J., and W. Ulrich. 2012. Statistical challenges in null model analysis.
464 *Oikos* 121:171–180.

465 Graham, C. H., and P. V. A. Fine. 2008. Phylogenetic beta diversity: linking
466 ecological and evolutionary processes across space in time. *Ecology Letters*
467 11:1265–1277.

468 Grainger, T. N., and B. Gilbert. 2016. Dispersal and diversity in experimental
469 metacommunities: linking theory and practice. *Oikos* 125:1213–1223.

470 Gravel, D., F. Massol, E. Canard, D. Mouillot, and N. Mouquet. 2011. Trophic theory
471 of island biogeography. *Ecology Letters* 14:1010–1016.

472 Hanski, I. 2009. The theories of island biogeography and metapopulation dynamics.
473 Pages 186–213 in J. B. Losos and R. E. Ricklefs, eds. *The theory of island*
474 *biogeography revisited*. Princeton University Press, Princeton.

- 475 Hanski, I., A. Moilanen, T. Pakkala, and M. Kuussaari. 1996. The quantitative
476 incidence function model and persistence of an endangered butterfly
477 metapopulation. *Conservation Biology* 10:578–590.
- 478 Helm, A., I. Hanski, and M. Pärtel. 2006. Slow response of plant species richness to
479 habitat loss and fragmentation. *Ecology Letters* 9:72–77.
- 480 Holt, R. D. 2009. Toward a trophic island biogeography. Pages 143–185 *in* J. B.
481 Losos and R. E. Ricklefs, eds. *The theory of island biogeography revisited*.
482 Princeton University Press, Princeton.
- 483 Hubbell, S. P. 2001. *The unified neutral theory of biodiversity and biogeography*.
484 Princeton University Press, Princeton.
- 485 ———. 2009. Neutral theory and the theory of island biogeography. Pages 263–292
486 *in* J. B. Losos and R. E. Ricklefs, eds. *The theory of island biogeography*
487 *revisited*. Princeton University Press, Princeton.
- 488 Hui, C., and M. A. McGeoch. 2014. Zeta diversity as a concept and metric that unifies
489 Incidence-based biodiversity patterns. *The American Naturalist* 184:684–694.
- 490 Jacquet, C., D. Mouillot, M. Kulbicki, and D. Gravel. 2017. Extensions of Island
491 Biogeography Theory predict the scaling of functional trait composition with
492 habitat area and isolation. *Ecology Letters* 20:135–146.
- 493 Jost, L. 2007. Partitioning diversity into independent alpha and beta components.
494 *Ecology* 88:2427–2439.
- 495 Kadmon, R., and O. Allouche. 2007. Integrating the effects of area, isolation, and

496 habitat heterogeneity on species diversity: a unification of island biogeography
497 and niche theory. *The American Naturalist* 170:443–454.

498 Kalyuzhny, M., and N. M. Shnerb. 2017. Dissimilarity-overlap analysis of community
499 dynamics: opportunities and pitfalls. *Methods in Ecology and Evolution* 0:1–10.

500 Kessler, D. A., and N. M. Shnerb. 2015. Generalized model of island biodiversity.
501 *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics* 91:1–11.

502 Ko, C. Y., O. J. Schmitz, and W. Jetz. 2016. The limits of direct community modeling
503 approaches for broad-scale predictions of ecological assemblage structure.
504 *Biological Conservation* 201:396–404.

505 Kraft, N. J. B., L. S. Comita, J. M. Chase, N. J. Sanders, N. G. Swenson, T. O. Crist, J.
506 C. Stegen, et al. 2011. Disentangling the drivers of diversity along latitudinal and
507 elevational gradients. *Science* 333:1755–1758.

508 Latombe, G., C. Hui, and M. A. McGeoch. 2017. Multi-site generalised dissimilarity
509 modelling: using zeta diversity to differentiate drivers of turnover in rare and
510 widespread species. *Methods in Ecology and Evolution* 8:431–442.

511 Legendre, P. 2014. Interpreting the replacement and richness difference components
512 of beta diversity. *Global Ecology and Biogeography* 23:1324–1334.

513 Legendre, P., and M. De Cáceres. 2013. Beta diversity as the variance of community
514 data: dissimilarity coefficients and partitioning. *Ecology Letters* 16:951–963.

515 Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F.
516 Hoopes, R. D. Holt, et al. 2004. The metacommunity concept: a framework for

517 multi-scale community ecology. *Ecology Letters* 7:601–613.

518 Leprieux, F., P. A. Tedesco, B. Hugueny, O. Beauchard, H. H. Dürr, S. Brosse, and T.

519 Oberdorff. 2011. Partitioning global patterns of freshwater fish beta diversity

520 reveals contrasting signatures of past climate changes. *Ecology Letters* 14:325–

521 334.

522 Lim, J. Y., and C. R. Marshall. 2017. The true tempo of evolutionary radiation and

523 decline revealed on the Hawaiian archipelago. *Nature* 543:710–713.

524 MacArthur, R. H., and E. O. Wilson. 1967. *The theory of island biogeography*.

525 Princeton University Press, Princeton.

526 Marion, Z. H., J. A. Fordyce, and B. M. Fitzpatrick. 2017. Pairwise beta diversity

527 resolves an underappreciated source of confusion in calculating species turnover.

528 *Ecology* 98:933–939.

529 Miller, E. T., D. R. Farine, and C. H. Trisos. 2017. Phylogenetic community structure

530 metrics and null models: a review with new methods and software. *Ecography*

531 40:461–477.

532 Morrison, L. W. 2017. Insular plant turnover across a 22-year interval: a critical

533 retrospective of the roles of pseudoturnover and cryptoturnover. *Journal of*

534 *Biogeography* 44:1007–1017.

535 Nekola, J. C., and P. S. White. 1999. The distance decay of similarity in biogeography

536 and ecology. *Journal of Biogeography* 26:867–878.

537 O’Dwyer, J. P., J. K. Lake, A. Ostling, V. M. Savage, and J. L. Green. 2009. An

538 integrative framework for stochastic, size-structured community assembly.

539 Proceedings of the National Academy of Sciences 106:6170–6175.

540 O’Dwyer, J. P., A. Rominger, and X. Xiao. 2017. Reinterpreting maximum entropy in

541 ecology: a null hypothesis constrained by ecological mechanism. Ecology

542 Letters 20:832–841.

543 Ojima, M. N., and L. Jiang. 2017. Interactive effects of disturbance and dispersal on

544 community assembly. Oikos 126:682–691.

545 Patiño, J., R. J. Whittaker, P. A. V. Borges, J. M. Fernández-Palacios, C. Ah-Peng, M.

546 B. Araújo, S. P. Ávila, et al. 2017. A roadmap for island biology: 50 fundamental

547 questions after 50 years of The Theory of Island Biogeography. Journal of

548 Biogeography 44:963–983.

549 Pigot, A. L., and R. S. Etienne. 2015. A new dynamic null model for phylogenetic

550 community structure. Ecology Letters 18:153–163.

551 Podani, J., and D. Schmera. 2011. A new conceptual and methodological framework

552 for exploring and explaining pattern in presence - absence data. Oikos 120:1625–

553 1638.

554 Qian, H., S. Chen, L. Mao, and Z. Ouyang. 2013. Drivers of β -diversity along

555 latitudinal gradients revisited. Global Ecology and Biogeography 22:659–670.

556 Qian, H., X. Wang, and Y. Zhang. 2012. Comment on “Disentangling the drivers of

557 diversity along latitudinal and elevational gradients.” Science 335:1573–1573.

558 Rodrigues, J. F. M., and J. A. F. Diniz-Filho. 2017. Dispersal is more important than

559 climate in structuring turtle communities across different biogeographical realms.
560 *Journal of Biogeography* 44:2109–2120.

561 Rosindell, J., and L. J. Harmon. 2013. A unified model of species immigration,
562 extinction and abundance on islands. *Journal of Biogeography* 40:1107–1118.

563 Rosindell, J., S. P. Hubbell, F. He, L. J. Harmon, and R. S. Etienne. 2012. The case
564 for ecological neutral theory. *Trends in Ecology and Evolution* 27:203–208.

565 Rosindell, J., and A. B. Phillimore. 2011. A unified model of island biogeography
566 sheds light on the zone of radiation. *Ecology Letters* 14:552–560.

567 Santos, A. M. C., R. Field, and R. E. Ricklefs. 2016. New directions in island
568 biogeography. *Global Ecology and Biogeography* 25:751–768.

569 Segre, H., R. Ron, N. De Malach, Z. Henkin, M. Mandel, and R. Kadmon. 2014.
570 Competitive exclusion, beta diversity, and deterministic vs. stochastic drivers of
571 community assembly. *Ecology Letters* 17:1400–1408.

572 Si, X., A. Baselga, F. Leprieur, X. Song, and P. Ding. 2016. Selective extinction
573 drives taxonomic and functional alpha and beta diversities in island bird
574 assemblages. *Journal of Animal Ecology* 85:409–418.

575 Simberloff, D. S. 1969. Experimental zoogeography of islands: a model for insular
576 colonization. *Ecology* 50:296–314.

577 Simberloff, D. S., and E. O. Wilson. 1969. Experimental zoogeography of islands: the
578 colonization of empty Islands. *Ecology* 50:278–296.

579 Simberloff, D. S., and E. O. Wilson. 1970. Experimental zoogeography of islands: a

580 two-year record of colonization. *Ecology* 51:934–937.

581 Stegen, J. C., A. L. Freestone, T. O. Crist, M. J. Anderson, J. M. Chase, L. S. Comita,
582 H. V. Cornell, et al. 2013. Stochastic and deterministic drivers of spatial and
583 temporal turnover in breeding bird communities. *Global Ecology and*
584 *Biogeography* 22:202–212.

585 Steinbauer, M. J., R. Field, J. Grytnes, P. Trigas, C. Ah-Peng, F. Attorre, H. J. B.
586 Birks, et al. 2016. Topography-driven isolation, speciation and a global increase
587 of endemism with elevation. *Global Ecology and Biogeography* 25:1097–1107.

588 Tuomisto, H. 2010*a*. A diversity of beta diversities: straightening up a concept gone
589 awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity.
590 *Ecography* 33:2–22.

591 ———. 2010*b*. A diversity of beta diversities: straightening up a concept gone awry.
592 Part 2. Quantifying beta diversity and related phenomena. *Ecography* 33:23–45.

593 Ulrich, W., A. Baselga, B. Kusumoto, T. Shiono, H. Tuomisto, and Y. Kubota. 2017.
594 The tangled link between β - and γ -diversity: a Narcissus effect weakens
595 statistical inferences in null model analyses of diversity patterns. *Global Ecology*
596 *and Biogeography* 26:1–5.

597 Valente, L. M., A. B. Phillimore, and R. S. Etienne. 2015. Equilibrium and
598 non-equilibrium dynamics simultaneously operate in the Galápagos islands.
599 *Ecology Letters* 18:844–852.

600 Veech, J. A., and T. O. Crist. 2007. Habitat and climate heterogeneity maintain

601 beta-diversity of birds among landscapes within ecoregions. *Global Ecology and*
602 *Biogeography* 16:650–656.

603 Wang, Y., Y. Bao, M. Yu, G. Xu, and P. Ding. 2010. Nestedness for different reasons:
604 the distributions of birds, lizards and small mammals on islands of an inundated
605 lake. *Diversity and Distributions* 16:862–873.

606 Wang, Y., V. Millien, P. Ding, and A. Algar. 2016. On empty islands and the
607 small-island effect. *Global Ecology and Biogeography* 25:1333–1345.

608 Warren, B. H., D. Simberloff, R. E. Ricklefs, R. Aguilée, F. L. Condamine, D. Gravel,
609 H. Morlon, et al. 2015. Islands as model systems in ecology and evolution:
610 prospects fifty years after MacArthur-Wilson. *Ecology Letters* 18:200–217.

611 Weigelt, P., W. Jetz, and H. Kreft. 2013. Bioclimatic and physical characterization of
612 the world’s islands. *Proceedings of the National Academy of Sciences*
613 110:15307–15312.

614 Weigelt, P., M. J. Steinbauer, J. S. Cabral, and H. Kreft. 2016. Late Quaternary
615 climate change shapes island biodiversity. *Nature* 532:99–102.

616 Whittaker, R. H. 1960. Vegetation of the Sisiyou Mountains, Orgeon and California.
617 *Ecological Monographs* 30:279–338.

618 Whittaker, R. H. 1965. Dominance and diversity in land plant communities. *Science*
619 147:250–260.

620 Whittaker, R. J., K. A. Triantis, and R. J. Ladle. 2008. A general dynamic theory of
621 oceanic island biogeography. *Journal of Biogeography* 35:977–994.

622 Wu, L., X. Si, R. K. Didham, D. Ge, and P. Ding. 2017. Dispersal modality
623 determines the relative partitioning of beta diversity in spider assemblages on
624 subtropical land-bridge islands. *Journal of Biogeography* 44:2121–2131.

625 Xiao, X., D. J. McGlinn, and E. P. White. 2015. A strong test of the Maximum
626 Entropy Theory of Ecology. *The American Naturalist* 185:E70–E80.

627 Xiao, X., J. P. O’Dwyer, and E. P. White. 2016. Comparing process-based and
628 constraint-based approaches for modeling macroecological patterns. *Ecology*
629 97:1228–1238.

630 Zillio, T., I. Volkov, J. R. Banavar, S. P. Hubbell, and A. Maritan. 2005. Spatial
631 scaling in model plant communities. *Physical Review Letters* 95:1–4.

632

Table 1. Models fitted to the presence-absence patterns of 93 bird species in 42 islands in the Thousand Island Lake, China.

Model	ΔAIC	AUC	Number of parameters	R^2		
				Richness	Dissimilarity	Turnover
non-neutral.GLM	-2627	0.95	97	0.91	0.37	0.53
neutral.GLM	-172	0.61	5	0.90	0.01	0.45
neutral.exponential	-8	0.57	2	0.41	0.01	0.17
neutral.inverse ratio	1587	0.60	1	0.69	0.04	0.46
Total observation = 3906						

Note: For the neutral exponential model: $c = \exp(-aI)$ and $e = \exp(-bA)$. For the neutral inverse ratio model: $c = \frac{a}{I}$ and $e = \frac{b}{A}$. Dissimilarity is the sum of turnover component and nestedness component. ΔAIC are calculated by subtracting the AIC of the null model (GLM with only one intercept) from the AIC of the fitted model. AUC are calculated with a Mann-Whitney U statistic.

Figure legends

Figure 1. Expected pairwise beta diversity of two identical islands (same colonization and extinction rates), conditioning on both islands having species ($i > 0, j > 0$). Results are derived from the joint presence-absence distribution of two islands and shown for different mainland pool sizes. Blue dashed line in panel (a) is the analytical solution $E(Jaccard.dissimilarity) = \frac{2\theta}{1+2\theta}$ for the case of at least one island having species ($N > 0$). Relative extinction rate is the ratio of extinction rate and colonization rate.

Figure 2. Expected pairwise Jaccard beta diversities of two islands differing in colonization and/or extinction rates, conditioning on both islands having species ($i > 0, j > 0$). The blue dashed line in panel (a), (e) and (i) are the analytical solutions of $E(Jaccard.dissimilarity) = \frac{(1+z)\theta}{1+(1+z)\theta}$, conditional on at least one island having nonzero species. Expectations are provided for different levels of z , i.e. the number of times relative extinction rate on the second island exceeds that on the first. The black dashed lines in panel (d), (h) and (l) represent a 1:1 ratio. Different lines in a graph represent different mainland pool sizes (for color legend see Fig. 1).

Figure 3. Empirical evaluation of the theoretically fitted alpha and beta diversity patterns for the Thousand Islands Lake bird dataset. (a) - (l) are predictions of three neutral models, (m) - (p) are predictions of a non-neutral model. Solid black lines

661 represent a 1:1 relationship and dashed red lines are fits of ordinary linear regressions,
662 Point densities from high to low are color-coded from red to blue.

Figure 1

Figure 2

Figure 3