The scent of the fly

Paul G. Becher¹, Sebastien Lebreton¹, Erika A. Wallin², Erik Hedenström², Felipe Borrero³, Marie Bengtsson¹, Volker Jörger⁴, Peter Witzgall^{1*}

¹Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Box 102, 23053 Alnarp, Sweden

²Department of Chemical Engineering, Mid Sweden University, Holmgatan 10, 85170 Sundsvall, Sweden

³Biological Control Laboratory, Colombian Corporation of Agricultural Research, AA 240142 Las Palmas, Bogota, Colombia

⁴Staatliches Weinbauinstitut, Merzhauserstr. 119, 79100 Freiburg, Germany

*Correspondence: Peter Witzgall, SLU, Box 102, 23053 Alnarp, Sweden
phone +46 70 2426939, peter.witzgall@slu.se

Paul G. Becher - paul.becher@slu.se

Sebastien Lebreton - sebastien.lebreton@univ-amu.fr

Erika A. Wallin - erika.wallin@miun.se

Erik Hedenström - erik.hedenstrom@miun.se

Felipe Borrero - fborrero@corpoica.org.co

Marie Bengtsson - marie.bengtsson@slu.se

Volker Jörger - volker.joerger@freenet.de

Abstract

(Z)-4-undecenal (Z4-11Al) is the volatile pheromone produced by females of the vinegar fly *Drosophila melanogaster*. Female flies emit Z4-11Al for species-specific communication and mate-finding. A sensory panel finds that synthetic Z4-11Al has a characteristic flavour, which we perceive even at the small amounts produced by one female fly. Since only females produce Z4-11Al, and not males, we can reliably distinguish between single *D. melanogaster* males and females, according to their scent. Females release Z4-11Al at 2.4 ng/h and we readily sense 1 ng synthetic Z4-11Al in a glass of wine (0.03 nmol/L), while a tenfold concentration is perceived as a loud off-flavour. This corroborates the observation that a glass of wine is spoiled by a single *D. melanogaster* fly falling into it, which we here show is caused by Z4-11Al. The biological

role of Z4-11Al or structurally related aldehydes in humans and the basis for this semiochemical convergence remains yet unclear.

Keywords

pheromone, semiochemical, odorant, off-flavour, olfaction, wine

Introduction

All living things communicate with chemicals. Unlike sounds or sights, semiochemicals interconnect species across the kingdoms, and enable information exchange between animals, plants and microorganisms (Schultz and Appel 2004). A fascinating, recurrent observation is that the same compound is bioactive in different species and context. Evolutionary convergence may result from the widespread occurrence or even physico-chemical properties facilitating information transmission, but is first of all thought to reflect the biological significance of chemicals, including the underlying biochemical pathways and precursors.

Citrus fruit is a preferred oviposition substrate for the vinegar fly *Drosophila melanogaster* (Dweck et al. 2013) provided that yeast is present (Becher et al. 2012). Citrus peel and brewer's yeast both produce linalool (Chisholm et al. 2003; Carrau et al. 2005), which the flies perceive via several odorant receptors (Ors), including DmelOr69a (Münch et al. 2016; Lebreton et al. 2017).

Linalool, commonly found in the headspace of foliage, flowers and fruit, is bioactive in many animals. Plant-produced linalool enhances mate-finding in several phytophagous insects, while other species release linalool as a sex pheromone component (Hefetz et al. 1979; Aldrich et al. 1986; Leal et al. 1993; Yang et al. 2004). Herbivory, on the other hand, upregulates linalool production in plants, which may protect against further infestation (Mithöfer and Boland 2012). The (R) and (S) enantiomers of linalool differentially attract pollinators and herbivores, for feeding and oviposition (Reisenman et al. 2010; Saveer et al. 2012; Raguso 2016), and enantiomeric changes during phenological development modulate our perception of flower aroma (Pragadheesh et al. 2017). In mammals, linalool induces psychopharmacological effects via glutamate receptors (Elisabetsky et al. 1995; Nakamura et al. 2009), perception via Ors produces a sweet, floral note and makes a prominent contribution to the bouquet of flowers, fruit and wine, where both grape and yeast are a source of linalool (Lewinsohn et al. 2001; Carrau et al. 2005; Swiegers et al. 2005).

The response to food and mate olfactory cues is strongly interconnected in *Drosophila* (Lebreton et al. 2015; Gorter et al. 2016; Das et al. 2017) and DmelOr69a is also tuned to the newly identified female pheromone (Z)-4-undecenal (Z4-11Al), in addition to food odorants (Lebreton et al. 2017). Curiously, Z4-11Al is also found in citrus essential oil (Chisholm et al. 2003). Unsaturated aldehydes are prominent constituents of a range of food aromas, including fruit, wine, meat and fish (e.g. Varlet et al. 2006; Cullere et al. 2007, Perez-Cacho and Rouseff 2008; Shi et al. 2013; Yang et al. 2008). Moreover, Z4-11Al is an anal gland volatile in the rabbit (Goodrich et al. 1978).

While collecting volatiles from *D. melanogaster* flies, we discovered that we can reliably distinguish single male from female flies by their scent, which is strongly reminiscent of synthetic Z4-11Al. We then employed a sensory panel to verify whether we can indeed discern single flies, and whether the newly discovered pheromone Z4-11Al contributes to the scent of the female fly.

Materials and methods

Chemicals

Isomeric and chemical purity of synthetic Z4-11Al were 98.6% and >99.9%, respectively, according to gas chromatography coupled to mass spectrometry (6890 GC and 5975 MS, Agilent Technologies, Santa Clara, CA, USA). Ethanol (redistilled, >99.9% purity; Merck, Darmstadt, Germany) was used as solvent.

Sensory evaluation

Eight members (six men and two women) of the sensory panel for organoleptic tests for the wine-growing area of Baden evaluated the odor of *D. melanogaster* and synthetic Z4-11Al. Members of this panel have been trained and selected for the official quality assessment of wines produced in Baden, at the Federal Institute for Viticulture, Freiburg, Germany. Each test comprised three glasses, control and two treatments, which were presented in random order. The panel was asked to score odor intensity, ranging from 1 (weak, silent) to 9 (strong, loud) and to comment on odor quality. The local human subjects committee approved sensory evaluation of Z4-11Al by sniffing. The first test compared the odor from single male and female flies. Flies were kept during 5 min in empty wine tasting glasses (215 ml) and were released shortly before tests. The second test compared a glass impregnated with fly odor and Z4-11Al (10 ng in 10 µl ethanol), which was applied to an empty glass, the solvent was allowed to evaporate during 2 min. Next, 10 ng Z4-11Al or a female fly were added to a glass filled with either water or white

wine (dry Pinot blanc, Freiburg 2013, Staatsweinkellerei Freiburg). The fly was removed after 5 min, prior to testing. Finally, 1 or 5 ng Z4-11Al was added to wine.

Statistical analysis

Odor panel data was analyzed using one-tailed analysis of variance (ANOVA) followed by a Tukey test. Normality was tested using Shapiro-Wilk and homoscedasticity was tested using Levene's test. All analysis were carried out using SPSS v. 20 (IBM Corp, 2011).

Results

D. melanogaster females (Fig. 1) produce a distinctive scent. The sensory panel found the odor of single female flies to be stronger and qualitatively clearly different from male flies (Fig. 2a).

Chemical analysis has shown earlier that Z4-11Al and its precursor, the cuticular hydrocarbon (Z,Z)-7,11-heptacosadiene, are produced by female flies, not by males (Billeter et al. 2009; Lebreton et al. 2017). Our panel tests established that synthetic Z4-11Al has a distinctive odor (Fig. 2b). Moreover, a female fly and 10 ng Z4-11Al were found to be similar, with respect to odor quality and intensity, when presented in an empty glass, in water or wine (Fig. 2c,d,e). Since 10 ng Z4-11Al was assessed as slightly louder than the odor of a fly, we compared Z4-11Al at 1 ng and 5 ng, showing that as little as 1 ng Z4-11Al in a wine glass (corresponding to ca. 5 ng/L or 0.03 nmol/L at the time of application) was clearly perceptible (Fig. 2f). Even at small amounts, Z4-11Al was perceived as a somewhat unpleasant off-flavour.

The detection threshold for Z4-11Al is apparently similar in flies and men, since we clearly sense Z4-11Al released from a single fly (Fig. 2a). Chemical analysis found that *D. melanogaster* females released Z4-11Al at a rate of 2.4 ng/h and solvent extracts of fly cuticula contained 0.3 ng Z4-11Al/female (Lebreton et al. 2017).

Discussion

Sensory evaluation confirmed that we sensitively smell Z4-11Al, the female-produced pheromone of the fruit fly *D. melanogaster* (Lebreton et al. 2017) and that we reliably distinguish single female from male flies. This supports the observation that one fly spoils a glass of wine, after falling into it - provided it is of the female sex. Other fly volatiles may contribute to our perception of fly odor. However, Z4-11Al is the most abundant

compound which is released by females only, other volatiles are found in both sexes (Lebreton et al. 2017).

A straightforward explanation for convergent perception of Z4-11Al is, however, not at hand. The occurrence of Z4-11Al in nature is probably incompletely known and its possible role in humans remains unclear. Z4-11Al may merely be reminiscent of other food aldehydes, or it might avert ingestion of fruit infested with vinegar flies, which may be contaminated with microbes vectored by flies.

A range of mammalian Ors is responsive to aldehydes (e.g. Benbernou et al. 2007; Saito et al. 2009; Nara et al. 2011), including human OR1A1 and OR2W1 (Geithe et al. 2017 a,b). Perception of Z4-11Al effects the heart rate of the rabbit, where it has been found in the anal gland (Goodrich et al. 1978). The positional isomer (*E*)-2-undecenal is a bovid body odor (Gikonyo et al. 2002) and olfactory sensory neurons of ticks (Acari, Ixodidae) tuned to aldehydes afford indirect evidence for aldehydes as vertebrate signals (Steullet and Guerin 1994). A characteristic scent which is reminiscent of tangerine emanates from colonies of crested auklet, a monogamous seabird. Two unsaturated aldehydes, the chain-shortened analog (*Z*)-4-decenal (Z4-10Al) and (*Z*)-2-decenal are main odor-active constituents. In crested auklet, Z4-10Al likely plays a role as an ectoparasite repellent and a signal of mate quality (Douglas et al. 2001; Hagelin et al. 2003; Caro and Balthazart 2010).

Z4-11Al is also part of clementine aroma (Chisholm et al. 2003) and it may play a dual role as social signal and food cue, not only in flies, but also in other animals. The olfactory sense in animals plays a key role during habitat adaptation. Tuning of Ors to habitat cues will create a bias for mate-finding signals that match or are structurally similar to habitat odorants (Endler 1992). This idea yields a tentative scenario for the convergence of semiochemicals. Insects and other animals feed on fruit, containing associated yeasts that facilitate digestion of plant materials, provide nutrients and protection of food from antagonistic microorganisms. Animal-produced compounds sharing structural motifs may have secondarily been adopted as mating signals, via established sensory channels dedicated to habitat and food odorants.

The olfactory system of *Drosophila* is conveniently accessible to experimental investigation and current research extends beyond the Or ligand repertoire (Münch and Galizia 2016) to neural circuits underlying odor-mediated behavior (Kohl et al. 2013; Auer and Benton 2016; Seki et al. 2017), chemical ecology (Depetris-Chauvin et al. 2015; Mansourian and Stensmyr 2015) and phylogenetic diversification (Shiao et al. 2015; Arguello et al. 2016; Ramasamy et al. 2016). A future challenge is to extend functional, behavioral, ecological and phylogenetic studies to include vertebrates, towards an understanding of the chemical vocabulary that interconnects us with other living things.

Acknowledgements

We thank the members of the sensory panel (Freiburg, Germany) for evaluating the fly scent and Cyrus Mahmoudi (comgraphix.de, Germany) for sharing a fruit fly photograph. Supported by the Linnaeus initiative "Insect Chemical Ecology, Ethology and Evolution" (Formas, SLU).

Data accessibility. Data is completely included in the figure.

Authors' contributions. P.G.B. sensed the fly scent, P.G.B., S.L., M.B. and V.J. conceived the idea and contributed to the experiment, F.B. calculated statistics, E.W. and E.H. synthesized the test chemical, P.W. supervised the project and wrote the manuscript, all authors contributed to and approved the final version of the manuscript.

Competing interests. Authors declare no competing interests.

References

- Aldrich JR, Lusby WR, Kochansky JP (1986) Identification of a new predaceous stink bug pheromone and its attractiveness to the eastern yellowjacket. *Cell Molec Life Sc* 42:583-585. (doi:10.1007/BF01946714)
- Arguello JR, Cardoso-Moreira M, Grenier JK, Gottipati S, Clark AG, Benton R (2016) Extensive local adaptation within the chemosensory system following *Drosophila melanogaster*'s global expansion. *Nature Comm* 7:11855 (doi:10.1038/ncomms11855)
- Auer TO, Benton R (2016) Sexual circuitry in *Drosophila*. *Curr Op Neurobiol* 38:18-26. (doi:10.1016/j.conb.2016.01.004)
- Becher PG, Flick G, Rozpedowska E, Schmidt A, Hagman A, Lebreton S, Larsson MC, Hansson BS, Piskur J, Witzgall P, Bengtsson M (2012) Yeast, not fruit volatiles mediate attraction and development of the fruit fly *Drosophila melanogaster*. *Funct Ecol* 26:822-828 (doi:10.1111/j.1365-2435.2012.02006.x)
- Benbernou N, Tacher S, Robin S, Rakotomanga M, Senger F, Galibert F (2007) Functional analysis of a subset of canine olfactory receptor genes. *J Heredity* 98:500-505. (doi:10.1093/jhered/esm054)
- Billeter JC, Atallah J, Krupp JJ, Millar JG, Levine JD (2009) Specialized cells tag sexual and species identity in *Drosophila melanogaster*. *Nature* 461:987-U250. (doi:10.1038/nature08495)
- Caro SP, Balthazart J (2010) Pheromones in birds: myth or reality? *J Comp Physiol A* 196:751-766. (doi:10.1007/s00359-010-0534-4)
- Carrau FM, Medina K, Boido E, Farina L, Gaggero C, Dellacassa E, Versini G, Henschke PA (2005) De novo synthesis of monoterpenes by *Saccharomyces cerevisiae* wine yeasts. *FEMS Microbiol Lett* 243:107-15. (doi:10.1016/j.femsle.2004.11.050)
- Chisholm MG, Jell JA, Cass DM. 2003 Characterization of the major odorants found in the peel oil of *Citrus reticulata* Blanco cv. Clementine using gas chromatography-olfactometry. *Flavour Fragrance J* 18:275-281. (doi:10.1002/ffj.1172)
- Cullere L, Cacho J, Ferreira V (2007) An assessment of the role played by some oxidation-related aldehydes in wine aroma. *J Agric Food Chem* 55:876-881. (doi:10.1021/jf062432)
- Das S, Trona F, Khallaf MA, Schuh E, Knaden M, Hansson BS, Sachse S. 2017. Electrical synapses mediate synergism between pheromone and food odors in *Drosophila melanogaster*. *Proc Natl Acad Sc USA* 114:9962-9971. (doi:10.1073/pnas.1712706114)
- Depetris-Chauvin A, Galagovsky D, Grosjean Y (2015) Chemicals and chemoreceptors: ecologically relevant signals driving behavior in *Drosophila*. *Front Ecol Evol* 3:41. (doi:10.3389/fevo.2015.00041)
- Douglas H, Jones T, Conner W (2001) Heteropteran chemical repellents identified in the citrus odor of a seabird (crested auklet: *Aethia cristatella*): evolutionary convergence in chemical ecology. *Naturwissenschaften* 88:330-332. (doi:10.1007/s001140100236)

216 Dweck HK, Ebrahim SA, Kromann S, Bown D, Hillbur Y, Sachse S, Hansson BH, Stensmyr
217 MC (2013) Olfactory preference for egg laying on citrus substrates in *Drosophila*.
218 Curr Biol 23:2472–80. (doi: 10.1016/j.cub.2013.10.047)

219 Elisabetsky E, Marschner J, Souza DO (1995) Effects of linalool on glutamatergic system
220 in the rat cerebral cortex. Neurochem Res 20:461-465. (doi:10.1007/BF00973103)

221 Endler JA (1992) Signals, signal conditions, and the direction of evolution. Am Naturalist
222 139:S125-S153. (doi:10.1086/285308)

223 Geithe C, Noe F, Kreissl J, Krautwurst D (2017a) The broadly tuned odorant receptor
224 OR1A1 is highly selective for 3-methyl-2, 4-nonanedione, a key food odorant in aged
225 wines, tea, and other foods. Chemical Senses 42:181-193.
226 (doi:10.1093/chemse/bjw117)

227 Geithe C, Protze J, Kreuchwig F, Krause G, Krautwurst D (2017b) Structural determinants
228 of a conserved enantiomer-selective carvone binding pocket in the human odorant
229 receptor OR1A1. Cell Molec Life Sci 74:4209-4229. (doi:10.1007/s00018-017-2576-
230 z)

231 Gikonyo NK, Hassanali A, Njagi PG, Gitu PM, Midiwo JO (2002) Odor composition of
232 preferred (buffalo and ox) and nonpreferred (waterbuck) hosts of some savanna
233 tsetse flies. J Chem Ecol 28:969-981. (doi:10.1023/A:1015205716921)

234 Goodrich BS, Hesterman ER, Murray KE, Mykytowycz R, Stanley G, Sugowdz G. 1978
235 Identification of behaviorally significant volatile compounds in the anal gland of the
236 rabbit, *Oryctolagus cuniculus*. J Chem Ecol 4:581-594. (doi:10.1007/BF00988922)

237 Gorter JA, Jagadeesh S, Gahr C, Boonekamp JJ, Levine JD, Billeter JC (2016) The
238 nutritional and hedonic value of food modulate sexual receptivity in *Drosophila*
239 *melanogaster* females. Sci Rep 6:19441. (doi:10.1038/srep19441)

240 Hagelin JC, Jones IL, Rasmussen LEL (2003) A tangerine-scented social odour in a
241 monogamous seabird. Proc R Soc B 270:1323-1329. (doi: 10.1098/rspb.2003.2379)

242 Hefetz A, Batra SWT, Blum MS (1979) Linalool, neral and geranial in the mandibular
243 glands of *Colletes* bees - an aggregation pheromone. Cell Molec Life Sc 35:319-320.
244 (doi:10.1007/BF01964324)

245 Kohl J, Ostrovsky AD, Frechter S, Jefferis GS (2013) A bidirectional circuit switch reroutes
246 pheromone signals in male and female brains. Cell 155:1610-1623.
247 (doi:10.1016/j.cell.2013.11.025)

248 Leal WS, Sawada M, Matsuyama S, Kuwahara Y, Hasegawa M (1993) Unusual periodicity
249 of sex pheromone production in the large black chafer *Holotrichia parallela*. J Chem
250 Ecol 19:1381-1391. (doi:10.1007/BF00984883)

251 Lebreton S, Trona S, Borrero-Echeverry F, Bilz F, Grabe V, Becher PG, Carlsson MA,
252 Nässel DR, Hansson BS, Sachse S, Witzgall P (2015) Feeding regulates sex
253 pheromone attraction and courtship in *Drosophila* females. Sci Rep 5:13132
254 (doi:10.1038/srep13132)

255 Lebreton S, Borrero-Echeverry F, Gonzalez F, Solum M, Wallin E, Hedenström E, Hansson
256 BS, Gustavsson A-L, Bengtsson M, Birgersson G, Walker WB, Dweck H, Becher PG,
257 Witzgall P (2017) A *Drosophila* female pheromone elicits species-specific long-range

258 attraction via an olfactory channel with dual specificity for sex and food. BMC Biology
259 15:88. (doi:10.1186/s12915-017-0427-x)

260 Lewinsohn E, Schalechet F, Wilkinson J, Matsui K, Tadmor Y, Nam KH, Amar O,
261 Lastochkin E, Larkov O, Ravid U, Hiatt W, Gepstein S, Pichersky E (2001) Enhanced
262 levels of the aroma and flavor compound S-linalool by metabolic engineering of the
263 terpenoid pathway in tomato fruits. Plant Physiol 127:1256-1265.
264 (doi:10.1104/pp.010293)

265 Mansourian S, Stensmyr MC (2015) The chemical ecology of the fly. Curr Op Neurobiol
266 34:95-102. (doi:10.1016/j.conb.2015.02.006)

267 Mithöfer A, Boland W (2012) Plant defense against herbivores: chemical aspects. Annu
268 Rev Plant Biol 63:431-450. (doi:10.1146/annurev-arplant-042110-103854)

269 Münch D, Galizia CG. 2016 DoOR 2.0 - comprehensive mapping of *Drosophila*
270 *melanogaster* odorant responses. Sci Rep 6:21841. (doi:10.1038/srep21841)

271 Nakamura A, Fujiwara S, Matsumoto I, Abe K (2009) Stress repression in restrained rats
272 by (R)-(-)-linalool inhalation and gene expression profiling of their whole blood cells.
273 J Agric Food Chem 57:5480-5485. (doi:10.1021/jf900420g)

274 Nara K, Saraiva LR, Ye X, Buck LB (2011) A large-scale analysis of odor coding in the
275 olfactory epithelium. J Neurosci 31:9179-9191. (doi:10.1523/jneurosci.1282-11.201)

276 Perez-Cacho PR, Rouseff RL (2008) Fresh squeezed orange juice odor: a review. Critical
277 Rev Food Sci Nutr 48:681-695. (doi: 10.1080/10408390701638902)

278 Pragadheesh VS, Chanotiya CS, Rastogi S, Shasany AK (2017) Scent from *Jasminum*
279 *grandiflorum* flowers: investigation of the change in linalool enantiomers at various
280 developmental stages using chemical and molecular methods. Phytochemistry
281 140:83-94. (doi:10.1016/j.phytochem.2017.04.018)

282 Raguso RA (2016) More lessons from linalool: insights gained from a ubiquitous floral
283 volatile. Curr Op Plant Biol 32:31-36. (doi:10.1016/j.pbi.2016.05.007)

284 Ramasamy S, Ometto L, Crava CM, Revadi S, Kaur R, Horner DS, Pisani D, Dekker T,
285 Anfora G, Rota-Stabelli O (2016) The evolution of olfactory gene families in
286 *Drosophila* and the genomic basis of chemical-ecological adaptation in *Drosophila*
287 *suzukii*. Gen Biol Evol 8:2297-2311. (doi:10.1093/gbe/evw160)

288 Reisenman CE, Riffell JA, Bernays EA, Hildebrand JG (2010) Antagonistic effects of floral
289 scent in an insect-plant interaction. Proc R Soc B 277:2371-2379.
290 (doi:10.1098/rspb.2010.0163)

291 Saito H, Chi Q, Zhuang H, Matsunami H, Mainland JD (2009) Odor coding by a
292 mammalian receptor repertoire. Science Signal 2(60):ra9.
293 (doi:10.1126/scisignal.2000016)

294 Saveer AM, Kromann S, Birgersson G, Bengtsson M, Lindblom T, Balkenius A, Hansson BS,
295 Witzgall P, Becher PG, Ignell R (2012) Floral to green: mating switches moth
296 olfactory coding and preference. Proc R Soc B 279:2314-2322.
297 (doi:10.1098/rspb.2011.2710)

298 Schultz JC, Appel HM (2004) Cross-kingdom cross-talk: hormones shared by plants and
299 their insect herbivores. Ecology 85:70-77. (doi:10.1890/02-0704)

- Seki Y, Dweck HK, Rybak J, Wicher D, Sachse S, Hansson BS (2017) Olfactory coding from the periphery to higher brain centers in the *Drosophila* brain. BMC Biol 15:56 (doi:10.1186/s12915-017-0389-z)
- Shi X, Zhang X, Song S, Tan C, Jia C, Xia S (2013) Identification of characteristic flavour precursors from enzymatic hydrolysis-mild thermal oxidation tallow by descriptive sensory analysis and gas chromatography-olfactometry and partial least squares regression. J Chromatogr B 913:69-76. (doi:10.1016/j.jchromb.2012.11.032)
- Shiao MS, Chang JM, Fan WL, Lu MY, Notredame C, Fang S, Kondo R, Li WH (2015) Expression divergence of chemosensory genes between *Drosophila sechellia* and its sibling species and its implications for host shift. Gen Biol Evolution 7:2843-2858. (doi:10.1093/gbe/evv183)
- Steullet P, Guerin PM (1994) Identification of vertebrate volatiles stimulating olfactory receptors on tarsus I of the tick *Amblyomma variegatum* Fabricius (Ixodidae). 1. Receptors within the Haller's organ capsule. J Comp Physiol A 174:27-38. (doi:10.1007/BF00192003)
- Swiegers JH, Bartowsky EJ, Henschke PA, Pretorius IS (2005) Yeast and bacterial modulation of wine aroma and flavour. Austral J Grape Wine Res 11:139-173. (doi:10.1111/j.1755-0238.2005.tb00285.x)
- Varlet V, Knockaert C, Prost C, Serot T (2006) Comparison of odor-active volatile compounds of fresh and smoked salmon. J Agric Food Chem 54:3391-3401. (doi:10.1021/jf053001p)
- Yang Z, Bengtsson M, Witzgall P (2004) Host plant volatiles synergize response to sex pheromone in codling moth, *Cydia pomonella*. J Chem Ecol 30:619-629. (doi:10.1023/B:JOEC.0000018633.94002.af)
- Yang DS, Shewfelt RL, Lee KS, Kays SJ (2008) Comparison of odor-active compounds from six distinctly different rice flavor types. J Agric Food Chem 56:2780-2787. (doi:10.1021/jf072685t)

Legends

Fig. 1 Fruit fly *D. melanogaster* female with exposed ovipositor on blueberry (Photo by Cyrus Mahmoudi).

Fig. 2 Sensory evaluation of fly odor and synthetic (Z)-4-undecenal (Z4-11Al). Odor intensity scale ranges from 1 (weak) to 9 (strong), symbols show evaluation by individual test panel members, mean intensity ratings followed by different letters are significantly different ($p < 0.001$). Olfactory intensity of (a) the odor of a single *D. melanogaster* male and female fly adsorbed during 5 min in an empty wine glass ($F=96.711$), (b) 10 ng synthetic Z4-11Al and solvent (ethanol) ($F=106.732$), (c) 10 ng Z4-11Al and the odor of a single *D. melanogaster* female fly in an empty glass ($F=34.720$), (d) in a glass with

337 water ($F=16.689$), (e) in a glass with wine ($F=12.952$), (f) 1 ng and 5 ng Z4-11Al in a
338 glass with wine ($F=110.694$).