Specific miRNA-GPCR networks regulate Sox9a/Sox9b activities to promote gonadal renewal in zebrafish

Xinlu Du^{1*}, Huiping Guo^{1*}, Ying Zhang¹, Jiacheng Wu¹, Minyou Li^{1,3,#}, Xianxian Hua^{2,#}, Jizhou Yan^{1,3,#}

1. Department of Developmental Biology, College of Fisheries and Life Science, Shanghai Ocean University;

2. Department of Cancer Biology, University of Pennsylvania School of Medicine

3. Key Laboratory of Exploration and Utilization of Aquatic Genetic Resources, Ministry of Education.

* The authors are regarded as joint First Authors

Correspondence to: jyan2@shou.edu.cn, huax@mail.upenn.edu, myli@shou.edu.cn

Key words: miRNA network, GPCR, Sox9, gonadal rejuvenation, zebrafish.

Running title: miRNA-GPCR networks and zebrafish gonadal renewal

Total characters: 6582, including abstract 169

Abstract

Fertility and endocrine function rely on a tightly regulated synchronicity within the hypothalamic-pituitary gonadal (HPG) axis. FSH/cAMP/MAPK/ Sox9 axis signaling and its regulated specific miRNAs are thought to regulate vertebrate gonadal development and sex differentiation, and yet the regulatory networks are largely unknown. Here we construct small RNA and mRNA libraries from sexually matured ovary and testis of zebrafish to identify specific miRNA-target pairs. Integration of Targetscan prediction and *in vivo* induced gene expression highlight four specific miRNAs that conditionally target three G protein-coupled receptor (GPCR) –Sox9 signaling genes, and implicate two regulatory circuits of miR430a-Sox9a in the testis and miR218a-Sox9b in the ovary. Co-injected Sox9a-miR430a mixture increases the proportion of spermatogonia but degenerates primary oocyte, while Sox9b-miR218a mixture induces renewal of ovarian follicles. Co-immunoprecipitation and mass-spectrometry analyses further reveal that miR430a and Sox9a synergistically activate testicular PKC/Rock1 signals while miR218a and Sox9b constrict ovary PKC/PI3K/Rock1 signaling. These results clarify specific miRNAs-GPCR regulatory networks of Sox9a/Sox9b switch, and also provide mechanistic insight into gonadal rejuvenation and plasticity.

Introduction

Although cell phenotypic changes often occur in stem cell-based differentiation, dedifferentiation or transdifferentiation, both stem cells and differentiated cells are able to reversibly dedifferentiate and transform into cells of different lineages under certain conditions. This phenomenon, called environmentally guided cell plasticity is the basis of tissue homeostasis, tissue regeneration, cancer cell recover (Galliot & Ghila, 2010), and sex reversal (Sun, Zhang et al., 2013). One excellent system to study cell plasticity is the vertebrate gonad, where distinct populations of somatic and germ stem cells give rise to different sex-dependent morphotypes: male testiculogenesis and female folliculogenesis (Chen, Zheng et al., 2013, Gassei & Schlatt, 2007). The early gonad is an undifferentiated primordium composed of bipotential somatic stem cells: precursors for supporting cells and steroid-secreting cells. When the primordial germ cells (PGCs) reside in the gonadal ridge, the supporting cell precursors develop into either testis-specific Sertoli cells or ovary-specific follicle (granulosa) cells. Meanwhile the immigrating mesonephric cells give rise to peritubular myoid (PM) cells, endothelial cells that form the male-specific vasculature, and fetal steroidogenic-Leydig cells (or ovarian theca

cells in female) (McClelland, Bowles et al., 2012, Wilhelm, Palmer et al., 2007). In parallel to the differentiation of somatic precursors, PGCs differentiate into oogonia in ovarian follicles or into spermatogonia in Sertoli cells-dominant testis cord.

Histologically, testiculogenesis with cord formation and Leyding cell differentiation supports spermatogenesis while folliculogenesis accompanies ootidogenesis, where the ovarian follicle surrounding the oocyte develops from a primordial follicle to a preovulatory one. There is a close anatomical relationship between the development of the genital ridge and the hormone excretory system during early ontogeny of all vertebrates, including fish. In both teleosts and mammals the testis contains Sertoli and Leydig cells in addition to germ cells, and the ovary consists of the thecal cells and granulosa cells surrounding the ovum. The gonadal sex differentiation involves the same sequential cellular events for spermatogenesis and folliculogenesis, in which ovarian development is the default pathway. Without Sox9-related signaling intervention, the bipotential gonads follow the ovarian pathway (McClelland et al., 2012, Sun et al., 2013).

An important difference is that the mammalian gonads are terminally developed into either testis or ovary by Sry-Sox9 genetic intervention, while fish gonads often retain the ability to change, making them sequential hermaphrodites. Despite the vast diversity of primary sex-determining mechanisms in vertebrate species, Sox9-related signals appear to be a conserved mechanism to switch on testis-differentiation pathway, where the expressions of *sox9* and its downstream genes (*amh* and *cyp19a1*) are proposed to be regulated by gonadotropins and its second messenger cAMP through MAPK signaling pathways (Kobayashi, Chang et al., 2005, Taieb, Grynberg et al., 2011). Intricately, activation of Sox9-MAPK pathways are also linked to chondrocyte differentiation (Murakami, Kan et al., 2000). Accumulating evidence implicates versatile functions of Sox9 in cell fate specification, stem cell biology, human diseases and tissue regeneration (Jo, Denduluri et al., 2014). Therefore, identifying partner factors, signaling pathways, post-transcriptional modifications, and integrative signaling pathways could facilitate us to better understand Sox9's versatility in cell fate switch and adaptive plasticity.

Zebrafish has two co-orthologs of *sox9* on its duplicated chromosomes: *sox9a* and *sox9b*. The pair of *sox9* genes show distinct and overlapping functions in zebrafish embryonic development (Rodriguez-Mari, Yan et al., 2005, Yan, Willoughby et al., 2005). In the adults, *sox9a* is expressed in many tissues including testis, whereas *sox9b* expression is restricted to previtellogenic oocytes of the ovary (Chiang, Pai et al., 2001).

Correspondingly two Sox9 downstream genes, *amh* and *cyp19a1a*, are also respectively expressed in the juvenile testis and ovary (Rodriguez-Mari et al., 2005). Therefore, zebrafish bipotential gonad is a unique system to explore more informative *sox9* axis signaling pathways controlling gonadal sexual fate. MicroRNAs (miRNAs) represent a second regulatory network that regulates gene expression in various biological processes by inducing degradation or translational inhibition of their target mRNAs (Tao, Sun et al., 2016). Based on miRNA expression in the gonads at the time of sexual differentiation, a number of miRNAs and the potential targets are identified. Some specific miRNAs show regulatory activities in gonad development and/or sex determination (Kang, Cui et al., 2013, Xiao, Zhong et al., 2014). Even though miR-124 was reported to be sufficient to induce the repression of both SOX9 translation and transcription in ovarian cells (Real, Sekido et al., 2013), no the specific miRNA-*sox9* axis regulatory networks have been identified so far (Tao et al., 2016). In this study we performed genome-wide search for specific miRNA-target pairs, and investigated physiological interaction effects between specific miRNAs and the target genes. Particularly, we focused on specific miRNA- Sox9a/Sox9b networks in regulation of testiculogenesis and folliculogenesis.

Results

Identification of genes related to sex-specific fate during the process of gonadal differentiation maturation

To identify genes functioning in folliculogenesis and testiculogenesis, we dissected the freshly matured gonads at 5mpf (months post-fertilization), and separated the ovary and testis for RNA deep sequencing analyses. Out of 26,459 genes (Table S1), 25,609 genes were expressed in either or both sex gonads. 18,993 genes exhibited 2 fold or more expression changes (Table S2), and 729 genes were significantly differentially expressed in testis or ovary ($P < 0.05$) (Table S3), including 421 ovary-, and 308 testis-highly expressed (Table S4). As the predominant expression of these induced genes synchronized with the reproductive growth cycle (testicular or follicle differentiation), we categorized 729 biased genes as sex-preferential effectors and/or regulators.

Our initial goal was to clarify the proposed gonadotropins-cAMP-MAPK- Sox9 signaling pathways during sex gonadal differentiation, Gonadotropins including follicle-stimulating hormone (FSH) and luteinizing

hormone (LH), are glycoprotein peptide hormones. Gonadotropin receptors are coupled to the G-protein system. We searched the two transcriptome datasets for GPCR-*sox9* axis genes. These included 92 GPCR, 35 cAMP, 34 MAPKs, and 21 p53 orthologs as well as 30 sex determinant genes. The vast majority of the GPCR-*sox9* axis genes (201/212) did not show significant differential expression between testis and ovary (Table S5). This result conformed to the phenomena that miRNAs only modestly downregulate the mRNA level of their target genes (Selbach, Schwanhauser et al., 2008).

Identification of specific miRNAs and GPCR-*sox9* axis target genes for gonadal differentiation

To identify specific miRNAs-target pairs in reproductive cycle, we analyzed two small RNA libraries, which were simultaneously constructed from the same ovary or testis total RNA samples as the mRNA libraries were made. After aligning the small RNA sequences with the miRBase (zebrafish miRNA database), bioinformatic analysis identified 350 unique mature miRNAs, and 346 miRNA precursors. 314 miRNAs were found in either of the two libraries, of which 51 miRNAs showed 2-fold higher transcription in testis (refer to testis miRNAs) and 106 miRNAs were preferentially transcribed in ovary (ovary miRNAs).

Using TargetScan to search for the target pairs between 157 sex-biased miRNAs and 209 GPCR-*sox9* axis genes, we found that 97 miRNAs targeted 183 GPCR-*Sox9* axis genes, including those multiple and cross targets (Table S7). For example *sox9a* was targeted by 25 miRNAs and *sox9b* was targeted by 20 miRNAs.

Four specific miRNAs represent putative regulators in GPCR-*sox9*-axis

To derive a short-list of specific miRNAs as putative regulators in GPCR-*sox9* axis, we selected the representative miRNAs-target pairs according to the criteria in Table 1. Combined with the expression pattern of endogenous pre-miRNAs (Fig.S1), four potentially interesting miRNAs were accordingly selected as the most overrepresented miRNAs: miR430a (multiple splicing isoforms and 17 multiple targets in GPCR-*sox9* axis of signaling cascades, not directly targeting *sox9*), miR218a (50 targets, not directly targeting *sox9* but targeting *vasa*), miR734 (95 targets, directly targeting *sox9a*), and miR141 (72 targets, directly targeting *sox9b*) (Table S7).

Table 1 A short list of overrepresented miRNAs

Criteria#	Ovary	Testis
-----------	-------	--------

Highly and differentially transcribed (expression abundance ≥ 100 , log2 ratio ≥ 1.5)	miR-23b, -724, -181a-5p, -135c, -200c-3p, -218b, R-218a, -200a-3p, -21, -455-5p, -144-3p, -455-2-5p, -194a, -146a, -181c-5p, -200a-5p, -181b-5p, -216a, -429a, -200b-3p, -34a, -223, -122, -184, -192, -181a-2-3p, -let-7i, -34c-5p, -34b, -146b, -34c-3p, -430b-5p, -430b-3p, -430a-3p	miR-723-3p, -2187-5p, -735-3p, -2189, -499-5p, -132-3p, -193b-3p, -7a, -458-3p, -7b
Capable of targeting multiple genes (target genes ≥ 25 for ovary miRNAs; target genes ≥ 14 for testis miRNAs)	miR-218a, -737, -734, -724, -181a, -144, -200b, -129, -139, -455, -459, -27a, -223, -216a, -122, -194a, -181a, -216b, -2192, -192, -205, -17a, -34b	miR-203a, -128, -9, -182, -2187, -2189, -499, -723, -124, -219, -134, -203a, -128, -124, -499, -2189
Commonly prominent in both gonads	miR-141, -363, -23a	miR-141, -363, -23a
Non target*	miR-301c, -301b	miR-301c, -301b

123 #, 1) miR-430a has multiple versions, such as miR-430a-3p, 5p; -4-5p; 11-17-5p, etc. Different versions of
124 miRNAs are indistinguishable in Targetscan. 2) The *vasa* gene is targeted by miR218a. *vasa* expression is
125 seen in germ cells, specifically the germline stem cells (GSC's) of female ovaries and in the early stages of
126 spermatogenesis in the male testis. 3) The miR-430/427/302 family is crucial for vertebrate embryogenesis by
127 controlling germ layer specification (Rosa, Spagnoli et al., 2009).

128 *, non-target miRNAs as negative control.

129 Interaction modes between specific miRNAs and GPCR-*sox9* targets in testis and ovary

130 To investigate the physiological interactions between four specific miRNAs and GPCR-*sox9* axis targets, we
131 devised an *in vitro* gonadal microinjection strategy to enforce exogenous expression of Sox9a/Sox9b and four
132 specific miRNAs in the gonads, and evaluated their effects on the expression of GPCR-*sox9* axis targets
133 (Table S8). When the Targetscan prediction and RT-PCR examinations were integrated together, four
134 interaction modes emerged (Table 2) : 1) a negative specific miRNA-target pair: a predicted target was
135 downregulated in both gonads by a specific miRNA; 2) a positive specific miRNA-target pair: a predicted
136 target was upregulated in both gonads by a specific miRNA; 3) a conditionally specific miRNA-target pair: a
137 predicted target was upregulated by a specific miRNA in one sex gonad but downregulated in the other gonad;
138 and 4) an indirect specific miRNA-target: a gene without specific miRNA binding sites was downregulated or
139 upregulated. Totally 92.3% (12 of 13) of predicted targets of GPCR-*sox9* genes were correctly scored as direct
140 targets in testis and/or ovary by at least one of four specific miRNAs. Fig.1 showed crossing alignment

between four specific miRNAs and 15 putative target genes.

Table 2 Interactions between specific miRNAs and GPCR-Sox9 axis*

Over-expression	<i>gpr157</i>	<i>gpr173</i>	<i>grk1a</i>	<i>grk7a</i>	<i>lgr4</i>	<i>grk5l</i>	<i>gpr161</i>	<i>grk4</i>	<i>gpr146</i>	<i>gpr137c</i>	<i>grk6</i>	<i>amh</i>	<i>sox9a</i>	<i>sox9b</i>	<i>cyp19a1a</i>
Testis microinjection															
sox9a	29.10 ±5.8	62.49 ±3.99	36.40±8.9	1612.38 ±412.3	12.81 ±0.63	8.51 ±0.94	103.96 ±13.01	-2.73 ±0.69	77.45 ±3.99	36.86 ±8.4	23.05 ±3.2	3.10 ±1.1	9042.18 ±2068.0	-11.52 ±8.4	-1.78 ±0.1
sox9b	1.27 ±0.41	-1.46 ±0.29	5.02 ±0.74	208.28 ±36.60	-7.64 ±1.85	11.91 ±6.98	22.82 ±9.8	18.54 ±5.2	2.07 ±0.25	5.00 ±0.19	3.27 ±0.9	2.24 ±0.43	135.88± 20.28	1095.05 ±62.07	9.73 ±2.33
miR-430a-3p	-1.18 ±0.26	1.55 ±0.04	1.56 ±0.29	-2.03 ±0.27	-5.10 ±0.26	-4.87 ±0.27	2.37 ±0.10	-11.51 ±4.60	1.06 ±0.10	1.31 ±0.06	-2.14 ±0.27	-1.49 ±0.12	35.10 ±2.1	-72.18 ±18.4	-79.74 ±4.6
miR-218a	-1.30 ±0.04	1.11 ±0.11	-1.66 ±0.01	22.24 ±0.33	1.02 ±0.03	-1.38 ±0.14	1.53 ±0.28	-1.10 ±0.23	-1.64 ±0.80	-1.24 ±0.03	-1.31 ±0.15	-1.50 ±0.19	-3.78 ±0.80	1.09 ±0.12	-2.17 ±0.16
miR-141-3p	-2.21 ±0.09	-4.15 ±0.26	-3.20 ±0.42	ND	-4.34 ±0.32	1.37 ±0.04	-2.84 ±0.47	15.91 ±2.85	-2.90 ±0.03	-2.68 ±0.12	-3.27 ±0.30	91.76 ±11.07	14.08 ±1.42	-457.87 ±41.48	-1.68 ±0.01
miR-734	-1.24 ±0.02	-3.38 ±0.34	-1.15 ±0.02	ND	-3.23 ±0.26	-1.84 ±0.21	-1.62 ±0.16	19.28 ±4.23	-3.75 ±0.17	-2.90 ±0.33	-2.59 ±0.12	117.69 ±21.36	-6.47 ±0.55	-166804 ±5099.0	2.80 ±0.28
Ovary microinjection															
sox9a	3.80 ±0.17	1.86 ±0.78	3.05 ±0.29	6.01 ±0.47	1.26 ±0.01	2.26 ±0.22	8.46 ±1.91	2.98 ±0.03	7.64 ±2.21	18.79 ±3.42	2.96 ±0.32	10.26 ±0.77	954.24 ±622.2	5.02 ±0.21	1.29 ±0.18
sox9b	14.61 ±7.37	28.32 ±3.20	26.77 ±5.26	54.88 ±4.76	10.02 ±1.52	16.63 ±6.86	169.36 ±29.59	5.52 ±0.12	45.68 ±4.42	74.58 ±34.84	27.50 ±2.41	7.25 ±3.57	984.32 ±40.25	1506.45 ±450.2	9.46 ±5.64
miR-430a-3p	-7.50 ±0.42	-7.10 ±0.14	-9.22 ±1.73	-37.29 ±3.35	-7.34 ±0.25	-8.32 ±0.32	-8.53 ±0.55	-13.60 ±1.22	-7.25 ±1.21	-4.44 ±0.19	-6.41 ±0.12	3.82 ±0.44	-9.10 ±2.32	-2.19 ±0.13	-9.59 ±0.35
miR-218a	-9.13 ±0.76	-1.29 ±0.26	-3.00 ±0.72	2.61 ±0.20	-1.28 ±0.15	-5.41 ±0.05	-4.07 ±1.24	-8.90 ±1.01	-1.30 ±0.61	-3.34 ±0.09	-2.09 ±0.16	1.25 ±0.09	-1.87 ±0.10	1.19 ±0.06	-5.46 ±0.42
miR-141-3p	-6.62 ±0.68	-35.00 ±6.40	-24.51 ±6.21	ND	-35.95 ±2.32	-1.37 ±0.04	-6.83 ±0.40	-4.97 ±0.91	-12.32 ±0.01	-1.59 ±0.15	-15.20 ±2.14	1.74 ±0.31	5.47 ±1.35	-82.96 ±25.89	2.45 ±0.16
miR-734	3.64 ±0.17	-5.20 ±1.41	-160.54 ±3.11	ND	-5.23 ±0.50	8.07 ±0.06	1.09 ±0.06	1.18 ±0.18	-2.99 ±0.11	-1.02 ±0.07	-1.98 ±0.36	7.12 ±1.46	19.15 ±3.12	-9.35 ±2.91	1.56 ±0.17

*, 27 Predicted targets in bold; 17 negative specific miRNA-target pairs in bold and underlined; 2 positive specific miRNA-target pairs in bold and italicized; 8 conditional specific miRNA-target pairs and one undetected (ND) in bold only. The remaining is indirect specific miRNA-targets.

Fig.1. Interaction modes between four specific miRNAs and GPCR-Sox9a/b targets

Specific miRNAs-GPCR network regulate Sox9a/Sox9b signaling at post-transcription and protein level

Previous study has identified several *sox9* downstream genes (*cyp19a*, *cyp19b*, *amh*) as potential targets of miR430 family during sexual transformation of the rice field eel (Gao, Guo et al., 2014). Here we found that miR734 and miR141 formed target-pairs or conditional target pairs with *sox9a*, *amh* and/or *sox9b*, while miR430a and miR218a only showed indirect interactions with these genes and *cyp19a1a* (Fig.1). Referred to *sox9a* and *sox9b* transcription, increased miR141 and miR734 showed similar regulatory functions as miR430a (Table S8). Additionally three miR430a direct target genes (*lgr4*, *grk5l*, *grk4*) were also pair-targets or conditional pair-targets for miR141, miR734 and miR218a (Fig.1). Thus, the four miRNAs may coordinate to regulate Sox9a/Sox9b activity through GPCR-signaling networks.

We then investigated the regulatory modes between the four specific miRNAs and two Sox9 isoforms. Sustained Sox9b expression increased the transcription of four pre-miRNAs in both ovary and testis, whereas enforced Sox9a only induced pre-miR430a transcription in testis (Fig.S2). Similarly, overexpressed Sox9a and Sox9b activated all tested GPCR-*sox9* genes in the ovary whereas a few genes were suppressed in testis.

164 When mature miR430a oligos were injected into the testis, *sox9a* transcript was increased but *sox9b* transcript
165 was strongly reduced. In the ovary, elevated miR430a somewhat reduced transcripts of *sox9a*, *sox9b* and
166 *cyp19a1a*. In contrast to miR430a-Sox9a reciprocal activation in the testis, miR218a downregulated
167 transcripts of *sox9a* and *cyp19a1a* but modestly upregulated *sox9b* transcript in both testis and ovary.
168 These results implicated two transcriptional regulatory circuits: miR430a-Sox9a, and miR218a-Sox9b.
169 Relative to miR218a-Sox9b's extensive and ovary-biased regulation, miR430a-Sox9a synergic regulation
170 was specific in the testis. When we constructed a GFP reporter for 3'UTR of *sox9a* and *sox9b*, and carried out
171 GFP reporter assays in zebrafish embryonic fibroblast cells (Pac2), we found that miR430a and miR218a not
172 only regulates the two Sox9 genes' transcription, but also confines the two proteins' subcellular distribution
173 (Fig.S4, S5).

174 **Gonadal microinjection of specific miRNAs-Sox9 mixture cause renewal of gonocytes**

175 To evaluate two miRNA-Sox9 regulatory circuits and its role in the gonadal differentiation, we conducted a
176 combinatory microinjection into mature gonads (5mpf), and then compared cellular events of somatic cells
177 and germline cells (Fig.2A, 2B). Enforced Sox9a-miR430a mixture expression increased the proportion of
178 mitotic condensed chromatin nuclear cell (Cn) and undifferentiated gonocytes (Gc), and increased
179 degeneration of primary oocytes (Fig.2Aa, d). Co-injected Sox9a- miR141-miR734 cocktails increased
180 proportion of spermatocytes (Sc), and also induced ovary-testis transition (indicated by oocyte degeneration
181 and multinucleate cell generation, Fig.2Ab, 2Ae). Conversely, enforced Sox9b- miR218a expression increased
182 each stage of follicles (oocytogenesis), but reduced testis Cn and Sc with increase of deforming Gc (Fig. 2Ac,
183 2Af). We didn't found consistent roles of other Sox9a/ Sox9b-miRNAs mixtures in testiculogenesis and
184 folliculogenesis (Fig.2C, Fig.S3).

185 Since Sox9a, Sox9b and pre-miRNAs transcripts were differentially transcribed in the testis and ovary
186 (Fig.S1), and naturally reduced with age, we posited that exogenous specific miRNAs-coupled Sox9a or
187 Sox9b may induce certain "rejuvenation" effects on the old gonads. To explore this possibility, we tested
188 1.5-2 years old of zebrafish. As expected, Sox9a-miR430a mixture significantly increased the proportion of
189 Gc and Cn whereas Sox9b-miR218a mixture increased all stages of follicles. Sox9b+miR430a increased the
190 testis Cn and Sc (Fig.2Bc, 2Cc) as well as the primordial follicles (Fig.2Be). Compared to the injection into

191 the young fish (5 months olds) (Fig.2A), Sox9b-miR218a mixture (Fig.2A-2B) more significantly increased
192 the ratio of the primordial follicles in the old fish.

Fig.2. Effects of exogenous Sox9a/Sox9b coupled specific miRNAs on testiculogenesis and folliculogenesis. (A) Histological observations on testiculogenesis and folliculogenesis in mature zebrafish (5 mpf). Based on cell morphology, the germline cells were classified into three categories: gonocytes (Gc), mitotic condensed chromatin nuclear cell (Cn), and mature sex gamete (Gm). Gc includes spermatogonia, oogonia or other undifferentiated PGCs. (Cn) includes spermatocytes, or primary oocytes (pOc), Gamete includes spermatids or spermocyte (Sc), secondary oocyte within five stages of follicles (I-V). Degenerating oocytes (dO) were indicated by arrow. Some enlarged gonocytes (indicated by arrow head) looked like early perinucleolar oocytes (or regarded as deforming gonocytes). (B) Histological observations on testiculogenesis and folliculogenesis in old fish (1.5 years old). a-c show testiculogenesis; d-f show various stages of folliculogenesis: stage Ia, Ib, II, III, and IV oocytes. (C) Statistical analyses on cell count of germ cells at various stages of spermatogenesis and oogenesis. The proportion of each category was compared among different specific miRNA-Sox9a/b treatments for 5mf fish (a, b) and 1-1.5 years old fish (c). * ($p<0.05$), ** ($p<0.01$). Scale bar =50 μ m (testis) and 500 μ m (ovary).

To further test the rejuvenation effects, we performed the same specific miRNA-Sox9a/b microinjection into

the old gonads of *vasa* promoter-driven GFP transgenic medaka. Germ cell-specific gene, *vasa*, is most abundantly expressed in mitotic germ cells (oogonia and spermatogonia) and reduced in meiotic germ cells (Tanaka, Kinoshita et al., 2001, Yuan, Li et al., 2014). As shown in Fig.S3B, miR430a-Sox9a increased the testis GFP-positive cells; miR430a-Sox9b increased the ovarian primordial follicles; and miR218-Sox9b increased all stages of follicles (mostly of primordial follicles). These results reaffirmed that miR430a-Sox9a coordination is sufficient to induce spermatogenesis while miR218a-Sox9b could promote renewal of ovarian follicles.

MiR430a and miR218a conversely regulate chemical modifications of Sox9a/Sox9b proteins

Since *grk4* and *grk5l* encode the members of G protein-coupled receptor kinase subfamily, our proposed specific miRNA-GPCR networks may regulate Sox9a and Sox9b protein modifications. Presumably covalent modifications in Sox9a and Sox9b proteins would change the molecular-weight sizes and/or band patterns. Immunoblotting analyses showed that specific Sox9a and Sox9b antibodies recognized several bands, which molecular-weight sizes were bigger than the predicted; and the band patterns were changed after cotransfections of miR430a or miR218a (Fig.3A). In the transfected Pac2 cells, the presumed protein modifications (Fig.3.B) were not significant as those occurred in the microinjected gonadal tissues.

Fig.3. Immunoblotting analyses of endogenous and exogenous Sox9a/Sox9b protein modifications. (A) The testes were microinjected with pSox9a-myc (Sox9a) or pSox9a-myc and miR430a mix (9a+miR430a); the ovaries were injected with pSox9b-myc (Sox9b) or pSox9b-myc and miR218a mix (9b+miR218a). (B). In the *in vitro* cultured Pac2 cells were transfected with pSox9a-myc, pSox9a-myc/miR430a mix, pSox9b-myc, pSox9b-myc/miR218a mix, or empty-control. (C). Sex gonads were microinjected as described in (A). The expression of each signaling component was examined. Relative signal intensity was calculated compared to total Akt.

Co-immunoprecipitation (Co-IP) coupled-mass spectrometer analysis indicated several reverse changes in covalent modifications (Table 3). First, enforced Sox9a-injected gonadal tissues showed enriched Sox9a peptide phosphorylation, acetylation and ubiquitination. On co-injection with miR430a, Sox9a phosphorylations and ubiquitination decreased, but oxidation increased. Meanwhile phosphorylation, ubiquitination and carbamidomethyl at Sox9b and Sox8b peptides were increased by enforced miR430a-Sox9a mixture. Second, enforced Sox9b induced extensive Sox9b acetylation while addition of miR218a retained Sox9b acetylation, but increased oxidation and phosphorylation at Sox9b, Sox9a and Sox8a peptides. Third, both Sox9a and Sox9b peptides were highly acetylated, and mostly accompanied with other modifications. Statistical analysis indicated that the putative enzymes responsible for significantly enriched phosphorylations, and acetylation, included phosphatidylinositol 3-kinase, rho-associated protein kinase, mitogen-activated protein kinase-activated protein kinase 2a, protein kinase C, novel protein similar to vertebrate carnitine acetyltransferase (CRAT), n-alpha-acetyltransferase 35, and acetyl-CoA acetyltransferase. We then verified whether the identified kinases linked to our proposed miR430a/miR218a-GPCR-Sox9 networks. Enforced Sox9b increased Rock1, p-PI3K and p-PKC in the ovary while co-injected miR218a suppressed Rock1 and p-PKC, partially suppressed p-PI3K, and increased p-Akt, suggesting that miR218 could modulate Sox9b activations (Fig.3C). In the testis, Sox9a did not activate Rock1, p-PI3K and p-PKC; However, Sox9a and miR430a mixture activated p-PKC and Rock1 signaling.

Table 3 CoIP-Mass spectrometer analyses on chemical modifications in Sox9a and Sox9b peptides

Enforced	Sequence	Matched	Modifications *	Putative enzymes
----------	----------	---------	-----------------	------------------

expression#		protein		
Sox9a	NGQSESEDGSEQTHI SPNAIFK	Sox9a	3xPhospho [S4; S6; S10]; 1xAcetyl [K22]	Phosphatidylinositol 3-kinase, Rho-associated protein kinase, Serine/threonine-protein kinase SBK1 , Tyrosine-protein kinase BAZ1B (Fragment)
	TLGKLWRLNEVEK	Sox9a	1xAcetyl [T/K]; 1xPhospho [T1]; 1xGG [K14]	
Sox9a+ miR430a	NKPHVKRPMNAFM VWAQAARR	Sox9a	Acetyl [N]; 1xOxidation [M9]	Novel protein similar to vertebrate carnitine acetyltransferase (CRAT) ; N-alpha-acetyltransferase 35, Mitogen-activated protein kinase-activated protein kinase 2a.
	ADLKR	Sox9a, Sox9b,	1xAcetyl [N-Term]	
	TDLPVCSKADLKR	Sox9b	1xPhospho [T1]; 1xAcetyl [T/K]	
	VSGSGKSKPHVK	Sox9b, Sox8b	1xPhospho [S7]; 1xAcetyl [N-Term]; 2xGG [K6; K8]	
Sox9b	VSGSGKSKPHVK	Sox9b	1xAcetyl [K6]	Acetyl-CoA acetyltransferase
	LRVQHKK	Sox9b	2xAcetyl[K6; K7]	
	DAVSQVLK	Sox9b		
Sox9b+ miR218a	RPMNAFMVWAQAA RRK	Sox9a, Sox9b, Sox10, Sox8a	2xOxidation [M3; M7]	Acetyl-CoA acetyltransferase, Protein kinase C.
	DAVSQVLK	Sox8a, Sox9b		
	VSGSGKSKPHVKRP MNAFMVWAQAAR	Sox9b, Sox8, Sox9a	Acetyl [K]; 1xPhospho [S7]; 2xOxidation [M15; M19]	
	ADLKRER	Sox9b	1xAcetyl [K4]	
	VNGSSKNKPHVKRP MNAFMVWAQAAR	Sox9a	Acetyl [K]; 1xOxidation [M19]	

254
255 #, Sex gonads from 20 Fish (2 months post fertilization, uncertain male or female) were divided four groups
256 and separately injected by 1) Sox9a-Myc plasmid; 2) Sox9a-Myc plasmid+miR430a+miR141; 3) Sox9b-Myc
257 plasmid; and 4) Sox9b-Myc plasmid+miR218a. Two more injections were repeated in the first week. Two
258 weeks later, the gonads were collected for co-immunoprecipitation with Sox9a-Myc antibody mix (group 1

and group2) or Sox9b-Myc antibody mix (group3 and 4).
 *,3xPhospho [S4; S6; S10], three phosphorylayions at 4th, 6th, 10th serine risidues of the peptide; 1xAcetyl [K22], one acetylation at 22nd lysine residue of the peptide; 1xGG [K14], one ubiquitination at 14th lysine residue of glyglylys peptide; 1xOxidation [M9], one oxidation at 9th methaline residue of the peptide.

Although specific antibodies are required to verify the exact modifications in Sox9a and Sox9b proteins, together with the results shown in Fig.3, Fig.S2, Table2 and Table 3, we can make following conclusions: 1) Sox9b up-regulated GPCR genes (*lgr4*, *grk5l*, *grk4*), and extensively activate PKC/PI3K/Rock1 signaling pathways and Sox9a/b modifications while miR218a could restrain such activations possibly by phosphorylation of Sox9b through PI3K-pAkt pathway. 2) In the testis, miR430a and Sox9a reciprocally regulate GPCR genes (*lgr4*, *grk5l*, *grk4*) and activate p-PKC-MAPK/Rock1 pathways to repress Sox9b activities.

Discussion

Inducing cell phenotypic conversions usually require multi-purpose signaling inputs and major remodeling of gene transcription (Zhang, Stokes et al., 2011). Specific miRNAs and targets have been identified to account for cell identity changes of skin stem cells (Zhang et al., 2011), mesenchymal progenitors (Huang, Zhao et al., 2010), and cardiomyocytes (Xin, Olson et al., 2013). Functionally phenotypic conversion of sex gonadal cells appear more intriguing, because it involves coordination of multiple somatic and germline lineages in order to successfully commit testiculogenesis or folliculogenesis. As scientists are trying to translate the information gained from basic scientific studies on ovarian organogenesis to the development of *in vitro* strategies to derive stem-cell based ‘prosthetic ovary’ (Truman, Tilly et al., 2017), an attractive possible alternative step in this process is discovery of new ways to drive oogenesis *in vivo*. In the present study, we find that specific miRNAs- GPCR networks could modulate and renew spermatogenesis and folliculogenesis.

First, we provide molecular basis for miRNAs-posttranscriptional regulation of Sox9 switch in the gonadal sex differentiation. In mammals, sex differentiation is controlled by two groups of sex-determining genes that promote one gonadal sex and antagonize the opposite one (Rodriguez-Mari et al., 2005). Zebrafish has no sex chromosome but has two isoforms of *sox9* gene: *sox9a* and *sox9b* (Chiang et al., 2001). Our *in vitro* cell

transfections and in vivo gonadal microinjections indicated that miR430a-Sox9a network could induce spermatogenesis while miR218a-Sox9b network promote folliculogenesis. We established miR430a/miR218a-GPCR-Sox9a/Sox9b switch regulatory model. In this model, miR430a and miR218a convergently target *lgr4*, *grk5l* and *grk4* GPCR genes, and conditionally regulate PI3K/ PKC/Rock signaling pathways and Sox9a/Sox9b activities in two sex gonads. MicroRNA-340 inhibits invasion and metastasis of cancer cells by downregulating ROCK1 in (Maskey, Li et al., 2017), however miR430a and Sox9a mixture coordinately activate Rock1 and PKC pathways to induce spermatogenesis. In the ovary Sox9b and miR218a together could restrain PI3K/ PKC/Rock signaling to increase primordial follicle reserve. This repression is important because over-activation of PI3K/ mTORC1 signaling pathways could accelerate depletion of primordial follicles, and lead to premature activation of the entire pool of primordial follicles (Adhikari, Zheng et al., 2010, Reddy, Liu et al., 2008).

GPCRs (G protein-coupled receptors) were found to play crucial roles in maintenance of planarian germ cell plasticity (Saber, Jamal et al., 2016). We identified three putative GPCR molecules and several modifier enzymes responsible for Sox9a and Sox9b posttranscriptional modifications (Table 3). Given that *Lgr4*, *Grk5l* and *Grk4* could respectively link to gonadotropins(Hsu, Liang et al., 1998), mTORC1 activity (Burkhalter, Fralish et al., 2013), as well as SUMO modification (Yu, Yin et al., 2017), our finding not only supplements the previously described composition and function of G protein-coupled receptor signalsomes (Bar Oz, Kumar et al., 2016, Luttrell, 2005, Spiegelberg, 2013), but also clarifies the signal transduction of gonadotropins /GPCR/MAPK/ Sox9a/b switch pathways.

Second, our analysis on the global mRNA-miRNA expression profiles that concur in the same sex gonads represents the first effort to systematically identify the gonadotropins-GPCR-Sox9 regulatory networks in testiculogenesis and folliculogenesis. We extend the previous finding that miRNAs could target multiple genes and form a complex miRNA regulatory network (Tao et al., 2016), and summarize four interaction modes between specific miRNAs and the target genes, *i.e.*, downregulatory target-pair, upregulatory target-pair, conditionally regulatory target-pair, and indirectly regulatory target-pair.

Previous studies have revealed dual regulatory module of miRNAs from repression to activation at transcriptional and translational levels (Vasudevan, Tong et al., 2007). The miR-430 allows the post-transcriptional activation of *nanos1* in germline cells, but induces deadenylation and translational

repression in somatic cells (Mishima, Giraldez et al., 2006). Now we present the other mechanism that miR430a could modulate expression and intercellular localization of Sox9a and Sox9b proteins through chemical modifications, such as phosphorylation, acetylation, oxidation and ubiquitination (Table 3). It has been found that these types of posttranscriptional modifications could modulate the stability, intracellular localization, and the overall activity of Sox9 (Bar Oz et al., 2016, Jo et al., 2014, Xu, Paige et al., 2010). Finally, we initiated gonadal injections of the specific miRNA-target cocktail to rejuvenate gonadal reserve. Replacement therapy with hormones, and stem cells, blood from aged male, and blood mononuclear cells have been tested to promote ovarian follicle dynamics, spermatogenesis, and tissue rejuvenation (Bukovsky, 2015, Moss, Crosnoe et al., 2013, Niikura, Niikura et al., 2010, Tilly & Telfer, 2009, Zouboulis & Makrantonaki, 2012). However our knowledge of how these factors and cells repair aged reproductive functions still remains limited. Although adult mouse ovaries retain proliferative germ cells resembling male spermatogonial stem cells *in vitro* (Tilly & Telfer, 2009), it is desirable to induce oogenesis under physiological conditions. In the present study, we found that gonadal injection of miR430a-Sox9a mixture could increase testicular gonocytes, and miR218a-Sox9b mixture could increase primordial follicles in old ovaries. Although it is unknown about the source origins from proliferation of reserved stem cell or dedifferentiation of adult gonadal cells, miR430a-Sox9a and miR430a-Sox9b at least could expand the testis spermatogonia reserve and ovarian follicle reserve respectively. If equivalent efficacy can be found in human gonads, miRNA-based testis rejuvenation and prevention of premature ovarian failure may one day be realized. The miRNA-target pair data reported here will help identify more potential therapeutic targets and promote to develop new therapeutic interventions for aging-related ovarian failure and age-related testicular regression.

Materials and methods

Fish husbandry

Zebrafish and *vasa*-GFP transgenic medaka were bred and maintained as previously described (Li, Guan et al., 2013, Wang, He et al., 2012, Yuan et al., 2014) . All experiments with zebrafish and medaka were approved by the laboratory animal care and use committee of Shanghai Ocean University, and performed in accordance with “Guide for the Care and Use of Laboratory Animals” (NIH).

Gonadal microinjection

343 A simple and feasible method for body surface injection into sex gonad has been described (personal data).
 344 Briefly, we dissected ten adult zebrafish (five each sex), measured the size ratio of gonad to the body, and
 345 figured out the integrated surface projection of the sex gonads, and determined the injection area. Based on the
 346 relative size and location of the sex gonads, injection site was about 1 mm upward from the intersect between
 347 pectoral fin upper horizontal and the front end vertical line of pelvic fin. By this way overexpression
 348 plasmids (100ng), and/or specific miRNAs (20pmol each oligos) (Table S10) were microinjected into sex
 349 gonads with lipohigh transfection reagent (Sangon Biotech, Shanghai). Microinjection efficiency was
 350 confirmed by western blotting analyses of enforced Sox9a-Myc and Sox9b-Myc (Fig.S6).

351 **Data Availability Statement:** All relevant data are within the paper and its supporting information files. All
 352 RNA sequencing data are available at the NCBI short read archive: under study number PRJNA293388
 353 (SRP062685) for mRNA transcriptome; under study number PRJNA294493 (SRP063086) for small RNA.

354

355 **Acknowledgements**

356 We thank Ying Peng for her technical assistance in immunoblotting analyses. This work was supported in part
 357 by Shanghai Universities First-class Disciplines Project of Fisheries to JY, and National Natural Science
 358 Foundation of China (31672700) to ML. The funders had no role in study design, data collection and analysis,
 359 decision to publish, or preparation of the manuscript.

360

361 **Author contributions**

362 Conceived and designed the experiments: JY. Performed the experiments: XD, HG, YZ, JW. Analyzed the data: JY, XH,
 363 XD, HG, YZ, JW. Wrote the paper: JY, XH, ML.

364

365 **Competing Interests:** The authors have declared that no competing interests exist.

366

367 **References**

368 Adhikari D, Zheng W, Shen Y, Gorre N, Hamalainen T, Cooney AJ, Huhtaniemi I, Lan ZJ, Liu K (2010) Tsc/mTORC1 signaling
 369 in oocytes governs the quiescence and activation of primordial follicles. *Hum Mol Genet* 19: 397-410
 370 Bar Oz M, Kumar A, Elayyan J, Reich E, Binyamin M, Kandel L, Liebergall M, Steinmeyer J, Lefebvre V, Dvir-Ginzberg M
 371 (2016) Acetylation reduces SOX9 nuclear entry and ACAN gene transactivation in human chondrocytes. *Aging Cell* 15:

499-508

Bukovsky A (2015) Novel methods of treating ovarian infertility in older and POF women, testicular infertility, and other human functional diseases. *Reprod Biol Endocrinol* 13: 10

Burkhalter MD, Fralish GB, Premont RT, Caron MG, Philipp M (2013) Grk5l controls heart development by limiting mTOR signaling during symmetry breaking. *Cell Rep* 4: 625-32

Chen SR, Zheng QS, Zhang Y, Gao F, Liu YX (2013) Disruption of genital ridge development causes aberrant primordial germ cell proliferation but does not affect their directional migration. *BMC Biol* 11: 22

Chiang EF, Pai CI, Wyatt M, Yan YL, Postlethwait J, Chung B (2001) Two sox9 genes on duplicated zebrafish chromosomes: expression of similar transcription activators in distinct sites. *Dev Biol* 231: 149-63

Galliot B, Ghila L (2010) Cell plasticity in homeostasis and regeneration. *Mol Reprod Dev* 77: 837-55

Gao Y, Guo W, Hu Q, Zou M, Tang R, Chi W, Li D (2014) Characterization and differential expression patterns of conserved microRNAs and mRNAs in three genders of the rice field eel (*Monopterus albus*). *Sex Dev* 8: 387-98

Gassei K, Schlatt S (2007) Testicular morphogenesis: comparison of in vivo and in vitro models to study male gonadal development. *Ann N Y Acad Sci* 1120: 152-67

Hsu SY, Liang SG, Hsueh AJ (1998) Characterization of two LGR genes homologous to gonadotropin and thyrotropin receptors with extracellular leucine-rich repeats and a G protein-coupled, seven-transmembrane region. *Mol Endocrinol* 12: 1830-45

Huang J, Zhao L, Xing L, Chen D (2010) MicroRNA-204 regulates Runx2 protein expression and mesenchymal progenitor cell differentiation. *Stem Cells* 28: 357-64

Jo A, Denduluri S, Zhang B, Wang Z, Yin L, Yan Z, Kang R, Shi LL, Mok J, Lee MJ, Haydon RC (2014) The versatile functions of Sox9 in development, stem cells, and human diseases. *Genes Dis* 1: 149-161

Kang L, Cui X, Zhang Y, Yang C, Jiang Y (2013) Identification of miRNAs associated with sexual maturity in chicken ovary by Illumina small RNA deep sequencing. *BMC Genomics* 14: 352

Kobayashi A, Chang H, Chaboissier MC, Schedl A, Behringer RR (2005) Sox9 in testis determination. *Ann N Y Acad Sci* 1061: 9-17

Li M, Guan G, Hong N, Hong Y (2013) Multiple regulatory regions control the transcription of medaka germ gene vasa. *Biochimie* 95: 850-7

Luttrell LM (2005) Composition and function of g protein-coupled receptor signalsomes controlling mitogen-activated protein kinase activity. *J Mol Neurosci* 26: 253-64

Maskey N, Li D, Xu H, Song H, Wu C, Hua K, Song J, Fang L (2017) MicroRNA-340 inhibits invasion and metastasis by downregulating ROCK1 in breast cancer cells. *Oncol Lett* 14: 2261-2267

McClelland K, Bowles J, Koopman P (2012) Male sex determination: insights into molecular mechanisms. *Asian J Androl* 14: 164-71

Mishima Y, Giraldez AJ, Takeda Y, Fujiwara T, Sakamoto H, Schier AF, Inoue K (2006) Differential regulation of germline mRNAs in soma and germ cells by zebrafish miR-430. *Curr Biol* 16: 2135-42

Moss JL, Crosnoe LE, Kim ED (2013) Effect of rejuvenation hormones on spermatogenesis. *Fertil Steril* 99: 1814-20

Murakami S, Kan M, McKeehan WL, de Crombrughe B (2000) Up-regulation of the chondrogenic Sox9 gene by fibroblast growth factors is mediated by the mitogen-activated protein kinase pathway. *Proc Natl Acad Sci U S A* 97: 1113-8

Niikura Y, Niikura T, Wang N, Satirapod C, Tilly JL (2010) Systemic signals in aged males exert potent rejuvenating effects on the ovarian follicle reserve in mammalian females. *Aging (Albany NY)* 2: 999-1003

Real FM, Sekido R, Lupianez DG, Lovell-Badge R, Jimenez R, Burgos M (2013) A microRNA (mmu-miR-124) prevents Sox9 expression in developing mouse ovarian cells. *Biol Reprod* 89: 78

Reddy P, Liu L, Adhikari D, Jagarlamudi K, Rajareddy S, Shen Y, Du C, Tang W, Hamalainen T, Peng SL, Lan ZJ, Cooney AJ,

416 Huhtaniemi I, Liu K (2008) Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool.
417 *Science* 319: 611-3

418 Rodriguez-Mari A, Yan YL, Bremiller RA, Wilson C, Canestro C, Postlethwait JH (2005) Characterization and expression
419 pattern of zebrafish Anti-Mullerian hormone (Amh) relative to sox9a, sox9b, and cyp19a1a, during gonad development.
420 *Gene Expr Patterns* 5: 655-67

421 Rosa A, Spagnoli FM, Brivanlou AH (2009) The miR-430/427/302 family controls mesendodermal fate specification via
422 species-specific target selection. *Dev Cell* 16: 517-27

423 Saberi A, Jamal A, Beets I, Schoofs L, Newmark PA (2016) GPCRs Direct Germline Development and Somatic Gonad
424 Function in Planarians. *PLoS Biol* 14: e1002457

425 Selbach M, Schwanhauser B, Thierfelder N, Fang Z, Khanin R, Rajewsky N (2008) Widespread changes in protein
426 synthesis induced by microRNAs. *Nature* 455: 58-63

427 Spiegelberg BD (2013) G protein coupled-receptor signaling and reversible lysine acetylation. *J Recept Signal Transduct*
428 *Res* 33: 261-6

429 Sun D, Zhang Y, Wang C, Hua X, Zhang XA, Yan J (2013) Sox9-related signaling controls zebrafish juvenile ovary-testis
430 transformation. *Cell Death Dis* 4: e930

431 Taieb J, Grynberg M, Pierre A, Arouche N, Massart P, Belville C, Hesters L, Frydman R, Catteau-Jonard S, Fanchin R, Picard
432 JY, Josso N, Rey RA, di Clemente N (2011) FSH and its second messenger cAMP stimulate the transcription of human
433 anti-Mullerian hormone in cultured granulosa cells. *Mol Endocrinol* 25: 645-55

434 Tanaka M, Kinoshita M, Kobayashi D, Nagahama Y (2001) Establishment of medaka (*Oryzias latipes*) transgenic lines with
435 the expression of green fluorescent protein fluorescence exclusively in germ cells: a useful model to monitor germ cells
436 in a live vertebrate. *Proc Natl Acad Sci U S A* 98: 2544-9

437 Tao W, Sun L, Shi H, Cheng Y, Jiang D, Fu B, Conte MA, Gammerdinger WJ, Kocher TD, Wang D (2016) Integrated analysis
438 of miRNA and mRNA expression profiles in tilapia gonads at an early stage of sex differentiation. *BMC Genomics* 17: 328

439 Tilly JL, Telfer EE (2009) Purification of germline stem cells from adult mammalian ovaries: a step closer towards control
440 of the female biological clock? *Mol Hum Reprod* 15: 393-8

441 Truman AM, Tilly JL, Woods DC (2017) Ovarian regeneration: The potential for stem cell contribution in the postnatal
442 ovary to sustained endocrine function. *Mol Cell Endocrinol* 445: 74-84

443 Vasudevan S, Tong Y, Steitz JA (2007) Switching from repression to activation: microRNAs can up-regulate translation.
444 *Science* 318: 1931-4

445 Wang X, He H, Tang W, Zhang XA, Hua X, Yan J (2012) Two origins of blastemal progenitors define blastemal regeneration
446 of zebrafish lower jaw. *PLoS One* 7: e45380

447 Wilhelm D, Palmer S, Koopman P (2007) Sex determination and gonadal development in mammals. *Physiol Rev* 87: 1-28

448 Xiao J, Zhong H, Zhou Y, Yu F, Gao Y, Luo Y, Tang Z, Guo Z, Guo E, Gan X, Zhang M, Zhang Y (2014) Identification and
449 characterization of microRNAs in ovary and testis of Nile tilapia (*Oreochromis niloticus*) by using solexa sequencing
450 technology. *PLoS One* 9: e86821

451 Xin M, Olson EN, Bassel-Duby R (2013) Mending broken hearts: cardiac development as a basis for adult heart
452 regeneration and repair. *Nat Rev Mol Cell Biol* 14: 529-41

453 Xu G, Paige JS, Jaffrey SR (2010) Global analysis of lysine ubiquitination by ubiquitin remnant immunoaffinity profiling.
454 *Nat Biotechnol* 28: 868-73

455 Yan YL, Willoughby J, Liu D, Crump JG, Wilson C, Miller CT, Singer A, Kimmel C, Westerfield M, Postlethwait JH (2005) A
456 pair of Sox: distinct and overlapping functions of zebrafish sox9 co-orthologs in craniofacial and pectoral fin
457 development. *Development* 132: 1069-83

458 Yu CQ, Yin LQ, Tu ZT, Liu DW, Luo WP (2017) The regulatory role of dopamine receptor D1 on PP2A via SUMO-1
459 modification. *Eur Rev Med Pharmacol Sci* 21: 3270-3276

460 Yuan Y, Li M, Hong Y (2014) Light and electron microscopic analyses of Vasa expression in adult germ cells of the fish
 461 medaka. *Gene* 545: 15-22
 462 Zhang L, Stokes N, Polak L, Fuchs E (2011) Specific microRNAs are preferentially expressed by skin stem cells to balance
 463 self-renewal and early lineage commitment. *Cell Stem Cell* 8: 294-308
 464 Zouboulis CC, Makrantonaki E (2012) Hormonal therapy of intrinsic aging. *Rejuvenation Res* 15: 302-12
 465
 466