1 Identification of a Novel Neutralizing and Two Non-Neutralizing Epitopes on Epstein-Barr Virus gp350 Protein

2

3 Lorraine Z Mutsvunguma^a, Anne Barasa^{a,b}, Charles Warden^c, Joslyn Foley^a, Murali Muniraju^a, David H
4 Mulama^{a,d}, Peng Ye^a, Hanjun Qin^c, Jinhui Wang^c, Xiwei Wu^c, Waithaka Mwangi^e, and Javier Gordon
5 Ogembo^{a,#}

6

7 ^aDepartment of Immuno-Oncology, Beckman Research Institute of City of Hope, Duarte, California, USA

8 ^bDepartment of Human Pathology, University of Nairobi, Nairobi, Kenya

9 ^cIntegrative Genomics Core, Beckman Research Institute of City of Hope, Duarte, California, USA

10 ^dDepartment of Biological Sciences, Masinde Muliro University of Science and Technology, Kakamega, Kenya

11 ^eDepartment of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University,
12 Manhattan, Kansas, USA

13

14 Running Head: Novel neutralizing and non-neutralizing gp350 epitopes

15 #Address correspondence to Javier Gordon Ogembo, jogembo@coh.org

16

17 Abstract word count: 250

18 Text word count:

Abstract

Prevention of Epstein-Barr virus (EBV) primary infection has focused on generating neutralizing antibodies (nAbs) targeting the major envelope glycoprotein gp350/220 (gp350). To date, eight gp350 epitopes have been identified, but only one has elicited nAbs. In this study, we generated 23 hybridomas that produced anti-gp350 antibodies. We compared the candidate anti-gp350 antibodies to nAb-72A1 by: (1) testing their ability to detect gp350 using ELISA, flow cytometry, and immunoblot; (2) sequencing their heavy and light chain complementarity-determining regions (CDRs); (3) measuring the ability of each monoclonal antibody (mAb) to neutralize EBV infection *in vitro*; and (4) mapping the gp350 amino acids bound by the mAbs using RepliTope peptide microarrays. Eight antibodies recognized both denatured and non-denatured gp350, whereas five failed to react with denatured gp350 but recognized native gp350, suggesting they recognized conformational epitope(s). Sequence analysis of the heavy and light chain variable regions of the hybridomas identified 15 mAbs with novel CDR regions unique from those of nAb-72A1. Seven of the new mAbs neutralized EBV *in vitro*, with HB20 and HB17 reducing EBV infection by 40% and >60%, and >30% and 80%, at 10 µg/ml and 50 µg/ml, respectively. Epitope mapping identified nine epitopes and defined their core residues, including two unique immunodominant epitopes, ²⁵³TPIPGTGYAYSLRLTPRPVSRFL₂₇₅ and ⁸⁷⁵LLLLVMADCAFRRLSTSHTYTTPPY₈₉₉, and a novel nAb epitope ³⁸¹GAFASNRFTDIT₃₉₂. This study provides comprehensive *in vitro* mapping of the exact residues defining nine epitopes of EBV gp350. Our findings will inform novel strategies to design optimal EBV vaccines capable of conferring broader protection against the virus.

Importance

Neutralizing antibodies (nAbs) directed against Epstein-Barr virus envelope glycoprotein gp350/220 (gp350) are generated in humans upon infection or immunization, and are thought to prevent neonatal infection. However, clinical use of exogenous nAbs (passive immunization) is limited to a single study using the only well-characterized nAb, 72A1. The gp350 ectodomain contains at least eight unique B-cell binding epitopes; two of these epitopes are recognized by nAb-72A1. The exact amino acid residues of the other six epitopes and their role in generating nAbs has not been elucidated. We used our 15 newly generated and fully characterized monoclonal antibodies and a peptide-overlapping RepliTope array to provide a comprehensive map of the core amino acid residues that define epitopes of gp350 and to understand their role in generating nAbs. These results will inform design of better-targeted gp350 peptide vaccines that contain only protective epitopes, which will focus the B-cell response to produce predominantly nAbs.

Keywords: EBV, glycoprotein, gp350, antibodies, epitopes, amino acid residues, vaccines

Introduction

Epstein-Barr virus (EBV) infection is the causal agent of acute infectious mononucleosis (9, 13). Persistent EBV infection in immunodeficient individuals is associated with numerous epithelial and lymphoid malignancies, such as nasopharyngeal carcinoma, gastric carcinoma, Burkitt lymphoma, Hodgkin lymphoma, and post-transplant lymphoproliferative diseases (PTLD) (24). Pre-existing antibodies provide the primary defense against viral infection. Prophylactic prevention of EBV primary infection has mainly focused on blocking the first step of viral entry by generating neutralizing antibodies (nAbs) that target EBV envelope glycoproteins. Five glycoproteins in particular—gp350/220 (gp350), gp42, gH, gL, and gB—are required for efficient infection of permissible host cells and have emerged as potential prophylactic targets (2, 3, 5, 20).

EBV predominantly infects epithelial cells and B cells, reflecting the viral tropism and the cellular ontogeny for EBV-associated malignancies (4). There are two schools of thought on how the initial EBV transmission into the human host cells occurs. In the first infection model, the incoming virus engages with ephrin receptor A2 via heterodimeric gH/gL, which triggers gB fusion with the epithelial cell membrane and entry of the virus into the cytoplasm (4). This interaction is thought to occur in the oral mucosa, where the virus undergoes lytic replication to release virions that subsequently infect B cells. In the alternative model, the incoming virus binds to the host cell via complement receptor type 1 (CR1)/CD35 (17) and/or CR2/CD21 through its major immunodominant glycoprotein, gp350 (6). The interaction between gp350 and CD35 and/or CD21 triggers viral adsorption, capping, and endocytosis into the B cell (31), which subsequently leads to the heterotrimeric viral glycoproteins complex, gp42/gH/gL, binding to HLA class II molecules to activate gB membrane fusion and entry. Because these two models are not necessarily mutually exclusive, and given that both gp350 and gH/gL complex are important in initiating the first viral contact with host cells, use of nAbs that target either gp350 or gH/gL complex, or both, may potentially block incoming virus at the oral mucosa.

Nearly all EBV-infected individuals develop nAbs directed to the ectodomains of these glycoproteins (25, 36). These antibodies can prevent neonatal infection, can protect against acute infectious mononucleosis in adolescents, and can protect against several human lymphoid and epithelial malignancies associated with EBV infection (7, 14, 23, 28). Although numerous monoclonal antibodies (mAbs) have been generated against EBV gp350 (11, 21, 34), only two murine mAbs, the non-neutralizing 2L10 and the neutralizing 72A1, have been extensively characterized and made commercially available (11, 34). Importantly, nAb-72A1 conferred short-term clinical protection against EBV transmission after transplantation in pediatric patients in a small phase I clinical trial (8).

EBV gp350 is the most immunogenic envelope glycoproteins on the virion. It is a type 1 membrane protein that encodes for 907 amino acid (aa) residues. A single splice of the primary transcript deletes 197 codons and joins gp350 codons 501 and 699, in frame, to generate the gp220 messenger RNA. Both gp350 and gp220 are comprised of the same 18-aa residue at the C terminus that is located within the viral membrane, a 25-aa residue at the transmembrane-spanning domain, and a large highly glycosylated N-terminal ectodomain, aa 1–841 (32). The first 470 aa of gp350 are sufficient for binding CD21 in B cells, as demonstrated by a truncated gp350 (aa 1–470) blocking the binding of EBV to B cells and reducing viral infectivity (8). The gp350-binding domain on CD21 maps to N-terminal short consensus repeats (SCRs) 1 and 2, which also bind to a bioactive fragment of complement protein 3 (C3d) (16, 18). A soluble truncated EBV gp350 fragment (aa 1–470) and soluble CD21 SCR1 and SCR2 can block EBV infection and immortalization of primary B cells (32). However, gp350 binding to CD35 is not restricted to N-terminal SCRs; it binds long homologous repeat regions as well as SCRs 29–30 (22).

The gp350 ectodomain is heavily glycosylated, with both N- and O-linked sugars, which accounts for over half of the molecular mass of the protein. Currently there is only one crystal structure available for gp350, comprised of a truncated structure between 4–443 aa, with at least 14 glycosylated arginines coating the protein

with sugars, with the exception of a single glycan-free patch (30). Mutational studies of several residues in the glycan-free patch resulted in the loss of CD21 binding (30), suggesting that binding of CD35 and CD21 by gp350 is mediated within this region.

There are at least eight unique CD21 binding epitopes located at the N-terminus of the gp350 ectodomain (35); at least one of these epitopes (aa 142–161) is capable of eliciting nAbs (32, 35). The aa residues 142–161 are also the binding site for nAb-72A1 (11, 30). Using gp350 synthetic peptides binding to CD21 on the surface of a B cell line, an additional gp350 epitope was identified in the C-terminal region of gp350 (aa 822-841), suggesting it is involved in EBV invasion of B cells (35). The role of other epitopes in eliciting nAbs has not been fully investigated. Furthermore, the exact aa residues that comprise the core binding sites for epitopes capable of eliciting neutralizing and non-nAbs have not been determined. Mapping the EBV gp350 protein residues that define immunodominant epitopes, identifying the critical aa residues of the known and unknown epitopes, and defining their roles in generating neutralizing and non-nAbs will guide rational design and construction of an efficacious EBV gp350-based vaccine that would focus B-cell responses to the protective epitopes.

In this study, we generated 23 hybridomas producing antibodies against EBV gp350. To assess their clinical potential and utility in informing future prophylactic and therapeutic vaccine design, we: (1) tested the ability of the antibodies produced by the new hybridomas to detect gp350 protein by enzyme-linked immunosorbent assay (ELISA), flow cytometry, and immunoblot; (2) sequenced the unique complementarity-determining regions (CDRs) of the heavy and light chains of all 23 hybridomas to identify novel mAbs; (3) measured the efficacy of each mAb to neutralize EBV infection *in vitro*; and (4) used RepliTope peptide microarrays to identify gp350 core aa residues recognized by neutralizing and non-neutralizing mAbs. Using the newly generated antibodies, we identified a new epitope bound preferentially by nAbs, distinct from the

canonical neutralizing epitope bound by nAb-72A1, as well as two immunodominant epitopes bound by both neutralizing and non-nAbs.

Results

Characterization of new anti-gp350 mAbs. We generated and biochemically characterized new EBV gp350-specific mAbs, and evaluated their ability to neutralize EBV infection. In addition, we used the antibodies to map immunodominant epitopes on the EBV gp350 protein. To generate hybridomas, we immunized BALB/c mice with purified UV-inactivated EBV, boosted them with virus-like particles (VLPs) that incorporate the EBV gp350 ectodomain on the surface to enrich for production of anti-gp350 antibodies, then isolated splenocytes from the immunized mice and fused them with myeloma cells. We used indirect ELISA to screen supernatants from the hybridomas for specificity against purified EBV gp350 ectodomain protein (aa 4–863) and identified 23 hybridomas producing gp350-specific antibodies.

We determined the isotypes of the new antibodies to be IgG1 (n=14), IgG2a (n=5), IgG2b (n=1), a mixture of IgG1 and IgG2b (n=1), and a mixture of IgG1 and IgM (n=2). We found that all 23 hybridomas that produced antibodies (designated HB1–23) recognized the gp350 antigen in an initial ELISA screening using unfractionated and unpurified hybridoma supernatants (data not shown). We used affinity purification with protein A followed by SDS-PAGE to confirm the purity of all 23 antibodies. When we re-evaluated quantified amount of the purified antibodies (10 µg/ml) using indirect ELISA, all of the 23 antibodies had ELISA signals two times greater than those of phosphate buffered saline (negative control), and were considered as positive or specific to gp350. Of these, five (HB4, HB5, HB7, HB13, and HB14) demonstrated binding affinity equal to or greater than that of the positive control, nAb-72A1 (**Fig. 1A**). This difference in binding of the 23 antibodies could be due to differential exposure of cognate epitopes on gp350 in the assay performed.

Determining the nature of the binding between an antibody and its target antigen is an important consideration for the performance and specificity of an antibody, as it can involve the recognition of a linear or conformational epitope (26). We characterized the antibodies using immunoblot analysis of denatured gp350 antigen expressed from Chinese hamster ovary (CHO) cells, and showed that 16 of the antibodies reacted to both the 350 kDa and the 220 kDa splice variant. In contrast, HB2, HB3, HB6, HB7, HB13, HB20, and HB21 failed to recognize either of the denatured isoforms of gp350 (**Fig. 1B**). We further characterized the antibodies using flow cytometric analysis of CHO cells stably expressing gp350 on the cell surface, and revealed that HB1, HB2, HB3, HB5, HB6, HB9, HB11, HB12, HB15, HB17, HB19, HB20, and HB21 antibodies readily recognized gp350 (**Fig. 1C**). Given that HB2, HB3, HB20, and HB21 detected gp350 by flow cytometry, but not by immunoblot, suggests that these four antibodies recognized conformational epitopes (native) on gp350, whereas HB5, HB9, HB11, HB15, HB17, and HB19 recognized both linear and conformational epitopes (**Fig. 1B-C**). The observation that all 23 anti-gp350 antibodies recognized the gp350 antigen either by indirect ELISA, flow cytometry, or immunoblot assay suggests that we successfully produced antibodies that are specific to EBV gp350 protein.

Analysis of the variable heavy and variable light chain sequences. We determined the sequences of the heavy and light chain variable region genes (V_H and V_L , respectively) of the 23 new anti-gp350 antibodies, as well as nAb-72A1, and compared the sequences to published nAb-72A1 sequences (10, 33). The sequence of the CDR of this antibody was recently determined and published, revealing two unique IgG1 heavy chains and two unique light chains, one kappa and one lambda (10, 33). We used PCR to amplify the genes encoding the V_H and V_L chain regions in cDNA generated from the 23 hybridoma cells, as well as from HB168 (nAb-72A1). The PCR products presented distinct bands at approximately 350–400 bp and 450–500 bp for V_H and V_L , respectively (data not shown). We sequenced purified fragments using Illumina MiSeq, followed by *in silico* analysis and identified CDRs for both V_H and V_L (**Fig. 2**). We identified two V_H and V_L sequences of nAb-

72A1 as >94% identical to the previously published sequences (10), suggesting that nAb-72A1 exists as a mixed antibody, instead of the reported mAb (33). Similar to nAb-72A1, HB4, HB13, HB15, and HB23 hybridomas each produced a mixture of two antibodies, with unique sequences of the V_H chain showing at >5% frequencies, suggesting that they are not mAbs (**Table 1**). We were unable to identify coding sequences for V_L chains for HB7, HB9, and HB17, unless the frequencies were lowered to >1% (**Table 1**); in this case, the identified coding V_L chain sequences were identical.

Our analysis and comparison of the V_H and V_L chain gene sequences of the 23 hybridomas compared to HB168 (nAb-72A1) showed unique sequences within the CDR 1–3 regions. Only HB8 and HB18 had identical V_H and V_L chain gene sequences, suggesting that the two are the same clone isolated separately; therefore, HB18 was excluded from subsequent experiments. One of the two HB15 antibodies had identical V_H and V_L gene sequences to that of HB10; however, based on the previous characterization, the presence of the additional antibody in HB15 was sufficient to confer subtle differences in biochemical characterizations for gp350 between the two antibodies. Thus, sequence analysis (**Fig. 2**) demonstrated that we generated 15 unique anti-gp350 mAbs, with distinct biochemical properties and sequence identities from the commercially available nAb-72A1.

Neutralization assay. We evaluated the ability of the 15 mAbs (10 µg/ml or 50 µg/ml) to neutralize purified eGFP-tagged AGS-EBV infection of the Raji B cell line *in vitro* following standardized procedures (25) and determined the percentage of eGFP⁺ cells using flow cytometry as described (17, 18). We used the nAbs 72A1 and anti- gH/gL (E1D1) as positive controls, whereas the non-neutralizing mAb 2L10 was used as a negative control. Because HB4, HB7, HB13, HB15, HB16, HB19, HB21, and HB23 were confirmed to be mixtures based on isotyping or sequence data, we eliminated them from further consideration in the neutralization assay. We considered an antibody to be a neutralizer if it inhibited EBV infection >20% at 10 µg/ml and >60% at 50 µg/ml. Several mAbs inhibited EBV infection in a dose-dependent manner. HB20 and

HB17 were the most effective in neutralizing EBV infection of Raji cells *in vitro*, whereby they reduced infection by 40% and >60%, and >30% and 80%, at 10 µg/ml and 50 µg/ml, respectively (**Fig. 3**). The HB9 and HB10 antibodies prevented EBV infection of Raji cells by ~25% at 10 µg/ml and ~60% at 50 µg/ml. The HB11 antibody neutralized <20% at 10 µg/ml, but showed a dose-dependent increase at 50 µg/ml by neutralizing EBV infection by 60%. By the set neutralization parameters, HB1–3, HB5–8, HB12, HB14, and HB22 did not neutralize EBV infection of Raji cells. In comparison, both nAb-72A1 and nAb-E1D1 neutralized EBV infection by >70% and 40%, respectively, at 10 µg/ml. The nAb-72A1 neutralized 100% of EBV infection at 50 µg/ml, whereas nAb- E1D1 neutralized infection by >60% at 50 µg/ml. As expected, the negative control, mAb-2L10, did not neutralize viral infection at either concentration. From the neutralization assay results, we organized the remaining 15 mAbs into two distinct groups, neutralizers (+) and non-neutralizers (-) as summarized in **Table 2**.

Epitope Mapping. We used the 15 new anti-gp350 mAbs (neutralizers vs. non-neutralizers) to identify most, if not all, of the relevant immunodominant aa residues targeted by both the neutralizing and non-nAbs. We used a RepliTope approach, in which overlapping peptides [15-mer with 11 aa overlap] that cover the complete sequence of gp350 (aa 1–907) were immobilized on microarray slides and probed with the purified anti-gp350 antibodies in an ELISA format. The nAb-72A1 was as positive control, because the cognate epitopes bound by the antibody have previously been reported.

We showed that two epitopes, ²⁵³TIPIGTGYAYSLRLTPRPVSRFL₂₇₅ and ³⁹³VSGLGTAPKTLIITRTATNATTT₄₁₅, were both bound by all 15 mAbs, as well as nAb-72A1 and 2L10, regardless of their neutralizing or non-neutralizing capabilities (**Fig. 4**). This consensus suggests that these epitopes are immunodominant. Several of the 15 mAbs (HB2, HB3, HB8, HB11, HB12, HB14, HB17, and HB22), as well as nAb-72A1, bound to ³⁴¹ANSPNVTVTAFWAWPNNTE₃₅₉. Two epitopes, aa 341–359 and aa 393–415, were found within the previously identified single epitope II, which is encoded by nucleotides

between 3,186 bp and 3,528 bp, corresponding to aa 326–439 of gp350 (37). Two mAbs, HB1 and HB10, bound ₆₀₅**TTTPNATGPTVGETSPQA**₆₂₃, an epitope located within the gp350 (aa 501–699) splice region that is involved in generation of the 220 kDa splice variant. A total of eight mAbs (HB1–3, HB8, HB10–12, and HB22) also bound to the region between ₁**MEAALLVCQYTIQSLIHLTGEDPG**₂₄, which includes a region homologous to C3d, another molecule known to interact with CD21 (15, 33). Two epitopes common between most neutralizing and non-nAbs, ₈₂₁**PPSTSSKLRPRWTFTSPPV**₈₃₉ and ₈₇₅**LLLLVMADCAFRNLSTSHTYTPPY**₈₉₉, were located upstream and downstream, respectively, of the transmembrane domain on the C-terminus of gp350. Epitope aa 821–839 is located within the previously identified epitope I, which is located between aa 733–841 (37). Furthermore, epitope aa 821–839 is potentially involved in EBV infection of B cells (35). However, our study could not identify two epitopes located at aa 282–301 and aa 194–211, which were previously shown to be involved in the binding of nAb-72A1 and CD21, respectively (30, 33, 35). Similar to previously reported data, we showed that nAb-72A1 bound ₁₄₅**EMQNPVYLIPETVPYIKWDN**₁₆₄, one of the neutralizing epitopes on gp350 (30, 33, 35).

Analysis of V_H-V_L sequences from the HB168 (nAb-72A1) hybridoma by our group and others revealed that the hybridoma produces two antibodies: one that is gp350-specific and another that recognizes mineral oil-induced plasmacytoma (MOPC)(10). To further interrogate gp350 for additional neutralizing epitopes, we used the gp350-specific nAb-72A1 V_H-V_L sequence to generate chimeric (mouse/human) recombinant antibodies. Similarly, we used the V_H-V_L sequence for the HB20 antibody, which our neutralization analysis above showed to be one of the most effective nAbs, to generate chimeric antibodies. We generated a negative control chimeric recombinant antibody using V_H-V_L sequences from the gp350-specific but non-neutralizing HB5 antibody described above (**Fig. 3**).

We performed comparative analysis of the epitope binding pattern using the chimeric recombinant antibodies and revealed a novel epitope, ₃₈₁**GAFASNRFTDIT**₃₉₂, which was bound by HB20 and nAb-72A1,

but not by HB5, and is distinct from the ₁₄₅EMQNPVYLIPETVPYIKWDN₁₆₄ epitope, which is bound only by the nAb-72A1 (**Fig. 5**).

Discussion

Despite more than 50 years of EBV vaccine research, few candidates have demonstrated even partial clinical efficacy, and none have been efficacious enough to elicit sterilizing immunity and be licensed (3). Antibodies, whether elicited in the host naturally or via passive immunization, provide an effective first-line of defense against viral infection. Several studies have indicated that the major immunodominant glycoprotein EBV gp350 is an ideal target for EBV nAbs production. Although the ectodomain of EBV gp350 (aa 1–841) contains at least eight unique CD21 binding epitopes, only one of these epitopes (aa 142–161) is capable of eliciting nAbs (30, 32, 35). Although the aa residues that constitute the other epitopes and their role in generating nAbs have not been elucidated, this information would be valuable in the precise design of effective EBV peptide vaccines. To date, nAb-72A1 remains the only EBV antibody with proven clinical prophylactic efficacy; it confers short-term protection by reducing and delaying EBV infection onset in immunized pediatric transplant patients (21).

In this study we generated 23 hybridomas producing gp350-specific antibodies and characterized their ability to bind gp350 protein (**Fig. 1**). Out of the 23 hybridomas, we determined that 15 were monoclonal and novel based on their V_H and V_L CDR sequences, compared to the reported sequence of nAb-72A1 (22) (**Fig. 2**).

Following confirmation that the new mAbs recognized EBV gp350 antigen and contained unique V_H-V_L sequences, our further characterization revealed that mAbs HB9, HB10, HB11, HB17, and HB20 inhibited EBV infection in a dose-dependent manner, with HB17 and HB20 being the best neutralizers (**Fig. 3**). Thus, our study provides new neutralizing and non-nAbs against EBV infection, which could potentially inform future EBV vaccine research.

Various methods, including lectin/ricin immune-affinity assay (34), purified mAbs (1, 22), purified soluble gp350 mutants, synthetic peptides (15, 16, 33), cell binding assays (35), and crystal structure of partial gp350 protein (aa 4–443) (30), have been used to identify the critical gp350 epitopes responsible for its interaction with the CD35 and CD21 cellular receptors (*for a detailed summary review see Table 3*). We used an epitope mapping assay to identify a total of nine epitopes, including two new epitopes, and one bound by all mAbs, including nAb-72A1 and 2L10, regardless of the mAb neutralizing capability to EBV infection. This suggests it is an immunodominant epitope on the gp350 protein. We further identified portions of four of the previously described epitopes, including the only currently recognized neutralizing epitope, and defined their exact aa residues. However, we could not identify two previously reported epitopes located at aa 282–301 and aa 194–211, which have been reported to be involved in the binding of nAb-72A1 or CD21, respectively (30, 33, 35) (**Fig. 4**).

Our comparative epitope mapping analysis results identified a novel neutralizing epitope, ³⁸¹GAFASNRTFDIT₃₉₂, which was bound preferentially by the nAbs HB20 and nAb-72A1, but not the non-neutralizing mAb HB5 (**Fig. 5**). The new neutralizing epitope is distinct from the reported canonical nAb-72A1-binding epitope aa 145–164, on gp350, suggesting that epitope 381–392 is a novel epitope on gp350 that might be capable of eliciting nAbs.

In conclusion, in this study we generated 15 novel EBV gp350-specific mAbs, characterized their binding to gp350, determined their neutralization activity against EBV infection *in vitro*, mapped their cognate epitopes, and defined the exact aa residues they recognize on gp350. We confirmed six of the eight previously described epitopes responsible for generating neutralizing and non-nAbs and defined the exact aa residues they bind. This study also confirmed that the binding epitopes on gp350 that elicit nAbs are between aa 4–443 (30). We identified an additional neutralizing epitope and two new non-neutralizing epitopes, with one located downstream of the gp350 ectodomain (aa 1–841). The newly developed mAbs will be useful research tools for

informing future vaccine development, diagnosis of viral infection, or therapeutic/prophylactic management of post-transplant lymphoproliferative diseases, either individually, in combination with nAb-72A1, or with other mAbs such as neutralizing anti-gH/gL (E1D1).

Materials and Methods

Cells and viruses. EBV-AGS, a human gastric carcinoma cell line infected with a recombinant Akata virus expressing enhanced fluorescent green protein (eGFP) was a kind gift of Dr. Liisa Selin (University of Massachusetts Medical School). Anti-EBV gH/gL (E1D1) hybridoma cell line was a kind gift of Dr. Lindsey Hutt-Fletcher (Louisiana State University Health Sciences Center). Chinese hamster ovary cells (CHO); human embryonic kidney cells expressing SV-40 T antigen (HEK-293T); HEK-293 6E suspension cells; EBV-positive Burkitt lymphoma cells (Raji); myeloma cells (P3X63Ag8.653); and anti-EBV gp350 nAb-72A1 hybridoma cells (HB168) were purchased from American Type Culture Collection (ATCC). EBV-AGS cells were maintained in Ham's F-12 media supplemented with 500 µg/ml neomycin (G418, Gibco). Raji, P3X63Ag8.653, and HB168 hybridoma cells were maintained in RPMI 1640. CHO and HEK-293T cells were maintained in DMEM. HEK-293 6E cells were maintained in FreeStyle F17 Expression Medium supplemented with 0.1% Pluronic F-68. All culture media were supplemented with 10% fetal bovine serum (FBS), 2% penicillin-streptomycin, and 1% l-glutamine, with the exception of Freestyle F17 expression medium.

Antibodies and plasmids. Primary antibodies: EBV anti-gp350 nAb (72A1) and anti-gH/gL (E1D1) were purified from the supernatant of HB168 and E1D1 hybridoma cell lines, respectively using Capture™ Protein A Maxiprep spin columns (Takara). Anti-gp350/220 mAb (2L10) was purchased from Millipore Sigma.

Secondary antibodies: Horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG for immunoblot or ELISA were purchased from Bio-Rad. Alexa Fluor® (AF) 488-conjugated goat anti-mouse IgG (H+L) for flow

cytometry was purchased from Life Sciences Tech. Goat anti-mouse IgG (H+L) secondary antibody and DyLight 650 for epitope mapping were purchased from Thermo Fisher Scientific.

The construction of the pCI-puro vector and pCAGGS-gp350/220-F has been described (17, 20).

Virus production and purification. eGFP-tagged EBV was produced from the EBV-infected AGS cell line as described (29). Briefly, EBV-AGS cells were seeded to 90% confluency in T-75 flasks in Ham's F-12 medium containing G418 antibiotic. After the cells reached confluency, G418 media was replaced with Ham's F-12 medium containing 33 ng/ml 12-O-tetradecanoylphorbol-13-acetate (TPA) and 3 mM sodium butyrate (NaB) to induce lytic replication of the virus. Twenty-four h post-induction, the media was replaced with complete Ham's F-12 media without G418, TPA, or NaB and cells were incubated for 4 days at 37°C. The cell supernatant was collected, centrifuged, and filtered using a 0.8-μm filter to remove cell debris. The filtered supernatant was ultra-centrifuged using a Beckman-Coulter type 19 rotor for 70 min at 10,000 rpm to pellet the virus. EBV-eGFP virus was titrated in both HEK-293T cells and Raji cells, and stocks were stored at -80°C for subsequent experiments.

Generation and purification of gp350 virus-like particles. To generate gp350 VLPs, equal amounts (8 μg/plasmid) of the relevant plasmids (pCAGGS-Newcastle disease virus [NDV]-M, -NP, and gp350 ectodomain fused to fusion protein cytoplasmic and transmembrane domains) were co-transfected into 80% confluent CHO cells seeded in T-175cm² flasks; supernatant from transfected cells was collected and VLPs were purified and composition characterized as previously described (18).

Production of hybridoma cell lines. Seven days prior to immunization, two eight-week-old BALB/c mice were bled for collection of pre-immune serum. The mice were immunized with purified UV-inactivated EBV three times (Day 0, 21, and 35) and boosted every 7 days three times (Day 42, 49, and 56) with VLPs incorporating gp350 on the surface after Day 35. The mice were sacrificed and their splenocytes were isolated, purified, and used to fuse with P3X63Ag8.653 myeloma cells at a ratio of 3:1 in the presence of polyethylene

glycol (PEG, Sigma). Hybridoma cells were seeded in flat-bottom 96-well plates and selected in specialized hybridoma growth media with HAT (Sigma) and 10% FBS.

Indirect ELISA. Hybridoma cell culture supernatant from wells that had colony-forming cells were tested for antibody production by indirect ELISA. Briefly, immunoplates (Costar 3590; Corning Incorporated) were coated with 50 μ l of 0.5 μ g/ml recombinant EBV gp350/220 (Millipore) protein diluted in phosphate buffered saline (PBS, pH 7.4) and incubated overnight at 4°C. After washing three times with PBS containing 0.05% (v/v) Tween 20 (washing buffer), plates were blocked with 100 μ l washing buffer containing 2% (w/v) bovine serum albumin (BSA), incubated for 1 h at room temperature, and washed as above. 100 μ l of hybridoma supernatant was added to each well (in triplicate) and incubated for 2 h at room temperature. PBS and nAb-72A1 were added as negative and positive controls, respectively. The plates were washed as described above, followed by incubation with goat anti-mouse IgG horseradish peroxidase-conjugated secondary antibody (1:2,000 diluted in PBS) at room temperature for 1 h. The plates were washed again and the chromogenic substrate 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS, Life Science Technologies) was added. The reaction was stopped using ABTS peroxidase stop solution containing 5% sodium dodecyl sulfate (SDS) in water. The absorbance was read at an optical density of 405 nm using an ELISA reader (Molecular Devices). Three independent experiments were performed.

Antibody purification, quantification, and isotyping. Hybridoma cells from selected individual positive clones were expanded stepwise from 96-well plates to T-75 flasks. At confluence in T-75 flasks, supernatant from individual clones was collected, clarified by centrifugation (4,000 g, 10 min, 4°C), and filtered through a 0.22- μ m-membrane filter (Millipore). Antibodies were further purified using Capturem™ Protein A Maxiprep (Takara) and stored in PBS (pH 7.4) at 4°C. Antibodies were analyzed by SDS-PAGE to determine purity. Bicinchoninic acid assay (BCA assay; Thermo Fisher Scientific) was conducted to determine the concentration

of purified antibodies. Isotype identification was performed with the Rapid ELISA mouse mAb isotyping kit (Thermo Fisher Scientific).

RNA extraction, cDNA synthesis, and sequencing of the variable region of the mAbs. Total RNA was extracted from 1×10^6 hybridoma cells using the RNeasy Mini Kit (Qiagen). Each hybridoma clone cDNA was synthesized in a total volume of 20 μ l using Tetro Reverse Transcriptase (200 u), RiboSafe RNase Inhibitor, Oligo(dT)18 primer, dNTP mix (10 mM each nucleotide), and 100–200 ng RNA. Reverse transcription was performed at 45°C for 30 min, and terminated at 85°C for 5 min. The cDNA was stored at -20°C. Immunoglobulin (Ig) V_H and V_L were amplified using the mouse Ig-specific primer set purchased from Novagen (12). The V_H and V_L genes were amplified in separate reactions and PCR products were visualized on 1% agarose gel.

The V_H and V_L amplicons were sequenced using an Illumina MiSeq platform: duplicate 50 μ l PCR reactions were performed, each containing 50 ng of purified cDNA, 0.2 mM dNTPs, 1.5 mM MgCl₂, 1.25 U Platinum Taq DNA polymerase, 2.5 μ l of 10x PCR buffer, and 0.5 μ M of each primer designed to amplify the V_H and V_L. The amplicons were purified using AxyPrep Mag PCR Clean-up kit (Thermo Fisher Scientific). The Illumina primer PCR PE1.0 and index primers were used to allow multiplexing of samples. The library was quantified using ViiA™ 7 Real-Time PCR System (Life Technologies) and visualized for size validation on an Agilent 2100 Bioanalyzer (Agilent Technologies) using a high-sensitivity cDNA assay. The sequencing library pool was diluted to 4 nM and run on a MiSeq desktop sequencer (Illumina). The 600-cycle MiSeq Reagent Kit (Illumina) was used to run the 6 pM library with 20% PhiX (Illumina), and FASTQ files were used for data analysis(19).

Chimeric mAb construct generation. To generate chimeric mAbs, the V_H and V_L sequences were cloned into the dual-vector system pFUSE CHIg/pFUSE CLIg (InvivoGen), which express the constant region of the heavy and light chains of human immunoglobulins, respectively (Genewiz). The constructs were transiently

transfected into HEK-293 6E cells. The supernatants were collected at 72 h post-transfection and IgG was purified using protein A/G affinity chromatography.

Immunoblot analysis. CHO cells were cultured and stably co-transfected with pCAGGS-gp350/220 F and pCI-puro vector containing a puromycin resistance gene. Forty-eight h post-transfection, DMEM media containing 10 µg/ml of puromycin was added to enrich for cells expressing gp350 protein. Puromycin-resistant clones were expanded, followed by flow cytometry sorting using nAb-72A1 to a purity >90%. EBV gp350-positive CHO cells were harvested and lysed in radioimmunoprecipitation assay buffer (RIPA) followed by centrifugation at 15,000 g for 15 min on a benchtop centrifuge. The supernatants were collected and heated at 95°C for 10 min in SDS sample buffer containing β-mercaptoethanol, then separated using SDS-PAGE. Proteins were transferred onto a nitrocellulose membrane using an iBlot™ Transfer System (Thermo Fisher Scientific) followed by incubation in blocking buffer (1% BSA; 20 mM Tris-HCl, pH 7.5; 137 mM NaCl; and 0.1% Tween-20 [TBST]) for 1 h. The blots were incubated in TBST containing purified anti-gp350 antibodies (1:50) overnight at 4°C. After three washes with TBST, the blots were incubated with a goat anti-mouse conjugated to horseradish peroxidase (1:2,000) in TBST for 1 h. After three washes, the antibody-protein complexes were detected using the Amersham ECL Prime Western Blotting Detection Reagent (GE Healthcare). All experiments were independently repeated three times.

Flow cytometry. To assess the ability of purified anti-gp350 mAb to detect surface expression of EBV gp350 protein by flow cytometry, CHO cells that stably express EBV gp350 were harvested and stained with purified anti-gp350 (10 µg/ml), followed by AF488 goat anti-mouse IgG secondary antibody. Flow cytometric analysis was performed on a C-6 FC (BD Biosciences) and data was analyzed using FlowJo Cytometry Analysis software (FlowJo, LLC) as described (18). All experiments were independently repeated three times.

EBV neutralization assay. Purified individual anti-gp350 mAbs were incubated with purified AGS-EBV-eGFP (titer calculated to infect at least 20% of HEK293 cells seeded in 100 µl of serum-free DMEM) for

2 h at 37°C. To represent EBV infection of B cells, the pre-incubated anti-gp350 mAbs/AGS-EBV-eGFP were used to infect 5×10^5 Raji cells seeded in a 96-well plate. Anti-gp350 neutralizing 72A1 and non-neutralizing 2L10 mAbs served as positive and negative controls, respectively. Plates were incubated at 37°C and the number of eGFP+ cells was determined using flow cytometry 48 h post-infection. All dilutions were performed in quintuplicate and the assays repeated three times for Raji cells. Antibody EBV neutralization activity was calculated as: % neutralization = $(EBV_{alone} - EBV_{mAb}) / (EBV_{alone}) \times 100$.

Epitope mapping. Anti-gp350 mAbs were incubated with a multi-well EBV GP350/GP340 RepliTope (JPT) peptide microarray displaying 224 peptides (15-mers with 11 aa overlap) in 3x7 subarrays. Briefly, anti-gp350 mAbs were diluted in blocking solution (TBS-T and 2% BSA) to a final concentration of 10 µg/ml and incubated with the microarray slide for 1 h at 30°C with shaking. Slides were washed 5 times with wash buffer (TBS-T), followed by incubation with 1 µg/ml secondary antibody Dylight 650 (Thermo Fisher Scientific) for 1 h at 30°C. After washing 5 times with wash buffer and 2 times with distilled water, microarray slides were dried by centrifugation. Detection was performed using the Agilent DNA microarray scanner.

Acknowledgments: We are grateful to Lindsey Hutt-Fletcher for providing the EBV gH/gL (E1D1) hybridoma cell line, Liisa Selin for providing EBV Akata virus expressing eGFP, and Chao Guo for technical assistance with the RepliTope protocol.

This work was supported by the National Institute of Health (NIH) grant R21CA205106 and a generous philanthropic donation from the V Foundation to J.G.O. Research reported in this publication also included work performed in the City of Hope Integrative Genomics Core, Flow Cytometry Core, and Animal Resource Center supported by the National Cancer Institute of the National Institutes of Health under award number P30CA033572. The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health. We thank Dr. Sarah T. Wilkinson for editing of the manuscript and for her insightful feedback and discussion.

Competing Interests: The Authors declare that they have no competing interests.

References

1. **Alfarano, C., C. E. Andrade, K. Anthony, N. Bahroos, M. Bajec, K. Bantoft, D. Betel, B. Bobechko, K. Boutilier, E. Burgess, K. Buzadzija, R. Caverio, C. D'Abreo, I. Donaldson, D. Dorairajoo, M. J. Dumontier, M. R. Dumontier, V. Earles, R. Farrall, H. Feldman, E. Garderman, Y. Gong, R. Gonzaga, V. Grytsan, E. Gryz, V. Gu, E. Haldorsen, A. Halupa, R. Haw, A. Hrvojic, L. Hurrell, R. Isserlin, F. Jack, F. Juma, A. Khan, T. Kon, S. Konopinsky, V. Le, E. Lee, S. Ling, M. Magidin, J. Moniakis, J. Montojo, S. Moore, B. Muskat, I. Ng, J. P. Paraiso, B. Parker, G. Pintilie, R. Pirone, J. J. Salama, S. Sgro, T. Shan, Y. Shu, J. Siew, D. Skinner, K. Snyder, R. Stasiuk, D. Strumpf, B. Tuekam, S. Tao, Z. Wang, M. White, R. Willis, C. Wolting, S. Wong, A. Wrong, C. Xin, R. Yao, B. Yates, S. Zhang, K. Zheng, T. Pawson, B. F. Ouellette, and C. W. Hogue.** 2005. The Biomolecular Interaction Network Database and related tools 2005 update. *Nucleic acids research* **33**:D418-424.
2. **Balfour, H. H., Jr.** 2014. Progress, prospects, and problems in Epstein-Barr virus vaccine development. *Curr Opin Virol* **6C**:1-5.
3. **Cohen, J. I.** 2015. Epstein-barr virus vaccines. *Clin Transl Immunology* **4**:e32.
4. **Connolly, S. A., J. O. Jackson, T. S. Jardetzky, and R. Longnecker.** 2011. Fusing structure and function: a structural view of the herpesvirus entry machinery: A structural view of herpesvirus entry machinery. *Nat Rev Microbiol* **9**:369-381.
5. **Cui, X., Z. Cao, Q. Chen, S. Arjunaraja, A. L. Snow, and C. M. Snapper.** 2016. Rabbits immunized with Epstein-Barr virus gH/gL or gB recombinant proteins elicit higher serum virus neutralizing activity than gp350. *Vaccine*.
6. **Fingerroth, J. D., J. J. Weis, T. F. Tedder, J. L. Strominger, P. A. Biro, and D. T. Fearon.** 1984. Epstein-Barr virus receptor of human B lymphocytes is the C3d receptor CR2. *Proc Natl Acad Sci U S A* **81**:4510-4514.
7. **Gu, S. Y., T. M. Huang, L. Ruan, Y. H. Miao, H. Lu, C. M. Chu, M. Motz, and H. Wolf.** 1995. First EBV vaccine trial in humans using recombinant vaccinia virus expressing the major membrane antigen. *Dev Biol Stand* **84**:171-177.
8. **Haque, T., I. Johannessen, D. Dombagoda, C. Sengupta, D. M. Burns, P. Bird, G. Hale, G. Mieli-Vergani, and D. H. Crawford.** 2006. A mouse monoclonal antibody against Epstein-Barr virus envelope glycoprotein 350 prevents infection both in vitro and in vivo. *J Infect Dis* **194**:584-587.
9. **Henle, G., and W. Henle.** 1979. The virus as the etiologic agent of infectious mononucleosis, p. 297-320, *The Epstein-Barr Virus*. Springer.
10. **Herrman, M., J. Muhe, C. Quink, and F. Wang.** 2015. Epstein-Barr Virus gp350 Can Functionally Replace the Rhesus Lymphocryptovirus Major Membrane Glycoprotein and Does Not Restrict Infection of Rhesus Macaques. *J Virol* **90**:1222-1230.

11. **Hoffman, G. J., S. G. Lazarowitz, and S. D. Hayward.** 1980. Monoclonal antibody against a 250,000-dalton glycoprotein of Epstein-Barr virus identifies a membrane antigen and a neutralizing antigen. *Proc Natl Acad Sci U S A* **77**:2979-2983.
12. **Jones, S. T., and M. M. Bendig.** 1991. Rapid PCR-cloning of full-length mouse immunoglobulin variable regions. *Biotechnology (N Y)* **9**:579.
13. **Luzuriaga, K., and J. L. Sullivan.** 2010. Infectious mononucleosis. *N Engl J Med* **362**:1993-2000.
14. **Moutschen, M., P. Leonard, E. M. Sokal, F. Smets, M. Haumont, P. Mazzu, A. Bollen, F. Denamur, P. Peeters, G. Dubin, and M. Denis.** 2007. Phase I/II studies to evaluate safety and immunogenicity of a recombinant gp350 Epstein-Barr virus vaccine in healthy adults. *Vaccine* **25**:4697-4705.
15. **Nemerow, G. R., R. A. Houghten, M. D. Moore, and N. R. Cooper.** 1989. Identification of an epitope in the major envelope protein of Epstein-Barr virus that mediates viral binding to the B lymphocyte EBV receptor (CR2). *Cell* **56**:369-377.
16. **Nemerow, G. R., C. Mold, V. K. Schwend, V. Tollefson, and N. R. Cooper.** 1987. Identification of Gp350 as the Viral Glycoprotein Mediating Attachment of Epstein-Barr-Virus (Ebv) to the Ebv/C3d Receptor of B-Cells - Sequence Homology of Gp350 and C3-Complement Fragment C3d. *Journal of Virology* **61**:1416-1420.
17. **Ogembo, J. G., L. Kannan, I. Ghiran, A. Nicholson-Weller, R. W. Finberg, G. C. Tsokos, and J. D. Fingerhuth.** 2013. Human complement receptor type 1/CD35 is an Epstein-Barr Virus receptor. *Cell reports* **3**:371-385.
18. **Ogembo, J. G., M. R. Muraswki, L. W. McGinnes, A. Parcharidou, R. Sutiwisesak, T. Tison, J. Avendano, D. Agnani, R. W. Finberg, and T. G. Morrison.** 2015. A chimeric EBV gp350/220-based VLP replicates the virion B-cell attachment mechanism and elicits long-lasting neutralizing antibodies in mice. *Journal of Translational Medicine* **13**:50.
19. **Pei, J., B.-H. Kim, and N. V. Grishin.** 2008. PROMALS3D: a tool for multiple protein sequence and structure alignments. *Nucleic Acids Research* **36**:2295-2300.
20. **Perez, E. M., J. Foley, T. Tison, R. Silva, and J. G. Ogembo.** 2016. Novel Epstein-Barr virus-like particles incorporating gH/gL-EBNA1 or gB-LMP2 induce high neutralizing antibody titers and EBV-specific T-cell responses in immunized mice. *Oncotarget*.
21. **Qualtiere, L. F., J. F. Decoteau, and M. Hassan Nasr-el-Din.** 1987. Epitope mapping of the major Epstein-Barr virus outer envelope glycoprotein gp350/220. *J Gen Virol* **68 (Pt 2)**:535-543.
22. **Qualtiere, L. F., J. F. Decoteau, and M. H. Nasreldin.** 1987. Epitope Mapping of the Major Epstein-Barr-Virus Outer Envelope Glycoprotein Gp350/220. *Journal of General Virology* **68**:535-543.
23. **Rees, L., E. J. Tizard, A. J. Morgan, W. D. Cubitt, S. Finerty, T. A. Oyewole-Eletu, K. Owen, C. Royed, S. J. Stevens, R. C. Shroff, M. K. Tanday, A. D. Wilson, J. M. Middeldorp, P. L. Amlot, and N. M. Steven.** 2009. A phase I trial of epstein-barr virus gp350 vaccine for children with chronic kidney disease awaiting transplantation. *Transplantation* **88**:1025-1029.
24. **Rickinson, A. B., and E. Kieff.** 2007. Epstein-Barr Virus, p. 2680-2700. *In* D. Knipe and P. Howley (ed.), *Fields Virology*, Fifth ed, vol. 2. Lippincott Wilkins and Williams, Philadelphia.
25. **Sashihara, J., P. D. Burbelo, B. Savoldo, T. C. Pierson, and J. I. Cohen.** 2009. Human antibody titers to Epstein-Barr Virus (EBV) gp350 correlate with neutralization of infectivity better than antibody titers to EBV gp42 using a rapid flow cytometry-based EBV neutralization assay. *Virology* **391**:249-256.
26. **Sela, M., B. Schechter, I. Schechter, and F. Borek.** 1967. Antibodies to Sequential and Conformational Determinants. *Cold Spring Harbor Symposia on Quantitative Biology* **32**:537-545.

27. **Sitompul, L. S., N. Widodo, M. S. Djati, and D. H. Utomo.** 2012. Epitope mapping of gp350/220 conserved domain of epstein barr virus to develop nasopharyngeal carcinoma (npc) vaccine. *Bioinformation* **8**:479-482.
28. **Sokal, E. M., K. Hoppenbrouwers, C. Vandermeulen, M. Moutschen, P. Leonard, A. Moreels, M. Haumont, A. Bollen, F. Smets, and M. Denis.** 2007. Recombinant gp350 vaccine for infectious mononucleosis: a phase 2, randomized, double-blind, placebo-controlled trial to evaluate the safety, immunogenicity, and efficacy of an Epstein-Barr virus vaccine in healthy young adults. *The Journal of infectious diseases* **196**:1749-1753.
29. **Speck, P., and R. Longnecker.** 1999. Epstein-Barr virus (EBV) infection visualized by EGFP expression demonstrates dependence on known mediators of EBV entry. *Arch Virol* **144**:1123-1137.
30. **Szakonyi, G., M. G. Klein, J. P. Hannan, K. A. Young, R. Z. Ma, R. Asokan, V. M. Holers, and X. S. Chen.** 2006. Structure of the Epstein-Barr virus major envelope glycoprotein. *Nature structural & molecular biology* **13**:996-1001.
31. **Tanner, J., J. Weis, D. Fearon, Y. Whang, and E. Kieff.** 1987. Epstein-Barr virus gp350/220 binding to the B lymphocyte C3d receptor mediates adsorption, capping, and endocytosis. *Cell* **50**:203-213.
32. **Tanner, J., Y. Whang, J. Sample, A. Sears, and E. Kieff.** 1988. Soluble gp350/220 and deletion mutant glycoproteins block Epstein-Barr virus adsorption to lymphocytes. *J Virol* **62**:4452-4464.
33. **Tanner, J. E., M. Coincon, V. Leblond, J. Hu, J. M. Fang, J. Sygusch, and C. Alfieri.** 2015. Peptides designed to spatially depict the Epstein-Barr virus major virion glycoprotein gp350 neutralization epitope elicit antibodies that block virus-neutralizing antibody 72A1 interaction with the native gp350 molecule. *J Virol* **89**:4932-4941.
34. **Thorley-Lawson, D. A., and K. Geilinger.** 1980. Monoclonal antibodies against the major glycoprotein (gp350/220) of Epstein-Barr virus neutralize infectivity. *Proc Natl Acad Sci U S A* **77**:5307-5311.
35. **Urquiza, M., R. Lopez, H. Patino, J. E. Rosas, and M. E. Patarroyo.** 2005. Identification of three gp350/220 regions involved in Epstein-Barr virus invasion of host cells. *The Journal of biological chemistry* **280**:35598-35605.
36. **Weiss, E. R., G. Alter, J. G. Ogembo, J. L. Henderson, B. Tabak, Y. Bakis, M. Somasundaran, M. Garber, L. Selin, and K. Luzuriaga.** 2017. High Epstein-Barr Virus Load and Genomic Diversity Are Associated with Generation of gp350-Specific Neutralizing Antibodies following Acute Infectious Mononucleosis. *J Virol* **91**.
37. **Zhang, P. F., M. Klutch, G. Armstrong, L. Qualtiere, G. Pearson, and C. J. Marcus-Sekura.** 1991. Mapping of the epitopes of Epstein-Barr virus gp350 using monoclonal antibodies and recombinant proteins expressed in *Escherichia coli* defines three antigenic determinants. *J Gen Virol* **72** (Pt 11):2747-2755.

Figure legends:

FIG 1 Specificity of anti-gp350 antibodies. (A) ELISA screening of hybridoma (HB) supernatants for anti-gp350-specific antibodies. Soluble EBV gp350 protein was used as the target antigen at 0.5 µg/ml. nAb-72A1 at 10 µg/ml and 1x phosphate buffered saline (PBS) were used as positive and negative (not shown) controls,

respectively. Bound antibodies were detected using HRP-conjugated anti-mouse IgG (1:2,000). Twenty-three HB clones with ELISA signals two times greater than those of PBS control were considered as positive hybridomas. (B) Determining specificity of anti-gp350-producing hybridoma supernatants by immunoblotting with gp350-transfected stable CHO lysate. (C) Flow cytometric analysis of surface expression of gp350 protein on gp350-expressing CHO cells. Cells were stained with anti-gp350 mAb (1:250), followed by secondary goat anti-mouse conjugated to AF488.

FIG 2 PROMALS3D multiple sequence alignment of (A) V_H and (B) V_L regions of 15 mAbs and nAb-72A1. The highly variable complementarity determining regions (CDR) 1–3, indicated by black boxes, define the antigen binding specificity. The conserved framework regions (FR) 1–4 flank the CDRs. Consensus amino acids (aa) are in bold and upper case. Consensus-predicted secondary structure (ss) symbols: alpha-helix, h; beta-strand, e.

FIG 3 Neutralization activity of novel anti-gp350 mAbs against EBV-eGFP in Raji cells. EBV-eGFP was pre-incubated with 15 anti-gp350 mAbs at (A) 10 µg/ml and (B) 50 µg/ml, followed by incubation with Raji cells for 48 h. EBV-eGFP⁺ cells were enumerated using flow cytometry. Anti-gp350 (nAb-72A1) and anti-gH/gL (E1D1) mAbs served as positive controls and non-neutralizing anti-gp350 (2L10) mAb served as negative control.

FIG 4 Identification of nine gp350 epitopes using 15 neutralizing and non-neutralizing mAbs. Residues in bold represent the gp220 splice variant region. Residues in red represent RepliTope-identified epitopes and exact residues. Italic residues represent canonical neutralizing epitope, underlined residues represents epitope bound by all assayed mAb.

561

562 FIG 5 Identification of novel gp350-neutralizing epitope by epitope mapping of neutralizing (nAb-72A1 and
563 HB20) vs. non-neutralizing (HB5) anti-gp350 mAbs. Residue positions of nAb-72A1 (red), HB20 (blue), and
564 HB5 (green) are indicated. Bold black residues represent splice variant region, underlined residues represent
565 epitopes bound by nAb-72A1, HB20, and HB5. Italic residues represent epitopes bound by nAbs.
566 Red residues represent epitopes bound by nAbs-72A1, blue residues represent residues bound by HB20 and
567 green represents residues bound by HB5.

568

Table 1. Summary of Illumina Dual Demultiplex of V_H and V_L regions.

Sample	Chain	Starting Pairs	PEAR Merged Reads	Length Filtered Reads	Primer Matched Reads	3x Reads	>5% Unique Coding	>5% Unique Non-Coding	
HB1	HEAVY	51,641	51,210	46,655	32,482	22,725	1	0	
	LIGHT	280,048	279,012	100,725	68,041	58,570	1	1	
HB2	HEAVY	22,793	22,621	16,475	11,415	7,429	1	0	
	LIGHT	167,230	166,496	161,764	132,752	115,886	1	1	
HB3	HEAVY	26,382	26,162	25,542	16,910	11,709	1	0	
	LIGHT	12,681	12,609	11,753	9,809	8,023	1	1	
HB4	HEAVY	38,811	38,238	17,151	11,957	7,217	2	0	\$
	LIGHT	179,249	129,752	111,996	78,392	66,419	1	1	
HB.5	HEAVY	42,951	42,173	35,793	25,842	17,173	1	0	
	LIGHT	176,073	175,267	168,806	141,712	127,045	1	1	
HB6	HEAVY	26,142	25,981	22,245	15,658	10,453	1	0	
	LIGHT	171,996	171,370	167,730	138,348	122,397	1	1	
HB7	HEAVY	32,443	32,094	25,449	17,615	11,836	1	0	
	LIGHT	67,031	63,924	37,344	26,271	22,378	1	1	*
HB8	HEAVY	140,091	103,349	92,744	58,292	31,583	1	0	
	LIGHT	151,244	115,527	102,439	82,154	70,803	1	1	
HB9	HEAVY	37,057	36,473	19,544	11,585	7,358	1	0	
	LIGHT	409,432	310,529	136,074	106,820	90,063	1	2	*
HB10	HEAVY	38,181	37,981	26,043	17,104	11,391	1	0	
	LIGHT	114,255	112,498	106,914	84,368	75,370	1	1	
HB11	HEAVY	22,225	21,841	6,956	4,408	2,465	1	0	
	LIGHT	106,465	102,278	65,332	50,232	44,527	1	0	
HB12	HEAVY	83,044	82,355	46,350	30,276	20,886	1	0	
	LIGHT	53,098	47,336	15,823	7,560	5,845	1	1	
HB13	HEAVY	81,451	80,372	47,995	32,216	20,139	2	0	\$
	LIGHT	27,314	24,774	8,987	5,457	4,104	2	1	
HB14	HEAVY	76,299	75,357	28,309	19,104	12,939	1	0	
	LIGHT	153,011	149,264	48,474	29,710	25,133	1	1	
HB.15	HEAVY	26,551	26,410	16,387	11,434	7,002	2	0	\$
	LIGHT	78,525	77,778	43,509	29,504	24,731	1	1	
HB16	HEAVY	54,249	53,943	9,517	7,128	4,179	1	0	
	LIGHT	42,048	40,351	30,602	22,758	18,251	2	1	\$
HB17	HEAVY	111,614	110,882	81,428	50,844	35,949	1	0	
	LIGHT	102,490	100,488	83,925	65,925	57,727	1	1	*
HB18	HEAVY	211,215	155,410	146,256	91,009	50,308	1	0	
	LIGHT	212,261	161,879	155,235	123,096	105,959	1	1	
HB19	HEAVY	109,692	82,221	20,546	12,587	7,274	1	1	
	LIGHT	70,828	69,744	62,572	48,354	42,051	1	1	
HB20	HEAVY	15,781	15,632	12,789	7,757	4,852	1	0	
	LIGHT	135,527	133,208	118,513	90,717	78,701	1	1	
HB21	HEAVY	15,312	15,202	8,577	5,645	3,420	1	0	
	LIGHT	102,450	100,171	89,059	68,552	60,500	1	1	
HB22	HEAVY	217,959	156,488	154,008	95,755	50,245	1	0	
	LIGHT	205,334	156,986	143,386	108,728	85,136	1	0	
HB23	HEAVY	196,390	143,929	123,028	71,076	39,358	2	0	\$
	LIGHT	158,594	120,140	115,476	90,004	78,787	2	0	

72A1	HEAVY	213,480	158,199	156,215	107,395	68,486	2	0	\$
	LIGHT	187,216	140,964	132,945	105,783	91,208	1	1	

*= hybridoma with V_L chain sequences identified with >1% frequency, \$ = hybridoma with more than one unique, plausible-coding V_H chain sequence with > 5% frequency. The term “unique” refers to unique sequence counts (so, identical sequences found in a substantial frequency of merged reads, not necessarily unique compared to other samples).

Table 2: Summarized biochemical and functional characterizations of anti-gp350 antibodies.

Antibody	IgG sub-class	Light chain	ELISA binding to purified EBV gp350/220	FACS (CHO Cells)	Western blot	Neutralization activity
HB1	IgG1	K	+	+	+	-
HB2	IgG2a	K	+	+	-	-
HB3	IgG2a	K	+	+	-	-
HB4	IgG1	K	+	-	+	ND
HB5	IgG2a	K	+	+	+	-
HB6	IgG1	K	+	+	-	-
HB7	IgG1	K	+	-	-	-
HB8	IgG1	K	+	-	+	-
HB9	IgG2a	K	+	+	+	+
HB10	IgG1	K	+	-	+	+
HB11	IgG1	K	+	+	+	+
HB12	IgG1	K	+	+	+	-
HB13	IgG1	K	+	-	-	ND
HB14	IgG1	K	+	-	+	-
HB15	IgG1	K	+	+	+	ND
HB16	IgG1/ IgM	K	+	-	+	ND
HB17	IgG2b	K	+	+	+	+
HB19	IgG1/ IgM	K	+	+	+	ND
HB20	IgG2a	K	+	+	-	+
HB21	IgG1/IgG2b	K	+	+	-	ND
HB22	IgG1	K	+	-	+	-
HB23	IgG1	K	+	+	+	ND
72A1	IgG1	K	+	+	+	+
E1D1 (anti gH/gL)	IgG1	K	-	-	-	+

ND=Not determined, + = positive, - = negative, ELISA = enzyme-linked immunosorbent assay, κ = Kappa

581 Table 3. Summary of gp350 epitope mapping over time using various methodologies.

Method	mAbs/protein/ peptides	Number of epitopes identified	Reference
Competitive binding assay: Tagged mAb vs. untagged mAb	Newly generated mAbs	8 epitopes – Sequence not defined	Qualtiere et al., 1987 (21)
Binding studies: Determine the effects of anti-gp350 mAbs on gp350 binding to CR2	Newly generated mAbs	2 possible regions identified by sequence alignment to C3d sequence: 1. aa 21–28 2. aa 372–378	Nemerow et al., 1987 (16)
Peptide digest and immunoprecipitation	Truncated and mutant protein; mAbs (72A1 and BOS-1)	Narrowed down to the first 470 residues	Tanner et al., 1988 (32)
Binding studies	Peptide and protein	2 sequences defined: 1. aa 21–28 2. N-terminus of gp350	Nemerow et al., 1989 (15)
Dot Blot immunoassay: Purified truncated protein incubated with mAbs	Protein – 8 clones overlapping N- and C-terminal portions of protein; mAbs from Qualtiere <i>et al.</i> , 1987	3 sequences defined: 1. aa 310–325 2. aa 326–439 3. aa 733–841	Zhang et al., 1991 (37)
Peptide cell binding assay to 2 CR2-positive (Raji and Ramos) and 1 -negative (P3HR-1) cell lines	Synthesized peptides covering gp350 (907 aa)	7 regions, 3 identified: 1. aa 142–161 2. aa 282–301 3. aa 822–841	Urquiza et al., 2005 (35)
Crystal structure and binding studies	Mutant proteins; mAbs 72A1	3 epitopes (based on 72A1 binding and gp350 4-443) 1. aa 16-29 2. aa 142-161 3. aa 282-301	Szakonyi et al., 2006 (30)
Structural docking studies and antigenicity mapping	gp350 and CR2 crystal structure alignment/docking	Single epitope (based on gp350 aa 1–470) 1. aa147–165	Sitompul et al., 2012 (27)

Structural alignment: computer modeling of gp350 and 72A1 and docking studies	Peptides (used in immunization); mAb (72A1)	4 epitopes: identified 1. aa 14-20 2. aa 144-161 3. aa 194-211 4. aa 288-291	Tanner et al., 2015 (33)
--	--	--	-----------------------------

582

		FR-1										FR-2											
Conservation:		6	9	96	6	69666	6	9	666	69	6	966	69	99669	9	6							
72A1	1	QAVLTQESALTTSPGETVTLTCRSST	GAVT--TSNYANWVQEKPDHLFTGLIG										GTNNRVPGVVPARFSGSLIGDKAALTIT										78
HB1	1	-----KFMSTSVGDRVSVTCKASQ	NV-----GTNVAWYQQKPGQSPKALIYSTSSRYTGVPDRFAGSGSGTDYTLTIS										STSSRYTGVPDRFAGSGSGTDYTLTIS										68
HB2	1	-----AILSASPGEKVTMTCRATSS	-----VNVMHWYQQKPGSSPKPWIYATSNLASGVVPARFSGSGSGTSYSLTIS										ATSNLASGVVPARFSGSGSGTSYSLTIS										67
HB3	1	-----SSLSASLGDRVTISCRASQD	-----IGNYLNWYQQKPDGTIKLLIYYTSRLHSGVPSRFSGSGSGTDYSLTIS										YTSRLHSGVPSRFSGSGSGTDYSLTIS										68
HB5	1	-----AILSVPSPGERVSFSCRASQS	-----IGTSHHWYQQRTNDSPRLLIKYASESISGIPPRFSGSGSGTDFTLSIN										YASESISGIPPRFSGSGSGTDFTLSIN										68
HB6	1	-----AILSASPGEKVTMTCRATSS	-----VNVMHWYQQKPGSSPKPWIYATSNLASGVVPARFSGSGSGTSYSLTIS										ATSNLASGVVPARFSGSGSGTSYSLTIS										67
HB7	1	-----KFMSTSVGDRVNITCKASQSV	-----GNVAVWFQQKPGQSPKLLIYSASNRYTGIPDRFTGSGSGTDFTLTCTN										SASNRYTGIPDRFTGSGSGTDFTLTCTN										68
HB8	1	-----LSLPVSLGDQASISCRSSQS	SIVHSNNGNTYLEWYLQKAGQSPKLLIYKVSNNRFSGVDPDRFGGSGSGTDFTLKIS										YKVSNNRFSGVDPDRFGGSGSGTDFTLKIS										73
HB9	1	-----LSLPVSLGDQASISCRSSQS	SIVHSNNGNTYLEWYLQKAGQSPKLLIYKVSNNRFSGVDPDRFGGSGSGTDFTLKIS										YKVSNNRFSGVDPDRFGGSGSGTDFTLKIS										73
HB10	1	-----SSLSASLGDRVTISCRASQD	-----IGNYLNWYQQKPDGTVKLLIYYTSRLHSGVPSRFSGSGSGTDYSLTIS										YTSRLHSGVPSRFSGSGSGTDYSLTIS										68
HB11	1	-----AIMSASLGEEKVTMSCRASSS	-----V-NFMNWYQQKSDDSPKLLIYYISNLAAPGVVPARFSGSGSGNSYSLTIS										YISNLAAPGVVPARFSGSGSGNSYSLTIS										67
HB12	1	-----LSLPVSLGDQASISCRSSQS	SLVHSNNGNTYLEWYLQKPGQSPKLLIYKVSNNRFSGVDPDRFSGSGSGTDFTLKIS										YKVSNNRFSGVDPDRFSGSGSGTDFTLKIS										73
HB14	1	-----LSLPVSLGDQASISCRSSQS	SIVHDNGNTYLEWYLQKPGQSPKLLIYKVSNNRFSGVLDKFSGSGSGTDFTLKIS										YKVSNNRFSGVLDKFSGSGSGTDFTLKIS										73
HB17	1	-----SSLSASLGDRVTISCRASQD	-----IGNYLNWYQQKPDGTIKLLIYYTSRLHSGVPSRFSGSGSGTDYSLTIS										YTSRLHSGVPSRFSGSGSGTDYSLTIS										68
HB20	1	-----SSLSASLGDRVTISCRASQD	-----IGNYLNWYQQKPDGTVKLLIYYTSRLHSGVPSRFSGSGSGTDYSLTIS										YTSRLHSGVPSRFSGSGSGTDYSLTIS										68
HB22	1	-----LSLPVSLGDQASISCRSSQS	SIVHSNNGNTYLEWYLQKPGQSPKLLIYKVSNNRFSGVDPDRFSGSGSGTDFTLRIS										YKVSNNRFSGVDPDRFSGSGSGTDFTLRIS										73
Consensus aa:		hsh S.G -phoho CR Tsps.....ssh hpW@bQK sspo.+ .LIb.hSp.hsGVPsRFoGSGSGsc@oLpIo																					
Consensus ss:		eeee		eeeeeeee				ee		eeeeee				eeee		eeee				eeee		eeee	

CDR-1

CDR-2

		FR-3						FR-4					
Conservation:		6	99	6	9	9	6	699	9969	699999999			
72A1	79	GAQTEDEAIYFC	VLWHS-NHWV	FGGGTKLTVL	-----							109	
HB1	69	NVQSEDLAEYFC	QQYNT-YPYTF	GGGTRLDIKRADAAPT	TV							107	
HB2	68	RVEAEDAATYYC	QQWSSNPP-TF	GAGTKLELKRADAAPT	TV							106	
HB3	69	NLEEEDIATYFC	QQGNTLPP-TF	GGGTKLEIKRADAAPT	TV							107	
HB5	69	SVESEDIADYHC	QQSNSWPMLTF	GAGTKLELKRADAAPT	TV							108	
HB6	68	RVEAEDAATYYC	QQWSSNPP-TF	GAGTKLELKRADAAPT	TV							106	
HB7	69	NMQSEDLADYFC	QQYSS-YPLTF	GAGTKLELKRADAAPT	TV							107	
HB8	74	RVEAEDLGVIYFC	FQGSHPYTF	GGGTKLEIKRADAAPT	TV							112	
HB9	74	RVEAEDLGVIYFC	FQGSHPYTF	GGGTKLEIKRADAAPT	TV							112	
HB10	69	NLEEEDIATYFC	QQGNTLPP-TF	GGGTKLEIKRADAAPT	TV							107	
HB11	68	GMEGEDAATYYC	QQFTSSPSWTF	GGGTKLEIKRADAAPT	TV							107	
HB12	74	RVEAEDLGVIYFC	SQSTH-VPLTF	GSGTKLEIK-----								104	
HB14	74	RVEAEDLGIYFC	FQGSHPVPTF	GGGTKLEIK-----								104	
HB17	69	NLEEEDIATYFC	QQGNTLPP-TF	GGGTKLEIKRADAAPT	TV							107	
HB20	69	NLEQEDIATYFC	QQGNALPP-TF	GGGTKLEIKRADAAPT	TV							107	
HB22	74	RVEAEDLGVIYFC	FQGSHPVWTF	GGGTKLEIKRADAAPT	TV							112	
Consensus aa:		.hps ED ht. Y@C.Q .sp.s.. TFGtG TKL- LKRADAAPT											
Consensus ss:		hhheeeeeeeee						ee		eeeeee			

CDR-3

1	MEAALLVCQY	TIQSLIHLTG	EDPGFFNVEI	PEFPFYPTCN	VCTADVNVTI	NFDVGGKKHQ
61	LDLDFGQLTP	HTKAVYQPRG	AFGGSENATN	LFLLELLGAG	ELALTMRSKK	LPINVTTGEE
121	QQVSLESVDV	YFQDVFGTMW	CHHAEMQNPV	YLIPETVPYI	KWDNCNSTNI	TAVVRAQGLD
181	VTLPLSLPTS	AQDSNFSVKT	EMLGNEIDIE	CIMEDGEISQ	VLPGDNKFNI	TCSGYESHVP
241	SGGILTSTSP	VATPIPGTGY	AYSLRLTPRP	VSRLGNNSI	LYVFYSGNGP	KASGGDYCIQ
301	SNIVFSDEIP	ASQDMPTNTT	DITYVGDNAT	YSVPMVTSED	ANSPNVTVTA	FWAWPNNTET
361	DFKCKWTLTS	GTPSGCENIS	GAFASNRTFD	IT/VSGLGTAP	KTLIITRTAT	NATTTTHKVI
421	FSKAPESTTT	SPTLNTTGFA	DPNTTTGLPS	STHVPTNLTA	PASTGPTVST	ADVTSPPTAG
481	TTSGASPVTP	SPSPWDNGTE	SKAPDMTSST	SPVTTPTPNA	TSPTPAVTTP	TPNATSPTPA
541	VTTPTPNATS	PTLGKTSPTS	AVTTPTPNAT	SPTLGKTSPT	SAVTTPTPNA	TSPTLGKTSP
601	TSAVTTPTPN	ATGPTVGETS	PQANATNHTL	GGTSPTPVVT	SQPKNATSAV	TTGQHNITSS
661	STSSMSLRPS	SNPETLSPST	SDNSTSHMPL	L TSAHPTGGE	NITQVTPASI	STHHVSTSSP
721	APRPGTTSQA	SGPGNSSTST	KPGEVNVTKG	TPPQNATSPQ	APSGQKTAVP	TVTSTGGKAN
781	STTGGKHTTG	HGARTSTEPT	TDYGGDSTTP	RPRYNATTYL	PPSTSSKLRP	RWTFTSPPVT
841	TAQATVPVPP	TSQPRFSNLS	MLVLQWASLA	VLTL LLLVM	ADCAFRRNLS	TSHTYTTPPY
901	DDAETYV					

1	MEAALLVCQY	TIQSLIHLTG	EDPGFFNVEI	PEFPFYPTCN	VCTADVNVTI	NFDVGGKKHQ
61	LDLDFGQLTP	HTKAVYQPRG	AFGGSENATN	LFLLELLGAG	ELALTMRSKK	LPINVTTGEE
121	QQVSLESVDV	YFQDVFGTMW	CHHAEMQNPV	YLIPETVPYI	KWDNCNSTNI	TAVVRAQGLD
181	VTLPPLSLPTS	AQDSNFSVKT	EMLGNEIDIE	CIMEDGEISQ	VLPGDNKFNI	TCSGYESHVP
241	SGGILTSTSP	VATPIPGTGY	AYSLRLTPRP	VSREFLGNNI	LYVFYSGNGP	KASGGDYCIQ
301	SNIVFSDEIP	ASQDMPTNTT	DITYVGDNAT	YSVPMVTSED	ANSPNVTVTA	FWAWPNNTET
361	DFKCKWTLTS	GTPSGCENIS	GAFASNRTFD	ITVSGLGTAP	KTLIIITRTAT	NATTTTHKVI
421	FSKAPESTTT	SPTLNTTGFA	DPNNTTGLPS	STHVPTNLTA	PASTGPTVST	ADVTSPTPAG
481	TTSGASPVTP	SPSPWDNGTE	SKAPDMTSST	SPVTTPTPNA	TSPTPAVTTP	TPNATSPTPA
541	VTTPTPNATS	PTLGKTSPTS	AVTTPTPNAT	SPTLGKTSPT	SAVTTPTPNA	TSPTLGKTSP
601	TSAVTTPTPN	ATGPTVGETS	PQANATNHTL	GGTSPTPVVT	SQPKNATSAV	TTGQHNITSS
661	STSSMSLRPS	SNPETLSPST	SDNSTSHMPL	L TSAHPTGGE	NITQVTPASI	STHHVSTSSP
721	APRPGTTSQA	SGPGNSSTST	KPGEVNVTKG	TPPQNATSPQ	APSGQKTAVP	TVTSTGGKAN
781	STTGGKHTTG	HGARTSTEPT	TDYGGDSTTP	RPRYNATTYL	PPSTSSKLRP	RWTFTSPVVT
841	TAQATVPVPP	TSQPRFSNLS	MLVLQWASLA	VLTL LLLLVM	ADCAFRRNLS	TSHTY TTPPY
901	DDAETYV					