1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

**Fusion expression and anti-*Aspergillus flavus* activity of a novel inhibitory
protein DN-AflR**

Yuan Liang, Qing Kong,[#] Yao Yao, Shujing Xu, Xiang Xie

School of Food Science and Engineering, Ocean University of China, Qingdao,
Shandong, China

Running Title: Fusion expression of DN-AflR

[#]: Address correspondence to Qing Kong, kongqing@ouc.edu.cn.

23

24 **ABSTRACT** The regulatory gene (*aflR*) of aflatoxin encodes AflR, a positive
 25 regulator that activates transcriptional pathway of genes in aflatoxin biosynthesis.
 26 New L-Asp-L-Asn (DN) extracted from *Bacillus megaterium* inhibited the growth of
 27 *A. flavus* had been elucidated in our laboratory. The genes encoding DN and binuclear
 28 zinc finger cluster protein of AflR were fused, then fusion protein could compete with
 29 the AflS-AflR complex for the AflR binding site and significantly improve anti-*A.*
 30 *flavus* activity of DN. The fusion gene *dn-aflR* was cloned into pET32a and
 31 recombinant plasmid was introduced into *Escherichia coli* BL21. The highest
 32 expression was observed after 10 h induction and purified by affinity chromatography
 33 column. Compared with DN, the novel fusion protein DN-AflR significantly inhibited
 34 the growth of *A. flavus* and biosynthesis of aflatoxin B₁. This study promoted the use
 35 of competitive inhibition of fusion proteins to reduce the expression of regulatory
 36 genes in the biosynthetic pathway of aflatoxin. Moreover, it provided more supports
 37 for deep research and industrialization of such novel, anti-*A. flavus* bio-inhibitors.

38 **IMPORTANCE** Aflatoxin contamination has seriously influence on export of
 39 agricultural products, income of farmers and economic development. Biological
 40 methods, especially using antagonistic microorganisms to inhibit aflatoxin
 41 biosynthesis gradually become the hot spot in recent years. DN (L-Asp-L-Asn) from
 42 *Bacillus megaterium*, which could inhibit growth of *Aspergillus flavus* and synthesis
 43 of aflatoxin, has been identified. In this report, we fused the genes encoding inhibitory
 44 peptides (DN) and specific zinc finger cluster protein, and expressed the novel anti-*A.*

flavus protein in *Escherichia coli*. Compared with DN, the inhibitory ability of novel protein has been improved significantly. This research showed fusion expression of anti-fungal proteins, such as DN-AflR, is a promising method to economically improve the inhibitory activity of bio-inhibitors for *A. flavus*.

KEYWORDS AflR, *Aspergillus flavus*, *dn-aflR*, fusion expression

INTRODUCTION

Aflatoxin is one of the most potent naturally occurring toxic and carcinogenic compound, which is a mycotoxin that poses a serious threat to human health (1). Aflatoxin contamination has seriously influence on export of agricultural products, income of farmers and economic development (2, 3). Biological methods, especially using antagonistic microorganisms to inhibit aflatoxin biosynthesis gradually become hot spot in recent years (4). Palumbo et al. isolated one strain of *Bacillus* from almonds, which could inhibit the growth of *Aspergillus flavus* (5). Early studies found that *Mycobacterium smegmatis* and *Rhodococcus erythropolis* could produce F420 H₂-dependent reductases to degrade aflatoxin (6, 7). Some studies isolated antifungal compounds from *Bacillus* and verified their inhibitory effects on the growth of *A. flavus* (8, 9). Above all, antagonistic microorganisms could produce metabolites or enzymes to inhibit expression of regulatory genes, or degrade aflatoxin.

Aflatoxin biosynthetic pathway has been studied for years and is one of the best understood fungal secondary metabolic pathways. The whole-genome sequencing of

A. flavus has been accomplished, and we could better control aflatoxin contamination through deep research of the regulatory genes and mechanisms. Up to now, at least 34 genes have been identified as members of the aflatoxin pathway gene cluster. On the 82-kb biosynthetic gene cluster of aflatoxin, *aflR* and *aflS* (formerly known as *aflJ*) are genes involved in pathway regulation (10, 11). *aflR* is necessary for transcription of some genes in *Aspergillus* gene cluster (12-14). This gene encodes a specific DNA binding protein (AflR) containing 444 amino acids, and the 29th to 56th amino acids constitute a binuclear zinc finger cluster protein with sequence Cys-Xaa2-Cys-Xaa6-Cys-Xaa6-Cys-Xaa2-Cys-Xaa6-Cys, which is the key region of AflR to activate gene transcription and determines AflR-binding specificity (15). AflR, which possesses DNA-binding and activation domains typical of the GAL4-type family of positive regulatory proteins in yeast and fungi (16). A common feature of fungal gene clusters, including those for secondary metabolites, is the presence of specific regulatory genes, which have been found to encode members of the zinc binuclear cluster protein family typified by GAL4 (17-19). AflS was found to be involved in the regulation of transcription. Between *aflR* and *aflS* is intergenic region, where promoter region is located. AflS-AflR complex binds to AflR binding site and activates aflatoxin biosynthesis. Studies suggested that failure to product aflatoxin due possibly to alternation in the interaction between AflS and AflR (10, 20). Bio-inhibitors inhibited expression of *aflR* or *aflS* mainly by acting on the intergenic region, thereby prevented the formation of AflS-AflR complex, so that AflS-AflR complex could not bind to AflR binding site and activate biosynthetic pathway of

aflatoxin.

Recently, we identified DN (L-Asp-L-Asn) from *Bacillus megaterium*, which could inhibit growth of *A. flavus* and synthesis of aflatoxin. To improve the inhibitory effect of DN, we transformed the genes encoding DN and GAL4-type zinc finger cluster protein (specifically binds to the AflR binding site) by gene fusion. We hypothesized the fusion protein could compete with AflS-AflR complex by acting on AflR binding sites to inhibit the activation of AflS-AflR complex and improve anti- *A. flavus* activity of DN.

RESULTS

Effects of DN on mycelia morphology of *A. flavus*. (i) TEM. In control group, the internal structure of cells grew normally and various organelles were clearly visible (Fig. 2A and B). However, cell structures of mycelia treated by DN were obviously abnormal: organelles were degenerated, and vacuole became significantly larger, expanded and fractured (Fig. 2C and D).

(ii) SEM. Mycelia of control group were integrity, showing straight, neatly arranged and smooth surface (Fig. 2E and F). While mycelia treated with DN grew abnormally: mycelia were rough and had uneven thickness, and appeared to be distorted and broken (Fig. 2G and H).

Expression and identification of fusion protein. The expected DNA and vector fragments were seen in 1% agarose electrophoresis after digesting with EcoR I and

Hind III. Colony PCR and DNA sequencing showed that target fragment (*dn-afIR*) was successful inserted into the vector. The molecular weight of empty plasmid (Trx-His-pET32a) and fusion protein (Trx-His-DN-AfIR) were predicted to be 19 kDa and 22 kDa. Expression vector with fusion gene had a distinct protein band (24 kDa, Fig. 3A, lanes 1 and 2). The empty plasmid did not express the band at the corresponding position and had its own specific band (Fig. 3A, lane 3). Results illustrated successful expression of the fusion protein DN-AFLR. The supernatant after ultrasonic breaking (Fig. 3B, lane 1) and the precipitate after 8 mM urea dissolved (Fig. 3B, lane 2) both showed obvious protein bands. The result was consistent with predicted weight of expected protein before purification (Fig. 3A). There were no obvious protein bands in wash buffer (Fig. 3B, lanes 3-5), and there was only one obvious protein band in elution buffer (Fig. 3B, lanes 6 and 7), which corresponded to the target protein. ImageJ software (<https://imagej.nih.gov/ij/>) was used to compare the density of the bands on the gel, the fusion protein accounts for approximately 57.4% of the total cellular protein, the relative content of fusion protein was about 600 $\mu\text{g ml}^{-1}$.

Inhibitory effect of DN-AfIR on growth of *A. flavus*. Plate treated by DN had only a small inhibition zone of 5.0 ± 0.2 mm (Fig. 4A, plate a). The control group (Fig. 4A, plate c) and 200 mM imidazole treated group (Fig. 4A, plate b) showed no significant zone of inhibition. Diameter of inhibition zone treated with fusion protein was as high as 21.3 ± 0.2 mm (Fig. 4A, plates d-f). Inhibitory ability of DN-AfIR was much stronger than that of DN. The MICs of DN-AfIR and DN were 150 $\mu\text{g ml}^{-1}$ and

400 $\mu\text{g ml}^{-1}$, respectively. MFC of DN-AflR was 400 $\mu\text{g ml}^{-1}$. Results of two repeated experiments were consistent (Table 1). The MIC and MFC of fusion protein were lower than that of DN, indicating that the fusion protein had stronger inhibitory effect on growth of *A. flavus* at a lower concentration.

Inhibitory effect of fusion protein on aflatoxin B₁ biosynthesis. Two groups of experiments were performed using DN-AflR and DN with final concentrations of 30, 60, and 90 $\mu\text{g ml}^{-1}$. The residual rate of aflatoxin B₁ after fusion protein treatment was 24.37%, 14.21% and 6.74%, respectively, and the residual rate of aflatoxin B₁ after DN treatment was 44.39%, 22.19% and 10.99%, respectively (Fig. 4B). Compared with DN, fusion protein had stronger inhibitory effect on biosynthesis of aflatoxin B₁ ($P<0.05$). At 30 $\mu\text{g ml}^{-1}$ of DN-AflR, more than 75% aflatoxin B₁ was degraded.

Physical properties and molecular structure of DN-AflR. Amino acid sequence, isoelectric point and other basic informations were analyzed by ExPASy (<http://web.expasy.org/protparam/>; Table 2). Secondary structure of protein was predicted by SPLIT (<http://splitbioinf.pmfst.hr/split/4/>; Fig. 5A). Spatial structure was predict by SWISS-MODEL. The fusion protein DN-AflR carries a positive charge, has a relative strong hydrophobic effect, and contains the characteristic secondary structure of the protein. By analyzing the predicted structures of DN-AflR (Fig. 5B) and GAL4 (Fig. 5C), DN-AflR has the zinc finger DNA-binding functional structure (specifically binds to AflR binding site) (Fig. 5D).

DISCUSSION

At present, more and more microorganisms and their metabolites were reported to inhibit the growth of *Aspergillus flavus* and degrade aflatoxins (5, 26). Albert et al. transformed the laccase gene of *T. versicolor* into recombinant *A. niger* by gene cloning, and the inhibition rate on aflatoxin of laccase (118 U L^{-1}) after recombinant expression as high as 55% (27). Douillard et al. demonstrate that novel *L. lactis* fusion partner expression vectors allow high-level expression of soluble heterologous proteins (28). Study showed that Fh8 tag fusion expression could significantly improve the ability of *E. coli* to express soluble exogenous proteins (29). Genetic engineering could highly express the active proteins and other metabolites in prokaryotic or eukaryotic hosts, which is the great way to cut down costs.

However, there was few report on the inhibition of growth of *A. flavus* by molecular modification of positive regulatory genes in aflatoxin biosynthesis. Fusion expression has been continuously applied to the process of expressing recombinant protein in order to improve functions of the active protein (30, 31). TEM and SEM are effective applications to analyze the characteristics and morphologies of samples. SEM showed that DN caused changes in mycelial morphology of *A. flavus*. TEM showed the ruptured vacuoles accounted for most of cell space and destroyed the function and balance of other organelles, which affected normal growth of the whole cell (Fig. 2). So that DN had the ability of inhibiting the growth of *A. flavus*.

In order to improve the anti-*A. flavus* effect of DN, we fused the genes encoding DN and sequences of zinc finger cluster protein (specifically binds to the AflR binding site), then successfully constructed a recombinant plasmid dn-aflR-pET32a

(Fig. 1). In the process of *E. coli* expression, when expressed in the host system at a high level, the recombinant protein was easy to form inclusion bodies (32). This may be due to the fact that during the expression, the protein folded too fast, while insufficient supply of enzymes or co-factors made it cannot form the correct secondary bond. According to the reports (33), we used 16°C, 160 rpm as the conditions for inducing expression. In general, target proteins with a molecular weight of less than 5 kDa or more than 100 kDa could not be expressed easily. The smaller molecular weight of protein, the easier it was to be degraded. In order to avoid degradation of target protein due to its low molecular weight, we used the pET32a vector to express the active protein, and its Trx and His tag significantly improved the stability of expression, and enhanced purification of protein, respectively. Because target protein contained the His tag, result of SDS-PAGE showed that molecular weight of fusion protein was 24 kDa (Fig 3), which was a little more than the predicted molecular weight by the theoretical calculations (20 kDa). Compared with DN, the fusion protein DN-AflR had stronger inhibitory effect on growth of *A. flavus* (Fig. 4A). Simultaneously, it also had significant advantages in inhibiting aflatoxin B₁ biosynthesis (Fig. 4B), especially, under 30 µg mL⁻¹ concentration of DN-AflR, more than 75% aflatoxin B₁ was inhibited. To get a deeper comprehension of DN-AflR, its physical properties and molecular structures were predicted. Outer membrane of most cells is negatively charged, while most anti-fungal proteins are positively charged and they can bind to the cells surface through electrostatic attraction (34). The α-helical peptides with higher hydrophobic properties have a higher ability to lyse membranes,

and the inhibitory activity of some peptides would even disappear with reduced hydrophobicity (35). Through the predicted results, the fusion protein DN-AflR carries positive charge and has a relative strong hydrophobic effect, and contains the secondary structure of helix. Especially, DN-AflR contains the zinc finger cluster protein structure (compare with GAL4), which make it could compete with AflS-AflR complex by acting on AflR binding sites to inhibit the activation of aflatoxin. These properties confirmed that DN-AflR had high inhibitory ability on growth of *A. flavus* and biosynthesis of aflatoxin.

To the best of our knowledge, this is the first report about fusing the genes encoding inhibitory peptides and specific zinc finger cluster protein, and fusion expressed the novel anti-*A. flavus* protein. Results showed the modified novel protein reduced aflatoxin biosynthesis and growth of *A. flavus*, which could control aflatoxin contamination. Compared with DN, the inhibitory ability of novel protein has been improved significantly.

This work promoted deep researches for fusion expression of anti-fungal proteins. Furthermore, this study also provided more theoretical basis and technical support for the further biocontrol of *A. flavus* and a new idea about enhancing the anti-*A. flavus* ability of inhibitory substances.

MATERIALS AND METHODS

Materials, strains, and culture conditions. *A. flavus* NRRL3357, preserved in

School of Food Science and Engineering, Ocean University of China, was maintained at 4°C on potato dextrose agar (PDA; Bio-way technology, Shanghai, China). For liquid culture, *A. flavus* was transferred into a 250 ml Erlenmeyer flask containing 100 ml of MM medium at 28°C. *Escherichia coli* DH5α and *E. coli* BL21 (Ruibio Biotech, Beijing, China) were used as hosts for plasmid amplification and genes expression, respectively. *E. coli* was grown in LB medium (10 g l⁻¹ Tryoton, 5 g l⁻¹ Yeast Extract, 10 g l⁻¹ NaCl) containing 50 μl ml⁻¹ ampicillin (Solarbio, Beijing, China) at 37°C with shaking at 180 rpm.

Effects of DN on mycelia morphology of *A. flavus*. (i) Preparation of spores suspension. Spores of *A. flavus* were washed off with sterile distilled water containing 0.1% Tween-80 from PDA medium and filtered with a cotton slag to prepare spores suspension. The number of spores was counted by haemocytometer and diluted to the desired concentration.

(ii) TEM and SEM. 0.8 mg ml⁻¹ DN (synthesized in GL Biochem, Shanghai, China) and 10⁵ spores ml⁻¹ were added in 50 ml MM medium and cultivated at 28°C with 180 rpm for 48 h. Control group didn't contain DN. Then the mycelia were fixed with 2.5% glutaraldehyde, dehydrated in ethanol and embedded in an epoxy resin to be observed by Transmission Electron Microscope (TEM; JEOL, Tokyo, Japan) (21), and the mycelia were also observed with Scanning Electron Microscope (SEM; JEOL, Tokyo, Japan) after dispersion by ultrasonic wave (22).

Expression and identification of fusion protein. (i) Expression vector and its

construction and transformation into *E. coli* BL21. The fusion gene *dn-aflR* containing genes encoding DN and zinc binuclear cluster protein (Cys-Xaa2-Cys-Xaa6-Cys-Xaa6-Cys-Xaa2-Cys-Xaa6-Cys, based on AAM03003.1, NCBI) was synthesized by Hongxun Biotech (Suzhou, China), and amplified with PCR (forward primer 5'-GAATTCGACAACGACAACT-3', reverse primer 5'-AAGCTTACAGGCAAGACCA-3'). Sequences for restriction site EcoR I were sequentially incorporated at the 5'-end and Hind III restriction site was in order added to the 3'-end, and extracted gene by Plasmid Mini Kit (OMEGA, Omega bio-tek, Shanghai, China). Restriction enzymes Hind III and EcoR I were used to digest the *dn-aflR* and pET32a (+), then the digested fragments were separated and identified on 1% agarose gel and recovered from gel using Agarose DNA Extraction Kit (OMEGA, Omega bio-tek, Shanghai, China). The digested fragments were ligated at 4°C with T4 DNA ligase (Thermo Fisher Scientific, New York, USA) (Fig. 1). Then recombinant plasmid *dn-aflR*-pET32a was transformed into *E. coli* DH5α and single colony was picked for colony PCR. The recombinant plasmid extracted from suspension and empty plasmid were both transformed into *E. coli* BL21 by heat shock, and cultivated on solid LB medium containing 50 μg ml⁻¹ ampicillin (23). After 16 h, single colony was inoculated into LB liquid medium and sent it to Ruibio Biotech (Beijing, China) for DNA sequencing.

(ii) Protein expression and analysis. A single colony (containing *dn-aflR*-pET32a) was inoculated into 10 ml LB medium and cultured at 37°C. Then overnight culture was inoculated into fresh 300 ml LB media in 1:5 ratio. When

optical density (OD₆₀₀) reached 0.6, the final concentration of 0.8 mM IPTG was added to induce the expression. After 10 h induction at 16°C under 160 rpm, the cells were washed with buffer (20 mM Tris, 30 mM NaCl, 10 mM imidazole, pH=7.5) and then lysed by sonication (JY92-IIN, Ningbo Scientz Biotech, Ningbo, China; ultrasonic power 300 w, ultrasound work 4 s, stop 6 s, 60 times). Supernatant was collected and identified by SDS-PAGE.

(iii) Protein purification. The presence of a His tag in the recombinant protein meant that the purification was performed with Ni²⁺-NTA affinity chromatography. Briefly, samples were passed through a column and extensively washed with the elution buffer (100 mM Tris base, 500 mM NaCl, and 200 mM Imidazole, pH=8). Purified fusion protein was demonstrated by SDS-PAGE and relative quantity was determined by Bradford protein assay in wave length of 595 nm by standard protein BSA (bovine serum albumin) (Solarbio, Beijing, China).

Effects of DN and DN-AflR on growth of *A. flavus*. (i) Anti-*A. flavus* activity.

Fusion protein was evaluated by agar disk diffusion experiment. The diameter of inhibition zone indicated the anti-*A. flavus* activity. 0.1 ml spores suspension (10⁵ spores ml⁻¹) was coated on PDA plate, and Oxford Cups filled with 100 µl DN (600 µg ml⁻¹) and DN-AflR (600 µg ml⁻¹) were put on it, respectively. The Oxford Cup on control plate was filled with 100 µl sterile water.

(ii) MIC and MFC. A series of solutions containing DN-AflR or DN at the concentrations of 30, 50, 100, 150, 400, 600 and 900 µg ml⁻¹ were prepared by serial dilution. 200 µl of above solutions was added into each well of 96-well plates

containing PDA medium, respectively. Pure PDA without DN-AflR and DN was the control. Then 5 μ l suspension (10^5 spores ml^{-1}) was added to each well and placed at 28°C for 48 h (24). By definition, the minimum concentration of compound that completely inhibited growth of *A. flavus* was the minimum inhibitory concentration (MIC) (non-visible *Aspergillus* hyphae); the culture was continued for 8 days to get the MFC, while the lowest fungicidal concentration (MFC) was the lowest concentration which no spore could germinate at.

Effects of DN and DN-AflR on AFB₁ biosynthesis. DN or DN-AflR (final concentrations of 30, 60 and 90 $\mu\text{g ml}^{-1}$) were added into 30 ml MM medium, respectively, and 100 μ l of suspension (10^5 spores ml^{-1}) was inoculated to each group and incubated at 28°C with 200 rpm. Each experiment was repeated 3 times. After culturing for 48 hours, samples were centrifuged at 8000 g for 20 min at room temperature. Five milliliters of supernatant were diluted with 5 ml ultrapure water. Next, 5 ml of the diluted sample was extracted in immune affinity columns (Huaan Magnech BioTech Co., Ltd., Beijing, China) and then eluted with 1 ml of methanol at a flow rate of 1 drop per second. The eluent was evaporated under a gentle stream of nitrogen at 45°C up to dryness condition, and then derivatized with 200 μ l hexane and 100 μ l trifluoroacetic acid (TFA) for 15 min. After being evaporated to dryness again, the eluent was redissolved in 200 μ l water–acetonitrile (85:15, v/v).

AFB₁ was analyzed according to retention time in HPLC system equipped with a ZORBAX Eclipse XDB-C18 column (4.6 $\times$ 150 mm, 5 μ m, Agilent, Palo Alto, CA, USA) and a 470 fluorescent detector (G1321A, Agilent, USA) (λ_{exc} 360 nm; λ_{em}

440 nm) using a mobile phase solvent of 10% acetonitrile, 40% methanol, and 50% water. The flow rate was 0.8 ml min⁻¹ and injection volume was 20 µl (25).

Statistical analysis. All data were presented as mean ± S.D. One-way analysis of variance (ANOVA) and Duncan's multiple comparison test were used and carried out with SPSS software (SPSS Inc., Chicago, IL, USA). *P*<0.05 was considered statistically significant.

ACKNOWLEDGEMENTS

We thank National Natural Science Foundation of China (31471657) and the research grant from Qingdao Municipal Science and Technology Project (16-6-2-39-nsh) for supporting this research.

REFERENCES

1. Cleveland TE, Yu JJ, Bhatngar D, Chen ZY, Brown RL, Chang KP, Cary GW. 2004. Progress in elucidating the molecular basis of the host plant–*Aspergillus flavus* interaction: a basis for devising strategies to reduce aflatoxin contamination in crops. *J Toxicol-Toxin Rev* 23:345-380.
2. Wu F, Guclu H. 2012. Aflatoxin regulations in a network of global maize trade. *PLoS One* 7:e45151.
3. Atehnkeng J, Donner M, Ojiambo PS, Ikotun B, Augusto J, Cotty PJ, Bandyopadhyay R. 2016. Environmental distribution and genetic diversity of vegetative compatibility groups determine biocontrol strategies to mitigate aflatoxin contamination of maize by *Aspergillus flavus*. *Microb Biotechnol*

329 9:75-88.

330 4. Mishra HN, Das C. 2010. A review on biological control and metabolism of
331 aflatoxin. Crit Rev Food Sci 43:245-264.

332 5. Palumbo JD, Baker JL, Mahoney NE. 2006. Isolation of bacterial antagonists of
333 *Aspergillus flavus* from almonds. Microb Ecol 52:45-52.

334 6. Taylor MC, Jackson CJ, Tattersall DB, French N, Peat TS, Newman J, Briggs LJ,
335 Lapalika GV, Campbell PM, Scott C, Russell RJ, Oakeshott JG. 2010.
336 Identification and characterization of two families of F420 H₂-dependent
337 reductases from Mycobacteria that catalyse aflatoxin degradation. Mol Microbiol
338 78:561-575.

339 7. Lapalika GV, Taylor MC, Warden AC, Scott C, Russell RJ, Oakeshott JG. 2012.
340 F420 H₂-dependent degradation of aflatoxin and other furanocoumarins is
341 widespread throughout the Actinomycetales. PLoS One 7:e30114.

342 8. Zhang T, Shi ZQ, Hu LB, Cheng LG, Wang F. 2008. Antifungal compounds from
343 *Bacillus subtilis* B-FS06 inhibiting the growth of *Aspergillus flavus*. World J
344 Microb Biot 24:783-788.

345 9. Bottone EJ, Peluso RW. 2003. Production by *Bacillus pumilus* (MSH) of an
346 antifungal compound that is active against *Mucoraceae* and *Aspergillus* species:
347 preliminary report. J Med Microbiol 52:69-74.

348 10. Cleveland TE, Yu JJ, Fedorova N, Bhatnagar D, Payne GA, Nierman WC,
349 Bennett JW. 2009. Potential of *Aspergillus flavus* genomics for applications in
350 biotechnology. Trends Biotechnol 27:151-157.

- 351 11. Yu JJ, Chang PK, Ehrlich KC, Cray JW, Bhatnagar D, Cleveland TE, Payne GA,
352 Linz JE, Woloshuk CP, Bennett JW. 2001. Clustered pathway genes in aflatoxin
353 biosynthesis. Appl Environ Microbiol 70:1253–1262.
- 354 12. Chang, PK, Ehrlich KC, Yu JJ, Bhatnagar D, Cleveland TE. 1995. Increased
355 expression of *Aspergillus parasiticus aflR*, encoding a sequence-specific
356 DNA-binding protein, relieves nitrate inhibition of aflatoxin biosynthesis. Appl
357 Environ Microbiol 61: 2372–2377.
- 358 13. Woloshuk CP, Foutz KR, Brewer JF, Bhatnagar D, Cleveland TE, Payne GA.
359 1994. Molecular characterization of *aflR*, a regulatory locus for aflatoxin
360 biosynthesis. Appl Environ Microbiol 60:2408–2414.
- 361 14. Matsushima K, Chang PK, Yu J, Abe K, Bhatnagar D, Cleveland TE. 2001.
362 Pre-termination in *aflR* of *Aspergillus sojae* inhibits aflatoxin biosynthesis. Appl
363 Microbiol Biot 55:585-589.
- 364 15. Burger G, Strauss J, Scazzocchio C, Lang BF. 1991. The pathway specific
365 regulatory gene of nitrate assimilation in *Aspergillus nidulans*, encodes a putative
366 GAL4-type zinc finger protein and contains four introns in highly conserved
367 regions. Mol Cell Biol 11:5746–5755.
- 368 16. Ehrlich KC, Montalbano BG, Cary JW. 1999. Binding of the C6-zinc cluster
369 protein, AFLR, to the promoters of aflatoxin pathway biosynthesis genes in
370 *Aspergillus parasiticus*. Gene 230:249-257.
- 371 17. Marmorstein R, Carey M, Ptashne M, Harrison SC. 1992. DNA recognition by
372 GAL4: structure of a protein-DNA complex. Nature 356:408-414.

- 373 18. Giniger E, Varnum SM, Ptashne M. 1985. Specific DNA binding of GAL4, a
374 positive regulatory protein of yeast. *Cell* 40:767-774.
- 375 19. Todd RB, Andranopoulos A. 1997. Evolution of a fungal regulatory gene family:
376 the Zn(II)₂Cys₆ binuclear cluster DNA binding motif. *Fungal Genet* 21:388-450.
- 377 20. Chang, PK. 2003. The *Aspergillus parasiticus* protein AFLJ interacts with the
378 aflatoxin pathway-specific regulator AFLR. *Molecul Genet Genomics*
379 268:711-719.
- 380 21. Vigneshwaran N, Ashtaputre NM, Varadarajan PV, Nachane RP, Paralikar KM,
381 Balasubramanya RH. 2007. Biological synthesis of silver nanoparticles using the
382 fungus *Aspergillus flavus*. *Mater Lett* 61:1413-1418.
- 383 22. Theis T, Marx F, Salvenmoser W, Stahl V, Meyer V. 2005. New insights into the
384 target site and mode of action of the antifungal protein of *Aspergillus giganteus*.
385 *Res Microbiol* 156:47-56.
- 386 23. Sambrook J, Russell DW. 2001. *Molecular Cloning: A Laboratory Manual*. third
387 ed. Vols. 1, 2 and 3. Cold Spring Harbor Laboratory Press, New York (2100 p.,
388 soft cover. SDS. ISBN-10 Cold Spring Harbor).
- 389 24. Wiegand I, Hilpert K, Hancock RE. 2008. Agar and broth dilution methods to
390 determine the minimal inhibitory concentration (MIC) of antimicrobial
391 substances. *Nat Protoc* 3:163–175.
- 392 25. Zhou GH, Chen YJ, Kong Q, Ma YX, Liu Y. 2017. Detoxification of Aflatoxin
393 B1 by *Zygosaccharomyces rouxii* with Solid State Fermentation in Peanut Meal.
394 *Toxins* 9:42-51.

- 395 26. Kong Q, Chi C, Yu JJ, Shan SH, Li QY, Li QT, Guan B, Nierman WC, Bennett
396 JW. 2014. The inhibitory effect of *Bacillus megaterium* on aflatoxin and
397 cyclopiazonic acid biosynthetic pathway gene expression in *Aspergillus flavus*.
398 Appl Microbiol Biot 98:5161-5172.
- 399 27. Alberts JF, Gelderblom WCA, Botha A, Vanzyl WH. 2009. Degradation of
400 aflatoxin B₁ by fungal laccase enzymes. Int J Food Microbiol 135:47-52.
- 401 28. Costa SJ, Almeida A, Castro A, Domingues L, Besir H. 2013. The novel Fh8 and
402 H fusion partners for soluble protein expression in *Escherichia coli*: a comparison
403 with the traditional gene fusion technology. Appl Microbiol Biot 97:6779-6791.
- 404 29. Douillard FP, O'Connell-Motherway M, Cambillau C, van Sinderen D. 2011.
405 Expanding the molecular toolbox for *Lactococcus lactis*: construction of an
406 inducible thioredoxin gene fusion expression system. Microb Cell Fact 10:66.
- 407 30. Schückel J, Rylott EL, Grogan G, Bruce NC. 2012. A gene-fusion approach to
408 enabling plant cytochromes p450 for biocatalysis. Chembiochem 13:2758-2763.
- 409 31. Terpe K. 2003. Overview of tag protein fusions: from molecular and biochemical
410 fundamentals to commercial systems. Appl Microbiol Biot 60:523–533.
- 411 32. Lilie H, Schwarz E, Rudolph R. 1998. Advances in refolding of proteins
412 produced in *E. coli*. Curr Opin Biotech 9:497-501.
- 413 33. Pal G, Srivastava S. 2014. Cloning and heterologous expression of p1nE, -F, -J
414 and -K genes derived from soil metagenome and purification of active plantaricin
415 peptides. Appl Microbiol Biot 98:1441-1447.
- 416 34. Carvalho AO, Gomes VM. 2009. Plant defensins-prospects for the biological

functions and biotechnological properties. Peptides 30:1007-1020.

35. Saberwal G, Nagaraj R. 1994. Cell-lytic and antibacterial peptides that act by perturbing the barrier function of membranes: facets of their conformational features, structure-function correlations and membrane-perturbing abilities. BBA-Rev Biomembranes 1197:109-131.

TABLE 1 The MICs and MBCs of fusion protein DN-AflR and DN

	Concentrations ($\mu\text{g ml}^{-1}$)							MIC	MBC
	30	50	100	150	400	600	900	($\mu\text{g/ml}$)	($\mu\text{g/ml}$)
DN-AflR	+	+	+	-	-	-	-	150	400
DN	+	+	+	+	+	-	-	600	>900
Negative control	+	+	+	+	+	+	+	/	/

Negative control: no inhibitory protein.

+: observed fungal growth.

-: no fungal growth observed.

/ : no inhibition.

454

455

456

457 **TABLE 2** The basic properties prediction of DN-AflR

Number of amino acids	36
Theoretical pI	8.33
Formula	C ₁₄₆ H ₂₅₀ N ₅₀ O ₅₄ S ₆
Total number of atoms:	506
Aliphatic index	40.83
Total number of negatively charged residues (Asp+Glu)	4
Total number of positively charged residues (Arg+Lys)	6
Grand average of hydropathicity (GRAVY)	-0.581

458 Basic informations were analyzed by ExPASy (<http://web.expasy.org/protparam/>).

459

460

461

462

463

464

465

466

467

468

469 Figure legends:

470 **FIG 1** Molecular modified fragments and characteristics of *E. coli* expression plasmid

471 *dn-aflR*-pet32a. (A) The chemical formula and base sequences of DN. (B) The
472 aflatoxin biosynthetic pathway and clustered genes, AflR binding site (black area) and
473 intergenic region. (C) Construction of expression plasmid. DNA fragment was
474 digested with Hind III and EcoR I, and inserted into downstream of the thioredoxin
475 (Trx) and then ligated with T4 DNA ligase. DN-AflR was expressed as a fusion
476 protein with Trx and 6His. Ar, an ampicillin resistance gene. (D) Base composition of
477 the inserted fusion gene (thick base) containing genes encoding DN (under line) and
478 zinc finger cluster protein (double line). 6His tag, Trx tag and unique restriction sites
479 (over line) were also indicated.

480 **FIG 2** Effects of DN on mycelia morphology of *A. flavus*. (A), (B), (C), (D): TEM;
481 (E), (F), (G), (H): SEM; (A), (B), (E), (F): Control groups; (C), (D), (G), (H): DN
482 treatment groups.

483 **FIG 3** SDS-PAGE analyzed the expression of DN-AflR. (A) M: molecular weight
484 marker (Takara). Lanes 1 and 2: protein bands of *dn-aflR*-pET32a expression. Lane 3:
485 protein band of empty pET32a expression. Lanes 4, 5 and 6: negative control (before
486 induced). The position of the fusion protein was indicated with an arrowhead. (B)
487 Analysis of purified fusion protein DN-AflR. M: marker. Lane 1: expressed
488 supernatant after ultrasonic breaking. Lane 2: precipitate after 8mM urea dissolved.

Lanes 3, 4 and 5: 50 mM imidazole buffer band, 100 mM imidazole buffer band and 300mM imidazole buffer band, respectively. Lanes 6 and 7: 200mM imidazole elution buffer band. The position of the purified protein was indicated with an arrowhead.

FIG 4 Anti- *A. flavus* activity and degradation of AFB₁. (A) Plate a: treated with DN. Plate b: treated with 200 mM imidazole. Plate c: control group with no treatment. Plates d, e and f: treated with DN-AflR, the circle diameters of them were around 24, 19 and 21mm, respectively. (B) The abscissa represented the final concentration of DN (solid line) and DN-AflR (dotted line), the ordinate (content of AFB₁ without any treatment as 100%. The residual rates of the final concentrations of inhibitory proteins in the three experimental groups, they were 30, 60 and 90 ug mL⁻¹, respectively.

FIG 5 Predicted structure of DN-AflR. (A) Secondary structure. Line 1: Transmembrane helix preference (THM index). Line 2: Beta preference (BET index). Line 3: Modified hydrophobic moment index (INDA index). (B) Spatial structure of DN-AflR. Marked position in circle: zinc finger structure. (C) Structure of GAL4. Marked position in circle: zinc finger structure. (D) Zinc finger structure.

511

512

513

514

515

516

517

D

T7 promoter → Lac operator Xba I

TAATACGACTCACTAGGGGAATTGTGAGCGGATAACAATTCCCCTCTAGAAATAATTTTGT

rbs Trx.Tag Msc I

TTAACTTTAAGAAGGAGATATACATATGAGC.....315bp....CTGGCCGGTTCTGGTTCTGGCCAT

His.Tag thrombin

ATGCACCATCATCATCATCATTCTTCTGGTCTGGTGCCACGCGGTTCTGGTAT

S.Tag Nsp V Bgl II Kpn I

GAAAGAAACCGCTGCTAAATTCGAACGCCAGCACATGGACAGCCCAGATCTGGGTACC

Nco I EcoR V BamHI EcoRI

GACGACGACGACAAGGCCATGGCTGATATCGGATCCGAATTCGACAACGACAAC

linker Asp Asn Asp Asn

TCTCCAGGTTCC TGTACGAGTTGTGCCAGCTCAAAAAGTGCATGCACCAAGGAGAAA

Cys Cys Cys

Hind III

CCGGCCTGTGCTCGGTGTATCGAACGTGGTCTTGCCTGTAAGCTTGAGCTCCGTCGACA

Cys Cys Cys

His.Tag

AGCTTGCGGCCGCACTCGAGCACCACCACCACCAC

518 **FIG 1** Molecular modified fragments and characteristics of *E. coli* expression plasmid

dn-aflR-pet32a. (A) The chemical formula and base sequences of DN. (B) The aflatoxin biosynthetic pathway and clustered genes, AflR binding site (black area) and intergenic region. (C) Construction of expression plasmid. DNA fragment was digested with Hind III and EcoR I, and inserted into downstream of the thioredoxin (Trx) and then ligated with T4 DNA ligase. DN-AflR was expressed as a fusion protein with Trx and 6His. Ar, an ampicillin resistance gene. (D) Base composition of inserted fusion gene (thick base) containing genes encoding DN (under line) and zinc finger cluster protein (double line). 6His tag, Trx tag and unique restriction sites (over line) were also indicated.

FIG 2 Effects of DN on mycelia morphology of *A. flavus*. (A), (B), (C), (D): TEM;
(E), (F), (G), (H): SEM; (A), (B), (E), (F): Control groups; (C), (D), (G), (H): DN
treatment.

FIG 3 SDS-PAGE analyzed the expression of DN-AflR. (A) M: molecular weight marker (Takara). Lanes 1 and 2: protein bands of dn-aflR-pET32a expression. Lane 3: protein band of empty pET32a expression. Lanes 4, 5 and 6: negative control (before induced). The position of the fusion protein was indicated with an arrowhead. (B) Analysis of purified fusion protein DN-AflR. M: marker. Lane 1: expressed supernatant after ultrasonic breaking. Lane 2: precipitate after 8 mM urea dissolved. Lanes 3, 4 and 5: 50 mM imidazole buffer band, 100 mM imidazole buffer band and 300 mM imidazole buffer band, respectively. Lanes 6 and 7: 200 mM imidazole elution buffer band. The position of the purified protein was indicated with an arrowhead.

FIG 4 Anti- *A. flavus* activity and degradation of AFB₁. (A) Plate a: treated with DN.

Plate b: treated with 200 mM imidazole. Plate c: control group with no treatment.

Plates d, e and f: treated with purified fusion protein DN-AflR, the circle diameter of

them was around 24, 19 and 21mm, respectively. (B) The abscissa represented the

final concentration of DN (solid line) and DN-AflR (dotted line), the ordinate (content

of AFB₁ without any treatment as 100%. The residual rates of the final concentrations

of inhibitory proteins in the three experimental groups, they were 30, 60 and 90 ug

mL⁻¹, respectively.

FIG 5 Predicted structure of DN-AflR. (A) Secondary structure. Line 1: Transmembrane helix preference (THM index). Line 2: Beta preference (BET index). Line 3: Modified hydrophobic moment index (INDA index). (B) Spatial structure of DN-AflR. Marked position in circle: zinc finger structure. (C) Structure of GAL4. Marked position in circle: zinc finger structure. (D) Zinc finger structure.