Understanding the regulation of STIM1-ORAI1 interaction using a reaction-diffusion model

Barbara Schmidt,^{1,2,3,*} Ivan Bogeski,^{3,4,†} Barbara A. Niemeyer,^{2,‡} and Heiko Rieger^{1,§}

¹*Department of Theoretical Physics, Saarland University, 66041 Saarbrücken, Germany*

²*Department of Molecular Biophysics, Saarland University, 66421 Homburg, Germany*

³*Department of Biophysics, Saarland University, 66421 Homburg, Germany*

⁴*Molecular Physiology, Institute of Cardiovascular Physiology,*

University Medical Center Georg-August-University, 37073 Göttingen, Germany

(Dated: April 23, 2018)

Release of Ca^{2+} from endoplasmatic reticulum (ER) Ca^{2+} stores causes stromal interaction molecules (STIM) in the ER membrane and ORAI proteins in the plasma membrane (PM) to interact and form the Ca^{2+} release activated Ca^{2+} (CRAC) channels, which represent a major Ca^{2+} entry route in non-excitabile cells and thus control various cell functions. Extracellular reactive oxygen species (ROS) can modulate the CRAC current I_{CRAC} via oxidation of ORAI1. We formulate a reaction-diffusion model to quantify the STIM1-ORAI1 interaction during CRAC channel formation and analyze different ORAI1 channel stoichiometries and different ratios of STIM1 and ORAI1 in comparison with experimental data. We incorporate the inhibition of ORAI1 channels by ROS into our model and calculate its contribution to the CRAC channel amplitude. We find that the possibility of reactions between CRAC channel subunits and established CRAC channels tunes the total amount of Ca^{2+} influx and determines which CRAC channel configuration is mostly preferred. High numbers of ROS-inhibited ORAI1 dimers (in comparison to non-inhibited ORAI1 dimers) are needed to induce a strong decrease of Ca^{2+} influx compared to the wildtype (WT) case.

Keywords: ORAI1; STIM1; CRAC channel; reaction-diffusion-model; stoichiometry; ROS

I. INTRODUCTION

Temporally and locally controlled changes of the intracellular Ca^{2+} concentration drive a plethora of cellular functions [1–3]. While electrically excitable cells utilize voltage gated Ca^{2+} channels to achieve rapid changes in intracellular Ca^{2+} concentration [2], immune cells require slower but long-lasting changes in intracellular Ca^{2+} concentration for activation and cytokine production [4, 5]. In these cells activation of T-cell receptors and other PLC coupled receptors results in a depletion of intracellular Ca^{2+} stores (endoplasmatic reticulum, ER). This results in an increased intracellular Ca^{2+} concentration, which by itself is not sufficient to trigger long-term immune cell activation and translocation of the nuclear factor of activated T-cells (NFAT). Thus extracellular Ca^{2+} needs to be ingested into the cell.

For long-lasting Ca^{2+} influx the information about the filling state of the ER has to be conveyed to ion channels in the plasma membrane, which then provide an entry pathway for extracellular Ca^{2+} [2, 6–10]. The drop in ER luminal Ca^{2+} is sensed by stromal interaction molecules (STIM), which undergo a conformational change, multimerize and move to regions near the plasma membrane, so called plasma membrane junctions (PMJ [11, 12]). Here STIM1 proteins trap ORAI1 ion channel

proteins diffusing within the plasma membrane [9, 12–14]. Depending on the STIM1-ORAI1 stoichiometry different ORAI1 conductance states are reached open and selectively conduct Ca^{2+} ions into the cell.

Although the electrophysiological correlate has been known as CRAC (Ca^{2+} release activated Ca^{2+}) current since 1991, the exact protein composition of the complex has been a matter of debate. Several reports pointed towards dimeric ORAI1 channels at rest while tetrameric and hexameric stoichiometries have been proposed to form the ion conduction pore. Indeed recent results obtained from using concatenated constructs as well as the crystal structure of purified *Drosophila* Orail point towards hexameric Orail channels as the predominant species underlying I_{CRAC} [15–17].

While the amplitude of I_{CRAC} is thus determined by the relative ratios of the STIM1 to ORAI1 protein levels [10, 18, 19], it is also modified by external factors. When immune cells enter an area of inflammation, they encounter environments rich in reactive oxygen species (ROS). Exposure to ROS prevents ORAI1 from being activated whereas preassembled STIM1-ORAI1 complexes are insensitive towards inhibition by ROS [20].

In [18] a Monod-Wynman-Chanceux model is used to analyze the dependence of I_{CRAC} on ORAI1 expression levels. In this model ORAI1 exists in two conformational states (open and closed) with four binding sites of STIM1 each, which results in ten channel states in total. Analysis of ORAI1 complexes with less binding sites disagreed with the experimental data. More binding sites have not been tested. With the assumption that STIM1 binds to ORAI1 with negative cooperativity the model provides a reasonable fit to I_{CRAC} . While for an ORAI1 expression

*Electronic address: schmidt@lusi.uni-sb.de

†Electronic address: ivan.bogeski@med.uni-goettingen.de

‡Electronic address: barbara.niemeyer@uks.eu

§Electronic address: h.rieger@mx.uni-saarland.de

level less than two (a. u.) the dominating state is the open ORAI1 conformation with four STIM1 bound OS_4 , for higher ORAI1 expression levels the open states with three and two STIM1 bound dominate and the state OS_4 is nearly not occupied anymore [18].

The stochastic reaction diffusion model of [21] assumes that the ORAI1 proteins are already accumulated into tetramers and that these ORAI1 complexes can trap one to four STIM1 dimers. This leads to four different CRAC channel states with different current capacities contributing to the calculation of I_{CRAC} and to CRAC channel currents reaching steady state values not earlier than about 300 s. Furthermore the numbers of different CRAC channel states in dependence of the cooperativity factors α and β is analyzed. It is found that for all tested configurations (α, β) the best ratio of STIM1 monomers to ORAI1 tetramers is 7.5. In addition the number of active channels depends not only on the total number of ORAI1 proteins but also on the chosen cooperativity. For negative cooperativity and a small number of ORAI1 proteins the tetrameric states are most dominant, for a higher number of ORAI1 proteins the single STIM1 bound states dominate.

In [22] the diffusional behavior of STIM1 and ORAI1 in PMJ regions is addressed. Single-particle tracking and photoactivation experiments are combined with Monte Carlo simulations to analyze STIM1 and ORAI1 diffusion at PMJ regions in resting cells as well as in activated cells. It is found that in resting cells STIM1 proteins follow Brownian motion and ORAI1 motility is subdiffusive. After ER-store depletion ORAI1 and STIM1 movement is mostly restricted to PMJ spots and the data show that both proteins move together as a complex. Furthermore, in activated cells more proteins are immobile than in resting cells.

The junctional regions between ER and plasma membrane are not static [23] but the ER is remodeled by STIM1 proteins. Activated STIM1 proteins are able to form elongated ER cisternae close to the plasma membrane, which increase in number and length during ER depletion [24, 25]. The PM junctions seem to be predetermined since the puncta structures appear repeatedly at the same spots [26].

In [27] an integrated particle system model is combined with stochastic modeling using a spatially heterogeneous Gillespie algorithm to track single particles while CRAC channel formation. The presented model allows to observe the dynamics of single molecules as in single molecule tracking experiments and allows to change rates of action according to the history of the molecules.

Using a similar setup as [21] the diffusion of STIM1 complexes and the four different channels states is analyzed. Therefore the size of the PMJ is enlarged and the channel states do not rest at one single PMJ spot. In the beginning none of the species show sub-diffusive behavior but after 30 s the CRAC channel states move sub-diffusively with decreased diffusion rate, which is a result of restricting these complexes to the small PMJ

spot. Together with this restriction the formation of STIM1-ORAI1 complexes results in differences of STIM1 movement before and after releasing Ca^{2+} from ER.

While previous studies have used a reaction-diffusion system to model I_{CRAC} [18, 21, 27, 28], these models were based on a tetrameric CRAC channel configuration and a single junctional interaction region.

Recent studies analyzing the CRAC channel stoichiometries conclude that activated ORAI1 channels are present as hexamers. Yen et al. expressed hexameric concatemers of human ORAI1 [15]. Their measured currents reproduce the characteristics of CRAC channel current which gives evidences that CRAC channels follow a hexameric stoichiometry. Single-molecule photobleaching experiments lead to the conclusion that ORAI1 forms dimers at resting state and that upon activation a mixture of dimers, tetramers and hexamers contributes to the total CRAC current [16]. The ORAI1 concatemer analysis of [17] finds that concatemers of different sizes (dimer to hexamer) all lead to significant Ca^{2+} influx. Substitution of non-conducting subunits into different places in the concatemers shows, that the conducting capability of the whole hexameric concatemer depends on the position of the substituted subunit. This leads to two different possibilities for the CRAC channels: a pure hexameric arrangement and an assembly as „trimer-of-dimers“.

The ROS-mediated CRAC channel inhibition is analyzed in [29]. Next to FRAP measurements to find differences in the diffusion parameter of inhibited and non-inhibited ORAI1, FRET measurements analyze the interaction strength between STIM1 and ORAI1 and between ORAI1 and ORAI1 for both WT and ROS-preincubated ORAI1. Altered diffusion and reaction behavior together with the findings of an intramolecular locking of the CRAC channels by H_2O_2 could explain the reduced CRAC current of ORAI1 channels under H_2O_2 influence.

The goal of the present study is to transfer these insights of CRAC channel stoichiometries into a new model for CRAC channel formation in order to analyze the different possibilities of ORAI1 channel configurations and their contribution to Ca^{2+} influx into the cell.

Therefore we combine dimeric, tetrameric and hexameric structures of ORAI1 channels and adjust the interaction regions to include as many junctional regions as can be observed in the experiment. Furthermore, we account for WT ORAI1 and ROS-preincubated ORAI1.

The paper is organized as follows: In section II the reaction-diffusion-model is introduced. We distinguish two scenarios: the base case scenario and the H_2O_2 scenario, which deals with a mixture of inhibited and non-inhibited ORAI1 channels. Consequently section III deals with the analytic, numeric and stochastic results. Finally, we discuss (IV) our results.

Figure 1: Schematic representation of CRAC channel formation

Figure 2: Schematic representation of CRAC channel formation under H_2O_2 pressure

II. METHODS

The formation of CRAC channels requires the interaction of two proteins, STIM1 and ORAI1. The STIM1 proteins are located at the ER membrane preassembled in form of dimers [30]. Similarly the ORAI1 proteins located at the plasma membrane are assumed as dimers as smallest unit. As soon as the STIM1 EF-hands, which reach into the ER lumen, sense a decrease in the luminal Ca^{2+} concentration, the STIM1 proteins unfold and start to multimerize. These STIM1 multimers accumulate at regions near the plasma membrane, the plasma membrane junctions (PMJ). Here they attach to the plasma membrane and trap the ORAI1 proteins. Already a channel consisting of four STIM1 and two ORAI1 (i. e. in our model one CRAC channel subunit) is functional although with a low conductance. Also tetrameric or hexameric CRAC channel structures have been observed [16]. These channels consisting of two or three CRAC channel subunits respectively yield an even higher Ca^{2+} influx per channel. Figure 1 schematically represents the different steps of CRAC channel formation. An external factor influencing the Ca^{2+} influx through CRAC channels is the presence of ROS within the extracellular space. ORAI1 proteins exposed to ROS become inactivated, which means they can still form CRAC channels but the conductance of these inhibited channels is small in comparison to non-inhibited channels or even zero (cf. figure 2).

Based on the experimental observation described above

we now formulate our theoretical model to analyze the ORAI1-STIM1 interaction during CRAC channel formation and the resulting CRAC current. In the following we assume that the ER store is already depleted, which is experimentally realized by adding $1 \mu M$ of the SERCA inhibitor thapsigargin. Then we describe CRAC channel formation by the following four steps:

1. Multimerization of the STIM1 proteins
2. Diffusion of STIM1 and ORAI1 proteins to the PMJ
3. Attachment of STIM1 proteins to a PMJ
4. Binding of STIM1 and ORAI1 proteins at the PMJ

The multimerization of the STIM1 proteins is given by:

$$S_{dimer} + S_{dimer} \xrightleftharpoons[k_{demulti}]{k_{multi}} S, \quad (1)$$

where S_{dimer} represents a STIM1 dimer (resting state configuration of STIM1) and S denotes a STIM1 multimer (tetramer), which is able to form a CRAC channel subunit when combined with an ORAI1 dimer. The rates k_{poly} and k_{depoly} define the rates of multimerization and demultimerization respectively. Since after ER store depletion a strong multimerization of STIM1 dimers is observed, one has $k_{multi} \gg k_{demulti}$. The stationary solution is given by:

$$S = \frac{1}{2} \frac{k_{multi}}{k_{demulti}} S_{dimer}^2. \quad (2)$$

The ORAI1 dimers diffuse within the cell membrane, the STIM1 dimers and tetramers diffuse within the ER plasma membrane. The diffusion rate within the PMJ is assumed to be smaller than outside the plasma membrane junction [22]. The different channel states ($|O\rangle$, $|OO\rangle$ and $|OOO\rangle$) and the STIM1 tetramers connected to the PMJ (S_{rest}) are assumed to rest at the PMJ. Furthermore the reactions

$$S + PMJ \xrightleftharpoons[k_{detach}]{k_{attach}} S_{rest} \quad (3)$$

describe the process of STIM1 tetramers attaching to the PMJ, which means these STIM1 tetramers S_{rest} are fixed at the PMJ and are able to trap an ORAI1 dimer nearby.

The core reaction scheme of CRAC channel formation is given by the following reactions, where S_{rest} denotes the concentration of STIM1 tetramers fixed at the PMJ, O denotes the concentration of ORAI1 dimers within the PMJ and Z_1, Z_2, Z_3 denote the concentration of the three different CRAC channel states:

$$S_{rest} + O \xrightleftharpoons[k_2]{k_1} Z_1 \quad (4)$$

$$Z_1 + Z_1 \xrightleftharpoons[k_4]{k_3} Z_2 \quad (5)$$

$$Z_2 + Z_1 \xrightleftharpoons[k_6]{k_5} Z_3 \quad (6)$$

Figure 3: Schematic representation of the reaction scheme

Resting STIM1 tetramers (S_{rest}) form together with free ORAI1 dimers (O) CRAC channel subunits (Z_1). Two of these subunits form an intermediate state (Z_2). The fully open CRAC channel (Z_3) is built out of an intermediate state and another subunit.

Since CRAC channel formation and disassembly seems to be highly dynamic ([18] and [29]) we analyze the influence of an additional reaction between a hexamer CRAC channel state Z_3 and a CRAC channel subunit Z_1 :

This reaction „destroys“ the fully open CRAC channel configuration and builds two intermediate CRAC channel states with lower Ca^{2+} conductivity. Therefore we will call this reaction „stealing mechanism“.

Figure 3 shows a schematic representation of the reaction scheme. The yellow boxes indicate reactions only allowed at the PMJ.

The reaction rates k_1 , k_3 and k_5 (on-rates) as well as the rates k_2 , k_4 and k_6 (off-rates) are not chosen to be independent but are connected through a cooperativity parameter (α respectively β) to account for the findings of [18]. In analogy to [21] the dependency of the rates is assumed to be:

$$\begin{aligned} k_{1+2n} &= \alpha^{2n} \cdot k_1, \quad n \in (0, 1, 2) \\ k_{2n} &= \beta^{2n} \cdot k_2, \quad n \in (0, 1, 2). \end{aligned} \quad (8)$$

As default we examine negative cooperativity ($0 < \alpha, \beta < 1$) since we assume, that it gets harder to bind more and more proteins to an already established STIM1-ORAI1 complex. Nevertheless within the numerical analysis we also examine positive cooperativity ($\alpha, \beta > 1$). Assuming that the reaction strength between a single CRAC channel subunit and a fully open CRAC channel is as strong as the reaction strength between a CRAC channel subunit and an intermediate state, we take for the rate for the stealing mechanism:

$$k_7 = k_5 = \alpha^2 \cdot k_1. \quad (9)$$

Because STIM1 multimerization, STIM1 attachment to the PM and STIM1-ORAI1 binding are more preferred than the respective backward reactions, we assume higher

parameter	value
k_{multi}	$4.8 \cdot 10^5 \frac{l}{mol \cdot s}$
k_{multi}	$0.01 \frac{l}{mol \cdot s}$
k_{attach}	$1.8 \cdot 10^6 \frac{l}{mol \cdot s}$
$k_{detacht}$	$0.3 \frac{l}{mol \cdot s}$
k_1	$1.2 \cdot 10^6 \frac{l}{mol \cdot s}$
k_2	$1 \frac{l}{mol \cdot s}$
k_3	$\alpha \cdot k_1$
k_4	$\beta \cdot k_2$
k_5	$\alpha^2 \cdot k_1$
k_6	$\beta^2 \cdot k_2$
k_7	$\alpha^2 \cdot k_1$
α	0.25
β	0.25

Table I: Reaction rate parameters for the reactions of the basic reaction scheme

rate constants for the forward reactions. Similar to [21] we assume that STIM1 multimerizing is slower than STIM1-ORAI1 binding. The process of attaching to the membrane is assumed to be very fast, since just one (instead of two) protein complex is involved. Furthermore, we assume strong negative cooperativity. With the default numerical values for the reaction rate parameters as given in table I the model provides a reasonable time dependence of I_{CRAC} .

The total Ca^{2+} influx into the cell, I_{CRAC} , is determined by the number of channels N_{Z_i} weighted with their respective conductivity c_i :

$$I_{CRAC} = c_1 \cdot N_{Z_1} + c_2 \cdot N_{Z_2} + c_3 \cdot N_{Z_3} \quad (10)$$

with $c_1 = 0.01$, $c_2 = 0.25$ and $c_3 = 1.0$. The number of channels can be calculated via the formula:

$$N_{Z_i} = Z_i \cdot A_{cell} \cdot p_{PMJ}, \quad (11)$$

where $A_{cell} \approx 2 \cdot 10^3 \mu m^2$ denotes the size of the cell surface and $p_{PMJ} \approx 15\%$ describes the fraction of the cell surface where ORAI1 proteins accumulate, the PMJ regions.

In the following we will analyze the reaction-diffusion model as follows: First, the core reactions without and with stealing mechanism at the PMJ will be analyzed analytically and numerically with special emphasis on the steady state values for the channels states and for I_{CRAC} . In a second step we will combine the reaction and diffusion part of the model into a stochastic reaction-diffusion model. Within this model the results of TIRF measurements are used to describe and quantify the changes in PMJ clusters.

To account for diffusion in the stochastic model the model area is discretized in a grid of 100×100 grid cells called subvolumes each representing an area of $\Delta l \times \Delta l = 0.1 \mu m \times 0.1 \mu m$. The diffusion of the proteins is treated as reactions with reaction rates $d = D/(\Delta l)^2$ assuming periodic boundary conditions. Numerical values for the diffusion constants are given in table II.

diffusion rate	value
D_O	$0.07 \mu\text{m}^2/\text{s}$ [13]
D_S	$0.1 \mu\text{m}^2/\text{s}$ [12]
D_{S_2}	$0.05 \mu\text{m}^2/\text{s}$ [12]
D_O^{PMJ}	$0.03 \mu\text{m}^2/\text{s}$ [22]
D_S^{PMJ}	$0.03 \mu\text{m}^2/\text{s}$ [22]
$D_{S_2}^{PMJ}$	$0.015 \mu\text{m}^2/\text{s}$ (*)

Table II: Diffusion constants for STIM1 and ORAI1 outside and within the PMJ, (*) assuming that as outside the PMJ the tetramers are half as fast as the dimers

The two-dimensional model area represents the cell membrane and the plasma membrane simultaneously. The interaction of ORAI1 and STIM1 is only possible at regions, where the ER membrane and the plasma membrane are close together, the plasma membrane junctions PMJ. Therefore, some of the subvolumes of our model area are labeled PMJ. Reaction (1) and diffusion is possible in all subvolumes, reactions (3) to (7) are just allowed at the PMJ.

To determine how many subvolumes must be PMJ we analyzed the ORAI1 cluster in HEK293 cells (i. e. HEK293 stably expressing STIM1 cells) via TIRF microscopy (cf. figure 4). After addition of the SERCA inhibitor thapsigargin ($1 \mu\text{M}$) leading to passive ER store depletion, the GFP-tagged ORAI1 proteins diffused within the cell membrane to the PMJs. Figure 4A shows a snapshot of two cells immediately after store depletion, figure 4C the same cells 300 s later. Analyzing in total 19 cells via TIRF microscopy and with the help of imageJ regarding number and mean size of ORAI1 clusters we find the following (cf. figure 5): Within the first 50 seconds the number of clusters rises to about 20 per cell and simultaneously the average size of the clusters increases rapidly from $0.3 \mu\text{m}^2$ to about $0.55 \mu\text{m}^2$. After this rapid rise the number of clusters doubles within the next 200 seconds until it reaches a value between 40 and 45. At the same time the average size of clusters drops slightly but then rises again to about $0.6 \mu\text{m}^2$.

In our analysis we increase the number of PMJ subvolumes from one to six within the first 50 seconds and from five to eleven within the next 150 seconds. Simultaneously we increase their size from $0.3 \mu\text{m}^2$ to $0.6 \mu\text{m}^2$ within the first 200 seconds in order to analyze the influence of the PMJ formation.

Analogously we analyzed cells preincubated with ROS (figures 4B and D). Since size and amount of ORAI1 cluster were similar between the two analysis, the same settings in the model are used.

To analyze the reaction-diffusion system described above we perform computer simulations using a Gillespie Monte Carlo algorithm [31] with an efficient implementation technique [32] optimizing the computation time. At the beginning of the simulation ($t = 0 \text{ s}$) all proteins are homogeneously distributed and the ER Ca^{2+} store is

Figure 4: TIRF images of HEK293 cells with GFP-tagged ORAI1, A: WT at $t = 0 \text{ s}$, B: ROS preincubated cells at $t = 0 \text{ s}$, C: WT at $t = 300 \text{ s}$, D: ROS preincubated cells at $t = 300 \text{ s}$

Figure 5: Cluster analysis of TIRF images, A: average number of ORAI1 clusters per cell over time, B: time dependence of the average size of ORAI1 clusters

assumed to be empty.

H_2O_2 scenario

We analyze two different scenarios: The base case scenario described above and the H_2O_2 scenario to analyze the influence of ROS. To account for the H_2O_2 preincubated cells we modify the reaction-diffusion scheme as follows:

- Additional to the CRAC channel states Z_1 , Z_2 and Z_3 , denoted for better readability as $|O\rangle$, $|OO\rangle$ and $|OOO\rangle$ respectively, composed of STIM1 and ORAI1 proteins, new channel states consisting of STIM1 and preincubated ORAI1 (O^H) or of STIM1 and a mixture of O and O^H were included: $|O^H\rangle$, $|OO^H\rangle$, $|O^H O^H\rangle$, $|OOO^H\rangle$, $|OO^H O^H\rangle$, $|O^H O^H O^H\rangle$.
- New reactions were introduced by exchanging the states O , $|O\rangle$, $|OO\rangle$ and $|OOO\rangle$ of reactions (4) to (7) by the new states in every possible combination.

- To model I_{CRAC} a new parameter δ was introduced quantifying the degree of inhibition. We assume that the conductance of CRAC channel states consisting of total ROS-inhibited ORAI1 is given by the fraction δ of the respective CRAC channel state of WT ORAI1. To model the conductance of mixed CRAC channel states (consisting of WT and ROS-inhibited ORAI1), we assume that each included WT dimer contributes to the conductance as before and each included ROS-inhibited dimer contributes with fraction δ of the WT case. This means that the intermediate state $|OO^H\rangle$ is weighted with the factor $(1 + \delta)/2$ and the mixed hexamer CRAC channel states $|OOO^H\rangle$ and $|OO^H O^H\rangle$ are weighted with the factors $(2 + \delta)/3$ and $(1 + 2\delta)/3$ respectively:

$$I_{CRAC}^{mix} = c_1 \cdot N_{Z_1} + c_2 \cdot N_{Z_2} + c_3 \cdot N_{Z_3} \\ + \delta (c_1 \cdot N_{|O^H\rangle} + c_2 \cdot N_{|O^H O^H\rangle} + c_3 \cdot N_{|O^H O^H O^H\rangle}) \\ + \frac{c_2}{2} (1 + \delta) \cdot N_{|OO^H\rangle} \\ + \frac{c_3}{3} (2 + \delta) \cdot N_{|OOO^H\rangle} + \frac{c_3}{3} (1 + 2\delta) \cdot N_{|OO^H O^H\rangle} \quad (12)$$

In [29] the interactions between STIM1 and ORAI1 proteins and between ORAI1 and ORAI1 proteins were analyzed. Foerster Resonance Energy Transfer (FRET) values of STIM1 proteins with ORAI1 proteins inhibited by H_2O_2 were significantly larger (0.24 ± 0.03) than FRET values of STIM1 proteins with WT ORAI1 (0.15 ± 0.01). In contrast, the ORAI1-ORAI1 subunit interaction was reduced by 46%. Fluorescence recovery after photobleaching (FRAP) measurements revealed a 1.7 times smaller rate of recovery for preincubated ORAI1 proteins [29]. Therefore the diffusion constant of O^H and the reaction rates $\tilde{k}_1$ and $\tilde{k}_2$ as well as the cooperativity parameters $\tilde{\alpha}$ and $\tilde{\beta}$ of the new reactions are defined as:

$$D_{O^H} = D_O/1.7 \quad (13) \\ \tilde{k}_i = k_i \cdot 1.6 \quad \text{for } i \in (1, 2) \\ \tilde{\alpha} = \alpha \cdot 0.46 \\ \tilde{\beta} = \beta \cdot 0.46.$$

III. RESULTS

A. Analytical and numerical analysis of the Base Case Scenario

Within the analysis of the base case scenario, we examine two reaction systems. The first reaction system consists of the core reactions only, the second reaction system includes the additional stealing mechanism. The core reaction system reads:

$$\begin{aligned} \frac{dS}{dt} &= -k_1 \cdot S \cdot O + k_2 \cdot Z_1 \\ \frac{dO}{dt} &= -k_1 \cdot S \cdot O + k_2 \cdot Z_1 \\ \frac{dZ_1}{dt} &= k_1 \cdot S \cdot O - k_2 \cdot Z_1 - k_3 \cdot \frac{Z_1^2}{2} \cdot 2 + 2 \cdot k_4 \cdot Z_2 \\ &\quad - k_5 \cdot Z_2 \cdot Z_1 + k_6 \cdot Z_3 \\ \frac{dZ_2}{dt} &= k_3 \cdot \frac{Z_1^2}{2} - k_4 \cdot Z_2 - k_5 \cdot Z_2 \cdot Z_1 + k_6 \cdot Z_3 \\ \frac{dZ_3}{dt} &= k_5 \cdot Z_2 \cdot Z_1 - k_6 \cdot Z_3 \end{aligned} \quad (14)$$

and when including the stealing mechanism, the right hand sides of (14) are extended by:

$$\begin{aligned} \frac{dZ_{1,3}}{dt} &\rightarrow \frac{dZ_{1,3}}{dt} - k_7 \cdot Z_1 \cdot Z_3 \\ \frac{dZ_2}{dt} &\rightarrow \frac{dZ_2}{dt} + 2 \cdot k_7 \cdot Z_1 \cdot Z_3. \end{aligned} \quad (15)$$

The total amounts of ORAI1 (in whichever CRAC channel configuration) and of STIM1 respectively have to be conserved, which means:

$$O^{tot} = O(t) + Z_1(t) + 2 \cdot Z_2(t) + 3 \cdot Z_3(t), \quad (16)$$

$$S^{tot} = S(t) + Z_1(t) + 2 \cdot Z_2(t) + 3 \cdot Z_3(t). \quad (17)$$

The second condition is always fulfilled when (16) is fulfilled, since the right hand sides of the differential equations for O and S are identical and we consider the following initial conditions:

$$\begin{aligned} S(0) &= S^{tot} \\ O(0) &= O^{tot} \\ Z_1(0) &= 0 \\ Z_2(0) &= 0 \\ Z_3(0) &= 0. \end{aligned} \quad (18)$$

Considering the reaction system without stealing (eqns (14)), the second equation of (14) can be rewritten as:

$$\frac{dZ_1}{dt} = - \left(\frac{dO}{dt} + 2 \cdot \frac{dZ_2}{dt} + 3 \cdot \frac{dZ_3}{dt} \right), \quad (19)$$

which results in

$$Z_1(t) = - (O(t) + 2 \cdot Z_2(t) + 3 \cdot Z_3(t) + c_0), \quad (20)$$

where c_0 is determined by the initial condition (18), which yields $c_0 = O^{tot}$. This is identical to the additional condition above (16).

The stationary state of the reaction system (14) obeys:

$$\begin{aligned} 0 &= -k_1 \cdot S^{stat} \cdot O^{stat} + k_2 \cdot Z_1^{stat} \\ 0 &= k_3 \cdot \frac{Z_1^{stat2}}{2} - k_4 \cdot Z_2^{stat} - k_5 \cdot Z_2^{stat} \cdot Z_1^{stat} \\ &\quad + k_6 \cdot Z_3^{stat} \\ 0 &= k_5 \cdot Z_2^{stat} \cdot Z_1^{stat} - k_6 \cdot Z_3^{stat} . \end{aligned} \quad (21)$$

These are three independent equations with five concentration parameters (S^{stat} , O^{stat} , Z_1^{stat} , Z_2^{stat} , Z_3^{stat} ,) and six reaction rates k_i .

With the cooperativity parameters α and β for the on- and off-rates and using the abbreviations

$$\begin{aligned} \gamma &= \frac{\alpha}{\beta} \\ k &= \frac{k_1}{k_2} , \end{aligned} \quad (22)$$

the equation system can be rewritten in terms of the stationary number of free ORAI1 and free STIM1 proteins:

$$\begin{aligned} Z_1^{stat} &= k \cdot S^{stat} \cdot O^{stat} \\ Z_2^{stat} &= 0.5 \cdot \gamma \cdot k \cdot S^{stat2} \cdot O^{stat2} \\ Z_3^{stat} &= 0.5 \cdot \gamma^3 \cdot k^2 \cdot S^{stat3} \cdot O^{stat3} . \end{aligned} \quad (23)$$

Together with the conservation laws (16) and (17) one gets a polynomial equation for Z_1^{stat} , which can be solved numerically. All other stationary concentrations then follow.

Similarly we analyzed the reaction system with stealing mechanism (eqns (15)): The additional stealing reaction (7) should help to regulate the formation of CRAC channels. The corresponding stationary equation system together with the two conservation laws (16) and (17) can be simplified to the following non-linear, non-homogeneous equation system:

$$\begin{aligned} O^{ges} &= O^{stat} + Z_1^{stat} + 2 \cdot Z_2^{stat} + 3 \cdot Z_3^{stat} \\ S^{ges} &= S^{stat} + Z_1^{stat} + 2 \cdot Z_2^{stat} + 3 \cdot Z_3^{stat} \\ Z_1^{stat} &= \frac{k_1}{k_2} S^{stat} \cdot O^{stat} \\ Z_2^{stat} &= \frac{k_3}{2k_4} (Z_1^{stat})^2 + \frac{k_7}{k_4} Z_1^{stat} \cdot Z_3^{stat} \\ Z_3^{stat} &= \frac{k_5 \cdot Z_1^{stat} \cdot Z_2^{stat}}{k_6 + k_7 Z_1^{stat}} , \end{aligned} \quad (24)$$

which is consistent with the basic solution for $k_7 = 0$.

Time evolution of the CRAC channel states

The reaction systems (14) and (15) can be solved numerically. In both reaction systems the stationary value of the CRAC channels states is reached after about two seconds. In experiments ([29]) it is observed that the

Figure 6: Time evolution of CRAC channel states Z_1 (blue), Z_2 (red) and Z_3 (green) according to the core reaction scheme (eq (14), solid lines) and the reaction system with stealing mechanism (eq (15), dashed lines) for the rate parameters as given in table I and for $O^{tot} = 120000$ and $S^{tot} = 240000$ for $t \in [0 \text{ s}, 5 \text{ s}]$.

delay between taspigargin addition and the steady state Ca^{2+} influx is larger than two seconds, which means that the delay is also influenced by the diffusion of STIM1 and ORAI1 proteins towards the PMJ. Figure 6 exemplifies the time evolution of all three CRAC channel states normalized to the total amount of ORAI1 (O^{tot}) within the first five seconds, with the parameters given in table I. The solid lines represent the core reaction system (14) and the dashed lines represent the system with stealing mechanism (15). In both cases the single subunit state Z_1 is rarely occupied (less than $2 \cdot 10^{-4}$), as on-reactions are more preferred than off-reactions, while Z_2 and Z_3 show more interesting features:

The time course of Z_2 and Z_3 is determined by k_1 and k_2 . Analyzing the base case scenario without stealing mechanism shows the following: The larger k_1 the steeper the rise of Z_2 within the first milliseconds, the larger k_2 the faster the following drop of Z_2 . Similarly the rise of Z_3 is determined by the magnitude of k_1 . Large k_2 yields a second (slower) rise of Z_3 within the next five seconds. Furthermore k_2 determines whether Z_2 or Z_3 dominates after these five seconds, i. e. the larger k_2 the more hexameric CRAC channel states are established and the less tetrameric states are found. Analysis of the base case scenario with stealing mechanism shows similar dependence of the time evolution of Z_2 and Z_3 . The main differences occur in the absolute values of these two states and that even for large k_2 the hexameric state is less occupied than the tetrameric state.

Influence of the rate parameters

In a second step we analysed the influence of the rate parameters (k_1 , k_2 , α and β) on the steady state values. Keeping all but the parameters displayed on the x-axis fixed as given in table I, figures 7 and 8 show the results

Figure 7: Graphic representation of the values of the steady state configuration (i. e. at $t = 300$ s) Z_1^{stat} (blue), Z_2^{stat} (red) and Z_3^{stat} (green) in dependence of the rate parameters k_1 (A) and k_2 (B) for the core reaction system (eq (14), solid lines) and with stealing mechanism (eq (15), dashed lines). If not varied as displayed on the x axis (log-scale!), the rate parameters are chosen as given in table I.

for varying the respective parameters. Solid lines indicate the core reaction system, while dashed lines indicate the inclusion of the stealing reaction.

Figure 7A shows the influence of the on-rate k_1 on the stationary solution of the reaction schemes. In both cases the behavior of Z_1 is similar: Only in the case when the on-rate is small (between 10^{-2} l/(mol · s) and 10^2 l/(mol · s)) a lot of dimeric CRAC channels states can be found. If the on-rate is even less than 10^{-3} l/(mol · s) no channels at all are formed. The behavior of intermediate CRAC channel states on changes of k_1 differs without (14) or with (15) stealing mechanism. In the second case a larger on-rate yields to more tetrameric CRAC channel states up to the value 0.22. In the first case just for on-rates between 10^{-1} l/(mol · s) to 10^5 l/(mol · s) a non negligible stationary value of Z_2 is reached. As already expected from the data of the time evolution analysis, the most dominant state of the core reaction system is the hexamer configuration for large on-rates. In case of core reactions with stealing mechanism the stationary value of Z_3 is smaller than the one of Z_2 for large on-rates, which means, that more tetramers than hexamers are formed.

Keeping the on-rate fixed and varying the off-rate (figure 7B) results always in no noticeable values of Z_1 . This is obvious from the data before, because in the examined interval the on-reactions are always more preferred than the off-reactions. In the core reaction scenario the tetramer configuration is more often adopted as the hexamer configuration as long as the off-rate is smaller than $10^{-3} \frac{1}{s}$ and then drops to almost zero for larger off-rates. In contrast to this the hexamer configuration starts at a value of 0.177 for small off-rates, rises to a value of 0.344 at $k_2 = 0.01$ and then slightly drops again to 0.289 for even larger off-rates. In the reaction system with stealing mechanism the tetramer configuration is always favored, while the gap between tetramer and hexamer configuration scales down from 0.2 for off-rates between 10^{-9} to 10^{-3} to 0.04 for larger off-rates.

Figure 8A shows the dependence of the CRAC channel states on the on-rate cooperativity parameter α . Only for small α (smaller than 0.05) compared to the off-rate

Figure 8: Graphic representation of the values of the steady state configuration Z_1^{stat} (blue, right y-axis), Z_2^{stat} (red) and Z_3^{stat} (green) in dependence of α and β for the core reaction system (eq (14), solid lines) and with stealing mechanism (eq (15), dashed lines). If not varied as displayed on the x axis, the rate parameters are chosen as given in table I.

Figure 9: Graphic representation of the values of the steady state configuration Z_1^{stat} (blue, right y-axis), Z_2^{stat} (red) and Z_3^{stat} (green) in dependence of the ratio STIM1 / ORAI1. Solid lines indicate the core reaction scheme (eq (14)), dashed lines show the corresponding values for the reaction scheme including the stealing mechanism (eq (15)). Rate parameters are chosen as given in table I.

cooperativity parameter β (0.25) the steady state values of Z_2 and Z_3 develop from zero to their respective constant value as in the analysis above. Z_1 drops from 1 ($\alpha = 0$) to 10^{-3} ($\alpha = 0.1$) and then to less than 10^{-5} for α rising until 0.2.

In contrast to this Z_i change more significantly on changes of the off-rate cooperativity parameter β , when α is fixed at 0.25 (cf. figure 8B). For the core reaction system Z_1 rises linear from zero to $2.5 \cdot 10^{-4}$ for β between zero and two. Similarly Z_2 rises from zero to 0.29 and Z_3 drops from 0.2 to 0.14 for β between 0.1 and two.

Dependency of I_{CRAC} of the ratio S^{ges}/O^{ges}

Major parameters influencing I_{CRAC} are the amount and ratio of STIM1 and ORAI1 proteins. Figure 9 shows

Figure 10: Graphic representation of the values of the steady state configuration Z_1^{stat} (blue, right y-axis), Z_2^{stat} (red) and Z_3^{stat} (green) in dependence of the ratio STIM1 / ORAI1 with additional increase of β (unbinding cooperativity parameter). Solid lines indicate the core reaction scheme (eq (14)), dashed lines show the corresponding values for the reaction scheme including the stealing mechanism (eq (15)). Constant parameters are chosen as s given in table I and β is varied linear in the interval [0.25, 6.25].

the results for the different stationary CRAC channel states normalized to O^{tot} . As soon as there are twice or more as many STIM1 dimers than ORAI1 dimers the stationary values of Z_i are constant. For ratios of STIM1 to ORAI1 below two we see a linear rise of the steady state values $Z_{2,3}$ from zero to their respective value at STIM1 / ORAI1 = 2 in both cases. Z_1 of the core reaction scheme first shows a steep linear and then slightly slower rise to its final value.

In comparison, figure 10 shows the stationary values of Z_i and I_{CRAC} normalized to O^{tot} in dependence of the ratio S^{tot}/O^{tot} with simultaneous rise of β (linear from 0.25 to 6.25). This scenario corresponds to the assumption that it gets more difficult for CRAC channel units to build up larger complexes as more and more STIM1 proteins arrive at the PMJ. In this case all Z_i first rise between $S^{tot}/O^{tot} = 0$ and $S^{tot}/O^{tot} = 2$ ($0.25 \leq \beta \leq 0.55$). Subsequently, Z_1 and Z_2 rise more slowly whereas Z_3 drops for larger β and larger ratios S^{tot}/O^{tot} . This results in both reaction systems (without stealing (14), solid line, and with stealing (15), dashed line) in a slight decrease of I_{CRAC} . A similar behavior is seen in [28].

B. Stochastic analysis of the Base Case Scenario

Analysis of I_{CRAC} in dependence of time and of the ratio STIM1 / ORAI1

Main aspects of the analysis of the base case scenario are the dependence of I_{CRAC} of time t and of the ratio of

Figure 11: Stochastic analysis of the temporal course of I_{CRAC} for different ratios of STIM1 / ORAI1. Solid lines indicate the core reaction scheme (eq (14)), dashed lines show the corresponding values for the reaction scheme including the stealing mechanism (eq (15)).

STIM1 proteins to ORAI1 proteins. Figure 11 shows the development of I_{CRAC} normalized to the total available amount of free ORAI1 dimers within the first 20 seconds for three different ratios STIM1 / ORAI1 for the core reaction scheme without (solid lines) and with (dashed lines) stealing mechanism. In contrast to the numerical analysis above the respective steady state value of I_{CRAC} is reached after about eight to ten seconds not already after about two seconds. This is because diffusion of ORAI1 and STIM1 proteins are now taken into account. For all three ratios of STIM1 / ORAI1 (STIM1 / ORAI1 = 1, 2 and 3 in black, blue and red respectively) the core reaction scheme without stealing mechanism leads to a higher amount of Ca^{2+} influx compared to the reaction scheme with stealing mechanism. In both reaction schemes, Ca^{2+} influx is highest for STIM1 / ORAI1 = 3. It is slightly lower for STIM1 / ORAI1 = 2 and significantly lower for STIM1 / ORAI1 = 1.

Figure 12 shows the steady state values of Z_i^{stat} and I_{CRAC}^{stat} normalized to O^{tot} in dependence of the ratio STIM1 / ORAI1. For both reaction schemes without and with stealing the numbers of tetramer and hexamer CRAC channel states rise linear from STIM1 / ORAI1 = 0 to STIM1 / ORAI1 $\simeq 2.5$ and then stay at a constant value except for statistical fluctuations. The number of CRAC channel subunits Z_1 is low in comparison to the number of higher CRAC channel states.

As in the numerical analysis the hexamer states are the most dominant for the base case scenario without stealing mechanism and the tetramer states are the most dominant for the base case scenario with stealing mechanism. This results in lower I_{CRAC} in the latter scenario.

In contrast to the numerical analysis above, the highest steady state value of I_{CRAC}^{stat} is reached at STIM1 / ORAI1 $\gtrsim 2.5$ and not at exactly STIM1 / ORAI1 = 2. The latter ratio is the minimum number of STIM1 tetramers needed to saturate all free ORAI1 dimers available and therefore is the theoretical minimal ra-

Figure 12: Stochastic analysis of the steady state values (i. e. at $t = 300$) of CRAC channel states Z_1^{stat} , Z_2^{stat} , Z_3^{stat} (blue, red, green respectively) and of I_{CRAC}^{stat} (black) in dependence of the ratio STIM1 / ORAI1. Solid lines indicate the core reaction scheme (eq (14)), dashed lines show the corresponding values for the reaction scheme including the stealing mechanism (eq (15)).

ratio at which the highest value of I_{CRAC} is reached. The stochastic analysis includes diffusion of STIM1 and ORAI1. In this case there is more than the minimum amount of STIM1 needed to saturate all ORAI1 dimers, because it is possible, that free STIM1 tetramers and free ORAI1 dimers are not at the same spot and therefore can not react with each other. Are there about 25% more STIM1 proteins available than the minimal theoretical amount needed the maximal Ca^{2+} influx is reached.

In the following we perform all analysis with the reactions of the core system with stealing mechanism.

Influence of the diffusion constant of ORAI1

Comparing WT ORAI1 channels with ROS preincubated ORAI1 channels one experimentally measured parameter concerns the diffusion constant of ORAI1. WT ORAI1 proteins are faster than ROS preincubated ORAI1 channels [29]. To analyze, whether this could influence the total Ca^{2+} influx into the cell, we altered the diffusion rate of the ORAI1 dimers outside and inside the PMJ areas in our model.

Figure 13 shows the amplitude of I_{CRAC}^{stat} as well as the number of active channels at steady state (i. e. at $t = 300$ s) normalized to O^{tot} in dependence of the diffusion constant of the ORAI1 dimers. The diffusion constant D_O outside the PMJ regions was varied between 0 and $0.5 \text{ mm}^2/\text{s}$. To account for the slower diffusion of ORAI1 within the PMJ, D_O^{PMJ} was set to $3 \cdot D_O/7$ in analogy to the difference in the measured values of [13] and [22] (compare table II). Apart from statistical fluctuations the number of active channels in the steady state and hence the value of I_{CRAC}^{stat} do not vary for different values of D_O . Consequently, the lowered diffusion constant alone can not explain the drastic decrease of Ca^{2+}

Figure 13: Stochastic analysis of the steady state values (i. e. at $t = 300$) of CRAC channel states Z_1^{stat} , Z_2^{stat} , Z_3^{stat} (blue, red, green respectively) and of I_{CRAC}^{stat} (black) in dependence of the diffusion constant D_O of the ORAI1 dimers according to the reaction scheme including the stealing mechanism (eq (15)).

Figure 14: Temporal course of I_{CRAC} according to eq (15). Faded lines indicate the parameters as given in tables I and II, bold lines indicate the modified parameters according to eq. (13) and $\bar{D}_O = D_O/1.7$

influx seen in experiments ([29]) between the H_2O_2 scenario and the WT scenario.

Influence of altered reaction rates

In a next step in [29] the interactions between STIM1 and ORAI1 proteins and between ORAI1 and ORAI1 proteins were analyzed. According to this we changed the reaction parameters as described in eq (13). Combined with the modified diffusion constant ($\bar{D}_O = D_O/1.7$) the temporal course of I_{CRAC} was examined. The results are shown in figure 14. The faded black, blue and red curves show the courses of WT ORAI1 for the three ratios of STIM1 / ORAI1 = 1, 2 and 3 respectively. These courses are being compared to the modified version (mod) of the base case scenario including the altered reaction rates and the altered

diffusion constant (black, blue and red respectively). The differences between the basic reaction scheme and the modified version are small.

To sum up, the analysis of the base case scenario leads to the following conclusions:

- I_{CRAC} strongly depends on the ratio of STIM1 / ORAI1. Ca^{2+} influx is highest for STIM1 / ORAI1 $\gtrsim 2.5$.
- The dominant channel form of the steady state value of I_{CRAC}^{stat} (at $t = 300$ s) depends on the possibility of single CRAC channel subunits to disrupt hexameric channels (what we call „stealing mechanism“). Are the subunits not able to disrupt the fully open CRAC channels (core reactions, eqns. (14)) the hexameric state is the most favored state. Including the stealing mechanism (eq. (15)) leads to a lower number of hexamers and a higher number of tetramers. The total amount of Ca^{2+} influx is decreased in this case.
- Altering the diffusion rate of free ORAI1 dimers does not result in significant changes of I_{CRAC}^{stat} .
- Altered reaction and diffusion rates according to [29] (eq. (13) and $\tilde{D}_O = D_O/1.7$) do not significantly change the temporal course of I_{CRAC} for different ratios of STIM1 / ORAI1.

C. Stochastic analysis of the H_2O_2 Scenario

In the next step we extended our model taking the findings of an intramolecular locking of ORAI1 channels by H_2O_2 ([29]) into account. We introduce a new particle species O^H , which represents ORAI1 proteins that reacted with ROS (cf. section II). Thus three parameters are of major interest:

- The ratio of the total amount of STIM1 proteins to the total amount of ORAI1 proteins (i. e. free O and O^H as well as all ORAI1 proteins in whichever CRAC channel configuration) S^{tot}/O^{tot} ,
- the ratio of ROS-inhibited ORAI1 (free and bound in CRAC channels) to the total amount of ORAI1 O^H/O^{tot} and
- the inhibition parameter δ (cf. eq. (12)).

As default values we choose $S^{tot}/O^{tot} = 2$, $O^H/O^{tot} = 0.95$ and $\delta = 0.1$.

To count the different channel states we summed up all different combinations occurring of each type of CRAC channel state, which means

$$\begin{aligned} Z_1 &= Z_1^0 + Z_1^H, \\ Z_2 &= Z_2^0 + Z_2^H + Z_2^{2H} \text{ and} \\ Z_3 &= Z_3^0 + Z_3^H + Z_3^{2H} + Z_3^{3H} \end{aligned} \quad (25)$$

where the numbers of H in the exponent indicate how many ORAI1 dimers that interacted with ROS are part of the respective CRAC channel.

First, we exemplarily analyzed the temporal course of all occurring Z_i^j at default values. Figure 15A shows the total number of single CRAC channel subunits in the H_2O_2 scenario (black) compared to the base case scenario (blue). In addition, the two possible configurations of CRAC channel subunits in the H_2O_2 scenario are shown in gray. We see that within the temporal course, the total numbers of subunit states stay constant in both scenarios except for statistical fluctuations. In contrast, the composition of Z_1^{tot} in the H_2O_2 scenario changes within time: The number of WT subunits (Z_1^0) increases, while the number of ROS-inhibited subunits (Z_1^H) decreases.

The analysis of the tetramer and the hexamer CRAC channel states shows similar features (figure 15B and C): The total numbers of the respective channel configurations stay constant in both scenarios, while the composition of Z_i^{tot} of the H_2O_2 scenario changes: For small t a lot of total ROS-inhibited channels can be found. For larger t the number of total ROS-inhibited channels decreases, while the numbers of mixed channel states and of homomeric WT channel states increase.

Comparing the total numbers of Z_i of the two different scenarios we see the following: There are less single CRAC channel subunits, roughly equal tetramer CRAC channels and more hexamer CRAC channel states in the base case scenario than in the H_2O_2 scenario. These differences as well as the dynamic changes in the composition of Z_i^{tot} occur due to the differences in the reaction rates between WT channels and partially or completely ROS-inhibited channels. While $\tilde{k}_1$ and $\tilde{k}_2$ are larger than k_1 and k_2 , the rates for reactions between higher CRAC channel states $\tilde{k}_3$ to $\tilde{k}_7$ are smaller than for simple WT CRAC channel states k_3 to k_7 (cf. eq. (13) and table I).

In the next step we varied these three parameters and analyzed the respective temporal courses of I_{CRAC} (figs 16A, 17A and 18A) and the steady state values of the different CRAC channel states Z_i as well as of I_{CRAC} normalized to O^{tot} (figs 16B, 17B and 18B).

First, we analyzed the ratio S^{tot}/O^{tot} while the other two parameters were fixed at their respective default values (figure 16B). As in the base case scenario, the highest Ca^{2+} influx is reached for $S^{tot}/O^{tot} \gtrsim 2.5$. Due to altered reaction parameters in the H_2O_2 scenario, less hexameric channels and more tetrameric and dimeric channels are formed. The absolute value of I_{CRAC} is significantly smaller ($\approx 35\%$) in the H_2O_2 scenario since the Ca^{2+} conductance of the ROS-inhibited ORAI1 channels (or channel parts) is just ten percent of the conductance of WT ORAI1 channels.

We compare the temporal course of I_{CRAC} between the two scenarios for different ratios of S^{tot}/O^{tot} (figure 16A, $S^{tot}/O^{tot} = 1$: black, $S^{tot}/O^{tot} = 2$: blue, $S^{tot}/O^{tot} = 3$: red). We see that in the base case scenario (solid lines) as well as in the H_2O_2 scenario (dashed lines) I_{CRAC} reaches a stable state after the first five seconds. In

Figure 15: A: Temporal courses of the number of CRAC channel subunits of the WT scenario (blue) and of the H_2O_2 scenario (black and gray) for $t \in [0 \text{ s}, 300 \text{ s}]$, B: temporal courses of tetramer CRAC channel states of the WT scenario (blue) and of the H_2O_2 scenario (black and gray) for $t \in [0 \text{ s}, 300 \text{ s}]$, C: temporal courses of hexamer CRAC channel states of the WT scenario (blue) and of the H_2O_2 scenario (black and gray) for $t \in [0 \text{ s}, 300 \text{ s}]$

Figure 16: A: Temporal course of I_{CCRAC}^{mix} (cf. eq(12)) for different ratios of S^{tot}/O^{tot} (base case scenario: solid lines, H_2O_2 scenario: dashed lines) with fixed $O^H/O^{tot} = 0.95$ and $\delta = 0.1$, B values of the different CRAC channel states Z_i and of I_{CCRAC}^{mix} at $t = 300 \text{ s}$ depending on the ratio of S^{tot}/O^{tot} (base case scenario: solid lines, H_2O_2 scenario: dashed lines)

Figure 17: A: temporal course of I_{CCRAC}^{mix} (cf. eq(12)) for different ratios of O^H/O^{tot} with fixed $S^{tot}/O^{tot} = 2$ and $\delta = 0.1$, B values of the different CRAC channel states Z_i and of I_{CCRAC}^{mix} at $t = 300 \text{ s}$ depending on the ratio O^H/O^{tot}

contrast to the base case scenario, in the H_2O_2 scenario this is not the final steady state value, since I_{CCRAC}^{mix} rises again after about 100 seconds. The first rise is due to diffusion of the proteins to the PMJ and the first CRAC channel formation. The second rise is due to the mixture of original and modified on- and off-rates and the high amount of ROS-inhibited ORAI1.

Analyzing the temporal course of I_{CCRAC}^{mix} at different ratios of O^H/O^{tot} (cf. figure 17A) at the fixed ratio $S^{tot}/O^{tot} = 2$ and for $\delta = 0.1$ shows, that the chosen

value of O^H/O^{tot} affects whether the value of I_{CCRAC}^{mix} at $t = 300 \text{ s}$ is the steady state value or not. Values of the case $O^H/O^{tot} = 0$ (black) correspond to the base case scenario. $O^H/O^{tot} = 1$ (green) correspond to all ORAI1 dimers interacted with ROS before building CRAC channels. $O^H/O^{tot} = 0.2$ (blue) connotes that 20 % of ORAI1 dimers interacted with ROS before building CRAC channels. In all three cases the value of I_{CCRAC} approached a stable after about five to ten seconds. In contrast at a ratio of $O^H/O^{tot} = 0.95$, which means that 95 % of the available ORAI1 interacted with ROS, I_{CCRAC}^{mix} rises continuously within the 300 seconds. This is due to the fact that a large amount of ORAI1 features the modified on- and off-rates of the H_2O_2 scenario.

Figure 17B shows the values of I_{CCRAC}^{mix} at $t = 300 \text{ s}$ and of Z_i normalized to O^{tot} in dependence of the ratio O^H/O^{tot} ($S^{tot}/O^{tot} = 2$, $\delta = 0.1$). The number of Z_1 increases, whereas the number of Z_3 decreases for higher O^H/O^{tot} . The number of Z_2 stays at similar level within the whole interval. This behavior combined with a ten percent contribution of Z_i to I_{CCRAC}^{mix} ($\delta = 0.1$) yields less I_{CCRAC}^{mix} the more ORAI1 dimers interacted with ROS (i. e. the larger the ratio O^H/O^{tot}). In total the final value of I_{CCRAC}^{mix} for $O^H/O^{tot} = 1$ is equal to ten percent of the value of I_{CCRAC}^{mix} for $O^H/O^{tot} = 0$, which is as expected: With $O^H/O^{tot} = 1$ equation (12) simplifies to

$$I_{CCRAC}^{mix}(O^H/O^{tot} = 1) = \delta \cdot (c_1 \cdot N_{Z_1} + c_2 \cdot N_{Z_2} + c_3 \cdot N_{Z_3}), \quad (26)$$

which is equal to multiplying equation (10) with δ .

Finally the influence of δ for fixed ratios $S^{tot}/O^{tot} = 2$ and $O^H/O^{tot} = 0.95$ is examined (figure 18B). In general the steady state value of I_{CCRAC}^{mix} increases with increasing δ , which is as expected, since the Ca^{2+} influx depends linear on δ (cf. equation 12).

Comparing different values of δ the temporal courses and absolute values of I_{CCRAC}^{mix} differ much from each other (figure 18A). Large δ leads to a faster increase of $I_{CCRAC}^{mix}(t)$, which is obvious from the data before (cf. figures 15B and C): In the first seconds mainly total ROS-inhibited CRAC channel states are formed. For large δ

Figure 18: A: temporal course of I_{CRAC}^{mix} (cf. eq(12)) for different values of δ with fixed $S^{tot}/O^{tot} = 2$ and $O^H/O^{tot} = 0.95$, B values of the different CRAC channels Z_i and of I_{CRAC}^{mix} at $t = 300$ s depending on δ

these states contribute strongly to I_{CRAC}^{mix} .

For $\delta = 0$ (black), which means CRAC channels including ROS preincubated ORAI1 dimers can not conduct any Ca^{2+} into the cell, I_{CRAC}^{mix} rises within the total analyzed time interval ($t \in [0 \text{ s}, 300 \text{ s}]$). A similar behavior is seen for small delta ($\delta = 0.1$, blue). In the case that channels including ROS preincubated ORAI1 contribute half as strong to I_{CRAC}^{mix} as the channels consisting of WT ORAI1 only ($\delta = 0.5$, red) the temporal course of I_{CRAC}^{mix} is almost stable after about 30 seconds. If the mixed and the WT channels contribute in equal manner to I_{CRAC}^{mix} ($\delta = 1$, green), the same rises to a maximum after 30 seconds and then drops continuously again.

These differences and especially these instabilities in the course of I_{CRAC} arise, because the compositions of the total numbers of CRAC channel states Z_i^{tot} are not stable either. Due to the different on- and off-rates as well as the different cooperativity parameters α and β for WT ORAI1 and WT CRAC channel states and for ROS preincubated ORAI1 and mixed CRAC channel states, the process of binding and unbinding of CRAC channel states stays dynamic. Therefore, also the predicted Ca^{2+} current, which is estimated by the numbers of different CRAC channel states is dynamic.

Summing up, the introduction of the new species, the ORAI1 dimers which reacted with ROS before building CRAC channels, leads to changes of the predicted Ca^{2+} influx into the cell:

- As CRAC channels including ROS preincubated ORAI1 dimers contribute less to the Ca^{2+} current than simple WT CRAC channels, the total I_{CRAC}^{mix} drops largely, when a lot of ROS preincubated ORAI1 dimers are taken into account.
- Introducing the new species O^H leads to dynamic binding and unbinding of the CRAC channel states and therefore the temporal course of I_{CRAC}^{mix} differs from the course seen in the base case scenario.
- Depending on the chosen set of parameters (S^{tot}/O^{tot} , O^H/O^{tot} and δ), I_{CRAC}^{mix} either continuously rises within the examined 300 seconds,

reaches a stable steady state value after less than 30 seconds, or drops again after reaching a maximum value.

Simultaneously, one feature of the base case scenario remains unchanged in the H_2O_2 scenario: If the number of available STIM1 dimers is at least 2.5 times larger than the total number of ORAI1 dimers, the highest values of I_{CRAC}^{mix} at $t = 300$ s are reached.

IV. DISCUSSION

With the help of a reaction-diffusion model we analyzed the STIM1-ORAI1 stoichiometries during CRAC channel formation, the influence of dynamic PM junctions and additionally focused on the effects of ROS inhibition.

Within the base case scenario, which correlates to WT ORAI1 in experiments we find that the highest value of the amplitude of I_{CRAC} is reached when the ratio of STIM1 / ORAI1 equals or is larger than 2 in the analytical and numerical analysis and equals or is larger than 2.5 in the stochastic analysis. The difference occurs, since in the stochastic case diffusion of the proteins is taken into account and the minimal number of STIM1 does not suffice the free ORAI1 dimers, since they are possibly not at the same spot.

Analyzing the numbers of active channels and the distribution of those, it is found that the most frequently occupied channel state is the hexameric channel state, followed by tetrameric channels for the core reaction system. Taking into account a further reaction between free ORAI1 dimers and fully open CRAC channel states (what we call „stealing mechanism“), the most dominant channel configuration is tetrameric followed by hexameric states. In this case the total amount of I_{CRAC} is lower than without stealing mechanism. The dimeric channel subunit plays a minor role in contributing to Ca^{2+} influx into the cell. This analysis indicates, that the most likely channel configuration and with that the amount of Ca^{2+} influx can be controlled by the interplay between single CRAC channel subunits and already established CRAC channels.

Analyzing variations of the diffusion constant of ORAI1 and changes in the reaction rate parameters (on- and off-rate as well as cooperativity parameters α and β) shows, that for small off-rates the tetramer CRAC channel state configuration is more often occupied than the hexamer configuration in both cases, without and with stealing mechanism. For larger off-rates the situation changes: Considering the core reactions only, the hexamer configuration is more likely than the tetramer configuration. Considering the core reaction system with stealing mechanism, the tetramer configuration is still more often occupied, but the gap between the tetramer and the hexamer configuration is much smaller. The influence of the on-rate cooperativity parameter α on the

different channel states is negligible in the examined parameter range. Large off-rate cooperativity parameter β leads to less hexamer CRAC channel states than small β and simultaneously more tetramer CRAC channel states.

Analytic and numeric analysis of the steady state value of I_{CRAC} in dependence of the on- and off-rates as well as the ratio of STIM1 / ORAI1 show, that increasing both the unbinding cooperativity and the ratio STIM1 / ORAI1 results in a slight decrease of I_{CRAC}^{stat} after the highest value (for STIM1 / ORAI1 = 2) is reached. This is in good agreement with the experimental results of [28].

In a further step the influence of ROS-inhibited channels was analyzed. Therefore the whole reaction-diffusion system was extended by a new type of species (the ROS preincubated ORAI1) and the formation of channel complexes was altered accordingly. Analysis show that still for the ratio of STIM1 / ORAI1 being equal or larger than 2.5 the largest Ca^{2+} influx is observed. In this case the maximum value of I_{CRAC} , which is only 35% of the maximum value of the base case scenario, is reached. The introduced inhibition parameter δ influences $I_{CRAC}(t)$ for given ratios STIM1 / ORAI1 and O^H/O^{tot} . For large amounts of O^H with respect to the total amount of ORAI1, I_{CRAC} drops drastically.

Alansary et al. [29] analyzed the influence of ROS on the amplitude of Ca^{2+} influx experimentally. In accordance to these experiments our analysis shows, that the drop in I_{CRAC} for preincubated ORAI1 is not only due to alterations in the diffusion constant of ORAI1 and in the ORAI1-ORAI1 and the ORAI1-STIM1 interaction, but that there is another mechanism needed to explain the strong decrease of I_{CRAC} . In [29] it is found that the interaction of two transmembrane domains lock the CRAC channel when preincubated with ROS. The introduction of the new channels into the model to account for this locking shows, that with this modification the drastic decrease of I_{CRAC} can be explained, while only changing the reaction and diffusion parameters according to the changes found in the experiment does not yield such a large decrease in I_{CRAC} .

Recent experiments, which analyze the ORAI1 stoichiometry of CRAC channels demonstrate that fully functional ORAI channels exist as hexamers, in accordance with the crystal structure [15, 17, 33]. Other experiments indicate that CRAC channels can occur as a mixture of hexameric, tetrameric and dimeric channel complexes or as hexamers with different conducting states [16, 34]. A very recent report provides evidence for the existence of different conductance states of hexameric channels due to the ability of STIM1 dimers to crosslink neighbouring ORAI1 hexamers to provide a more efficient activation for a given number of STIM1 molecules [35]. Until now, there has been no model taking these new findings into account. We therefore close this gap and present a reaction-diffusion model in which we include three types of channels and therefore three types of conducting states: A CRAC channel subunit with only one ORAI1 dimer bound to STIM1 proteins with low conduc-

tance, an intermediate tetrameric CRAC channel with medium conductance and a fully open (i. e. hexameric) CRAC channel state with full conductance. Our model would lead to similar results if we assume the different conductance states to be due to hexameric channels with different number of bound STIM1 molecules.

With the help of the introduced „stealing mechanism“ we can account for the dynamic reversible binding reactions between free STIM1 and ORAI1 as seen in [22]. In contrast to these findings in our model the free proteins remain within the junctional region.

Previous models [18, 21, 22, 27] all assume tetramers as highest ORAI1 complexes. All studies evolve the best ratio of single STIM1 monomers to single ORAI1 monomers to be roughly 2:1. Also in our model, which is adapted to tetrameric and hexameric CRAC channels, the ratio of STIM1 / ORAI1 must be at least two or larger to gain highest Ca^{2+} influx.

While the models of [21] and [27] assume a static area as PMJ region, we allow a dynamic grow of the PMJ regions as seen in the cluster analysis of TIRF experiments.

In contrast to the previous models we can distinguish between the reaction-diffusion analysis of WT ORAI1 with STIM1 and the reaction-diffusion analysis of ROS-inhibited ORAI1 with STIM1. We can tune the quantity of inhibited proteins and therefore can predict the magnitude of Ca^{2+} influx in dependence of the ratio of preincubated ORAI1 to the total amount of ORAI1.

In summary the described model opens the possibility to predict Ca^{2+} influx into cells while modifying reaction and diffusion parameters, the amount of ROS-inhibited ORAI1 proteins and the predefinition of PMJ regions. So far, assessing and tuning these parameters experimentally is hard. Therefore our model can reveal new insights in the complex and dynamic stoichiometry of STIM1 and ORAI1 proteins during CRAC channel formation.

Addressing the Orai-STIM interaction during CRAC channel formation may also rise the question of how different STIM and Orai homologs (i. e. STIM2, ORAI2 and ORAI3) may influence the stoichiometry of the CRAC channels and the Ca^{2+} influx. For example recent studies [36] find that the CRAC channels consisting of a mixture of ORAI1 and Orai3 show redox insensitivity. Further extensions of a model according to these homologs may offer new insights into the complex analysis of CRAC channel formation.

Author Contributions

BS performed calculations, performed the experiments, and analyzed the data. HR supervised and designed the theoretical aspects of the research, BAN advised on biological aspects and supervised the experiments. BS, BAN, IB and HR wrote the manuscript.

Acknowledgments

We thank Dalia Alansary for the help with the experimental work. This work was funded by the German Re-

search Foundation (DFG) within the Collaborative Research Center SFB 1027, projects A3 to HR and C4 to BN and IB.

- [1] Berridge M J, Lipp P and Bootman M D, The versatility and universality of calcium signalling, *Nature Reviews Molecular Cell Biology*, 1, 11-21 (2000)
- [2] Clapham, David E., Calcium Signaling, *Cell*, 131, 6, 1047-1058 (2007)
- [3] Carafoli E, Calcium signaling: A tale for all seasons, *PNAS*, 99, 3, 1114-1122 (2002)
- [4] Falcke M, Reading the patterns in living cells — the physics of Ca^{2+} signaling, *Advances in Physics*, 53, 3, 255-440 (2004)
- [5] Lewis R S, Calcium Signaling Mechanisms in T Lymphocytes, *Annu. Rev. Immunol.* 19, 497-521 (2001)
- [6] Parekh A B and Putney Jr. J W, Store-Operated Calcium Channels, *Physiological Reviews*, 85, 2, 757-810 (2005)
- [7] Hoth M and Penner R, Depletion of intracellular calcium stores activates a calcium current in mast cells, *Nature*, 355, 6358, 353-356, (1992)
- [8] Muik M, Frischauf I, Derler I, Fahrner M, Bergsman J, Eder P, Schindl R, Hesch C, Polzinger B, Fritsch R, Kahr H, Madl J, Gruber H, Groschner K and Romanin C, Dynamic Coupling of the Putative Coiled-coil Domain of ORAI1 with STIM1 Mediates ORAI1 Channel Activation, *The Journal of Biological Chemistry*, 283, 12, 8014-8022 (2008)
- [9] Luik R M, Wang B, Prakriya M, Wu M M and Lewis R S, Oligomerization of STIM1 couples ER calcium depletion to CRAC channel activation, *Nature*, 454(7203), 538-542 (2008)
- [10] Derler I, Jardin I and Romanin C, Molecular mechanisms of STIM/Orai communication, *Am J Physiol Cell Physiol*, 310, C643-C662 (2016)
- [11] Wu M M, Buchanan JA, Luik R and Lewis R S, Ca^{2+} store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane, *The Journal of Cell Biology*, 174, 6, 803-813 (2006)
- [12] Liou J, Fivaz M, Inoue T and Meyer T, Live-cell imaging reveals sequential oligomerization and local plasma membrane targeting of stromal interaction molecule 1 after Ca^{2+} store depletion, *PNAS*, 104, 22, 9301-9306 (2007)
- [13] Park C Y, Hoover P J, Mullins F M, Bachhawat P, Covington E D, Raunser S, Walz T, Garcia K C, Dolmetsch R E and Lewis R S, STIM1 Clusters and Activates CRAC Channels via Direct Binding of a Cytosolic Domain to Orai1, *Cell*, 136, 876-890 (2009)
- [14] Yuan J P, Zeng W, Dorwart M R, Choi Y-J, Worley P F and Muallem S, SOAR and the polybasic STIM1 domains gate and regulate Orai channels, *Nature Cell Biology*, 11, 3, 337-343 (2009)
- [15] Yen M, Lokteva L A and Lewis R S, Functional Analysis of Orai1 concatemers Supports a Hexameric Stoichiometry for the CRAC Channel, *Biophysical Journal*, 111, 1897-1907 (2016)
- [16] Li P, Miao Y, Dani A and Vig M, α -SNAP regulates dynamic, on-site assembly and calcium selectivity of Orai1 channels, *Molecular Biology of Cell*, 27(16):2542-2553 (2016)
- [17] Cai X, Zhou Y, Nwokonko R M, Loktionova N A, Wnag X, Xin P, Trebak M, Wang Y and Gill D L, The Orai1 Store-operated Calcium Channel Functions as a Hexamer, *The Journal of Biological Chemistry*, 291(50):25764-25775 (2016)
- [18] Hoover P J and Lewis R S, Stoichiometric requirements for trapping and gating of Ca^{2+} release-activated Ca^{2+} (CRAC) channels by stromal interaction molecule 1 (STIM1), *Proc. Natl. Acad. Sci. U.S.A.*, 108, 32, 13299-13304 (2011)
- [19] Scrimgeour N, Litjens T, Ma L, Barritt G J and Rychkov G Y, Properties of Orai1 mediated store-operated current depend on the expression levels of STIM1 and Orai1 proteins, *The Journal of Physiology*, 587, 12, 2903-0918 (2009)
- [20] Bogeski I, Kummerow C, Al-Ansary D, Schwarz E, Koehler R, Kozai D, Takahashi N, Peinelt C, Griesemer D, Bozem M, Mori Y, Hoth M and Niemeyer B A, Differential Redox Regulation of ORAI Ion Channels: A Mechanism to Tune Cellular Calcium Signaling, *Science Signaling*, 3, 115 (2010)
- [21] Peglow M, Niemeyer B A, Hoth M and Rieger H, Interplay of channels, pumps and organelle location in calcium microdomain formation, *New Journal of Physics*, 15, 27pp (2013)
- [22] Wu M M, Covington E D and Lewis R S, Single-molecule analysis of diffusion and trapping of STIM1 and ORAI1 at endoplasmatic reticulum-plasma membrane junctions, *Molecular Biology of the Cell*, 25(22):3672-85 (2014)
- [23] Shen W-W, Frieden Ma and Demareux N, Remodeling of the endoplasmic reticulum during store-operated calcium entry, *Biology of the Cell*, 103 (8), 365-380 (2011)
- [24] Saüc S, Bulla M, Nunes P, Orzi L, Marchetti A, Antigny F, Bernheim L, Cosson P, Frieden M and Demareux N, STIM1 traps and gates Orai channels without remodeling the cortical ER, *The company of Biologists*, 128, 1568-1579 (2015)
- [25] Wu M M, Buchanan JA, Luik R M, and Lewis R S, Ca^{2+} store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane, *The Journal of Cell Biology*, 174, 6, 803-813 (2006)
- [26] Malli R, Naghdi S, Romanin C and Graier W F, Cytosolic Ca^{2+} prevents the subplasmalemmal clustering of STIM1: an intrinsic mechanism to avoid Ca^{2+} overload, *Journal of Cell Science*, 121(019), 3133-3139 (2008)
- [27] Melunis J, Hershberg U, A spatially heterogeneous Gillespie algorithm modeling framework that enables individual molecule history and tracking, *Engineering Applications of Artificial Intelligence*, 62, 304 - 311 (2017)
- [28] Kilch T, Alansary D, Peglow M, Dörr K, Rychkov G, Rieger H, Peinelt C and Niemeyer B A, Mutations of the Ca^{2+} -sensing Stromal Interaction Molecule STIM1 regulate Ca^{2+} influx by altered oligomerization of STIM1

- and by destabilization of the Ca^{2+} channel Orai1 J. Biol. Chem. 288, 1653 (2013)
- [29] Alansary D, Schmidt B, Dörr K, Bogeski I, Rieger H, Kless A and Niemeyer B A, Thiol dependent intramolecular locking of Orai1 channels, Scientific Reports, 6, 33347 (2016)
- [30] Zhou Y, Wang X, Wang X, Loktionova N A, Cai X, Nwokonko R M, Vrana E, Wang Y, Rothberg B S and Gill D L, STIM1 dimers undergo unimolecular coupling to activate Orai1 channels, Nature Communications, 6:8395 (2015)
- [31] Gillespie D T, A General Method for Numerically Simulating the Stochastic Time Evolution of Coupled Chemical Reactions, Journal of Computational Physics 22, 403-434 (1976)
- [32] Gibson M A and Bruck J, Efficient Exact Stochastic Simulation of Chemical Systems with Many Species and Many Channels, J. Phys. Chem. A, 104, 1876-1889 (1999)
- [33] Hou X, Pedi L, Diver M M, and Long S B, Crystal structure of the calcium release-activated calcium channel Orai, Science, 338, 1308-1313 (2012)
- [34] Dynes J L, Amcheslavsky A and Cahalan M D, Genetically targeted single-channel optical recording reveals multiple Orai1 gating states and oscillations in calcium influx, PNAS 113 2, 440-445 (2016)
- [35] Zhou Y, Nwokonko R M, Cai X, Loktionova N A, Abdulqadir R, Xin P, Niemeyer B A, Wang Y, Trebak M and Gill D, Cross-linking of Orai1 channels by STIM proteins, PNAS, 115 15, E3398-E3407 (2018)
- [36] Saul S, Gibhardt C S, Schmidt B, Lis A, Pasieka B, Conrad D, Jung P, Gaupp R, Wönnenberg B, Diler E, Stanis H, Vogt T, Schwarz E C, Bischoff M, Herrmann M, Tschernig T, Kappl R, Rieger H, Niemeyer B A and Bogeski I, A calcium-redox feedback loop controls human monocyte immune responses: The role of ORAI Ca^{2+} channels, Science Signaling, 9, 418 (2016)