1 **Interruption of signaling pathways in lung epithelial cell by *Mycobacterium tuberculosis***

2
3 Shima Hadifar^{1,2}, Ava Behrouzi^{1,2}, Abolfazl Fateh^{1,2}, Shohreh Khatami³, Fatemeh Rahimi
4 Jamnani^{1,2}, Seyed Davar Siadat^{1,2}, Farzam Vaziri^{1,2*}

5
6 ¹ Department of Mycobacteriology and Pulmonary Research, Pasteur Institute of Iran, Tehran,
7 1316943551, Iran

8 ² Microbiology Research Center (MRC), Pasteur Institute of Iran, Tehran, 1316943551, Iran.

9 ³ Department of Biochemistry, Pasteur Institute of Iran, Tehran, 1316943551, Iran

10 * Corresponding author: farzam_vaziri@yahoo.com; f_vaziri@pasteur.ac.ir.

11
12 **Running title:** *M.tb*-lung epithelial cell interactions

13 14 **Abstract**

15 Alveolar epithelial cell (AEC) provides a replication niche for *Mycobacterium tuberculosis*
16 (*M.tb*). Based on the role of AEC in *M.tb* pathogenesis and existence of genetic diversity within
17 this bacterium, we investigated interactions between AEC II and two different *M.tb* lineages. We
18 have compared the transcriptome and cytokines/chemokines levels of A549 infected by *M.tb*
19 lineage 3 and 4 using qRT-PCR and ELISA arrays, respectively. We showed different *M.tb*
20 strains induced changes in different effectors that involved in TLRs and NF-κB signalling
21 pathways. We observed different reaction of the studied lineages specifically in pathogenesis,
22 immune evasion mechanism, IL-12/IFN-γ axis and autophagy. Similar behaviour was detected in
23 regarding to apoptosis, necroptosis, anti-inflammatory responses and canonical inflammasome.
24 Our findings contribute to elucidate more details in pathogenesis, immune evasion strategies,
25 novel target and druggable pathway for therapeutic intervention and host directed therapy in TB
26 infection. Also, different *M.tb* lineages-dependent host-pathogen interactions suggested using
27 only one standard strain (*e.g.* H37Rv) for this kind of research will be controversial.

28
29 **Keywords:** Alveolar epithelial cell/ *Mycobacterium tuberculosis*/ NF-κB signalling/ TLR
30 signalling/ WGS

31
32
33
34
35
36
37

Introduction

Mycobacterium tuberculosis (*M.tb*), a highly virulent respiratory pathogen, generally infects its host through the aerosol route. After inhalation, *M.tb* have the opportunity to interact with many cell types in human lung, especially professional phagocytic cells that reside within alveolar sacs, such as alveolar macrophages and dendritic cells which are investigated extensively (Bermudez, Sangari et al., 2002, Manca, Tsenova et al., 2001a, Sasindran & Torrelles, 2011, Van Golde, 1985). In recent studies, alveolar epithelium as nonprofessional phagocytic cells have been described to display capacity as a safe replication niche for intracellular *M.tb* (Chuquimia, Petursdottir et al., 2012, Garcia-Perez, Castrejon-Jimenez et al., 2012, Ryndak, Singh et al., 2015). Two types of alveolar epithelium described which including type I alveolar epithelial cells (AEC I), and type II alveolar epithelial cells (AEC II). AEC I, large squamous cells, lines 90 to 95% of the alveolar surfaces and AEC II, small cuboidal cells, covers a 7% of the alveolar surfaces (Castranova, Rabovsky et al., 1988, Fehrenbach, 2001, Williams, 2003). Both cells are contributing to respiratory tract defence, specially AEC II due to secrete a variety of active substances which participate in the formation of local immune responses (Lo, Hansen et al., 2008). However, human respiratory epithelial cells responses in mycobacterial infection were poorly investigated. Due to the significant role of innate immunity in early control of *M.tb* infection and providing a platform for adaptive immune response, this issue turns into an interesting field for profound studies (Sia, Georgieva et al., 2015). Epithelial cells as well as macrophage cells expressed pattern recognition receptors (PRRs) such as the TLRs, CLRs, and NLRs that have been implicated in interacting with pathogen-specific ligands (Ioannidis, Ye et al., 2013, Lala, Dheda et al., 2007, Lee, Yuk et al., 2009). Among the TLRs family, as one of the most important PRR families which are capable to recognizing mycobacterial ligands, TLR2, TLR4, and TLR9 have an important role in shape of immune response against tuberculosis (TB) (Kleinnijenhuis, Oosting et al., 2011). The TLRs family act through a MyD88-dependent and/or independent pathway to trigger of downstream signalling pathway and activation of the targets (Kawai & Akira, 2010). Activation of these pathways either lead to induction of pro- and anti-inflammatory cytokines by activation of NF- κ B and MAPKs pathways or induction of type I IFN genes as well as inflammatory cytokines by activation IRFs, NF- κ B and MAPKs pathways (Kawasaki & Kawai, 2014). Therefore, prior to activation of the adaptive immune response, innate mechanisms that occurred must be investigated for understanding of the clinical outcomes of mycobacterial infection; this will elucidate the ways which can increase the immune response against *M.tb* more efficiently.

M.tb strains based on their phylogeographic structure are grouped into different lineages. Lineage-specific characteristics of *M.tb* strains (transmission, growth, cytokine and IFN I gene induction) were reported previously (Carmona, Cruz et al., 2013, Coscolla & Gagneux, 2010, Mvubu, Pillay et al., 2017, Sarkar, Lenders et al., 2012). Lineage 4 sublineage L4.5 and Lineage 3 sublineage L3-CAS1-Delhi strains are the two dominant *M.tb* genotype which are circulating in capital of Iran (Stucki, Brites et al., 2016)(Unpublished data). The present study was conducted with the aim of comparing different lineages of *M.tb* strains (lineages 3 and 4) in interrupting of epithelial signalling pathways in A549 cell line, from TLRs and NF- κ B signalling pathways to specific cytokine secretion.

Results

Gene expression analysis

To determine the differential mRNA expression profile of genes which involves in TLRs and NF- κ B-signalling pathways during infection, the result of fold change/regulation of each gene was compared and up and down regulated genes were identified (Figure 1). Choose of time point of post infection (p.i.) was performed based on results of previous study; in which it has been proven that *M.tb* is found in endosome of infected cell after 3 days and induced cell death after 4-5 days p.i. (García-Pérez, Mondragón-Flores et al., 2003, McDonough & Kress, 1995). We also generated a volcano (Figure 2) and clustergram plots (Figure 3) by the mean log₂ fold change of genes in infected cell line to visualize gene expression under different treatment. After 72 hrs p.i., genes which were differentially expressed, sorted into functional groups and their fold regulation is illustrated in Table 1.

- TLRs group including CD180 (LY64; RP105), TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10.

-Negative Regulator genes in TLRs signalling including SARM1, SIGIRR and TOLLIP.

-The TIR domain-containing adaptors genes group such as MyD88, TICAM1 (TRIF), TICAM2 or TRAM, TIRAP/MAL.

-Kinases genes group including AKT1, RAF1, MAPK8 (JNK1), MAP2K3 (MEK3), MAP2K4 (JNKK1), MAP3K1 (MEKK), MAP3K7, MAP4K4, CHUK (IKK1), IRAK1, IRAK2, IRAK4, RIPK1, RIPK2, TBK1, BTK, EIF2AK2.

-Transcription factors group involved in NF- κ B pathway NFKB1, NFKB2, REL, RELA, RELB, ATF1, EGR1, ELK1, FOS, JUN, STAT1 and IKBKB or IKK2, IKBKG (NEMO), NFKBIA, NFKBIE, NFKBIL1.

- Interferon regulatory factor gene group such as IRF1, IRF3.

- Apoptosis related genes: AGT, BCL2A1 (BCL-X), BCL2L1, BIRC3 (c-IAP2), CASP8.

-TNF pathway genes group including TNFRSF10A, TNFRSF10B, TNFRSF1A, TNFSF10 (TRAIL), TRADD, TRAF2, TRAF3, TRAF6, BIRC2 (c-IAP1), BIRC3, CASP8 and CFLAR (FLIP), ICAM1, LTB, LTBR, CD83, TNFAIP2, TNFAIP3 (A20), TIFA, HMOX1(HO-1), FASLG.

-Chemokines and cytokines genes: MCP-1 (CCL2), RANTES (CCL5), MCSF (CSF1), GM-CSF(CSF2), GCSF (CSF3), CXCL10, CXCL2, CXCL3, CCL20, IFNA1, IFNB1, IFNG, IL8 (CXCL8), L12A, IL2, IL6, IL10, IL1A, IL1B, LTA, TNF α .

-Regulator genes of adaptive immunity: CD80, CD86, HSPD1, IFNG, IL10, IL12A, IL1B, IL2, MAP3K7, TRAF6.

-Other genes: CARD11 (CARMA1), CASP1, BCL3, CD40, EGFR, IL1R, MALT1, NOD1, PSIP1, RHOA, TIMP1, STX11, UBE2N, TAB1, PTGS2 (COX2), PPARA, PELI1, NR2C2,

NFRKB, MAPK8IP3, LY96 (MD2), LY86 (MD1), HSPD1, HSPA1A, HRAS, HMGB1, ECSIT, CLEC4E (MINCLE), CD14.

Cytokine/chemokine production patterns

Several cytokines and chemokines were down or up-regulated by the different strain lineages in infected A549 cells. The release of RANTES by A549 infected with lineage 4 strain was significantly increased in compare to lineage 3 strain ($p < 0.001$). Similarly, the release of IL-8 in lineage 4 strain-infected cells was higher when compared with lineage 3 strain ($p < 0.0001$). Further, the release of MCP-1 significantly was high in both lineages-infected cells. The secretion levels of TNF α , IL-6, IL-1B, IL-12, IL-17A, IL-10, MDC (CCL22), MIP-1 α , MIP-1 β and Eotaxin were mainly negative or very poor in both mock cell and infected cells (Figure 4).

Intracellular growth and intracellular internalization Index

The intracellular growth was evaluated by CFU counts. Figure 5A showed intracellular bacterial loads of different *M.tb* genotypes at different time points. We determined lineage 3 strain had a greater intracellular load at 72 hrs p.i in comparsion to lineage 4. The cell viability of infected cell was 96.24%. Figure 5B illustrates percentile of infected A549 cell line with each studied strains. The intracellular internalization index of each *M.tb* genotypes using Ziehl-Neelsen acid fast and auramine staining was showed in Figure 5C.

Discussion

Study of host-*M.tb* interactions to investigating specific host molecular signatures appear as new platform in controlling *M.tb* infection. In various *in vivo* and *in vitro* studies, interaction of *M.tb* strain with consideration of its genetic background has been investigated and a wealth of knowledge is generated (Carmona et al., 2013, Krishnan, Malaga et al., 2011, Manca, Reed et al., 2004, Mvubu et al., 2017, Sarkar et al., 2012, Subbian, Bandyopadhyay et al., 2013, Wiens & Ernst, 2016). The translational implications of diversity in *M.tb* lineages may be useful for improving and developing of diagnostic, prophylactic and therapeutic methods. However, significant lineage diversity exists in *M.tb* population; a few representatives of a lineage in different *M.tb* target cells (macrophage, AEC) have been studied. In the current study, we showed that infection of pulmonary epithelial cells by different *M.tb* strains induced changes in different effectors that involved in TLRs and NF- κ B signalling pathways.

Secretion of cytokines and chemokines from epithelial cells during an immune response to *M.tb* infection has been well documented (Li, Wang et al., 2012, Lin, Zhang et al., 1998, Sato, Tomioka et al., 2002, Sharma, Sharma et al., 2007). TNF α as a major proinflammatory mediator and pleiotropic cytokine is critical for establishment and maintenance of latent TB (Stenger, 2005). Expression of TNF α can affect expression of several chemokines such as CCL2 (MCP-1), CCL5 (RANTES), GM-CSF(CSF2) and surface adhesion genes such as ICAM1 in host cell (Lakshminarayanan, Beno et al., 1997). In addition, interactions between mycobacterial microbe-associated molecular pattern (MAMPs) and host PRRs can produce other chemokines and cytokines such as CXCL2, CXCL3, CCL2, CCL5 and IL8 (Mortaz, Adcock et al., 2015). In our study, lineage 4 strain induced up-regulation in all of aforementioned genes in A549 cell culture. On the contrary, lineage 3 induced down-regulation in TNF α and CSF2 and had no effect on

ICAM-1. The reduction in TNF α level in response to lineage 3 can be justified by downregulation of TLR2 and its downstream effectors by specific cell wall component of lineage 3 strain. However, it has been demonstrated that different lineages of *M.tb* stimulated different innate immune response but unable to show this response associated with specific cell-surface expressed lipids (Krishnan et al., 2011). Based on transcriptional profile of A549 infected with both strains and the role of ICAM1 in facilitating crosstalk between host and neutrophils, and also regard to the role of IL-8, CXCL2 and CCL2 cytokines in neutrophil activating, it has been expected that lineage 4 comparing to lineage 3 more efficiently stimulated a large number of neutrophil to migrate toward the infected host cell. Neutrophils contribute to the mycobacterial protection by attracting circulating DC and activating effector T cells (Blomgran & Ernst, 2011). Also, recent evidence suggested a suppressor function for PMN which can dampen acquired immunity to *M.tb* and consequently contribute to augment of TB pathology (Dallenga & Schaible, 2016). Interestingly, Hasan *et al.* showed that a high level of CCL2 was found to be associated with TB severity possibly due to increased systemic inflammation (Hasan, Cliff et al., 2009). Therefore, upregulation of CCL2 (at both transcriptional and translational level) in response to both lineages, specially lineage 4 can be so crucial in the pathogenesis of these lineages. Based on induced response by lineage 4 and regards to increase of G-CSF and its role in the induction of lung destruction, it can be suggested that infection by this genotype is potent to disseminate in lung. While the low level of TNF α and down-regulation in all types of CSF and CXCL10 in response to lineages 3 provides an advantage for the establishment and development of infection before encounter to adaptive immune response. Recently it has proven that RP105-mediated activation of Akt phosphorylation is essential to mycobacteria-infected macrophages to releasing TNF (Yu, Micaroni et al., 2015). Our data showed that lineage 3 (but not lineage 4) strain strongly reduced the expression of CD180/RP105, AKT1 and TNF α simultaneously; therefore, interruption of this innate immune signaling axis by this *M.tb* genotype can be proposed as a novel strain specific immune evasion strategy. Besides, Li *et al.* demonstrated that *M.tb* can inhibit the ERK1/2 signalling pathway, leading to the suppression of TNF α , IL-6, and the improvement of mycobacterial survival within macrophages (Li, Chai et al., 2015). It seems that in our study based on the secretion profile of TNF α and IL-6 in response to both lineages; this can be the strategy which *M.tb* used for promoting its survival in lung epithelial cell. This issue is more reliable about lineage 3 which down-regulated *raf-1*, the initial kinase in ERK1/2 cascade, significantly. Besides, lineage 3 comparing to lineage 4 has a higher level of intracellular growth and number of infected cells during 48 and 72 hrs p.i. Based on the intracellular bacillary growth level, an indicator of virulence of the *M.tb* strains, it seems lineage 3 is more virulent than lineage 4 and can replicate and survive in lung epithelial cell more efficiently. This characteristic can be another proof for selective advantage of lineage 3 for its survival in the epithelial cell.

In regards to RANTES and TNF α , some discrepancies between the results of qPCR and ELISA were detected. This difference between mRNA expression level and protein production could be explained by optimum p.i. time point which needed for peak production of the cytokine. Accordingly, it has been shown that peak production of a cytokine attained after 18 hrs p.i. in an infected epithelial cell by H37Rv strain (Lee, Shin et al., 2009). While in our study, the supernatant was collected for ELISA assay at 72 hrs p.i. Other speculation for this reduction was destabilization of mRNA transcripts and change in post-transcriptional regulator resulted from difference in cell-wall component of *M.tb* (Rajaram, Ni et al., 2011).

Differences in type of induced cell death (apoptosis or necroptosis) by *M.tb* strains may contribute to define outcome of infection. TNF α production can induce host necroptosis and apoptosis (Pasparakis & Vandenabeele, 2015). TNF-induced apoptosis occurs through RIPK1, TRADD, FADD, CASP8 and FLIP. It has been shown that RIPK1 normally induces cell survival through recruitment of a group of downstream genes including cIAP1/2, TAB2, TAK1, LUBAC, IKK complex which consequently promotes activation of MAPKs, NF- κ B and proinflammatory genes during TNF and its receptor (TNFRSF1A) interaction. This normal process depends upon ubiquitination of RIPK1 by cIAP1/2 protein as anti-apoptosis effector. On the other hand, any changes in optimal RIPK1 ubiquitination results in apoptosis or necroptosis. In addition, level of FLIP is important in the control of necroptosis and apoptosis by formation of heterodimers with caspase-8. If expression levels of FLIP(CFLAR) or cIAP (BIRC2/3) be low, RIPK1 able to interact with caspase 8 and apoptosis is induced (Newton & Manning, 2016, Pasparakis & Vandenabeele, 2015). In the current study, expression of TRADD, FADD, RIPK1 and CASP8 were down-regulated and no change was observed in expression of FLIP in response to lineage 3. In response to lineage 4, FADD was upregulated, TRADD was down-regulated and no change was detected in RIPK1, CASP8 and FLIP. Based on our results, we expected probability of apoptosis in infected cells by both lineages might be low. TRAIL (TNFSF10), a member of the TNF family of cytokines, considered as a potent inducer of apoptosis in various tumor cells and primary lung airway epithelial cells (Akram, Lomas et al., 2013, Walczak, Miller et al., 1999). This cytokine exerts its effect through FADD and CASP8/10/3. Also, apoptosis by mitochondrial pathways (Bid, Bax, BCL2/X and CASP8/3/9) can be induced by growth factor (Abdulghani & El-Deiry, 2010). In our study, lineage 3 induced upregulation in TRAIL, while its receptor and downstream effectors such as FADD and CASP8 were down-regulated. In mitochondrial pathway, expression of antiapoptotic effector, BCLx, was down-regulated and BCL2 was upregulated. The reduced level of BCLx can resulted from down-regulation of EGFR and IKKs expression level and consequently activated NF- κ B expression level. This reduction lead to the infected cell make sensitive to TRAIL markedly (Ravi & Bedi, 2002). However, it is expected that in the infected cell, the probability of apoptosis will be low, because expression of other anti-apoptotic gene (*bcl-2*) was increased and expression of CASP8 and FADD has been reduced significantly. In response to lineage 4 strain, BCLx and TRAIL were down-regulated, but its receptor, BCL2 and FADD were up-regulated and CASP8 was not changed. Therefore, the probability of apoptosis in the infected cell can also might be low. This finding is similar to most of the studies which showed that *M.tb* is capable of inhibit the various apoptotic models that were elucidated so far (Parandhaman & Narayanan, 2014).

Induction of necroptosis in response to TNF occurs through formation of the complex which containing RIPK1, RIPK3 and MLKL (Murphy, Czabotar et al., 2013). In response to lineage 3, expression of RIPK1 was down-regulated and no change was observed in cIAP1/2. In response to lineage 4, cIAP1/2 was up-regulated and no change was detected in RIPK1. Based on this expression profile, necroptosis may not be induced in response to lineage 3 because cIAP1/2 has normal performance and RIPK1 as one of key gene in induction of the phenomenon was down-regulated. Similarly, in response to lineage 4, necroptosis might not be induced because cIAP1/2 was up-regulated and RIPK1 expression level was not changed. Another proof in this regard is the significant reduction in the expression of TNFRSF1A, the receptor of TNF, in response to both lineages which will have significant effect on necroptosis. This observation is in contrast to Stutz's study who observed that *M.tb* infection of macrophages remodeled the intracellular signaling by upregulating TNFR1, whilst downregulating cIAP1, thereby establishing a strong

necroptotic response (Stutz, Ojaimi et al., 2017). Collectively, based on the host cellular target (AEC or macrophage) this clonal bacteria show significantly different reaction.

IL-1 α and IL-1 β cytokines have substantial role in inflammation, homeostasis and protection against *M.tb* infection (Garlanda, Dinarello et al., 2013). Mice deficient either IL-1 α or IL-1 β or both showed susceptibility to infection by *M.tb* strains (Mayer-Barber, Andrade et al., 2011). IL-1 β as well as IL-18 is synthesized as precursor (pro-IL-1 β). Activation of these cytokines is dependent on NLRP3 inflammasome pathway (Latz, Xiao et al., 2013). Involvement of TLR2/TLR6 in promoting IL-1 β production has been shown in macrophages during *M.tb* infection (Kleinnijenhuis, Joosten et al., 2009). In our study, IL-1 α was increased in response to both strains, while expression level of IL-1 β was reduced in response to lineage 3 strain and no change was observed for lineage 4. In response to lineage 4, expression level of TLR2 was not altered but TLR6 was down-regulated. In infected cell line by lineage 3, expression of TLR2, TLR6 and downstream effectors such as MYD88, IKKB, a kinase subunit of IKK complex, NF- κ B were decreased, and expression of IKK α as other kinase subunit of IKK complex and effector which negatively regulates canonical NF- κ B activation (Li, Lu et al., 2005) was up-regulated which can be one of the explanations for reduction of IL-1 β for lineage 3 strain. Role of type I IFNs in negative regulation of host resistance to TB infection in mouse models has been reported (Manca, Tsenova et al., 2001b). Expression of type I IFN can be mediated by TLR 3, TLR 4, TLR 7/8, TLR 9, MyD88 or TRIF adaptors, TBK1, IKK ϵ and a group of transcription factor as known IRFs, especially IRF3 genes (Akira, Uematsu et al., 2006, Fitzgerald, McWhirter et al., 2003). In the current study, lineage 4 down-regulated expression level of IFN- β . This reduction can be induced through several pathways: first, TRIF dependent TLR4 pathway that TLR4, TRIF and TBK1 were down-regulated while expression of IRF3 was up-regulated. Due to down-regulation of TBK1 as key factor which phosphorylates and activates IRF3 and upregulation of SARM1 as negative regulator of TLR, consequently end product will be reduced. In the second pathway, TLR3, TRIF and IRF3 were up-regulated while TBK1 was down-regulated. It has been shown induction of IFN- β gene is further mediated by IRF3 while IRF7 promotes induction of type I IFN genes (Sato, Hata et al., 1998). In response to lineage 3 IFNA, TLR4 and its downstream effectors except of IRF3 were reduced. Also, expression level of IL-10 was down-regulated (at both transcriptional and translational level) in response to both lineages. This reduction can be justified by down-regulation of IFN β in response to lineage 4. Reduced expression of IL10 and type I IFNs, showing that infection with both lineages cannot promote anti-inflammatory response in pulmonary epithelial cells. Furthermore, the reduced level of type I IFNs in response to both lineage strains can be considered an explanation for increase of IL-1 α in both infected cell (Mayer-Barber, Barber et al., 2010). IL-1 α can act through COX-2 and regulate *M.tb* replication during infection. COX2 level can affect level of IFN type I in a reciprocal manner. In our study, COX2 expression level was up-regulated in response to both lineages. In addition, up-regulation of COX2 as well as IL-8 contributes to attractant of neutrophils to infected cell (Xu, Reichner et al., 2008). This increased expression level can be other explanation for reduction of type I IFN in response to our studied strains. This result can be so controversial regarding to recent proposed of using interleukin-1 and type I interferon crosstalk for host-directed therapy against TB (Mayer-Barber, Andrade et al., 2014).

Activation of TLR4/TRIF signalling is one of check point in NLRP3 inflammasome activation. In addition, it has been shown that FADD and CASP8 have an important role in inflammasome pathway which including induction of pro-IL-1 β in TLR4/TRIF dependent pathway and Dectin-

1/mycobacterial MAMPs pathway and also in canonical inflammasome activation (Gringhuis, Kaptein et al., 2012, Gurung, Anand et al., 2014). In our study, TLR4/TRIF and downstream effector such as IKKs and IFN α were down-regulated in response to lineage 3. In response to lineage 4, TLR4 and IFN β were down-regulated, but TRIF and IKKs were up-regulated. Based on this profile, interaction of *M.tb* LPS with TLR4 cannot result in non-canonical inflammasome activation. Furthermore, based on expression of CASP8 and FADD in response to both lineages, canonical inflammasome activation may not be induced. Among of the genes involved in Dectin-1 dependent pathway, we evaluated MALT1 and CASP8 which expression level of MALT-1 was up-regulated and no change was observed in response to lineage 4 and 3, respectively. It seems that upregulation in expression level of MALT-1 cannot affect expression of IL-1 β and inflammasome activation. The expression level of CASP1, as a critical regulator of IL-1 β and IL-18 maturation, was up-regulated in response to lineage 3 strain. In our study, although the expression of CASP1 was increased, but maturation of this protein by TLR/TRIF-mediated induction of caspase-11 has not been induced. In addition, it has been reported that the conversions of procaspase-1 into caspase-1 (active form) can be suppressed by Bcl-2 production in human macrophages (Shimizu, Eguchi et al., 1996). In our study, expression of a member of BCL2 protein family (BCL2A1) was increased in response to lineage 3. This increased expression level can be one of the possible mechanisms of this phenomenon. Furthermore, it has been shown that *atg-16* (an autophagic protein) deleted mice produce high level of IL-1 β and IL-18 in response to MAMPs (Saitoh, Fujita et al., 2008). Also, depletion of autophagic proteins LC3B and Beclin 1 prompted high activation of caspase-1 and IL1 β and IL-18 production (Nakahira, Haspel et al., 2011). Based on the evidence and expression level of CASP1 and IL-1 β in our study, lineage 4 likely stimulates autophagic activity. Also, this activation can be explanation for failure in activation of CASP1. Besides, down-regulation of AKT as downstream mediator of class I PI3K signalling which acts through inhibiting mTORC1 or regulating Beclin1 in autophagic pathway, may promote autophagy activation (Bento, Empadinhas et al., 2015). HMGB1 as prototypic DAMPs have multifunctional role such as regulation of autophagy, chemotaxis, inflammatory response, DNA damage and innate and adaptive immunity (Sims, Rowe et al., 2010, Yanai, Matsuda et al., 2013). In our study, HMGB1 was down-regulated in response to both genotypes (especially lineage 3). The high level of down-regulation can be one of explanations for reduced probability of autophagy pathway induction for this lineage. Therefore, these results help us to understanding the molecular mechanisms of bacterial evasive strategies relating to inflammasome and autophagy activation; which may lead to development of potent therapies against TB.

The heterodimer of IL-12A (p35), as gene constitutively and ubiquitously expressed, and IL-12B (p40), expressed in specific cells, which known as IL-12 (IL-12p70) can act as growth factor for T cells and NK cells and promotes induction of IFN- γ (Vignali & Kuchroo, 2012). Activation of IRF pathways induces expression of IL-12A and IFNs type I, whereas activation of NF- κ B results in the induction of IL12B. In our study, based on ELISA assay, any change in IL-12 production was not identified in response to both lineages compared to the mock cell. So, this reduced level can be considered as another explanation for reduction of IFN type II in current study. In addition, this reduction in IL-12 level in response to lineage 3 can be justified by reduced expression level of IKK2, NFKB1-RELA (NF κ B) and increased expression of IKK1 and I κ B as inhibitor of NF κ B which may result in reduction in IL12B (not evaluated in our study) and reduced IRF pathways that lead to reduction in IL-12A and subsequently reduction in bioactive IL-12 level. In addition, TLR2 and TLR9 as important receptor for optimal

production of IL-12p40 (Bafica, Scanga et al., 2005) also was down-regulated in response to this lineage. IL-12 and IFN- γ production is essential to establishment of protective immunity in mycobacterial infection (Frucht, Fukao et al., 2001). Accordingly, the deficiency in IL-12p40 results in to inherent susceptibility to *M.tb* infection (Özbek, Fieschi et al., 2005). Based on protective role of IL-12 and IFN- γ , reduction in their expression level provide safe environmental for replication of both *M.tb* strains before encounter to adaptive immune response. However, profile of other cytokine and chemokine can cover IFN- γ reduction in infected cell by lineage 4. In contrast to our results, it has been shown that alveolar epithelial cells in response to infection by *M.tb* strains activated the IFN- γ response pathway (Sharma et al., 2007). Inconsistent results may be attributed to difference in genetic background of the studied strains. Collectively, it seems that inducing or blocking the IL-12/IFN- γ axis in mycobacterial disease is a strain-specific phenomenon. Another strain specific strategy which we observed is about MMP-1. Induction of this enzyme may result in cavity formation in lung tissue. This cavity can provide safe niche for replication and transmission of *M.tb* strains. In addition, *M.tb* suppressed expression of TIMP-1 as specific inhibitor of MMP (79). In our study, in response to lineage 3, expression of TIMP-1 was down-regulated. This reduced expression suggested that infection by lineage 3 strain promotes cavity formation in host and provide safe haven for its transmission. In addition, the current study is consistent with Ryndak's report (Ryndak et al., 2015) who showed that AEC II responsiveness to different *M.tb* strains stimulation and could provide a safe intracellular environment for bacterial replication and dissemination during primary infection. Furthermore, according to gene expression profile of an infected cell line in response to both lineages, it is likely that lineage 3 is more potent than lineage 4 strain to manipulate responsiveness of AEC II to induction for its progressive infection and transmission. This characteristic may provide a selective advantage for lineage 3 strain. On the contrary, previous study in co-culture model showed that primary human airway epithelium (PBEC) is inert to mycobacterial infection and no increase in expression of any genes was observed within 24 hrs (Reuschl, Edwards et al., 2017). Also the authors described *M.tb* strain have poor invasion and adherence to epithelial cells. These discrepant results are attributed to the use of the different genotype of *M.tb*, the different type of epithelial cells and different p.i. time.

In summary, results of the current study demonstrated the interruption of signalling pathways of the respiratory epithelium in the interaction by two different lineages of *M.tb* strains; which seems is only the tip of the iceberg. Further mechanistic exploration of interactions between alveolar epithelial cell and *M.tb* especially with consideration of genetic background of strains may be useful for elucidate more details in pathogenesis, immune evasion strategies, novel target and druggable pathway for therapeutic intervention and host directed therapy in TB infection. The last but not least is the significant strain-specific behaviour of this clonal bacterium in host-pathogen interactions, which brings us to this conclusion that using only one standard strain (e.g. H37Rv) for this kind of research will be controversial.

Materials and Methods

Bacterial strains and infection of A549 cell line

The *M. tb* strains, including the Lineage 4 sublineage L4.5 and Lineage 3 sublineage L3-CAS1-Delhi strains which were previously identified based on 24 loci MIRU-VNTR and Spoligotyping and confirmed by whole genome sequencing method (Unpublished data). Briefly, both strains were grown in Middlebrook 7H9 culture medium (Sigma Aldrich, St. Louis, MO, USA), supplemented with 10% ADC enrichment (Becton Dickinson, Oxford, UK) at 37°C until mid-log phase (OD600: 0.6-0.9), clumps were disaggregated using glass beads. The A549 (ATCC CCL-185) as the AEC II were cultured at 37°C in 5% CO₂ in Dulbecco's Modified Eagle Medium (DMEM) supplemented with penicillin, streptomycin, and 10% fetal bovine serum (FBS)(Gibco, Paisley, UK) in six well plates (Sorfa, Zhejiang, China). Once confluent (60-70%), A549 were infected in triplicates with each strain at an MOI of ~50:1(50 bacteria: cell) (Sequeira, Senaratne et al., 2014) and remained for 2 hrs. After the time, infected cells were washed extensively with warm DMEM to remove extracellular bacteria, then the infected cell line were maintained in supplemented DMEM (1% FBS) and incubated at 37°C in 5% CO₂ for 72 hrs.

Cell viability

To determine the viability of epithelial cells that infected with mycobacterial strains, trypan blue exclusion assay (Louis & Siegel, 2011) was done based on manufactures instructions (Sigma Aldrich, Germany).

Intracellular growth assay

To determine CFU/ml of the cultures, before infection, each bacterial standard concentration of the inoculum vial was plated for reconfirmation. Also, after 48 and 72 hrs infected cells were detached with Trypsin-EDTA 1X (Gibco) and washed with DMEM. The cell suspension was centrifuged to pellet cell. The pellet was lysed with 100 ml of 0.1% SDS and incubated at room temperature for 15 mins. Lysates were serially diluted and plated onto Middlebrook 7H11 agar at each time point and incubated at 37°C for 3-4 weeks.

Evaluation of intracellular internalization Index

To investigate internalization index, the A549 cells were seeded on a sterile glass coverslip and infected with studied *M.tb* strains. After 2 hours of infection, infected cells were washed extensively with warm DMEM to remove extracellular bacteria. Infected cell remained in DMEM media until 72 hrs. After that, for counting of internalized bacilli, Ziehl-Neelsen (ZN) acid fast and auramine staining were performed. At least 300 consecutive epithelial cells were counted and classified based on the number of intracellular bacteria.

RNA Extraction and cDNA Synthesis

Total RNA from infected and mock cells were extracted using a commercial RNA extraction kit (QIAGEN, USA, Cat.No.74104,) according to the manufacturer's instructions. The cDNA was synthesized using RT2 First Strand cDNA synthesis kit (QIAGEN, Cat.No.330401) according to the manufacturer's instructions.

Quantitative Real Time-PCR (qRT-PCR)

The qRT-PCR was performed by RT² Profiler™ PCR Array kits (QIAGEN) which including RT² Profiler™ PCR Array Human Toll-Like Receptor Signalling Pathway (QIAGEN, Cat.No. PAHS-018ZF-2) and RT² Profiler™ PCR Array Human NF-κB Signalling Pathway (QIAGEN, Cat.No. PAHS-025YF-2) according to the manufacturer's instructions. Each array contains 84 pathway-specific genes. Thirty nine genes were shared between the two arrays and a total of 129 genes were investigated. All assays were done in triplicate.

Cytokine/chemokine measurements

Supernatants of infected and mock cells were collected after 72 hrs and filtered through 0.22 μm Syringe filters (Sorfa) and frozen at -70°C until used. Determination of TNFα, IL1B, IL6, IL12, IL17A, IL8, MCP-1, RANTES, MIP-1α, MIP-1β, MDC and Eotaxin was performed by Multi-Analyte ELISArray Kit (QIAGEN, Cat. No. MEH-008A) and IL-10 was assessed by Single Analyte ELISArray Kit (QIAGEN, Cat. No. SEH00572A,) according to the manufacturer's instructions. All assays were done in duplicate.

Statistical analysis

For analysis of the threshold cycle (CT) values to calculate changes in gene expression and generates different plots, RT2 Profiler PCR Array Data Analysis Webportal was applied (<https://www.qiagen.com/ir/resources/geneglobe/>). The student's t test was performed for data comparisons. The comparison was performed based on differentially expressed genes with a fold change cut off of two. For pathways drawing, path Visio software (3.2.4) was used (Kutmon, van Iersel et al., 2015). GraphPad Prism 7 (GraphPad, La Jolla, CA, USA) was used for statistical analyses and drawing figures. Values of p< 0.05 were considered to be statistically significant.

Acknowledgments: We thank all the personnel of Mycobacteriology and Pulmonary Research Department, Pasteur Institute of Iran for their assistance in this project. The work was supported by a grant [Number 927] from Pasteur Institute of Iran.

Author contributions: F.V. and S.H. designed experiments. S.D.S. and S.K. directed research. F.V. and S.H. wrote the manuscript. S.H. and A.B. performed experiments. F.R.J. and A.F. analyzed data. F.V., S.D.S. and S.K.edited the manuscript.

Conflict of interest: The authors declare no competing interests.

References

- Abdulghani J, El-Deiry WS (2010) TRAIL receptor signaling and therapeutics. Expert opinion on therapeutic targets 14: 1091-1108
- Akira S, Uematsu S, Takeuchi O (2006) Pathogen recognition and innate immunity. Cell 124: 783-801
- Akram KM, Lomas NJ, Spiteri MA, Forsyth NR (2013) Club cells inhibit alveolar epithelial wound repair via TRAIL-dependent apoptosis. The European respiratory journal 41: 683-94
- Bafica A, Scanga CA, Feng CG, Leifer C, Cheever A, Sher A (2005) TLR9 regulates Th1 responses and cooperates with TLR2 in mediating optimal resistance to *Mycobacterium tuberculosis*. Journal of Experimental Medicine 202: 1715-1724
- Bento CF, Empadinhas N, Mendes V (2015) Autophagy in the fight against tuberculosis. DNA and cell biology 34: 228-42
- Bermudez LE, Sangari FJ, Kolonoski P, Petrofsky M, Goodman J (2002) The efficiency of the translocation of *Mycobacterium tuberculosis* across a bilayer of epithelial and endothelial cells as a model of the alveolar wall is a consequence of transport within mononuclear phagocytes and invasion of alveolar epithelial cells. Infection and immunity 70: 140-146
- Blomgran R, Ernst JD (2011) Lung neutrophils facilitate activation of naive antigen-specific CD4+ T cells during *Mycobacterium tuberculosis* infection. The Journal of Immunology 186: 7110-7119
- Carmona J, Cruz A, Moreira-Teixeira L, Sousa C, Sousa J, Osorio NS, Saraiva AL, Svenson S, Kallénius G, Pedrosa J (2013) *Mycobacterium tuberculosis* strains are differentially recognized by TLRs with an impact on the immune response. PLoS One 8: e67277
- Castranova V, Rabovsky J, Tucker J, Miles P (1988) The alveolar type II epithelial cell: a multifunctional pneumocyte. Toxicology and applied pharmacology 93: 472-483
- Chuquimia OD, Petursdottir DH, Rahman MJ, Hartl K, Singh M, Fernández C (2012) The role of alveolar epithelial cells in initiating and shaping pulmonary immune responses: communication between innate and adaptive immune systems. PLoS one 7: e32125
- Coscolla M, Gagneux S (2010) Does *M. tuberculosis* genomic diversity explain disease diversity? Drug Discovery Today: Disease Mechanisms 7: e43-e59
- Dallenga T, Schaible UE (2016) Neutrophils in tuberculosis—first line of defence or booster of disease and targets for host-directed therapy? Pathogens and disease 74
- Fehrenbach H (2001) Alveolar epithelial type II cell: defender of the alveolus revisited. Respiratory research 2: 33
- Fitzgerald KA, McWhirter SM, Faia KL, Rowe DC, Latz E, Golenbock DT, Coyle AJ, Liao S-M, Maniatis T (2003) IKKε and TBK1 are essential components of the IRF3 signaling pathway. Nature immunology 4: 491
- Frucht DM, Fukao T, Bogdan C, Schindler H, O'Shea JJ, Koyasu S (2001) IFN-γ production by antigen-presenting cells: mechanisms emerge. Trends in immunology 22: 556-560
- Garcia-Perez B, Castrejon-Jimenez N, Luna-Herrera J (2012) The role of non-phagocytic cells in mycobacterial infections. In Understanding tuberculosis-Analyzing the origin of *Mycobacterium tuberculosis* pathogenicity, InTech
- García-Pérez BE, Mondragón-Flores R, Luna-Herrera J (2003) Internalization of *Mycobacterium tuberculosis* by macropinocytosis in non-phagocytic cells. Microbial pathogenesis 35: 49-55

Garlanda C, Dinarello CA, Mantovani A (2013) The interleukin-1 family: back to the future. *Immunity* 39: 1003-18

Gringhuis SI, Kaptein TM, Wevers BA, Theelen B, Van Der Vlist M, Boekhout T, Geijtenbeek TB (2012) Dectin-1 is an extracellular pathogen sensor for the induction and processing of IL-1 β via a noncanonical caspase-8 inflammasome. *Nature immunology* 13: 246

Gurung P, Anand PK, Malireddi RS, Walle LV, Van Opdenbosch N, Dillon CP, Weinlich R, Green DR, Lamkanfi M, Kanneganti T-D (2014) FADD and caspase-8 mediate priming and activation of the canonical and noncanonical Nlrp3 inflammasomes. *The Journal of Immunology* 192: 1835-1846

Hasan Z, Cliff JM, Dockrell HM, Jamil B, Irfan M, Ashraf M, Hussain R (2009) CCL2 responses to *Mycobacterium tuberculosis* are associated with disease severity in tuberculosis. *PloS one* 4: e8459

Ioannidis I, Ye F, McNally B, Willette M, Flaño E (2013) Toll-like receptor expression and induction of type I and type III interferons in primary airway epithelial cells. *Journal of virology* 87: 3261-3270

Kawai T, Akira S (2010) The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nature immunology* 11: 373-384

Kawasaki T, Kawai T (2014) Toll-like receptor signaling pathways. *Frontiers in immunology* 5

Kleinnijenhuis J, Joosten LA, van de Veerdonk FL, Savage N, van Crevel R, Kullberg BJ, van der Ven A, Ottenhoff TH, Dinarello CA, van der Meer JW (2009) Transcriptional and inflammasome-mediated pathways for the induction of IL-1 β production by *Mycobacterium tuberculosis*. *European journal of immunology* 39: 1914-1922

Kleinnijenhuis J, Oosting M, Joosten LA, Netea MG, Van Crevel R (2011) Innate immune recognition of *Mycobacterium tuberculosis*. *Clinical and Developmental Immunology* 2011

Krishnan N, Malaga W, Constant P, Caws M, Chau TTH, Salmons J, Lan NTN, Bang ND, Daffe M, Young DB (2011) *Mycobacterium tuberculosis* lineage influences innate immune response and virulence and is associated with distinct cell envelope lipid profiles. *PloS one* 6: e23870

Kutmon M, van Iersel MP, Bohler A, Kelder T, Nunes N, Pico AR, Evelo CT (2015) PathVisio 3: an extendable pathway analysis toolbox. *PLoS computational biology* 11: e1004085

Lakshminarayanan V, Beno DW, Costa RH, Roebuck KA (1997) Differential regulation of interleukin-8 and intercellular adhesion molecule-1 by H₂O₂ and tumor necrosis factor- α in endothelial and epithelial cells. *Journal of Biological Chemistry* 272: 32910-32918

Lala S, Dheda K, Chang J-S, Huggett JF, Kim LU, Johnson MA, Rook GA, Keshav S, Zumla A (2007) The pathogen recognition sensor, NOD2, is variably expressed in patients with pulmonary tuberculosis. *BMC infectious diseases* 7: 96

Latz E, Xiao TS, Stutz A (2013) Activation and regulation of the inflammasomes. *Nature Reviews Immunology* 13: 397

Lee H-M, Shin D-M, Jo E-K (2009) *Mycobacterium tuberculosis* induces the production of tumor necrosis factor- α , interleukin-6, and CXCL8 in pulmonary epithelial cells through reactive oxygen species-dependent mitogen-activated protein kinase activation. *Journal of Bacteriology and Virology* 39: 1-10

Lee H-M, Yuk J-M, Shin D-M, Jo E-K (2009) Dectin-1 is inducible and plays an essential role for mycobacteria-induced innate immune responses in airway epithelial cells. *Journal of clinical immunology* 29: 795

Li J, Chai Q-Y, Zhang Y, Li B-X, Wang J, Qiu X-B, Liu CH (2015) *Mycobacterium tuberculosis* Mce3E suppresses host innate immune responses by targeting ERK1/2 signaling. The Journal of Immunology 194: 3756-3767

Li Q, Lu Q, Bottero V, Estepa G, Morrison L, Mercurio F, Verma IM (2005) Enhanced NF- κ B activation and cellular function in macrophages lacking I κ B kinase 1 (IKK1). Proceedings of the National Academy of Sciences of the United States of America 102: 12425-12430

Li Y, Wang Y, Liu X (2012) The role of airway epithelial cells in response to mycobacteria infection. Clinical and Developmental Immunology 2012

Lin Y, Zhang M, Barnes PF (1998) Chemokine production by a human alveolar epithelial cell line in response to *Mycobacterium tuberculosis*. Infection and immunity 66: 1121-1126

Lo B, Hansen S, Evans K, Heath JK, Wright JR (2008) Alveolar epithelial type II cells induce T cell tolerance to specific antigen. The journal of immunology 180: 881-888

Louis KS, Siegel AC (2011) Cell viability analysis using trypan blue: manual and automated methods. In Mammalian Cell Viability, pp 7-12. Springer

Manca C, Reed MB, Freeman S, Mathema B, Kreiswirth B, Barry CE, Kaplan G (2004) Differential monocyte activation underlies strain-specific *Mycobacterium tuberculosis* pathogenesis. Infection and immunity 72: 5511-5514

Manca C, Tsenova L, Bergtold A, Freeman S, Tovey M, Musser JM, Barry CE, 3rd, Freedman VH, Kaplan G (2001a) Virulence of a *Mycobacterium tuberculosis* clinical isolate in mice is determined by failure to induce Th1 type immunity and is associated with induction of IFN- α /beta. Proceedings of the National Academy of Sciences of the United States of America 98: 5752-7

Manca C, Tsenova L, Bergtold A, Freeman S, Tovey M, Musser JM, Barry CE, Freedman VH, Kaplan G (2001b) Virulence of a *Mycobacterium tuberculosis* clinical isolate in mice is determined by failure to induce Th1 type immunity and is associated with induction of IFN- α /beta. Proceedings of the National Academy of Sciences 98: 5752-5757

Mayer-Barber KD, Andrade BB, Barber DL, Hieny S, Feng CG, Caspar P, Oland S, Gordon S, Sher A (2011) Innate and adaptive interferons suppress IL-1 α and IL-1 β production by distinct pulmonary myeloid subsets during *Mycobacterium tuberculosis* infection. Immunity 35: 1023-1034

Mayer-Barber KD, Andrade BB, Oland SD, Amaral EP, Barber DL, Gonzales J, Derrick SC, Shi R, Kumar NP, Wei W (2014) Host-directed therapy of tuberculosis based on interleukin-1 and type I interferon crosstalk. Nature 511: 99

Mayer-Barber KD, Barber DL, Shenderov K, White SD, Wilson MS, Cheever A, Kugler D, Hieny S, Caspar P, Núñez G (2010) Cutting edge: caspase-1 independent IL-1 β production is critical for host resistance to *Mycobacterium tuberculosis* and does not require TLR signaling in vivo. The Journal of Immunology 184: 3326-3330

McDonough KA, Kress Y (1995) Cytotoxicity for lung epithelial cells is a virulence-associated phenotype of *Mycobacterium tuberculosis*. Infection and immunity 63: 4802-4811

Mortaz E, Adcock IM, Tabarsi P, Masjedi MR, Mansouri D, Velayati AA, Casanova J-L, Barnes PJ (2015) Interaction of pattern recognition receptors with *Mycobacterium tuberculosis*. Journal of clinical immunology 35: 1-10

Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang J-G, Alvarez-Diaz S, Lewis R, Lalaoui N, Metcalf D, Webb AI (2013) The pseudokinase MLKL mediates necroptosis via a molecular switch mechanism. Immunity 39: 443-453

Mvubu NE, Pillay B, McKinnon LR, Pillay M (2017) *Mycobacterium tuberculosis* strains induce strain-specific cytokine and chemokine response in pulmonary epithelial cells. Cytokine

Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, Englert JA, Rabinovitch M, Cernadas M, Kim HP, Fitzgerald KA, Ryter SW, Choi AM (2011) Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. Nat Immunol 12: 222-30

Newton K, Manning G (2016) Necroptosis and inflammation. Annual review of biochemistry 85

Özbek N, Fieschi C, Yilmaz BT, De Beaucoudrey L, Demirhan B, Feinberg J, Bikmaz YE, Casanova J-L (2005) Interleukin-12 receptor β 1 chain deficiency in a child with disseminated tuberculosis. Clinical Infectious Diseases 40: e55-e58

Parandhaman DK, Narayanan S (2014) Cell death paradigms in the pathogenesis of *Mycobacterium tuberculosis* infection. Frontiers in cellular and infection microbiology 4: 31

Pasparakis M, Vandenabeele P (2015) Necroptosis and its role in inflammation. Nature 517: 311

Rajaram MV, Ni B, Morris JD, Brooks MN, Carlson TK, Bakthavachalu B, Schoenberg DR, Torrelles JB, Schlesinger LS (2011) *Mycobacterium tuberculosis* lipomannan blocks TNF biosynthesis by regulating macrophage MAPK-activated protein kinase 2 (MK2) and microRNA miR-125b. Proceedings of the National Academy of Sciences 108: 17408-17413

Ravi R, Bedi A (2002) Requirement of BAX for TRAIL/Apo2L-induced apoptosis of colorectal cancers: synergism with sulindac-mediated inhibition of Bcl-xL. Cancer research 62: 1583-1587

Reuschl A-K, Edwards MR, Parker R, Connell DW, Hoang L, Halliday A, Jarvis H, Siddiqui N, Wright C, Bremang S (2017) Innate activation of human primary epithelial cells broadens the host response to *Mycobacterium tuberculosis* in the airways. PLoS pathogens 13: e1006577

Ryndak MB, Singh KK, Peng Z, Laal S (2015) Transcriptional profile of *Mycobacterium tuberculosis* replicating in type II alveolar epithelial cells. PLoS One 10: e0123745

Saitoh T, Fujita N, Jang MH, Uematsu S, Yang BG, Satoh T, Omori H, Noda T, Yamamoto N, Komatsu M, Tanaka K, Kawai T, Tsujimura T, Takeuchi O, Yoshimori T, Akira S (2008) Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1 β production. Nature 456: 264-8

Sarkar R, Lenders L, Wilkinson KA, Wilkinson RJ, Nicol MP (2012) Modern lineages of *Mycobacterium tuberculosis* exhibit lineage-specific patterns of growth and cytokine induction in human monocyte-derived macrophages. PLoS one 7: e43170

Sasindran SJ, Torrelles JB (2011) *Mycobacterium tuberculosis* infection and inflammation: what is beneficial for the host and for the bacterium? Frontiers in microbiology 2

Sato K, Tomioka H, Shimizu T, Gonda T, Ota F, Sano C (2002) Type II alveolar cells play roles in macrophage-mediated host innate resistance to pulmonary mycobacterial infections by producing proinflammatory cytokines. The Journal of infectious diseases 185: 1139-1147

Sato M, Hata N, Asagiri M, Nakaya T, Taniguchi T, Tanaka N (1998) Positive feedback regulation of type I IFN genes by the IFN-inducible transcription factor IRF-7. FEBS letters 441: 106-110

Sequeira PC, Senaratne RH, Riley LW (2014) Inhibition of toll-like receptor 2 (TLR-2)-mediated response in human alveolar epithelial cells by mycolic acids and *Mycobacterium tuberculosis* mce1 operon mutant. Pathogens and disease 70: 132-140

Sharma M, Sharma S, Roy S, Varma S, Bose M (2007) Pulmonary epithelial cells are a source of interferon- γ in response to *Mycobacterium tuberculosis* infection. Immunology and cell biology 85: 229-237

Shimizu S, Eguchi Y, Kamiike W, Matsuda H, Tsujimoto Y (1996) Bcl-2 expression prevents activation of the ICE protease cascade. *Oncogene* 12: 2251-2257

Sia JK, Georgieva M, Rengarajan J (2015) Innate immune defenses in human tuberculosis: an overview of the interactions between *Mycobacterium tuberculosis* and innate immune cells. *Journal of immunology research* 2015

Sims GP, Rowe DC, Rietdijk ST, Herbst R, Coyle AJ (2010) HMGB1 and RAGE in inflammation and cancer. *Annual review of immunology* 28: 367-88

Stenger S (2005) Immunological control of tuberculosis: role of tumour necrosis factor and more. *Annals of the rheumatic diseases* 64: iv24-iv28

Stucki D, Brites D, Jeljeli L, Coscolla M, Liu Q, Trauner A, Fenner L, Rutaihwala L, Borrell S, Luo T (2016) *Mycobacterium tuberculosis* lineage 4 comprises globally distributed and geographically restricted sublineages. *Nature genetics* 48: 1535

Stutz MD, Ojaimi S, Allison C, Preston S, Arandjelovic P, Hildebrand JM, Sandow JJ, Webb AI, Silke J, Alexander WS (2017) Necroptotic signaling is primed in *Mycobacterium tuberculosis*-infected macrophages, but its pathophysiological consequence in disease is restricted. *Cell Death & Differentiation*: 1

Subbian S, Bandyopadhyay N, Tsenova L, O'Brien P, Khetani V, Kushner NL, Peixoto B, Soteropoulos P, Bader JS, Karakousis PC (2013) Early innate immunity determines outcome of *Mycobacterium tuberculosis* pulmonary infection in rabbits. *Cell Communication and Signaling* 11: 60

Van Golde LM (1985) Synthesis of surfactant lipids in the adult lung. *Annual review of physiology* 47: 765-774

Vignali DA, Kuchroo VK (2012) IL-12 family cytokines: immunological playmakers. *Nat Immunol* 13: 722-8

Walczak H, Miller RE, Ariail K, Gliniak B, Griffith TS, Kubin M, Chin W, Jones J, Woodward A, Le T (1999) Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nature medicine* 5: 157

Wiens KE, Ernst JD (2016) The mechanism for type I interferon induction by *Mycobacterium tuberculosis* is bacterial strain-dependent. *PLoS pathogens* 12: e1005809

Williams MC (2003) Alveolar type I cells: molecular phenotype and development. *Annual review of physiology* 65: 669-695

Xu XJ, Reichner JS, Mastrofrancesco B, Henry WL, Albina JE (2008) Prostaglandin E2 suppresses lipopolysaccharide-stimulated IFN- β production. *The Journal of Immunology* 180: 2125-2131

Yanai H, Matsuda A, An J, Koshiba R, Nishio J, Negishi H, Ikushima H, Onoe T, Ohdan H, Yoshida N, Taniguchi T (2013) Conditional ablation of HMGB1 in mice reveals its protective function against endotoxemia and bacterial infection. *Proceedings of the National Academy of Sciences of the United States of America* 110: 20699-704

Yu C-H, Micaroni M, Puyskens A, Schultz TE, Yeo JC, Stanley AC, Lucas M, Kurihara J, Dobos KM, Stow JL (2015) RP105 engages phosphatidylinositol 3-kinase p110 δ to facilitate the trafficking and secretion of cytokines in macrophages during mycobacterial infection. *The Journal of Immunology* 195: 3890-3900

Table 1. The list of up-regulated, down-regulated and unchanged genes in A549 infected cells by two *M.tb* lineages in comparing to the uninfected cells at 72 hrs p.i.

Up-Down Regulation (comparing to control group)									
Symbol	Group1 (Sublineage L3-CAS1- Delhi)		Group2 (Sublineage L 4.5)		Symbol	Group1 (Sublineage L3- CAS1- Delhi)		Group2 (Sublineage L 4.5)	
	Fold Regulation	P Value	Fold Regulation	P Value		Fold Regulation	P Value	Fold Regulation	P Value
AGT	-2.0373	0.116117	-1.4306	0.25332	HRAS	3.2792	0	6.6807	3.00E-06
AKT1	-33.3589	0.016968	-6.7116	0.02503	HSPA1A	10.5805	0	4.0093	4.50E-05
ATF1	1.0070	0.830182	3.1748	0.00082	HSPD1	7.3445	0.000001	9.7585	4.00E-06
BCL2A1	3.1095	0	4.7131	0	HMOX1	-1.6208	0.052295	20.9179	0
BCL2L1	-24.0839	0.008385	-2.8350	0.02866	IFNA1	-2.1435	0.000673	1.1620	0.063039
BCL3	-12.0977	0.021229	-1.6170	0.16136	IFNB1	1.1865	0.031514	-4.7131	0.00015
BIRC2	-1.5728	0.227905	2.3027	0.03917	IFNG	-24.0284	0.000068	-9.4261	0.00009
BIRC3	1.2570	0.692297	4.0746	0.00077	IKBKB	-5.5919	0.024808	2.0801	0.00012
BTk	-4.2575	0.038407	-440.6023	0.00065	IKBKG	-10.8028	0.003	-1.2114	0.24391
CARD11	-9.7360	0.334011	1.0892	0.51556	ICAM1	-1.0000	0.83439	4.6161	5.60E-05
CASP1	2.4623	0.135402	-1.9009	0.20077	IL12A	-1.7411	0.000381	2.0562	0.000521
CASP8	-4.2575	0.038407	-1.1514	0.47084	IL1A	2.2038	0.000002	3.1602	8.00E-06
CCL2	2.0232	0.037661	5.9932	8.20E-05	IL1B	-2.0139	0.000848	-1.7371	0.0016
CD14	4.7240	0.000014	-1.217	0.13897	IL1R1	-10.7779	0.03236	-2.4680	0.08865
CD180	-24.0284	0.000068	-9.4261	0.00009	IL2	-24.0284	0.000068	-9.4261	0.00009
CD80	-3.6301	0.00021	-5.8563	0.00012	IL6	-1.8790	0.018595	-1.2924	0.12789
CD86	-24.0284	0.000068	-9.4261	0.00009	IL10	-9.7585	0.000538	-23.8624	0.000424
CLEC4E	2.4453	0.000053	2.2974	0.00027	IRAK1	1.3947	0.074013	1.1355	0.50341
CCL5	4.3570	0	1.9925	0	IRAK2	-2.4623	0	3.3173	0.00055
CD40	-1.3597	0.133717	1.3980	0.07428	IRAK4	-2.7258	0.002234	-2.2553	0.00358
CFLAR	-1.5440	0.068564	1.3755	0.08381	IRF1	-5.7094	0.008851	-1.8747	0.04961
CHUK	2.2868	0.00003	-1.3566	0.004702	IRF3	2.2243	0.000001	5.3270	0
CSF1	-4.0371	0.058892	1.2426	0.77318	JUN	3.1456	0.000007	12.5824	0
CSF2	-2.4453	0.00001	1.7053	8.00E-06	LTA	-2.4396	0.002354	-4.3469	0.00087
CSF3	-3.7235	0.00085	2.2868	0.000339	LY86	-1.6396	0.002234	-2.8945	0.00032
CXCL10	-2.1287	0.000737	1.0140	0.86249	LY96	-1.1487	0.077616	2.3839	1.60E-05
CCL20	2.3620	0.000169	3.0809	5.20E-05	LTBR	-4.3369	0.031	-1.1147	0.52452
CD83	-1.1045	0.153774	-1.3755	0.00658	MALT1	-1.0570	0.644815	2.0753	0.00519
CXCL2	2.9349	0.001168	5.3765	4.50E-05	MAP2K3	-2.0753	0.007287	1.7171	0.00323
CXCL3	2.1585	0.000652	3.5145	3.40E-05	MAP2K4	8.8152	0	13.3614	3.00E-06
CXCL8	6.3643	0.000015	20.9663	0	MAP3K1	4.5842	0.000008	6.6499	0

ECSIT	-1.7818	0.003502	-1.6208	0.00621	MAP3K7	1.5227	0.00486	2.8945	2.40E-05
EIF2AK2	6.3643	0	3.7235	0.00013	MAP4K4	12.1257	0	20.6299	3.60E-05
ELK1	-3.8996	0.00079	-3.7149	0.00085	MAPK8	-1.2864	0.000018	1.9009	0
EGFR	-3.5472	0.031121	1.2864	0.41303	MAPK8IP3	-13.1167	0.000157	-12.3234	0.00016
EGR1	-2.1435	0.000352	4.4898	1.00E-06	MYD88	-2.2658	0.015169	2.8024	0.05435
FASLG	-1.9954	0.396602	1.4109	0.65753	NFKB1	-2.6635	0.061876	2.4852	0.00735
FADD	-3.325	0.044358	36.7583	0.00131	NFKB2	-3.3326	0.000909	2.1735	0.00014
FOS	44.0173	0	31.0531	0.00055	NFKBIE	-2.5847	0.002523	2.3729	0.00017
HMGB1	-6.2046	0.000005	-2.6268	2.20E-05	NFKBIA	13.8646	0	2.4737	0.00392
PSIP1	-3.9724	0.040745	-2.0753	0.09633	TLR3	1.454	0	3.0596	0.01561
PELI1	12.7286	0	8.2059	4.00E-06	TLR4	-7.3785	0.000558	-9.6911	0.00048
PPARA	-3.1383	0	-3.8459	0	TLR5	1.0842	0.231596	-2.7511	0.00035
PRKRA	2.4794	0.000165	4.0652	0.00038	TLR6	-12.1538	0.005031	-5.2054	0.00785
PTGS2	40.8803	0	84.4485	1.00E-06	TLR7	-3.3173	0.000237	-8.7341	9.30E-05
RAF1	-2.6268	0.000002	1.7942	1.00E-06	TLR8	-11.6049	0.000082	-9.4261	0.00009
REL	-5.0865	0.044454	4.6268	3.70E-05	TLR9	-3.0455	0.000276	1.651	0.0011
RELA	7.8173	0.000002	18.0009	0.0001	TNF	-6.774	0	5.9244	0.00193
RELB	-4.5948	0.038984	-1.2512	0.397	TIFA	-1.3915	0.198832	-1.2058	0.36188
RIPK2	1.1947	0.108534	2.835	0.00208	TNFRSF10 B	-2.4737	0.047724	2.4284	0.00434
RHOA	-1.2454	0.048335	1.5263	0.00232	TNFRSF10 A	-8.7949	0.010248	2.1337	0.006051
RIPK1	-3.5554	0.03545	1.2002	0.64688	TNFRSF1 A	-7.7097	0.002488	-4.4076	0.00383
SARM1	1.6663	0.00034	4.357	0	TNFSF10	2.0753	0.000048	-2.5315	0.00036
STAT1	1.8747	0.012517	-1.1755	0.37894	TRADD	-5.8428	0.037202	-2.308	0.09517
SIGIRR	-6.3643	0.001906	-10.9536	0.00145	TRAF2	-21.6556	0.007317	-2.4116	0.0338
STX11	1.6663	0.008282	2.7574	0.00023	TOLLIP	-4.7349	0	-4.4691	0
TAB1	-6.5735	0.000005	-5.2295	6.00E-06	TRAF6	-1.3441	0.005839	2.514	0.00024
TBK1	-4.0093	0.004566	-2.1886	0.013833	TRAF3	-11.4188	0.004246	-1.9862	0.0309
TICAM1	-2.7132	0.000017	2.9828	1.00E-06	TNFAIP2	-16.374	0.01594	-8.1492	0.01961
TICAM2	4.7459	0.000001	5.0281	1.00E-06	TNFAIP3	-4.0558	0.019691	1.5018	0.10394
TIRAP	-9.8264	0.000079	-7.8717	8.90E-05	NFKBIL1	26.9709	0	2.514	0.00023
TIMP1	-2.0186	0.068338	-1.0281	0.72517	NFRKB	1.7291	0.000317	4.3169	2.70E-05
TLR1	1.1173	0.118853	2.395	0.02185	NR2C2	2.7195	0.000872	4.0746	0.00013
TLR10	-15.4907	0.000075	-9.4261	0.00009	NOD1	-2.4284	0.015252	-1.021	0.77791
TLR2	-8.9797	0.000092	1.651	0.0011					

691
692
693
694
695
696
697
698

Figures

Figure Legends:

Figure 1. The canonical pathway for “TLRs and NF- κ B Signalling” at 72 hrs p.i. in **A)** Sublineage L4.5 infected A549; **B)** Sublineage L3-CAS1-Delhi infected A549 cell line at MOI of 50. Red colour indicates genes which were up-regulated in infected cells in compare to the mock cells. Blue colour indicates genes which were down regulated in infected cells in compare to the mock cells. Purple colour indicates unchanged genes.

Figure 2. Volcano plot of genes involved in TLRs and NF- κ B signalling pathways. This plot displays the statistical significance versus fold-change on the y- and x-axes, respectively.

Figure 3. Clustergram plot of genes involved in TLRs and NF- κ B signalling pathways. This clustergram performs non-supervised hierarchical clustering of the entire dataset to display a heat map with dendrograms indicating co-regulated genes.

Figure 4. Evaluated cytokines and chemokines production infected cells with Sublineage L 4.5 (lineage 4) and L3-CAS1-Delhi (lineage 3) strains, at an MOI~50 :1 by ELISA assay. * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0001$. The results are shown as the mean + SD of duplicate measurements.

Figure 5. **A)** Intracellular growth of different studied strains of *M.tb* in A549 cell line; **B)** Graph shows the percentage of infected cells by two lineages; the intracellular internalization index was classified into 1–5 bacilli, 6–10 bacilli and more that 10 bacilli per cell; **C)** Auramine staining of the *M. tb*-infected cells.

TLRs

Group 1 vs. Control Group

Lineage3

Group 2 vs. Control Group

Lineage4

NFkB

Group 1 vs. Control Group

Group 2 vs. Control Group

TLRs

NFkB

bioRxiv preprint doi: <https://doi.org/10.1101/298862>; this version posted April 26, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

c