1 Identification of novel genes including *rpmF* and *yjjQ* critical for Type II 2 persister formation in *Escherichia coli*

3 Shuang Liu¹, Nan Wu¹, Shanshan Zhang¹, Yumeng Zhang¹, Wenhong Zhang^{1*} and
4 Ying Zhang^{1,2*}

5 ¹ Key Lab of Molecular Virology, Institute of Medical Microbiology, Department of
6 Infectious Diseases, Huashan Hospital, Fudan University, Shanghai, China,

7 ² Department of Molecular Microbiology and Immunology, Bloomberg School of Public
8 Health, Johns Hopkins University, Baltimore, MD, USA

9 * Correspondence:

10 Ying Zhang
11 yzhang@jhsph.edu

12 WenHong Zhang
13 zhangwenhong@fudan.edu.cn

14
15 **Keywords:** persister gene, knockout mutant, key time point, transcriptome, ribosomal
16 protein

17 Abstract

18 Persister cells, which are characterized by inactive metabolism and tolerance to antibiotics or
19 stresses, pose a significant challenge to the treatment of many persistent infections. Although
20 multiple genes have been reported to be involved in persister formation through transposon
21 mutant library screens, how persisters are formed during the natural process of persister
22 formation as the culture transitions from log phase to stationary phase is unclear. Here, using
23 *E. coli* as a model, we performed a comprehensive transcriptome analysis of gene expression
24 profiles of successive cultures of an *E. coli* culture at different critical time points, starting
25 from persister-free S1-nonexistence phase (3h) to persister appearing S2-emergence phase
26 (4h), and persister abundant stage S3-abundance phase (5h). The differentially expressed
27 genes (≥ 2 -fold) in persister appearing stage (S1 to S2 transition) and persister abundant stage
28 (S1 to S3) were compared, and 51 and 29 genes were identified to be up-regulated,
29 respectively. Importantly, 13 genes (*gnsA*, *gnsB*, *ybfA*, *yjjQ*, *ymdF*, *yhdU*, *csgD*, *yncN*, *rpmF*,
30 *ycdX*, *yohJ*, *ssrA*, *rbsD*) overlap in both persister S2-emergence phase and S3-abundance
31 phase, including a member of the trans-translation pathway (*ssrA*) as well as an orphan toxin
32 (*ycdX*), which are two well-known persister genes while the remaining 11 novel genes (*gnsA*,
33 *gnsB*, *ybfA*, *yjjQ*, *ymdF*, *yhdU*, *csgD*, *yncN*, *rpmF*, *yohJ*, *rbsD*) have not been reported
34 previously. Persister levels of 7 constructed knockout mutants ($\Delta gnsA$, $\Delta ybfA$, $\Delta yjjQ$, $\Delta yhdU$,
35 $\Delta csgD$, $\Delta yohJ$ and $\Delta rpmF$) and 10 overexpression strains (*gnsA*, *gnsB*, *ybfA*, *yjjQ*, *ymdF*,
36 *yhdU*, *csgD*, *rpmF*, *yohJ*, *rbsD*) in *E. coli* uropathogenic strain UTI89 were determined upon
37 treatment with different cidal antibiotics (ampicillin, levofloxacin and gentamicin).
38 Additionally, ranking of these overlapping genes according to their impact on persister levels

were also performed. Two genes (*rpmF* encoding 50S ribosomal subunit protein L32, and *yjjQ* encoding a putative LuxR-type transcription factor) showed the most obvious phenotype on persister levels in both knockout and overexpression studies, which suggests they are broad and key factors for persister formation. While previous studies cannot distinguish if a given persister gene is involved in persister formation or persister survival, our findings clearly identify novel persister forming genes and pathways involving a ribosome protein and a LuxR type transcription factor during the bona fide persister formation process and may have implications for developing improved treatment of persistent infections.

Introduction

A small percentage of bacterial cells can survive antibiotic or other stress-induced cell death by entering a transient dormant or slow-growing state (Kint et al., 2012; Lewis, 2010; Zhang, 2014), and this phenomenon is termed persistence and was first discovered by Hobby (1942) and Bigger (1944). Persistence is considered to be associated with chronic and recalcitrant bacterial infections which can ratchet up risks for antibiotic resistance and pose a grave threat to human health (Mulcahy et al., 2010; Zhang et al., 2012; Zhang, 2014; Van den Bergh and Michiels, 2016). Persister cells are believed to form either randomly or as an induced product (Balaban et al., 2004; Nierman et al., 2015; Van den Bergh et al., 2017). Persister cells can be divided into two types, type I persisters and type II persisters. Type I are generated from stationary phase while type II persisters are continuously generated through the whole bacterial growth stage as the culture grows from log phase to stationary phase (Balaban et al., 2004). Multiple molecular mechanisms have been proposed to be involved in the formation of persister (Li and Zhang, 2007; Dorr et al., 2009; Ma et al., 2010; Wang and Wood, 2011; Vega et al., 2012a; Li et al., 2013; Maisonneuve et al., 2013; Marques et al., 2014; Wu et al., 2015) (stringent response, toxin–antitoxin (TA) systems, SOS response, global regulators, signaling molecules, energy metabolism).

Although various persister pathways have been identified, they are mainly studied at one time point of bacterial growth stage, and little is known about the overall dynamic changes of gene expression profile associated with type II persister cell formation naturally during growth cycle from “nonexistence” to “emergence” and finally “abundance”, which we define them as S1, S2, S3, respectively, in this study. Here, using transcriptional analysis by RNA-seq, we provide a “dynamic evolution” of genes when type II persister cells are formed from non-persister cells. We identified many genes that have not been previously reported that are involved in type II persister formation. Fifty-three and thirty-two genes were found to have significantly different expression in S2 and S3 compared with S1, respectively. Specially, 13 genes overlap in both comparison groups. Apart from 4 genes (*gnsB*, *ymdF*, *yncN*, *ybfA*) with uncharacteristic functions and 2 genes (*ssrA*, *ycdX*) with definite effect on persistence (Li et al., 2013; Islam et al., 2015), the remaining 7 overlapping genes were mapped to ribosomal protein (*rpmF*), regulator of phosphatidylethanolamine synthesis (*gnsA*), membrane protein (*yohJ*, *yhdU*), DNA-binding transcriptional activator (*csgD*), transcription factor (*yjjQ*) and cytoplasmic sugar-binding protein (*rbsD*). We confirm two genes (*rpmF* and *yjjQ*) play crucial roles in persister formation under multiple antibiotic treatment and stress conditions. Together, our data provide a dynamic profile of genes involved in type II persister formation

82 which leads to novel insight on bona fide mechanisms of persister formation.

83 **Methods**

84 **Bacteria and culture conditions**

85 *E. coli* K12 strain W3110 from glycerol stocks at -80 °C were 1:1000 diluted in Luria-Bertani
86 (LB) broth (10 g Bacto-tryptone, 5 g yeast extract, and 10 g NaCl/liter) without antibiotics
87 and incubated at 37 °C for 16 hours. For transcriptomics, the overnight culture was diluted
88 1:10⁵ inoculated in 3.5 liters, 2 liters and 1 liter fresh LB medium for each time point (3 hrs, 4
89 hrs and 5 hrs) to satisfy the quantity requirement for library construction and grown at 37 °C
90 (100 rpm).

91 **RNA isolation and preparation of cDNA libraries and RNA-sequencing**

92 Samples of each time point were collected from three independent replicates. Briefly, cells
93 grown to the indicated time points were harvested by centrifugation at 4°C (4000g x 10 min)
94 in 50ml centrifuge tubes and cell cultures were kept on ice during the whole harvest process.
95 Samples were then preserved at -80 °C until RNA was extracted. Total RNA was extracted
96 from the three samples using TRIzol® reagent according to the manufacturer's instructions
97 (Invitrogen, USA). Integrity of RNA was determined by 2100 Bioanalyzer (Agilent
98 Technologies, USA). RNA was purified using oligo (dT) beads (Illumina, San Diego, CA,
99 USA). Fragmentation buffer was then added to cut the long mRNA into small pieces. The
100 mRNA short fragments were primed with random hexamer primers for synthesis of
101 single-stranded cDNAs which were adopted as the templates for synthesis of the second cDNA
102 strand. The double-stranded cDNAs were purified using the QiaQuick PCR extraction kit
103 (Qiagen). The purified cDNAs were then processed in end repair and appended with poly(A).
104 The appropriate cDNA fragments were obtained through agarose gel electrophoresis and
105 amplified using PCR. The cDNA libraries were sequenced using an Illumina HiSeq™ 2500
106 (OE Biotech Company, Shanghai, China) and produced 125 bp double-end reads.

107 **Transcriptome bioinformatic analysis**

108 The raw reads (FastQ format) were qualified by NGS QC Toolkit to abandon end joints,
109 low-quality bases and N-base which resulted in the “clean reads”. All the subsequent analysis
110 was based on the “clean reads”. Reads were mapped to the *E. coli str. K12 substr. W3110*
111 genome (NC_007779.1; NCBI) using Tophat (<http://tophat.cbcb.umd.edu/>). Then FPKM and
112 count values of the matched reads were obtained using eXpress (Mortazavi et al., 2008).
113 These values were normalized by DESeq (2012) R package (Siska and Kechris, 2017)
114 (estimate Size Factors). Subsequently, P values and fold changes were calculated using
115 nbinomTest. Differential expression genes (DEGs) with P value less than 0.05 were selected
116 for KEGG and COG analysis to determine the gene functions and pathways. Finally,
117 expression profile of DEGs in different samples was displayed using heat map.

118 **Validation of RNA-seq data by quantitative real-time PCR**

Thirty DEGs were chosen randomly and processed in qRT-PCR to validate the RNA-seq data. cDNAs in qRT-PCR were synthesized from the same RNA extractions used for the RNA-seq using a Prime Script RT Reagent Kit (TaKaRa, Japan) and then 1:5 diluted. *rrsB* was selected as the control gene. qRT-PCRs were carried out in ABI 7500 Real-Time instrument thermocycler (Applied Biosystems, Foster City, CA, USA). Relative fold changes were worked out using the $2^{-\Delta\Delta C_t}$ method.

Knockout mutant construction and persister assay

Deletion of persister genes made in the UTI89 background was achieved by using the λ Red recombination systems, as previously described by Datsenko and Wanner (Datsenko and Wanner, 2000). The deleted genes were stably replaced with a chloramphenicol resistance cassette. All mutants were confirmed with PCR and sequencing (Biosune, Shanghai, China). Persistence was measured by determining the bacterial survival as colony-forming units (CFUs) per 1mL after exposure to cidal antibiotics, i.e., 200 μ g/mL ampicillin, or 5 μ g/mL levofloxacin, or 40 μ g/mL gentamicin. Following overnight growth, 1ml cultures were transferred to a 1.5 ml Eppendorf tube and immediately treated with the above antibiotics and incubated at 37°C without shaking for different times. To confirm the persister levels of samples for transcriptome assay, 1 ml culture was withdrawn from the culture of each time point and treated with ampicillin (100 μ g/ml) for 3 hours. The cell viability was measured by samples withdrawn at the desired time points, washed and serially diluted in PBS, followed by inoculation onto LB agar without antibiotics. The CFU numbers were counted after overnight incubation at 37°C.

Construction of overexpression mutant strains

Eleven overlapping genes were amplified by PCR and the primers flanked by *Nco*I and *Eco*RI restriction enzyme sites are listed in Table S1. Following double-enzyme digestion, the resultant DNA fragments were cloned into the same restriction sites of plasmid vector pBAD202, resulting in a series of recombinant plasmids with kanamycin resistance cassette. After selection on kanamycin-containing agar plates and confirmation by DNA sequencing, the resulting constructs were transformed into uropathogenic *E. coli* strain UTI89 and designated as the overexpression mutant strains.

Measurement of cellular ATP concentrations

The cellular ATP concentrations of wild type and knockout mutants (or overexpression strains) were determined using an ATP assay kit according to the manufacturer's protocol.

Results

Determination of persister levels at three time points under ampicillin treatment

To obtain the gene expression profile of associated with persister formation, the emergence of persisters under ampicillin treatment at different time points was performed. To minimize the

influence of type I persister cells in the inoculum, we inoculated the overnight culture in fresh LB broth at a high dilution of $1:10^5$ and incubated in 37°C (100 rpm) for different times. To detect the persister levels, 1 ml bacterial culture from different ages was withdrawn and treated with ampicillin ($100\ \mu\text{g/ml}$) for the same length of time (3 hours). We found no viable cells or persisters existed if the age of cultures was 3hrs or younger. However, when the bacterial culture age was extended to 4 hrs, persister cells started to emerge, but with a number of less than 10 CFU/ml. As we prolonged the incubation time to 5 hrs, there was a surge in the number of persister cells (see Fig. S1). Based on the above results, we defined the three ampicillin-associated critical time points (3hrs, 4hrs and 5hrs) as S1 “nonexistence” to S2 “emergence” and S3 “abundance”, respectively. Bacteria of S1/S2/S3 were then analyzed using RNA-seq to identify the genes differentially expressed in persister emergence and abundance stages S2 and S3 compared with no persister stage S1.

Identification and verification of differentially expressed genes (DEGs) during persister formation

The up- or down-regulated genes (S2 vs. S1, and S3 vs. S1) whose p values were below 0.05 were identified (Table 2). Thirteen overlapping DEGs of S2/S1 and S3/S1 were chosen to validate the reliability of RNA-seq using quantitative real-time PCR (RT-PCR). As a result, all the genes we tested presented concordant expression patterns in RT-PCR and RNA-seq. The results presented here indicate the data from RNA-seq is credible in the follow-up analysis (see Table 1).

Core DEGs in S2 versus S1 and S3 versus S1

S1 “nonexistence” to S2 “emergence” and S3 “abundance” referred to the three different stages of persister formation, therefore DEGs of S2 vs. S1 and S3 vs. S1 would allow us to discover critical genes in the formation of persisters. Fifty-one and twenty-nine DEGs that showed significantly elevated levels (≥ 2 -fold increase) were identified based on the values of S2/S1 and S3/S1 comparison, respectively (see Table 2). Nine of the 51 genes up-regulated in S2 vs. S1 encoded uncharacterized or hypothetical proteins (*ydcA*, *ymdF*, *ybfA*, *gnsB*, *yncN*, *ybfH*, *yobB*, *ygeP*, *yhhP*, *ymcB*). Of the remaining genes, 7 were assigned to metabolism pathways according to the KEGG analysis and they are phosphatidylethanolamine synthesis (*gnsA*), fatty acid metabolism (*fadE*, *yfcY*), carbon metabolism (*idnK*), porphyrin and chlorophyll metabolism (*cobS*), amino sugar and nucleotide sugar metabolism (*glmU*) and pyruvate metabolism (*ldhA*), respectively. Other pathways such as trans-translation (*ssrA*), ATP-binding cassette (ABC) transporters (*rbsA*, *rbsD*), and two-component system (*torR*, *torT*, *degP*) were also involved in the formation of persisters. In addition, genes encoding toxin (*ydcX*), ribosomal protein or ribosomal associated proteins (*rpmF*, *yfiA*), phage shock protein (*pspC*), small heat shock proteins (*ibpB*, *ibpA*), membrane proteins (*yhdU*, *yohJ*, *yebN*, *yfdY*, *fxsA*, *yohK*, *yjfV*), transcriptional activators or factors (*yjjQ*, *csgD*, *arsR*, *pspC*, *ybeF*, *yagI*), many enzymes: ATPase (*zntA*), dehydrogenase (*idnD*), arsenate reductase (*arsC*), oxidoreductase (*ykgE*), transposase (*yhhI*) and fructose-6-phosphate aldolase 2 (*fsaB*). Moreover, fimbrial-like protein (*yfcV*, *yadK*), IS2 insertion element repressor (*insC*), cytochrome c-type protein (*torY*) also showed significant increase in S2 compared with S1

(see Fig. 1A). Of these genes, phage shock protein (*pspC*) has been reported to be involved in indole-mediated persister formation (Vega et al., 2012b). And *ycdX* also participates in persister formation as an orphan toxin (Islam et al., 2015), and *ssrA* involved in trans-translation has previously been shown to be involved in persister formation (Li et al., 2013).

During the period of “nonexistence to abundance”, the number of DEGs (≥ 2 -fold increase) was only 29 (see Table 3). Thirteen of them overlapped with those of S2/S1 while the other 16 genes encode proteins that are involved in arginine and proline metabolism (*argC*, *argG*, *argD*), glycerophospholipid metabolism (*glpD*), propanoate metabolism (*tdcD*), DNA-binding transcriptional regulator (*envR*), ABC transporters (*ccmD*), amino-acid transporter (*yeeF*), antimonite transporter (*arsB*), ribosome modulation factor (*rmf*), regulatory peptide (*mokB*), oxidoreductase (*ymjC*), membrane protein (*yohM*), inner membrane lipoprotein (*yghJ*), cytoplasmic ferritin iron storage protein (*ftn*) and uncharacterized protein (*yhfG*) (see Fig. 1B). As for the overlapping genes, 4 genes (*ymdF*, *ybfA*, *yncN*, *gnsB*) encode putative or uncharacterized proteins. Other overlapping genes were related to ribosomal protein (*rpmF*), toxin (*ycdX*), membrane proteins (*yohJ*, *yhdU*), DNA-binding transcriptional activator or transcriptional factors (*csgD*, *yjjQ*), trans-translation (*ssrA*), phosphatidylethanolamine synthesis (*gnsA*) and ABC transporters (*rbsD*). The 13 overlapping genes of S2/S1 and S3/S1 were supposed to be of significance for both “boot-up” and “shoot-up” of persister formation and required a thorough investigation. Apart from two genes (*ssrA* and *ycdX*) which have been known to participate in the persister formation (Li et al., 2013; Islam et al., 2015), the other 11 overlapping genes have not been reported to be related to persistence.

Persister levels of the overlapping gene knockout mutants in the presence of ampicillin, gentamicin and levofloxacin

Because *ssrA* and *ycdX* have previously been shown to be involved in persister formation (Li et al., 2013; Islam et al. 2015), we focused on assessing the impact of the knockout strains on persister levels of the remaining 11 of the 13 previously unreported overlapping genes under ampicillin treatment. However, only 7 knockout mutants ($\Delta gnsA$, $\Delta ybfA$, $\Delta yjjQ$, $\Delta yhdU$, $\Delta csgD$, $\Delta yohJ$ and $\Delta rpmF$) were successfully constructed presumably because some genes are either essential, or lack of sequence homolog (i.e. *yncN*) in UTI89 strain. We chose to construct the mutants in uropathogenic strain UTI89 because it is ureopathogenic strain that is conducive to further in-vivo experiments. Cells of wild type and knockout mutants from log phase ($\sim 10^8$ CFU/ml) were challenged with ampicillin (200 μ g/ml), gentamicin (40 μ g/ml) and levofloxacin (5 μ g/ml). Persister numbers were determined at 3 hrs, 5 hrs and 24 hrs, respectively. Of the 7 knockout mutants tested, *rpmF*, which encodes the 50S ribosomal subunit protein L32, manifested the most obvious phenotype in persister assays with all three antibiotics. After 24 hrs of ampicillin exposure, the $\Delta rpmF$ mutant had $\sim 10^5$ -fold decrease from the initial CFU numbers (10^8 CFU/ml) while UTI89 had 10^4 -fold decrease from 10^8 CFU/ml (see Fig. 2A). When exposed to gentamicin and levofloxacin, $\Delta rpmF$ also exhibited lower persister levels compared with UTI89 (10~100-fold change compared with UTI89) (see Fig. 2B and C). $\Delta ybfA$, also had dramatic decreased persister levels (10~1000-fold

decrease) in the presence of ampicillin as well as gentamicin compared with UTI89. While under levofloxacin treatment, *ΔybfA* showed a negligible effect on persister levels. *ΔgnsA*, though exhibited significantly lower persister levels compared with UTI89 (10~10³-fold decrease) when exposed to gentamicin and levofloxacin, showed a higher persister level (>10-fold increase) in the presence of ampicillin. *ΔyhdU* also showed a 100-fold increase in persister numbers when exposed to ampicillin. However, under gentamicin and levofloxacin treatment, *ΔyhdU* exerted little impact on persister numbers. Other knockout mutant strains, including *ΔcsgD* and *ΔyohJ*, exhibited similar persister levels to UTI89 under ampicillin and levofloxacin treatment. However, persister levels of *ΔcsgD* and *ΔyohJ* were dramatically decreased when exposed to gentamicin (~10⁶-fold decrease for *ΔcsgD* and ~10-fold decrease for *ΔyohJ*). However, when tested in the presence of levofloxacin and gentamicin, instead of higher persister level, *ΔyhdU* showed the similar persister number as UTI89 and *ΔgnsA* exhibited 10~100-fold reduction in persister level. We did not observe any alteration of persister numbers for *ΔyjjQ* compared with the wild type under our experiment conditions. The impact of *ybfA* exerted on persister formation was specific to ampicillin and gentamicin while *ΔcsgD* and *ΔyohJ* affected persister formation only in the presence of gentamicin. Contrary to our expectation, our data revealed two unique knockout mutants (*ΔgnsA* and *ΔyhdU*) which displayed 10~100-fold higher persister numbers than UTI89 when exposed to ampicillin. And *ΔgnsA* even exerted opposite effect on persister formation under different antibiotic treatment. This seemingly paradoxical result observed for *ΔgnsA* was not unexpected, instead, it may indicate that its role is to suppress persister formation. Overall, the data presented above revealed only one knockout mutant, *ΔrpmF* mutant exhibited a universal effect on persister formation for all the antibiotics tested.

Effect of overexpression of the 10 overlapping genes on persister levels

To further investigate whether these genes participate in persister formation, we constructed 10 overexpression strains and subjected them to treatment with ampicillin (200 μg/ml), levofloxacin (5 μg/ml) and gentamicin (50 μg/ml). UTI89 transformed with pBAD202 plasmid was used as a control. pBAD202 plasmids containing the 10 genes were constructed and 0.2% arabinose was added to induce the gene expression. Twenty-four hours after ampicillin treatment, 7 overexpression strains (*gnsA*, *gnsB*, *yhdU*, *csgD*, *yohJ*, *rpmF* and *rbsD*) had 10~10⁴-fold higher persister levels than the control strain (see Fig. 3A). Contrary to the RNA-seq data, two genes (*yjjQ* and *ymdF*) exhibited negative effect on persister formation (see Fig. 3A), while *ybfA* showed no influence on persister levels (see Fig. 3A). Collectively, the overexpression study under ampicillin treatment further confirmed the validity of the transcriptome assay, though two genes (*yjjQ* and *ymdF*) showed the opposite effect. These overexpression strains were also exposed to gentamicin and levofloxacin. Fig. 3B shows in the presence of gentamicin, 7 overexpression mutants (*gnsB*, *yhdU*, *csgD*, *rpmF*, *rbsD*, *yjjQ* and *ymdF*) displayed consistent effect with what they did when exposed to ampicillin. Overexpression of the 5 genes could lead to 10~100-fold increase in persister formation while *yjjQ* and *ymdF* still had negative effect on persister formation (100~1000-fold decrease compared with UTI89+pBAD202). *gnsA*, though could increase persister level by more than 100-fold, had negative effect on persister formation (~2.5-fold decrease) when exposed to gentamicin. Overexpression of *ybfA* which had little effect on persister level under ampicillin

treatment could increase persister numbers by 4-fold in gentamicin treatment. *yohJ* had little influence on persister formation. When exposed to levofloxacin, increased persister levels compared with the control strain were observed for only 3 overexpression strains (*gnsA*, *ybfA* and *ymdF*), with ~4-fold for *ybfA* and >10-fold for *gnsA* and *ymdF* (see Fig. 3C). Other genes, except for *yjjQ* and *yohJ* which exhibited >100-fold and ~4-fold decreased persister levels, had no obvious effect on persister formation. Notably, in the presence of levofloxacin, only two genes (*gnsA* and *yjjQ*) had the same effect on persister formation as observed in the presence of ampicillin. Collectively, our data suggest that of the genes *gnsA*, *gnsB*, *ybfA*, *yhdU*, *csgD*, *yohJ*, *rpmF* and *rbsD* conducive to persister formation, 5 genes (*gnsB*, *yhdU*, *csgD*, *rpmF* and *rbsD*) were specific to ampicillin and gentamicin, one gene (*gnsA*) was ampicillin- and levofloxacin-specific, one gene (*ybfA*) was gentamicin- and levofloxacin-specific and one gene (*yohJ*) was specific only to ampicillin. Of the genes (*yjjQ*, *ymdF* and *yohJ*) which exhibited negative role in persister formation, *yjjQ* had a general effect for all the three antibiotics, while *ymdF* was specific to ampicillin and gentamicin, and *yohJ* was specific to levofloxacin.

Ranking of the overlapping genes according to their impact on persister levels under antibiotic exposure

To determine the relative importance of the 10 overlapping genes (*gnsA*, *gnsB*, *ybfA*, *yjjQ*, *ymdF*, *yhdU*, *csgD*, *rpmF*, *yohJ*, *rbsD*) under exposure to all antibiotics, the results (≥ 5 -fold change) from knockout and overexpression assay were gathered and ranked, respectively. Among the 8 knockout mutants, *rpmF* was shown to play a key and broad role in the process of persister formation with 3 points under treatment with all the three antibiotics. The second most important genes were *ybfA* and *gnsA* with 2 scores. Although *gnsA* exhibited significant effect on persister formation under all the three antibiotics, its knockout mutant showed an enhancement in persister level under ampicillin treatment, which has an opposite effect in the presence of gentamicin and levofloxacin. *yohJ*, *csgD* and *yhdU* scored only 1 point, which suggests that their impact on persister formation is limited to only one antibiotic. Unfortunately, *yjjQ* scored 0 in its knockout mutant (see Table 4). Table 5 shows a distinct gene arrangement for the 10 overexpression strains. *yjjQ* overexpression strain showed a broad impact on persister formation and scored 3 points. Overexpression strains of seven genes (*ymdF*, *gnsA*, *gnsB*, *rpmF*, *yhdU*, *csgD* and *rbsD*) made 2 points. Among these genes, *ymdF* showed significant effects on persister levels upon treatment with all the three antibiotics. However, the effect *ymdF* exerted in the presence of ampicillin and gentamicin was opposite to that in the presence of levofloxacin. *yohJ* received 1 point only when exposed to ampicillin while overexpressing *ybfA* and *yncN* resulted in similar persister levels to the control strain. Furthermore, we also observed some gene knockout mutants exhibited a different persistence profile when compared with their overexpression strains.

Relative ATP concentration assay

Since ATP levels correlate with persister levels, we also analysed the ATP levels for knockout or overexpression strains. Figure 4A and B show the relative ATP levels for 6 knockout mutants ($\Delta csgD$, $\Delta gnsA$, $\Delta yohJ$, $\Delta ybfA$, $\Delta yhdU$, $\Delta rpmF$) and 9 overexpression strains (*yohJ*, *csgD*, *yhdU*, *gnsA*, *gnsB*, *rbsD*, *rpmF*, *yjjQ*, *ymdF*). Of the 6 knockout mutants, interestingly,

ATP production dramatically increased in the $\Delta rpmF$ (5.9-fold change). ATP produced in $\Delta yhdU$ was significantly lower than that in UTI89 (the ATP ratio was 0.42). The other 4 knockout mutants ($\Delta csgD$, $\Delta gnsA$, $\Delta ybfA$, $\Delta yohJ$) produced similar ATP levels as UTI89. Of the 9 overexpression strains, only $yjjQ$ overexpression exhibited a similar ATP concentration as UTI89+pBAD202 vector control strain and all the other 8 overexpression strains ($yohJ$, $csgD$, $yhdU$, $gnsA$, $gnsB$, $rbsD$, $rpmF$, $ymdF$) had lower ATP concentrations than UTI89+pBAD202, indicating their role in persistence.

Discussion

In this study, we completed the detection of dynamic transcriptional profiles in the process of persister formation for *E. coli* K12 strain W3110 using Illumina RNA sequencing technology. Numerous previous studies investigated mechanisms involved in drug specific persistence by comparison the differentially expressed genes of samples from “pre-drug treatment” and “post-drug treatment”. Samples used here were not treated by antibiotics but had specific antibiotic tolerance profiles. By drawing a comprehensive blueprint of the gene expression levels for three key points (S1 “nonexistence”, S2 “emergence” and S3 “abundance”) of persister formation under ampicillin treatment, we observed 51 and 29 genes were significantly up-regulated (>2-fold change) in S2 and S3 in comparison with S1. We mainly focused on the up-regulated genes and differences between S2 and S1 or S3 and S1 because the period from S1 to S2 (S3) was a “boot-up” (“shoot-up”) process and we believe the two periods (S1 to S2 and S1 to S3) are of critical importance. As stated above, different pathways were involved in persister “boot-up” and “shoot-up” stages. However, metabolic pathways, membrane proteins and transcription factors occupied the principal parts in both stages. This suggests during persister formation, there is a good chance that bacteria undergo considerable changes in their metabolic state as well as membrane characteristics. This concept appears to contradict the current acceptable view that persisters are predominantly dormant and in agreement with Orman and Brynildsen who claim “Dormancy is not necessary or sufficient for bacterial persistence”(Orman and Brynildsen, 2013). This is mainly because we are looking at genes involved in the actual process of persister formation rather than when persisters are already formed. Further experiments should be performed to detect the effects of these metabolic pathways on ATP production. Membrane and membrane proteins have previously been shown to participate in persistence (Cui et al., 2016; Guo et al., 2017), and our data indicate that their roles in persister formation may be far more important and complicated than previously thought, and changes to lowermembrane permeability to drugs has been noted in some dormant bacteria (Dick T, 2015)

By comparing the 51 DEGs in S2/S1 and the 29 DEGs in S3/S1, we observed 13 DEGs ($gnsA$, $gnsB$, $ybfA$, $yjjQ$, $ymdF$, $yhdU$, $csgD$, $yncN$, $rpmF$, $ydcX$, $yohJ$, $ssrA$, $rbsD$) that overlapped between S2/S1 and S3/S1. The 13 overlapping DEGs may play crucial roles throughout the persister formation process. It is surprising that only two previously known persister genes $ssrA$ (Li and Zhang, 2013) and $ydcX$ () are identified in our study despite many persister genes have been reported previously. This result further confirmed the significance of $ssrA$ and $ydcX$ in persister formation. However, this did not mean other persister genes were less important than them. A plausible explanation is that other previously identified

persister genes either display their effect on persister formation at different conditions rather than the natural persister formation as a function of ageing or time as in this study, or their influence is more on persister maintenance or survival rather than persister formation, as is investigated here.

We found that *rpmF* is involved in participation in multidrug tolerance. *rpmF*, which encodes 50S ribosomal subunit protein L32, is responsible for protein synthesis. Since persisters are considered slow-growing or in dormant state, and metabolically inactive, this prompted us to hypothesize that deletion of *rpmF* would have a significant effect on metabolic proteins, disturb the homeostasis of metabolism in bacteria leading to decreased persister levels. Another explanation is that deletion of *rpmF* would enhance the activity of drug or stress targets and increase the susceptibility of $\Delta rpmF$. The multidrug and multiple stress susceptibility of $\Delta rpmF$ have previously been shown to be hypersensitive to ampicillin, sulfamethoxazole, rifampicin and metronidazole (Tamae et al., 2008). However, subinhibitory concentrations of antibiotics were used in that study, but were not subjected to persister assays where the antibiotic concentrations are usually far more than MICs. Our study is the first to demonstrate the role of *rpmF* in persister formation. Since RpmF is a conserved protein, it is likely that such homologs play similar roles in persister formation in other bacteria. Future studies are needed to confirm this and address the mechanism of RpmF in persister formation. RpmF could serve as a good persister drug target for future drug development.

rpmF and *plsX* gene encoding a protein involved in membrane lipid synthesis and several fatty acid biosynthetic genes (*fabH*, *fabD* and *fabG*) are cotranscribed in *E. coli*. Organization of these genes into an operon may play a role in the coordinated regulation of the synthesis of ribosomes, cell membranes and fatty acid and lipid biosynthesis (Podkovyrov S1, Larson TJ. 1995) and link these processes to cellular metabolism. It is of interest to note that the same fatty acid synthesis operon *fabHGD* is activated by FadR but inhibited by ppGpp, a well-known molecule mediating persister formation (My L, et al. 2013). In addition, it has been shown that *rpmF* is involved in biofilm formation in *Actinobacillus pleuropneumoniae* (Grasteau A, et al., 2011) and that *rpmF* (L32) mutant is hypersensitive to ROS-generating agent hydroxyurea (HU) (Nakayashiki T, Mori H, 2013). These findings are consistent with our findings that RpmF involvement in persister formation may be mediated via ppGpp. Future studies are needed to address the detailed mechanism.

In summary, the data presented here provides a portrait of the overall profile of genes involved in type II persister formation. In particular, besides two known genes including a member of the trans-translation pathway (*ssrA*) and an orphan toxin (*ydcX*), 11 novel genes (*gnsA*, *gnsB*, *ybfA*, *yjjQ*, *ymdF*, *yhdU*, *csgD*, *yncN*, *rpmF*, *yohJ*, *rbsD*) previously not reported are identified in this study. Among them, two genes *rpmF* (encoding 50S ribosomal subunit protein L32) and *yjjQ* (a LuxR-type transcription factor) are identified as two key factors for persister formation. Our findings shed new light on mechanisms of type II persister formation and provide new therapeutic targets for intervention.

References

- Balaban, N.Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004). Bacterial persistence as a phenotypic switch. *Science* 305(5690), 1622-1625. doi: 10.1126/science.1099390.
- Cui, P., Niu, H., Shi, W., Zhang, S., Zhang, H., Margolick, J., et al. (2016). Disruption of Membrane by Colistin Kills Uropathogenic Escherichia coli Persisters and Enhances Killing of Other Antibiotics. *Antimicrob Agents Chemother* 60(11), 6867-6871. doi: 10.1128/aac.01481-16.
- Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* 97(12), 6640-6645. doi: 10.1073/pnas.120163297.
- Dorr, T., Lewis, K., and Vulic, M. (2009). SOS response induces persistence to fluoroquinolones in Escherichia coli. *PLoS Genet* 5(12), e1000760. doi: 10.1371/journal.pgen.1000760.
- Grasteau A, Tremblay YD, Labrie J, Jacques M. (2011). Novel genes associated with biofilm formation of Actinobacillus pleuropneumoniae. *Vet Microbiol.* 153:134-143.
- Guo, Y., Liu, X., Li, B., Yao, J., Wood, T.K., and Wang, X. (2017). Tail-Anchored Inner Membrane Protein ElaB Increases Resistance to Stress While Reducing Persistence in Escherichia coli. *J Bacteriol* 199(9). doi: 10.1128/jb.00057-17.
- Hobby, G.L., Meyer, K., and Chaffee, E. (1942). Observations on the mechanism of action of penicillin. *Proc Soc Exp Biol Med* 50, 281-285.
- Islam, S., Benedik, M.J., and Wood, T.K. (2015). Orphan toxin OrtT (YdcX) of Escherichia coli reduces growth during the stringent response. *Toxins (Basel)* 7(2), 299-321. doi: 10.3390/toxins7020299.
- Kim, J.S., and Wood, T.K. (2016). Persistent Persister Misperceptions. *Front Microbiol* 7, 2134. doi: 10.3389/fmicb.2016.02134.
- Kint, C.I., Verstraeten, N., Fauvart, M., and Michiels, J. (2012). New-found fundamentals of bacterial persistence. *Trends Microbiol* 20(12), 577-585. doi: 10.1016/j.tim.2012.08.009.
- Lewis, K. (2010). Persister cells. *Annu Rev Microbiol* 64, 357-372. doi: 10.1146/annurev.micro.112408.134306.
- Li, J., Ji, L., Shi, W., Xie, J., and Zhang, Y. (2013). Trans-translation mediates tolerance to multiple antibiotics and stresses in Escherichia coli. *J Antimicrob Chemother* 68(11), 2477-2481. doi: 10.1093/jac/dkt231.
- Li, Y., and Zhang, Y. (2007). PhoU is a persistence switch involved in persister formation and tolerance to multiple antibiotics and stresses in Escherichia coli. *Antimicrob Agents Chemother* 51(6), 2092-2099. doi: 10.1128/aac.00052-07.
- Ma, C., Sim, S., Shi, W., Du, L., Xing, D., and Zhang, Y. (2010). Energy production genes sucB and ubiF are involved in persister survival and tolerance to multiple antibiotics and stresses in Escherichia coli. *FEMS Microbiol Lett* 303(1), 33-40. doi: 10.1111/j.1574-6968.2009.01857.x.
- Maisonneuve, E., Castro-Camargo, M., and Gerdes, K. (2013). (p)ppGpp controls bacterial persistence by stochastic induction of toxin-antitoxin activity. *Cell* 154(5), 1140-1150. doi: 10.1016/j.cell.2013.07.048.
- Marques, C.N., Morozov, A., Planzos, P., and Zelaya, H.M. (2014). The fatty acid signaling molecule cis-2-decenoic acid increases metabolic activity and reverts persister cells to an antimicrobial-susceptible state. *Appl Environ Microbiol* 80(22), 6976-6991. doi: 10.1128/aem.01576-14.
- Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L., and Wold, B. (2008). Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* 5(7), 621-628. doi: 10.1038/nmeth.1226.
- Mulcahy, L.R., Burns, J.L., Lory, S., and Lewis, K. (2010). Emergence of Pseudomonas aeruginosa strains producing high levels of persister cells in patients with cystic fibrosis. *J Bacteriol* 192(23), 6191-6199. doi: 10.1128/jb.01651-09.

453 My L, Rekoske B, Lemke JJ, Viala JP, Gourse RL, Bouveret E. Transcription of the Escherichia coli Fatty Acid
454 Synthesis Operon fabHGD is Directly Activated by FadR and Inhibited by ppGpp. *J Bacteriol.* 2013 Aug;
455 195(16): 3784–3795.

456 Nakayashiki T, Mori H. (2013). Genome-wide screening with hydroxyurea reveals a link between nonessential
457 ribosomal proteins and reactive oxygen species production. *J Bacteriol.* 195:1226-1235.

458 Nierman, W.C., Yu, Y., and Losada, L. (2015). The In vitro Antibiotic Tolerant Persister Population in Burkholderia
459 pseudomallei is Altered by Environmental Factors. *Front Microbiol* 6, 1338. doi:
460 10.3389/fmicb.2015.01338.

461 Orman, M.A., and Brynildsen, M.P. (2013). Dormancy is not necessary or sufficient for bacterial persistence.
462 *Antimicrob Agents Chemother* 57(7), 3230-3239. doi: 10.1128/aac.00243-13.

463 Podkovyrov S, Larson TJ. (1995). Lipid biosynthetic genes and a ribosomal protein gene are cotranscribed. *FEBS*
464 *Lett.* 368:429-431.

465 Siska, C., and Kechris, K. (2017). Differential correlation for sequencing data. *BMC Res Notes* 10(1), 54. doi:
466 10.1186/s13104-016-2331-9.

467 Tamae, C., Liu, A., Kim, K., Sitz, D., Hong, J., Becket, E., et al. (2008). Determination of antibiotic hypersensitivity
468 among 4,000 single-gene-knockout mutants of Escherichia coli. *J Bacteriol* 190(17), 5981-5988. doi:
469 10.1128/jb.01982-07.

470 Van den Bergh, B., Fauvart, M., and Michiels, J. (2017). Formation, physiology, ecology, evolution and clinical
471 importance of bacterial persisters. *FEMS Microbiol Rev* 41(3), 219-251. doi: 10.1093/femsre/fux001.

472 Van den Bergh, B., and Michiels, J.E. (2016). Frequency of antibiotic application drives rapid evolutionary
473 adaptation of Escherichia coli persistence. 1, 16020. doi: 10.1038/nmicrobiol.2016.20.

474 Vega, N.M., Allison, K.R., Khalil, A.S., and Collins, J.J. (2012a). Signaling-mediated bacterial persister formation.
475 *Nat Chem Biol* 8(5), 431-433. doi: 10.1038/nchembio.915.

476 Vega, N.M., Allison, K.R., Khalil, A.S., and Collins, J.J. (2012b). Signaling-mediated bacterial persister formation.
477 *Nature Chemical Biology* 8(5), 431-433. doi: 10.1038/nchembio.915.

478 Wang, X., and Wood, T.K. (2011). Toxin-antitoxin systems influence biofilm and persister cell formation and the
479 general stress response. *Appl Environ Microbiol* 77(16), 5577-5583. doi: 10.1128/aem.05068-11.

480 Wu, N., He, L., Cui, P., Wang, W., Yuan, Y., Liu, S., et al. (2015). Ranking of persister genes in the same
481 Escherichia coli genetic background demonstrates varying importance of individual persister genes in
482 tolerance to different antibiotics. *Front Microbiol* 6, 1003. doi: 10.3389/fmicb.2015.01003.

483 Zhang, Y. (2014). Persisters, persistent infections and the Yin-Yang model. *Emerg Microbes Infect* 3(1), e3. doi:
484 10.1038/emi.2014.3.

485 Zhang, Y., Yew, W.W., and Barer, M.R. (2012). Targeting persisters for tuberculosis control. *Antimicrob Agents*
486 *Chemother* 56(5), 2223-2230. doi: 10.1128/aac.06288-11.

Table 1 Persister formation related gene transcripts with more than 2-fold up-regulation in S2 compared with S1 (p -value <0.05)*.

NO.	Gene ID	Gene name	Fold change	Gene description
1	12933889	<i>ydcA</i>	45.44144	Uncharacterized protein
2	12933782	<i>yjjQ</i>	22.85771	Putative transcription factor
3	12933901	<i>ydcX</i>	10.62014	Orphan toxin OrtT
4	12930553	<i>ymdF</i>	7.767768	Uncharacterized protein
5	12933458	<i>yhdU</i>	7.602703	Predicted membrane protein
6	12930933	<i>ybfA</i>	6.672799	Uncharacterized protein
7	12932386	<i>idnK</i>	6.291892	Pentose phosphate pathway Carbon metabolism
8	12933843	<i>gnsA</i>	6.216046	Predicted regulator of phosphatidylethanolamine synthesis
9	12931244	<i>gnsB</i>	5.887559	Protein GnsB
10	12934174	<i>yfcV</i>	5.534535	Uncharacterized fimbrial-like protein
11	12930501	<i>yncN</i>	4.272272	Hypothetical protein
12	12933181	<i>yohJ</i>	4.326215	Conserved inner membrane protein
13	12933927	<i>zntA</i>	3.713078	Lead, cadmium, zinc and mercury-transporting ATPase
14	12930672	<i>idnD</i>	3.620335	L-idonate 5-dehydrogenase (NAD(P)(+))
15	12932329	<i>arsR</i>	3.140484	Arsenical resistance operon repressor sequence-specific DNA binding transcription factor activity
16	12933817	<i>ykgE</i>	3.131381	Predicted oxidoreductase
17	12930745	<i>yadK</i>	3.087688	Uncharacterized fimbrial-like protein
18	12934032	<i>yebN</i>	3.006995	Conserved inner membrane protein
19	12930936	<i>ybfH</i>	2.8713	Hypothetical protein
20	12930776	<i>fadE</i>	2.87019	Fatty acid degradation Fatty acid metabolism
21	12931540	<i>yfcY</i>	2.823285	Valine, leucine and isoleucine degradation Fatty acid degradation
22	12932494	<i>torR</i>	2.789674	Two-component system
23	12934145	<i>fsaB</i>	2.738138	Fructose-6-phosphate aldolase 2
24	12934116	<i>yhhI</i>	2.71125	Predicted transposase

25	12934023	<i>insC</i>	2.667615	IS2 insertion element repressor InsA
26	12931070	<i>csgD</i>	2.621622	DNA-binding transcriptional activator
27	12934464	<i>rbsA</i>	2.598083	ABC transporters
28	12931901	<i>yfdY</i>	2.588437	Predicted inner membrane protein
29	12932146	<i>arsC</i>	2.586667	Arsenate reductase;
30	12933846	<i>torT</i>	2.487179	Periplasmic sensory protein associated with the TorRS two-component regulatory system
31	12933202	<i>fxsA</i>	2.485491	Inner membrane protein
32	12931368	<i>yobB</i>	2.467409	Uncharacterized protein YobB;
33	12933149	<i>glmU</i>	2.440769	Amino sugar and nucleotide sugar metabolism
34	12931356	<i>torY</i>	2.427427	Cytochrome c-type protein TorY;
35	12930508	<i>ldhA</i>	2.387548	Pyruvate metabolism
36	12932242	<i>rbsD</i>	2.384947	ABC transporters
37	12934234	<i>pspC</i>	2.338203	Transcriptional activator
38	12931784	<i>ybeF</i>	2.21888	Uncharacterized HTH-type transcriptional regulator YbeF;
39	12932035	<i>rpmF</i>	2.21342	50S ribosomal protein L32
40	12932624	<i>yfiA</i>	2.213267	Cold shock protein associated with 30S ribosomal subunit
41	12931638	<i>yjfV</i>	2.174975	Arsenical pump membrane protein
42	12931815	<i>ibpB</i>	2.160241	Small heat shock protein IbpB;
43	12933322	<i>ygeP</i>	2.115695	Uncharacterized protein YgeP;
44	12932504	<i>yagI</i>	2.105242	Uncharacterized HTH-type transcriptional regulator YagI;
45	12930217	<i>ssrA</i>	2.103644	Trans-translation
46	12931840	<i>yohK</i>	2.063606	Inner membrane protein YohK;
47	12933596	<i>ibpA</i>	2.062559	Small heat shock protein IbpA;
48	12933928	<i>yhhP</i>	2.058133	Uncharacterized protein
49	12931415	<i>cobS</i>	2.05437	Porphyrin and chlorophyll metabolism
50	12932416	<i>ymcB</i>	2.026644	Uncharacterized protein
51	12932000	<i>degP</i>	2.019632	Two-component system

* The bold fonts represent the 13 overlapping genes.

Table 2 Fold-changes of the expression levels of the 13 genes that overlap in S2:S1 and S3:S1 by RT-PCR

Gene name	Fold-change		RT-PCR primers
	S2:S1	S3:S1	
<i>yjjQ</i>	2.57	3.19	F: 5' CGC TCC GTC ATT CTC ACT CTT 3' R: 5' TTG CTG TGA ATT GCC AGT CG 3'
<i>ymdF</i>	6.85	8.73	F: 5' AAC CAT CGA GGC GGT TCC 3' R: 5' TTT GCC GTG ACT GCT CTT ACC 3'
<i>gnsA</i>	3.22	3.78	F: 5' AAA ACA AGC CGA AAC GGA AAT 3' R: 5' CAT TTT TTC TCG TGC GGT GAA 3'
<i>gnsB</i>	3.65	4.21	F: 5' CTT AAA AAC AAA AGC AGA AGC AGA T 3' R: 5' TTC AAG ACC CGT CAT TTT TTC C 3'
<i>rpmF</i>	5.34	6.15	F: 5' GAA TAA ACC AAC CCG TTC CAA AC 3' R: 5' ACC GTC GGC AGT GAT GTG G 3'
<i>yhdU</i>	2.72	2.31	F: 5' TAT TGG TGG CTC GTC GTT TTC 3' R: 5' TTG CGG CAT TTG TCT GTT TC 3'
<i>csgD</i>	3.43	5.14	F: 5' GAG ATC GCT CGT TCG TTG TTC 3' R: 5' CGC CTG AGG TTA TCG TTT GC 3'
<i>rbsD</i>	4.68	3.32	F: 5' CCA GGG TGT ACC TTC TTT TAT GC 3' R: 5' TTT CGT GGA GTT GCG GAT TA 3'
<i>yohJ</i>	4.09	2.76	F: 5' TTT GTG CCG ATT GGC GTA G 3' R: 5' CGT TCA CCG TGT ACC AGT TGC 3'
<i>ybfA</i>	2.90	2.22	F: 5' CGC CGT ACT TAT GCG GTT GC 3' R: 5' CGG TTT ATC GCT GGT TTT CGA 3'
<i>yncN</i>	5.76	6.93	F: 5' TCA GGT TTC ATG GGA GGC G 3' R: 5' CGA GTT GTT TCA GGA TTG CTT TAC 3'
<i>ycdX</i>	4.67	6.71	F: 5' TTA TCA CCT GGT TTC TTT CTC ACG 3' R: 5' TAA AAC AGT GAA AAC AAC AGC G 3'
<i>ssrA</i>	2.49	2.12	F: 5' TCG CAA ACG ACG AAA ACT ACG 3' R: 5' GAC GGA CAC GCC ACT AAC AA 3'

Table 3 Persister formation related gene transcripts with more than 2-fold up-regulation in S3 compared with S1 (p -value <0.05) *.

NO.	Gene ID	Gene name	Fold change	Gene description
1	12930501	<i>yncN</i>	15.67	Hypothetical protein
2	12931025	<i>rmf</i>	15.6	Ribosome modulation factor
3	12932035	<i>rpmF</i>	15.2	50S ribosomal protein L32
4	12933891	<i>mokB</i>	14.6	Regulatory peptide
5	12932090	<i>yhfG</i>	14.4	Uncharacterized protein YhfG
6	12930553	<i>ymdF</i>	10.1	Uncharacterized protein
7	12933901	<i>ydcX</i>	6.6	Orphan toxin OrtT
8	12933458	<i>yhdU</i>	5.5	Predicted membrane protein
9	12933782	<i>yjjQ</i>	5.0	LuxR-type transcription factor
10	12930933	<i>ybfA</i>	4.2	Uncharacterized protein
11	12932958	<i>envR</i>	4.0	DNA-binding transcriptional regulator
12	12930217	<i>ssrA</i>	3.8	Trans-translation
13	12932271	<i>glpD</i>	3.5	Glycerophospholipid metabolism
14	12931490	<i>ccmD</i>	3.2	ABC transporters
15	12933181	<i>yohJ</i>	3.06	Conserved inner membrane protein
16	12933196	<i>argC</i>	3.0	Arginine and proline metabolism 2-Oxocarboxylic acid metabolism Biosynthesis of amino acids
17	12932242	<i>rbsD</i>	2.9	ABC transporters
18	12933881	<i>ymjC</i>	2.6	Predicted oxidoreductase
19	12931423	<i>yeeF</i>	2.6	Predicted amino-acid transporter
20	12933985	<i>yghJ</i>	2.5	Predicted inner membrane lipoprotein
21	12931244	<i>gnsB</i>	2.5	Protein GnsB
22	12934127	<i>argG</i>	2.5	Alanine, aspartate and glutamate metabolism Arginine and proline metabolism Biosynthesis of amino acids
23	12933423	<i>tdcD</i>	2.4	Propanoate metabolism
24	12931961	<i>yohM</i>	2.1	Membrane protein conferring nickel and cobalt resistance
25	12934013	<i>ftn</i>	2.1	Cytoplasmic ferritin iron storage protein

26	12931748	<i>argD</i>	2.0	Lysine biosynthesis Arginine and proline metabolism 2-Oxocarboxylic acid metabolism Biosynthesis of amino acids
27	12931070	<i>csgD</i>	2.0	DNA-binding transcriptional activator
28	12933843	<i>gnsA</i>	2.0	Predicted regulator of phosphatidylethanolamine synthesis
29	12932331	<i>arsB</i>	2.0	Arsenite/antimonite transporter

* The bold fonts represent the 13 overlapping genes.

Table 4 Ranking of the top 7 persister genes according to their knockout mutant scores upon exposure to antibiotics ^a.

Knockout genes	Pathways or proteins	Scores for each antibiotic ^b				Sum
		Amp	Gen	Lev	Opposite	
<i>rpmF</i>	50S ribosomal protein L32	1	1	1		3
<i>ybfA</i>	Hypothetical protein	1	1	0		2
<i>gnsA</i>	predicted regulator of phosphatidylethanolamine synthesis	1	1	1	-1	2
<i>yohJ</i>	Conserved inner membrane protein	0	1	0		1
<i>csgD</i>	DNA-binding transcriptional activator in two-component regulatory system	0	1	0		1
<i>yhdU</i>	Predicted membrane protein	1	0	0		1
<i>yjjQ</i>	Putative transcription factor	0	0	0		0

^a Genes were ranked according to cell survival under exposure to three different antibiotics. The genes whose overexpression strains show difference (≥ 5 -fold) compared to the parent strain are scored as “1” point, whereas those mutants that show difference (< 5 -fold) were scored as “0”. All scores for a given mutant are calculated from the sum of different antibiotic exposures to obtain the ranking order of the persister genes.

^b The three abbreviations “Amp”, “Gen” and “Lev” are symbols for “Ampicillin”, “Gentamicin” and “Levofloxacin”, respectively. “Opposite” represents that a given mutant exhibited decreased persister levels in the presence of one or two antibiotics while showing increased persister numbers in other antibiotic exposures compared with UTI89.

Table 5 Ranking of the 10 persister related genes according to their overexpression scores upon exposure to antibiotics ^a.

Overexpression genes	Pathways or proteins	Scores for each antibiotic ^b				Sum
		Amp	Gen	Lev	Opposite	
<i>yjjQ</i>	Putative transcription factor	1	1	1		3
<i>ymdF</i>	Hypothetical protein	1	1	1	-1	2
<i>gnsA</i>	Predicted regulator of phosphatidylethanolamine synthesis	1	0	1		2
<i>gnsB</i>	Hypothetical protein	1	1	0		2
<i>rpmF</i>	50S ribosomal protein L32	1	1	0		2
<i>yhdU</i>	Predicted membrane protein	1	1	0		2
<i>csgD</i>	DNA-binding transcriptional activator in two-component regulatory system	1	1	0		2
<i>rbsD</i>	Predicted cytoplasmic sugar-binding protein	1	1	0		2
<i>yohJ</i>	Conserved inner membrane protein	1	0	0		1
<i>ybfA</i>	Hypothetical protein	0	0	0		0

^a Genes are ranked according to cell survival under exposure to three different antibiotics. The genes whose overexpression strains show difference (≥ 5 -fold) compared to the parent strain are scored as “1” point, whereas those mutants that show difference (< 5 -fold) are scored as “0”. All scores for a given mutant are calculated from the sum of different antibiotic exposures to obtain the ranking order of the persister genes.

^b The abbreviations “Amp”, “Gen” and “Lev” are symbols for “Ampicillin”, “Gentamicin” and “Levofloxacin”, respectively. “Opposite” represents that a given strain exhibited decreased persister levels in the presence of one or two antibiotics while showing increased persister numbers in exposure to other antibiotics compared with UTI89.

Table S1 Primers for amplification of the 10 overlapping persister genes^a

Gene name	Primers
<i>gnsA</i>	F: 5' CATG <u>CCA TGG</u> GTG AAT ATT GAA GAG TTA AAA AAA CAA G 3' R: 5' CCCG <u>GAA TTC</u> TCA CAT TAT TTT GAT TTT GAC ATC A 3'
<i>gnsB</i>	F: 5' CATG <u>CCA TGG</u> GTG AAT ATT GAA GAG TTA AAA AAA CAA G 3' R: 5' CCCG <u>GAA TTC</u> TCA CAT TAT TTT GAT TTT GAC ATC A 3'
<i>ybfA</i>	F: 5' CATG <u>CCA TGG</u> ATG GAA CTC TAC AAA GAA TAT CCT GC 3' R: 5' CCCG <u>GAA TTC</u> TCA GTA AAA ATC ACC AGT TGC CT 3'
<i>yjjQ</i>	F: 5' CATG <u>CCA TGG</u> ATG TTG CCA GGA TGC TGC A 3' R: 5' CCCG <u>GAA TTC</u> CTA TGA GTG CGA CAT TTC TCT TCT T 3'
<i>ymdF</i>	F: 5' CATG <u>CCA TGG</u> ATG GCA AAC CAT CGA GGC 3' R: 5' CCCG <u>GAA TTC</u> CTA GTT GTC GCT TTT ACC GTG AC 3'
<i>yhdU</i>	F: 5' CATG <u>CCA TGG</u> TTG GTT AAA CAT GCA CAG GCT C 3' R: 5' CCCG <u>GAA TTC</u> TCA CAT TCT GTC CTG AAA ATT CAG T 3'
<i>csgD</i>	F: 5' CGG <u>GGT ACC</u> ATG TTT AAT GAA GTC CAT AGT ATT CAT G 3' R: 5' CCCG <u>GAA TTC</u> TTA TCG CCT GAG GTT ATC GTT T 3'
<i>rpmF</i>	F: 5' CATG <u>CCA TGG</u> ATG GCC GTA CAA CAG AAT AAA CC 3' R: 5' CCCG <u>GAA TTC</u> TTA CTT AGC GAT GAC CTT GCG G 3'
<i>yohJ</i>	F: 5' CATG <u>CCA TGG</u> TTG TAT GAT GAA TCC ATC TCA TCT G 3' R: 5' CCCG <u>GAA TTC</u> TCA TTC TTC TGA TCC TTT CTG ACC T 3'
<i>rbsD</i>	F: 5' CATG <u>CCA TGG</u> ATG AAA AAA GGC ACC GTT CTT AA 3' R: 5' CCCG <u>GAA TTC</u> TCA GAA CGT CAC GCC AGC A 3'

^a The protective bases are in *italic*. **Bold** and underline marks the sequences of *NcoI* and *EcoRI* sites.

Figure 1 Persister associated DEG (≥ 2 -fold increase) distributions and percentage (A) S2:S1 and (B) S3:S1

Figure 2 Effect of the 7 overlapping gene knockout mutations on *E. coli* persister formation. Cultures of UTI89 and persister gene knockout mutants grown to log phase and were immediately treated with (A) ampicillin (200 µg/ml) for 24 hours, (B) gentamicin (40 µg/ml) for 5 hours, (C) levofloxacin (5 µg/ml) for 3 hours. Surviving bacteria were counted after 16h incubation at 37 °C on LB plates without antibiotics.

553

554

555

Figure 3 Effect of overexpression of the 10 overlapping genes on *E. coli* persister formation. Cultures of UTI89+pBAD202 and persister gene overexpression mutants grown to log phase and were immediately treated with (A) ampicillin (200 μ g/ml) for 24 hours, (B) gentamicin (40 μ g/ml) for 5 hours, (C) levofloxacin (5 μ g/ml) for 3 hours. Surviving bacteria were counted after 16h incubation at 37 $^{\circ}$ C on LB plates without antibiotics.

(A)

(B)

Figure 4 The relative ATP concentrations of 6 knockout mutants and 9 overexpression strains. Cultures of knockout mutants (A) and overexpression strains (B) grown to log phase and immediately subjected to ATP assay as described in Methods.

589 **Supplementary Figure 1**

590

591 **Figure S1 Different persister levels of *E. coli* at three time points.** Cultures of *E. coli*
592 W3110 grown to different time points (3h-S1, 4h-S2 and 5h-S3) were withdrawn and
593 immediately treated with ampicillin (100 µg/ml) for 3 hours. Surviving bacteria were counted
594 after 16h incubation at 37 °C on LB plates without antibiotics. As can be seen, no persisters
595 are formed at 3 hrs (S1) but persisters begin to appear at 4 hrs (S2) and shoot up considerably
596 at 5hrs (S3).

597

598

599