

Exploring the genetic correlations of antisocial behavior and life history traits

Jorim J. Tielbeek

VU University Medical Center Amsterdam

J.C. Barnes

University of Cincinnati

Arne Popma

VU University Medical Center Amsterdam

Tinca J.C. Polderman

VU University Amsterdam

James J. Lee

University of Minnesota

John B. Perry

Cambridge University

Danielle Posthuma

VU University Amsterdam

Brian B. Boutwell*

Saint Louis University

***Corresponding Author Email: brian.boutwell@slu.edu**

The authors declare no conflicts of interest. All results reported are original.

Abstract

Prior evolutionary theory provided reason to suspect that measures of development and reproduction would be correlated with antisocial behaviors in human and non-human species. Behavioral genetics has revealed that most quantitative traits are heritable, suggesting that these phenotypic correlations may share genetic etiologies. To date, few studies have capitalized on the advances in genome-wide analyses (GWAS) to test such predictions. We use prior GWAS (N=31,814) to estimate the genetic correlations between various measures of reproductive development and antisocial behavior. Our results revealed significant genetic correlations between antisocial behaviour and age of first birth ($r_g = -.64$, $p = .0008$) and number of children ever born ($r_g = 0.50$, $p = .0065$).

A moderate proportion of the variance in antisocial phenotypes is accounted for by genetic variation [1]. Phenotypic indicators of externalizing traits, moreover, are correlated with other forms of psychopathological behavior [2]. These findings suggest a general vulnerability underlying the spectrum of externalizing disorders, one underpinned by pleiotropic genetic influences. With the proliferation of population-based samples containing measured genes, researchers have investigated possible molecular correlates of antisocial behaviors. While some promising results emerged, so too did to the recognition of limitations [3]. Chief among them included the absence of theory driven hypothesis testing, the use of underpowered designs, and an absence of correction for multiple testing bias [4].

As a result, studies utilizing genome-wide techniques have become the preferred approach to unraveling the genetic architectures of human complex traits. Recent GWAS of antisocial phenotypes revealed a number of trait-relevant alleles that were nearly genome-wide significant [5]. More recent GWAS evidence has also revealed associations between a number of SNPs and phenotypes known to correlate with antisocial behaviors [6]. In particular, various indicators of reproductive behavior such as age at menarche, age at first sexual contact, and number of sexual partners have all been examined using genetically sensitive data, with results identifying a number of genome-wide significant SNPs [6].

What remains in need of further testing is whether the genetic underpinnings of antisocial behavior covary with the genetic underpinnings of reproductive indicators. For this topic, in particular, prior theoretical work has suggested that some forms of antisocial behavior may represent natural variation in broad reproductive strategies—captured broadly in the area of life history evolution—that have been shaped by natural selection over the course of human evolution [7]. This line of reasoning makes the explicit prediction that reproductive and

antisocial behavior should be correlated at a phenotypic level, and perhaps at a genetic level, such that alleles associated with antisocial behavior are also associated with higher reproductive output and more rapid physical maturation.

We combined GWAS data from the Broad Antisocial Behavior Consortium (BroadABC) with data from the Early Genetics and Life-course Epidemiology (EAGLE) Consortium. In total, we meta-analyzed genotypic and phenotypic data from 31,814 individuals across thirteen unique samples, making it the largest collective sample available to estimate the effects of genome-wide genetic variation on antisocial behavior (ASB). To the extent that alleles associated with antisocial behavior are related to alleles underpinning variation in reproductive outcomes, it could provide a better understanding of why the traits tend to correlate at the phenotypic level [7].

Method

In the current study, we used (cross-trait) LD-score regression to estimate the SNP-based heritability of ASB and reproductive traits and the genetic correlation between the traits explained by all SNPs [8]. LD-score regression disentangles the contribution of true polygenic signal and bias due to population stratification to the inflated test statistics in GWAS, and optionally calculates a genetic correlation (r_g) between traits [8]. An important condition is that the per-SNP heritability near a given SNP must not be confounded with the extent of that SNP's LD with neighbouring SNPs. This condition is likely to be violated in the case of a phenotype related to fitness producing a downwardly biased SNP-based heritability estimate [8]. Yet, it is still possible for the genetic correlation between two traits to be estimated accurately given the bias in the numerator (the genetic covariance) cancelling the bias in the denominator (the square root of the product of two heritability estimates).

Here, we estimated the genetic correlation of ASB with reproductive traits based on the summary statistics from the largest GWAS meta-analyses available. For reproductive traits, summary statistics were derived from the LD HUB database. Moreover, we employed PLINK to ‘clump’ the SNPs utilizing 1000 Genomes V3 reference panel for Europeans, with 0.1 as LD r^2 threshold, and 500 KB as physical distance threshold. Then, we examined whether the signs of the regression coefficients of the SNPs for ASB and age of first birth (yielding the highest r_g) were, more often than expected by chance, in the same direction. We tested this using a binomial test to verify whether the proportion of SNPs (yielding p-values $< .05$) with concordant sign was higher or lower than expected by chance (0.5).

For ASB we performed a GWA meta-analysis in order to obtain a large GWAS sample. We combined summary data from the publicly available EAGLE consortium (N= 18,988, [9]) with those from non-overlapping samples of the BroadABC (QIMR, TEDS, COGA, Yale-Penn, N=12,166), totalling 31,814 participants. The genetic correlation between these two meta-analysed datasets was .38 (SE= .48). To maximize sample size, we included studies with a broad range of antisocial measures, including both aggressive and non-aggressive domains of antisocial behaviour, and utilizing study-specific scales in different age groups [5, 9-10]. The meta-analysis was run using a fixed-effects model with z-scores weighted by sample size as implemented in the software METAL [11].

Results

Insert Table 1 about Here

First, we calculated the SNP-based heritability estimates of ASB and the life-history traits, after which the genetic correlations between the traits were computed. The estimated heritability for ASB utilizing LD-score regression was low—2.8% with a standard error of 1.5%—and

substantially higher estimates of the SNP-heritability of ASB have been reported elsewhere using genome-wide complex traits analyses (GCTA) tools. We identified four cohorts (including QIMR, ALSPAC and GENR samples) with ASB measures that previously performed GCTA, resulting in a sample-size weighted h^2 of 35% (Panel A in Table 1). These GCTA estimates are based on more homogeneous individual cohorts yielding smaller samples, which might explain the discrepancy in h^2 . The SNP heritabilities based on LD-score regression were 5.9% (SE=2.7%) and 5.2% (SE=2.7%) respectively for the EAGLE and BroadABC consortia. Another possibility is that LD-score regression is more sensitive to causal SNPs being in low LD with their neighbours [12].

Our main analysis (Panel B in Table 1) revealed significant ($\alpha < 0.0071$) genetic correlations of ASB with age of first birth ($r_g = -.64$, $p = .0008$) and number of children ever born ($r_g = 0.50$, $p = .0065$) as well as suggestive associations with parents', mothers' and fathers' age at death, but not with age at menarche or age at menopause ($r_g = .026$, $p = .76$; $r_g = -.10$, $p = .48$, respectively). Moreover, sign-tests revealed less SNPs yielding a p-value $< .05$ with the same direction of effect than expected by chance (proportion = .47, $p < .001$) for ASB and age of first birth. Ultimately, our findings coalesce with prior preliminary evidence concerning genetic correlations between antisocial traits and behavioral and reproductive outcomes [5].

Discussion

The present study expands on prior work [5], showing preliminary evidence of a genetic correlation between reproductively relevant traits and antisocial behavior. Our genetic correlation analyses demonstrate that alleles associated with higher reproductive output (i.e., faster life history styles) were positively correlated with alleles associated with antisocial behavior, whereas alleles associated with giving birth later in life were negatively associated

with alleles linked to antisocial behavior. It is important to acknowledge that this study is preliminary and not without limitations. Our correlations were somewhat higher than expected (ranging from 0.5 to 0.7). We hypothesize that with increasing samples and more homogeneous phenotypic measures these estimates will be more in the range of 0.3-0.5, as reported by previous studies [5].

Our genetic correlation analyses are based on, and limited by, relatively low SNP-based heritability estimates of ASB (2.9%), yielding a heritability Z score of 1.93. The low SNP-based h^2 estimate of ASB may be due to heterogeneity in measurement by meta-analyzing multiple cohorts or the tendency of the causal SNPs to be in low LD with their neighbors. Finally, although somewhat elevated, the magnitude of the genetic correlations observed herein is similar in magnitude compared to other related phenotypes, utilizing alternative samples [5,13]. These results are interesting, and the hypotheses tested were guided by prior theory [7], yet should be viewed only as preliminary. As sample sizes continue to grow, and a more diverse range of phenotypes become available for testing, additional insight should emerge regarding the molecular underpinnings for antisocial behavior and human reproductive strategy.

References

1. Polderman TJ, Benyamin B, De Leeuw CA, Sullivan PF, Van Bochoven A, Visscher PM, Posthuma D. Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nat Genet* 2015; **47**: 702-709.
2. Krueger RF, Hicks BM, Patrick CJ, Carlson SR, Iacono WG, McGue M. Etiologic connections among substance dependence, antisocial behavior and personality: Modeling the externalizing spectrum. *J Abnorm Psychol* 2002; **111**: 411.
3. Dick DM, Agrawal A, Keller MC, Adkins A, Aliev F, Monroe S, Sher KJ. Candidate gene–environment interaction research: Reflections and recommendations. *Perspec Psychol Sci* 2015; **10**: 37-59.
4. Chabris CF, Lee JJ, Cesarini D, Benjamin DJ, Laibson DI. The fourth law of behavior genetics. *Curr Dir Psychol Sci* 2015; **24**: 304-312.
5. Tielbeek JJ, Johansson A, Polderman TJ, Rautiainen MR, Jansen P, Taylor M, Viding E. Genome-Wide Association Studies of a Broad Spectrum of Antisocial Behavior. *JAMA Psychiatry* 2017.
6. Barban N, Jansen R, De Vlaming R, Vaez A, Mandemakers JJ, Tropf FC, Tragante V. Genome-wide analysis identifies 12 loci influencing human reproductive behavior. *Nat Genet* 2016; **48**: 1462-1472.
7. Boutwell BB, Barnes JC, Beaver KM, Haynes RD, Nedelec JL, Gibson CL. A unified crime theory: the evolutionary taxonomy. *Aggress Violent Behav* 2015; **25**: 343-353.
8. Bulik-Sullivan BK, Loh PR, Finucane HK, Ripke S, Yang J, Patterson N, Schizophrenia Working Group of the Psychiatric Genomics Consortium. LD Score regression

distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015; **47**: 291-295.

9. Pappa I, St Pourcain B, Benke K, Cavadino A, Hakulinen C, Nivard MG, Evans DM. A genome-wide approach to children's aggressive behavior: the EAGLE consortium. *Am J Med Genet B: Neuropsychiatr Genet* 2016; **171**: 562-572.
10. Tielbeek JJ, Medland SE, Benyamin B, Byrne EM, Heath AC, Madden PA, Verweij KJ. Unraveling the genetic etiology of adult antisocial behavior: a genome-wide association study. *PloS one* 2012; **7**: e45086.
11. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010; **26**: 2190-2191.
12. Lee JJ, Chow CC. Conditions for the validity of SNP-based heritability estimation. *Hum Genet* 2014; **133**: 1011-1022.
13. Hill WD, Marioni RE, Maghzian O, Ritchie SJ, Hagenaars SP, McIntosh AM, Deary IJ. A combined analysis of genetically correlated traits identifies 187 loci and a role for neurogenesis and myelination in intelligence. *Molec Psychiat* 2018; **1**.

Table 1

Panel A. Previously reported genome-wide complex trait analysis (GCTA) estimates on ASB.

Cohort, Trait	N	SNP h^2	SE	P
<i>QIMR</i> ¹ , adult ASB	4816	.55	.41	.07
<i>ALSPAC</i> ² (middle childhood/early adolescence)	5299	.08	.06	.10
<i>GENR</i> ² (6-year)	2101	.54	.19	.002
<i>NTR</i> ² (3-year)	908	.46	.35	.09

Estimates included above were derived from studies by ¹Tielbeek et al. (2012) and ²Papa et al. (2016). SNP h^2 is the estimation of narrow-sense heritability contributed by common SNPs.

Panel B. Genetic correlations across life history speed indicators (four reproductive traits and three aging traits) and broad antisocial behavior.

Phenotype	Sample Size	SNP h^2	r_g (SE)	Bivariate Intercept	P-value
<i>Age at Menarche</i>	182.416	.21	.026(.085)	-0,0083	.76
<i>Age at Menopause</i>	69.360	.13	-.10(.144)	-0,0016	.48
<i>Age of First Birth</i>	222.037	.063	-.64(.19)	-0,0003	.0008**
<i>Number of Children Ever Born</i>	318.863	.025	.50(.18)	-0,007	.0065**
<i>Parents' Age at Death</i>	45.627	.032	-.70(.29)	0,0093	.015*
<i>Mother's Age at Death</i>	52.776	.040	-.51(.26)	-0,0017	.045*
<i>Father's Age at Death</i>	63.775	.042	-.52(.24)	-0,0036	.026*

Note: *Nominally significant ($p < .05$) **Significant at the multiple-testing corrected p-value (.0071). GWAS summary statistics from our combined analyses were used to calculate the r_g 's with other traits. SNP h^2 is the estimation of narrow-sense heritability contributed by common SNPs. r_g is the genetic correlation and is calculated with the LDSC software package using pre-calculated LD scores from Bulik-Sullivan et al.