1 **Enterovirus 71 structural viral protein 1 promotes mouse Schwann cell**
2 **autophagy via endoplasmic reticulum stress-mediated peripheral myelin protein**
3 **22 upregulation**

4

5 **Running title:** VP1 in autophagy

6

7 Pei-qing LI¹, Si-da YANG^{2*}, Dan-dan HU², Dan WeEI³, Jing LU⁴, Huan-ying
8 ZHENG⁴, Shu-shan NIE¹, Guang-ming LIU¹, Hao-mei YANG¹

9

10 ¹Department of Pediatric Emergency, Guangzhou Women and Children's Medical
11 Center, Guangzhou Medical University, Guangzhou, Guangdong, China

12 ²Department of Pediatric Neurology, Guangzhou Women and Children's Medical
13 Center, Guangzhou Medical University, Guangzhou, China

14 ³Pediatric Intensive Care Unit, First Affiliated Hospital of Guangxi Medical
15 University, Nanning, China

16 ⁴ Guangdong Provincial Institution of Public Health, Guangdong Provincial Center
17 for Disease Control and Prevention, China

18

19 ***Corresponding Author**

20 Si-da Yang

21 Department of Pediatric Neurology, Guangzhou Women and Children's Medical
22 Center, Guangzhou Medical University, Guangzhou, China

23 Tel: +86-13808879852

24 E-mail: yangsida2013@126.com

25

26

27 **Abstract**

28 Enterovirus 71 (EV71) accounts for the majority of hand, foot and mouth disease-
 29 related deaths due to fatal neurological complications. The clinical observations and
 30 animal models found the early invasion of nervous system, and the demyelinating
 31 phenomenon was observed. As one of the receptors of EV71 structural viral protein 1
 32 (VP1), SCARB2 mainly exists on the myelin sheath. EV71 VP1 can promote viral
 33 replication through inducing autophagy in neuron cells. This study aims to investigate
 34 the role and mechanism of VP1 in autophagy of mouse Schwann cells (MSCs). An
 35 EV71 VP1-expressing vector (pEGFP-C3-VP1) was generated and transfected into
 36 MSCs. Transmission electron microscopy (TEM) and Western blot analysis of the
 37 autophagy marker microtubule-associated proteins 1A/1B light chain 3B (LC3B)
 38 were used to assess autophagy in the cells. Real-time PCR and immunofluorescent
 39 staining were performed to determine the expression of PMP22. Small interfering
 40 RNA against PMP22 was employed to investigate the role of PMP22 in MSCs
 41 autophagy. Selective endoplasmic reticulum (ER) stress inhibitor salubrinal (SAL)
 42 was employed to determine whether PMP22 is mediated by ER stress. Our results
 43 demonstrated that VP1 played a promotive role in MSC autophagy. Overexpression of
 44 VP1 upregulated PMP22. PMP22 deficiency downregulated LC3B and thus inhibited
 45 autophagy. Furthermore, PMP22 expression was significantly suppressed by SAL.
 46 VP1 promotes MSC autophagy through upregulating ER stress-mediated PMP22
 47 expression. VP1/ER stress/ PMP22 axis in autophagy may be a potential therapeutic
 48 target for EV71 infection-induced fatal neuronal damage.

49

50 **Key words:** Enterovirus 71 structural viral protein 1; autophagy; endoplasmic
51 reticulum stress; peripheral myelin protein 22

Introduction

Enterovirus 71, a single-stranded RNA virus, is one of the major causative pathogens of contagious hand, foot and mouth disease (HFMD)(1, 2) that mainly affects children under the age of 5. HFMD is an emerging public health issue worldwide, especially in Asia-Pacific countries(1, 3, 4). Although HFMD is commonly considered as a self-limited disease characterized by ulcerating vesicles in the mouth and viral rashes on hands and feet(5-7), a small proportion of cases are severe and even fatal due to cardiopulmonary or neurological complications(8, 9). EV71 infection accounts for at least 80% of severe cases and 90% of deaths in China according to the recent data (10). Increasing evidence indicates that EV71 may target human neurons in central nervous system (CNS), leading to neuronal degeneration and severe neurological disorders in fatal cases(11-13). Our previous study also showed that neuronal necrosis and neuronophagia were present in the brainstem in fatal EV71-infected cases(14). Despite these neurotropic characteristics of EV71 virus, the pathogenesis and molecular mechanisms of EV71-induced neuronal damage remain largely unknown.

Autophagy is an intracellular process that is mediated by a unique organelle named autophagosome and transports cytoplasmic components to the lysosomes for degradation(15, 16). The alteration of autophagy in the nervous system is associated with various neurodegenerative and neurometabolic disorders such as Alzheimer's disease and Niemann-Pick disease(17-19). Autophagy can be observed using transmission electron microscopy (TEM) and can be assessed by measuring the conversion of microtubule-associated protein 1 chain 3 (LC3) to phosphatidylethanolamine (PE)-conjugated LC3 (LC3-II) localized in autophagosomal membranes, which reflects the number of autophagosomes or the degree of autophagy(20-22). EV71 has been shown to induce autophagy in infected

human rhabdomyosarcoma and neuroblastoma cells(23, 24). Our previous study demonstrates that EV71 structural viral protein 1 (VP1) also induces autophagy in cultured primary EV71-infected brainstem neurons, which can be inhibited by endoplasmic reticulum (ER) stress inhibitor salubrinal (SAL)(25), suggesting an essential role of ER stress in VP1-induced autophagy.

ER stress is triggered by the accumulation of unfolded or misfolded proteins in ER(26, 27). Although the relationship between ER stress and autophagy is not yet fully understood, it is well established that there is a dynamic crosstalk between these two systems, and ER stress either stimulates or inhibits autophagy(26, 28, 29). Since ER stress and autophagy are commonly concurrent in some human pathologies, such as cardiovascular diseases, cancers, and neurodegenerative disorders(29-31), it is of great importance to identify ER stress-associated molecules as positive or negative regulators of autophagy. Peripheral myelin protein 22 (PMP22) is a transmembrane glycoprotein highly expressed in the myelinating Schwann cells of peripheral neurons, and majorly contributes to synthesis and function of myelin sheaths(32). In Schwann cells, newly synthesized PMP22 is transiently retained in ER and Golgi before transported to the plasma membrane(33, 34). Under pathological conditions, excessive mature or premature (unfolded or misfolded) PMP22 accumulates in ER and interacts with calnexin, a Ca^{2+} -binding chaperone, leading to ER retention and activation of ER stress(35, 36). However, it remains unknown whether the relationship between PMP22 and ER stress is associated with autophagy.

In this study, we hypothesize that PMP22 is a downstream effector of ER stress and triggers activation of autophagy in response to EV71 capsid protein VP1. To confirm this hypothesis, we transfected mouse Schwann cells (MSCs) with VP1-expressing vectors to explore its effect on MSC autophagy and PMP22 expression. Our results

showed that ER stress mediates the expression of PMP22 that is essential for MSC autophagy, suggesting an involvement of VP1/ER stress/PMP22 axis in the regulation of MSC autophagy. Targeting VP1/ER stress/PMP22 axis in autophagy may be a novel therapeutic strategy against EV71 infection-induced neuronal damage.

Results

Cloning and identification of VP1 cDNA.

To determine if VP1 cDNA was successfully cloned into pEGFP-C3 vector, we prepared plasmids from transformed bacteria and digested them with *Bam*HI and *Xho*I. The results of agarose electrophoresis showed that a band was located between 750 and 1000 bp following enzymatic digestion (Fig. 1), which is consistent with the size of VP1 cDNA (894 bp) based on the GenBank database. The sequencing results also indicated that the cloned fragment was identical to the VP1 cDNA sequence (supplementary Fig. 1), suggesting that VP1 cDNA was successfully cloned into the vector without any mutation.

Overexpression of VP1 activates MSC autophagy.

To examine whether VP1 has an effect on MSC autophagy, we analyzed the cellular and subcellular morphology of VP1-overexpressing MSCs using TEM. As shown in Fig. 2, VP1-overexpressing MSCs exhibited the features of autophagy such as swelling mitochondria, dilation and degranulation of rough ER, and vesicle-like dilation of Golgi(37), whereas the organelles in untransfected and GFP-transfected control MSCs were still morphologically normal. These results suggest that VP1 may activate autophagy in MSCs.

Overexpression of VP1 markedly upregulates PMP22 expression in MSCs

Our previous study indicates an essential role of ER stress in VP1-induced autophagy in primary cultured EV71-infected brainstem neurons(14). In combination with the findings that PMP22 is abundant in Schwann cells and is closely associated with ER stress activation(32, 35, 36), we hypothesize that PMP22 might correlate with VP1 and play an important role in VP1-induced autophagy. To test this hypothesis, we detected the mRNA and protein expression of PMP22 in VP1-overexpressing MSCs. As shown in Fig. 3A, the mRNA expression of PMP22 was dramatically elevated in VP1-overexpressing MSCs compared with GFP-transfected cells. Immunofluorescent staining assay also showed similar results (Fig. 3B). These data indicate that VP1 is an upstream regulator of PMP22, suggesting a possible involvement of PMP22 in VP1-mediated activation of MSC autophagy.

PMP22 is essential for MSC autophagy.

We next sought to investigate whether PMP22 is involved in MSC autophagy. PMP22 was knocked down by siRNA, which was confirmed by markedly decreased expression of PMP22 in siPMP22-transfected MSCs (Fig. 4A). Importantly, compared with siCtrl-transfected groups, knockdown of PMP22 significantly downregulated the expression of LC3 isoform LC3B-II, a gold standard autophagy marker (38), as shown in Fig. 4B and 4C. Consistently, the ratio of LC3B-II to LC3B-I in PMP22-deficient MSCs was also significantly lower than that in siCtrl-transfected groups (Fig. 4D). Furthermore, TEM images showed that there was no observable autophagic structure in siPMP22-transfected MSCs as compared with siCtrl-transfected cells (Fig. 5). Taken together, these data suggest that PMP22 is required for activation of autophagy in MSCs.

ER stress mediates PMP22 expression in MSCs.

Since PMP22 is closely associated with ER stress and both PMP22 and ER stress are essential for activation of autophagy, we further sought to clarify the relationship between PMP22 and ER stress in MSCs using selective ER stress inhibitor SAL. As shown in Fig. 6A, compared with the control groups, mRNA expression of PMP22 was significantly downregulated following SAL treatment. Consistently, markedly weak fluorescent staining of PMP22 was also observed in SAL-treated MSCs (Fig. 6B), suggesting that PMP22 expression in MSCs is mediated by ER stress. These results indicate that VP1/ER stress/PMP22 signaling axis is an important component in MSC autophagy.

Discussion

In the present study, we investigated the role and mechanism of EV71 capsid protein VP1 in MSC autophagy, and demonstrated for the first time that VP1 promotes MSC autophagy through ER stress-mediated PMP22 upregulation, suggesting VP1/ER stress/PMP22 axis as a novel potential target against EV71-induced neuronal disorder in severe HFMD cases.

EV71 possesses four structural proteins including VP1, VP2, VP3, and VP4. VP1 homodimers are the main component of the characteristic icosahedral capsid contributing to the pathogenicity and stability of EV71 virus to survive in the environment of the gastrointestinal tract(39, 40). In the present study, we demonstrated that VP1 plays a promotive role in MSC autophagy (Fig. 2), which is consistent with our previous findings(25). However, the effect of VP1-induced autophagy on MSC survival still remains unclear because autophagy plays dual roles in the nervous system. Excessive autophagy may be protective in chronic neurodegenerative diseases but detrimental in acute neural damages(18, 41). It has

been reported that inhibition of EV71-induced autophagy in human rhabdomyosarcoma cells inhibits cell apoptosis at autophagosome formation stage and autophagy execution stage, but promotes apoptosis at the autophagosome-lysosome fusion stage. Furthermore, the inhibition of autophagy in the autophagosome formation stage or apoptosis decreases the release of EV71 viral particles, which is an effective strategy against virus infection(42). On the other hand, EV71-induced autophagy promotes viral replication in human rhabdomyosarcoma and neuroblastoma cells, and aggravates physiopathological parameters including weight loss, disease symptoms, and mortality in mouse models(23, 24). Further *in vitro* and *in vivo* studies are required to clarify the exact role of VP-induced autophagy in neuron cells.

In the present study, we also found that VP1 overexpression upregulated an important ER stress-associated protein PMP22(35, 36) in MSCs (Fig. 3), suggesting an involvement of ER stress activation in VP1-induced autophagy. It is well-established that excessive or premature PMP22 retaining in the ER induces ER stress(35, 36). However, the effect of ER stress activation on PMP22 expression hasn't been investigated yet. Our data revealed for the first time that inhibition of ER stress significantly downregulated the expression of PMP22 in MSCs (Fig. 6), suggesting that PMP22 is a downstream effector of ER stress. It seems that there is a positive feedback loop between ER stress and PMP22 in MSCs. Furthermore, our results showed that, in PMP22-deficient MSCs, there was no morphological signs of autophagy and the autophagy marker LC3B-II was remarkably downregulated (Fig. 4 and 5), suggesting that PMP22 is essential for MSC autophagy. Interestingly, in an EV71-infected mouse model, VP1 was found co-localized with LC3 and/or autophagosome-like vesicles in neurons, and VP1 expression was positively

correlated with LC3-II expression, aggregation and autophagosome formation(24). Upregulation of LC3-II expression was also observed in VP1-transfected HEK293 cells(25). Considering the regulatory role of VP1 in both MSC autophagy and PMP22 expression (Fig. 3), we conclude that VP1/ER stress/PMP22 pathway may play an important role in activation of MSC autophagy. In summary, our data demonstrated that MSC autophagy can be activated by EV71 capsid protein VP1. Mechanistically, the expression of ER stress-associated protein PMP22 was significantly upregulated by VP1, suggesting that ER stress-mediated PMP22 upregulation is possibly responsible for VP1-induced autophagy activation. The VP1/ER stress/PMP22 axis may serve as a potential therapeutic target against EV71 infection.

Materials and Methods

Cell line and culture

Mouse Schwann cells (MSC) were purchased from ScienCell Research Laboratories (Carlsbad, CA, USA) and maintained in Schwann cell medium (ScienCell) containing penicillin (100U/mL)/ streptomycin (100 µg/mL) (Hyclone, Logan, UT, USA) in poly-L-lysine-coated (2 µg/cm²) flasks at 37°C in a humidified atmosphere of 5% CO₂.

Sample collection

EV71 were isolated from clinical specimens including throat, anal swabs and stools of HFMD patients with EV71 infection, and were provided by the Center for Disease Control and Prevention of Guangdong Province (Guangzhou, Guangdong, China). The patient was diagnosed by Guangxi Medical University (Nanning, Guangxi, China)

based on the pathological analysis by Forensic Identification Center, Zhongshan School of Medicine, Sun Yat-sen University (Guangzhou, Guangdong, China).

Gene cloning and transfection

Total RNA was extracted from EV71 using Trizol (Invitrogen, Carlsbad, CA, USA). The 894-bp VP1 cDNA was synthesized by reverse transcription polymerase chain reaction (RT-PCR) using the primer sets 5'-CCGCTCGAGGCCACCATGGGTGATGGAATTGCAGACATGA-3' (forward) and 5'-CGCGGATCCTAGTGTTGTTATTTGTCCCTACTTGTGC-3' (reverse) (Genewiz, Suzhou, Jiangsu, China). The PCR products were then subcloned into pEGFP-C3 (Green Fluorescent Protein, GFP) expression vector (Clontech, Terra., USA) and sequencing was performed by Sangon Biotech (Shanghai, China). The results were compared with VP1 cDNA sequence reported by GenBank database. Cells were transiently transfected with plasmids using Lipofectamine 2000 (Invitrogen) following the manufacturer's instruction.

Small interfering RNA (siRNA)

siRNA against PMP22 (siPMP22) was from Santa Cruz Biotechnology (Dallas, TX, USA) and transfected using siRNA transfection reagent (Santa Cruz Biotechnology). Scramble siRNA (siCtrl) was used as a negative control.

Quantitative real-time PCR (qPCR)

Total RNA was extracted from cells using Trizol (Invitrogen) following the manufacturer's instructions and was reversely transcribed into cDNA using reverse transcriptase (Promega, Madison, WI, UDA). Real-time PCR was performed using SYBR Green qPCR SuperMix (Invitrogen) and the primers as shown in Table 1, following the manufacturer's instruction. GAPDH was used as an internal control. Data were analyzed using the $2^{-\Delta\Delta C_t}$ method.

Western blot analysis

MSCs were lysed and the lysates were collected. Protein concentration was determined using BCA protein assay reagent (?). 50 ng of proteins were separated by 10% SDS-PAGE gel and transferred to polyvinylidene fluoride membranes. The membranes were then blocked with 5% nonfat milk powder in Tris-buffered saline and Tween 20 (TBST), and then incubated with anti-GAPDH (1:1000; Abcam, Cambridge, UK) or anti-LC3B (1:1000; Abcam) for 1–2 h at room temperature. Following 3 washes with cold TBST, the membranes were incubated with peroxidase-conjugated secondary antibody (1:4000; Thermo Fisher Scientific, Rockford, IL 61105 USA) for additional 1 h at room temperature. After 3 washes with TBST, the protein expression was detected using enhanced chemiluminescent development reagent (GE Healthcare, Little Chalfont, UK) and X-ray films.

Immunofluorescence staining

MSCs were seeded on sterile coverslips 48 h after transfection and incubated overnight at 37 °C. Cells were then fixed with 4% paraformaldehyde for 30 min, followed by incubation with 0.2% Triton-X 100 at 4 °C for 5 min. After phosphate-buffered saline (PBS) washes, cells were blocked with 10% normal goat serum (Jackson ImmunoResearch, West Grove, PA, USA) for 30 min and incubated with anti-PMP22 antibody (Abcam) overnight at 4 °C. Cells were then incubated with fluorescence-conjugated secondary antibodies (Thermo Fisher Scientific, Waltham, MA, USA) for 1 h at room temperature. The images of stained cells were captured with a Leica camera (Leica, Wetzlar, Germany).

Transmission electron microscopy (TEM) analysis

MSCs were prefixed with 2.5% glutaraldehyde for 2 h and postfixes with 1% osmic acid for additional 2 h at 4 °C, followed by gradient dehydration in 30%, 50%, and 70%

ethanol (10 min each), 80%, 90%, and 95% acetone (10 min each), and 100% acetone (10 min twice). Cells were then embedded in the resin and stained with lead citrate. The stained cells were observed and imaged under a Hitachi H-7500 transmission electron microscope (Hitachi, Tokyo, Japan).

Statistical analysis

All experiments were repeated at least three times. Data were expressed as the mean $\pm$ standard error (SE). Statistical significance was assessed using Student's *t* test or one-way ANOVA with SPSS16.0 statistical software (SPSS Inc, IL, USA). $P < 0.05$ was considered statistically significant.

Acknowledgements

None

Conflict of interests

All authors declare that they have no conflict of interests.

References

1. **Liu MY, Liu J, Lai W, Luo J, Liu Y, Vu GP, Yang Z, Trang P, Li H, Wu J.** 2016. Characterization of enterovirus 71 infection and associated outbreak of Hand, Foot, and Mouth Disease in Shawo of China in 2012. *Sci Rep* **6**:38451.
2. **Wang Y, Zou G, Xia A, Wang X, Cai J, Gao Q, Yuan S, He G, Zhang S, Zeng M, Altmeyer R.** 2015. Enterovirus 71 infection in children with hand, foot, and mouth disease in Shanghai, China: epidemiology, clinical feature and diagnosis. *Virol J* **12**:83.
3. **Aswathyraj S, Arunkumar G, Alidjinou EK, Hober D.** 2016. Hand, foot and mouth disease (HFMD): emerging epidemiology and the need for a vaccine strategy. *Med Microbiol Immunol* **205**:397-407.
4. **Shimizu H, Nakashima K.** 2014. Surveillance of hand, foot, and mouth disease for a vaccine. *Lancet Infect Dis* **14**:262-263.
5. **Zhang Y, Tan XJ, Wang HY, Yan DM, Zhu SL, Wang DY, Ji F, Wang XJ, Gao YJ, Chen L, An HQ, Li DX, Wang SW, Xu AQ, Wang ZJ, Xu WB.** 2009. An outbreak of hand, foot, and mouth disease associated with subgenotype C4 of human enterovirus 71 in Shandong, China. *J Clin Virol* **44**:262-267.
6. **Chong CY, Chan KP, Shah VA, Ng WY, Lau G, Teo TE, Lai SH, Ling AE.** 2003. Hand, foot and mouth disease in Singapore: a comparison of fatal and non-fatal cases. *Acta Paediatr* **92**:1163-1169.
7. **Sabanathan S, Tan le V, Thwaites L, Wills B, Qui PT, Rogier van Doorn H.** 2014. Enterovirus 71 related severe hand, foot and mouth disease outbreaks in South-East Asia: current situation and ongoing challenges. *J Epidemiol Community Health* **68**:500-502.
8. **Solomon T, Lewthwaite P, Perera D, Cardosa MJ, McMinn P, Ooi MH.** 2010. Virology, epidemiology, pathogenesis, and control of enterovirus 71. *Lancet Infect Dis* **10**:778-790.
9. **Gui J, Liu Z, Zhang T, Hua Q, Jiang Z, Chen B, Gu H, Lv H, Dong C.** 2015. Epidemiological Characteristics and Spatial-Temporal Clusters of Hand, Foot, and Mouth Disease in Zhejiang Province, China, 2008-2012. *PLoS One* **10**:e0139109.
10. **Xing WJ, Liao QH, Viboud C, Zhang J, Sun JL, Wu JT, Chang ZR, Liu FF, Fang VJ, Zheng YD, Cowling BJ, Varma JK, Farrar JJ, Leung GM, Yu HJ.** 2014. Hand, foot, and mouth disease in China, 2008-12: an epidemiological study. *Lancet Infectious Diseases* **14**:308-318.
11. **Hsueh C, Jung SM, Shih SR, Kuo TT, Shieh WJ, Zaki S, Lin TY, Chang LY, Ning HC, Yen DC.** 2000. Acute encephalomyelitis during an outbreak of enterovirus type 71 infection in Taiwan: report of an autopsy case with pathologic, immunofluorescence, and molecular studies. *Mod Pathol* **13**:1200-1205.
12. **Kuo RL, Kung SH, Hsu YY, Liu WT.** 2002. Infection with enterovirus 71 or expression of its 2A protease induces apoptotic cell death. *J Gen Virol* **83**:1367-1376.
13. **Li ML, Hsu TA, Chen TC, Chang SC, Lee JC, Chen CC, Stollar V, Shih SR.** 2002. The 3C protease activity of enterovirus 71 induces human neural cell apoptosis. *Virology* **293**:386-395.
14. **Yang SD, Li PQ, Li YM, Li W, Lai WY, Zhu CP, Tao JP, Deng L, Liu HS, Ma WC, Lu JM, Hong Y, Liang YT, Shen J, Hu DD, Gao YY, Zhou Y, Situ MX, Chen YL.** 2017. Clinical manifestations of severe enterovirus 71 infection and early assessment in a Southern China population. *BMC Infect Dis* **17**:153.
15. **Klionsky DJ.** 2007. Autophagy: from phenomenology to molecular understanding in less than a decade. *Nat Rev Mol Cell Biol* **8**:931-937.
16. **Mizushima N, Klionsky DJ.** 2007. Protein turnover via autophagy: implications for metabolism. *Annu Rev Nutr* **27**:19-40.

- 341 17. **Ebrahimi-Fakhari D, Wahlster L, Hoffmann GF, Kolker S.** 2014. Emerging role of
342 autophagy in pediatric neurodegenerative and neurometabolic diseases. *Pediatr Res*
343 **75**:217-226.
- 344 18. **Ginet V, Spiehlmann A, Rummel C, Rudinskiy N, Grishchuk Y, Luthi-Carter R, Clarke**
345 **PG, Truttmann AC, Puyal J.** 2014. Involvement of autophagy in hypoxic-excitotoxic
346 neuronal death. *Autophagy* **10**:846-860.
- 347 19. **Lee JH, Yu WH, Kumar A, Lee S, Mohan PS, Peterhoff CM, Wolfe DM, Martinez-**
348 **Vicente M, Massey AC, Sovak G, Uchiyama Y, Westaway D, Cuervo AM, Nixon RA.**
349 2010. Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted
350 by Alzheimer-related PS1 mutations. *Cell* **141**:1146-1158.
- 351 20. **Klionsky DJ, Abdelmohsen K, Abe A, Abedin MJ, Abeliovich H, Acevedo Arozena A,**
352 **Adachi H, Adams CM, Adams PD, Adeli K, Adhihetty PJ, Adler SG, Agam G, Agarwal**
353 **R, Aghi MK, Agnello M, Agostinis P, Aguilar PV, Aguirre-Ghiso J, Airolidi EM, Ait-Si-**
354 **Ali S, Akematsu T, Akporiaye ET, Al-Rubeai M, Albaiceta GM, Albanese C, Albani D,**
355 **Albert ML, Aldudo J, Algul H, Alirezaei M, Alloza I, Almasan A, Almonte-Beceril M,**
356 **Alnemri ES, Alonso C, Altan-Bonnet N, Altieri DC, Alvarez S, Alvarez-Erviti L, Alves S,**
357 **Amadoro G, Amano A, Amantini C, Ambrosio S, Amelio I, Amer AO, Amessou M,**
358 **Amon A, An Z, et al.** 2016. Guidelines for the use and interpretation of assays for
359 monitoring autophagy (3rd edition). *Autophagy* **12**:1-222.
- 360 21. **Tanida I, Ueno T, Kominami E.** 2008. LC3 and Autophagy. *Methods Mol Biol* **445**:77-
361 88.
- 362 22. **Benito-Cuesta I, Diez H, Ordonez L, Wandosell F.** 2017. Assessment of Autophagy in
363 Neurons and Brain Tissue. *Cells* **6**.
- 364 23. **Huang SC, Chang CL, Wang PS, Tsai Y, Liu HS.** 2009. Enterovirus 71-induced
365 autophagy detected in vitro and in vivo promotes viral replication. *J Med Virol*
366 **81**:1241-1252.
- 367 24. **Lee YR, Wang PS, Wang JR, Liu HS.** 2014. Enterovirus 71-induced autophagy
368 increases viral replication and pathogenesis in a suckling mouse model. *J Biomed Sci*
369 **21**:80.
- 370 25. **Hu DD, Mai JN, He LY, Li PQ, Chen WX, Yan JJ, Zhu WD, Deng L, Wei D, Liu DH, Yang**
371 **SD, Yao ZB.** 2017. Glucocorticoids Prevent Enterovirus 71 Capsid Protein VP1
372 Induced Calreticulin Surface Exposure by Alleviating Neuronal ER Stress. *Neurotox*
373 *Res* **31**:204-217.
- 374 26. **Rashid HO, Yadav RK, Kim HR, Chae HJ.** 2015. ER stress: Autophagy induction,
375 inhibition and selection. *Autophagy* **11**:1956-1977.
- 376 27. **Lin W, Popko B.** 2009. Endoplasmic reticulum stress in disorders of myelinating cells.
377 *Nat Neurosci* **12**:379-385.
- 378 28. **Lee WS, Yoo WH, Chae HJ.** 2015. ER Stress and Autophagy. *Curr Mol Med* **15**:735-
379 745.
- 380 29. **Cai Y, Arikath J, Yang L, Guo ML, Periyasamy P, Buch S.** 2016. Interplay of
381 endoplasmic reticulum stress and autophagy in neurodegenerative disorders.
382 *Autophagy* **12**:225-244.
- 383 30. **Zou X, Xu J, Yao S, Li J, Yang Y, Yang L.** 2014. Endoplasmic reticulum stress-mediated
384 autophagy protects against lipopolysaccharide-induced apoptosis in HL-1
385 cardiomyocytes. *Exp Physiol* **99**:1348-1358.
- 386 31. **Cheng X, Liu H, Jiang CC, Fang L, Chen C, Zhang XD, Jiang ZW.** 2014. Connecting
387 endoplasmic reticulum stress to autophagy through IRE1/JNK/beclin-1 in breast
388 cancer cells. *Int J Mol Med* **34**:772-781.
- 389 32. **Snipes GJ, Suter U, Welcher AA, Shooter EM.** 1992. Characterization of a novel
390 peripheral nervous system myelin protein (PMP-22/SR13). *J Cell Biol* **117**:225-238.

- 391 33. **Pareek S, Suter U, Snipes GJ, Welcher AA, Shooter EM, Murphy RA.** 1993. Detection
392 and processing of peripheral myelin protein PMP22 in cultured Schwann cells. *J Biol*
393 *Chem* **268**:10372-10379.
- 394 34. **Ryan MC, Notterpek L, Tobler AR, Liu N, Shooter EM.** 2000. Role of the peripheral
395 myelin protein 22 N-linked glycan in oligomer stability. *J Neurochem* **75**:1465-1474.
- 396 35. **Dickson KM, Bergeron JJ, Shames I, Colby J, Nguyen DT, Chevet E, Thomas DY,**
397 **Snipes GJ.** 2002. Association of calnexin with mutant peripheral myelin protein-22
398 ex vivo: a basis for "gain-of-function" ER diseases. *Proc Natl Acad Sci U S A* **99**:9852-
399 9857.
- 400 36. **Hara T, Hashimoto Y, Akuzawa T, Hirai R, Kobayashi H, Sato K.** 2014. Rer1 and
401 calnexin regulate endoplasmic reticulum retention of a peripheral myelin protein 22
402 mutant that causes type 1A Charcot-Marie-Tooth disease. *Sci Rep* **4**:6992.
- 403 37. **Shibutani ST, Yoshimori T.** 2014. A current perspective of autophagosome
404 biogenesis. *Cell Res* **24**:58-68.
- 405 38. **Barth S, Glick D, Macleod KF.** 2010. Autophagy: assays and artifacts. *J Pathol*
406 **221**:117-124.
- 407 39. **Shi M, Zhou Y, Cao L, Ding C, Ji Y, Jiang Q, Liu X, Li X, Hou X, Peng H, Shi W.** 2013.
408 Expression of enterovirus 71 capsid protein VP1 in *Escherichia coli* and its clinical
409 application. *Braz J Microbiol* **44**:1215-1222.
- 410 40. **Lal SK, Kumar P, Yeo WM, Kar-Roy A, Chow VTK.** 2006. The VP1 protein of human
411 enterovirus 71 self-associates via an interaction domain spanning amino acids 66-
412 297. *Journal of Medical Virology* **78**:582-590.
- 413 41. **Puyal J, Ginet V, Grishchuk Y, Truttmann AC, Clarke PG.** 2012. Neuronal autophagy
414 as a mediator of life and death: contrasting roles in chronic neurodegenerative and
415 acute neural disorders. *Neuroscientist* **18**:224-236.
- 416 42. **Xi XY, Zhang XY, Wang B, Wang T, Wang J, Huang H, Wang JW, Jin Q, Zhao ZD.** 2013.
417 The Interplays between Autophagy and Apoptosis Induced by Enterovirus 71. *Plos*
418 *One* **8**.

419

420

Figure legends

Figure 1. Cloning and identification of VP1 cDNA. Agarose electrophoresis for intact (lane 1) and restriction enzyme-digested (lane 2) VP1 cDNA cloning vector pEGFP-C3 plasmids. M: DNA marker.

Figure 2. The effect of VP1 overexpression on MSC autophagy. MSCs were transfected with pEGFP-C3-VP1 plasmids for 48 h. Untransfected and pEGFP-C3-transfected cells were used as blank and negative controls, respectively. Representative transmission electron microscopic images depict subcellular structures of MSCs. N: nucleus, M: mitochondrion, L: lysosome, AP: autophagosome, AL: autolysosome, DV: degradation vesicles, GA: Golgi apparatus, ER: endoplasmic reticulum, SV: secretory vesicles. MSC, mouse Schwann cell.

Figure 3. The effect of VP1 on PMP22 expression in MSCs. MSCs were transfected with pEGFP-C3-VP1 plasmids for 48 h. Untransfected and pEGFP-C3-transfected cells were used as blank and negative controls, respectively. A. The mRNA expression of PMP22 was detected by qPCR. Data are expressed as the mean $\pm$ SE; *** $P < 0.001$ vs. untransfected group; $n = 3$. B. Immunofluorescent staining for PMP22 in pEGFP-C3- or pEGFP-C3-VP1-transfected MSCs. GFP expression was used to monitor the transfection efficacy. Magnification: 400 $\times$. MSC, mouse Schwann cell; SE, standard error.

446

447 **Figure 4. The effect of PMP22 knockdown on autophagy marker LC3B-II in**
 448 **MSCs.** MSCs were transfected with siPMP22. Untransfected and scramble siRNA-
 449 trasfected cells were used as blank and negative controls, respectively. mRNA and
 450 proteins expression of LC3B were detected by real-time PCR (A) and Western blot
 451 assay (B), respectively. (C) Quantification of Western blot assay. (D) Ratio of LC3B-
 452 II to LC3B-I. Data are expressed as the mean $\pm$ SE; *** $P < 0.001$ vs. untransfected
 453 group; n = 3. MSC, mouse Schwann cell; SE, standard error.

454

455

456 **Figure 5. The effect of PMP22 knockdown on cellular and subcellular**
 457 **morphology of MSCs.** MSCs were transfected with siPMP22 for 48 h. Untransfected
 458 and scramble siRNA-trasfected cells were used as blank and negative controls (siCtrl),
 459 respectively. Representative transmission electron microscopic images depict
 460 subcellular structures of MSCs. N: nucleus, M: mitochondrion, L: lysosome, AP:
 461 autophagosome, AL: autolysosome, DV: degradation vesicles, GA: Golgi apparatus.
 462 MSC, mouse Schwann cell.

463

464

465 **Figure 6. The effect of ER stress activation on PMP22 expression in MSCs.** MSCs
 466 were treated with 15 μ M of selective ER stress inhibitor salubrinal (SAL) for 48 h.
 467 Untreated and DMSO-treated cells were used as blank and negative controls. (A) The
 468 mRNA level of PMP22 was detected by real-time PCR. Data are expressed as the
 469 mean $\pm$ SEM; *** $P < 0.001$ vs. untransfected group; n = 3. (B) Immunofluorescent
 470 staining (red) for PMP22 in DMSO- or SAL-treated MSCs. Furo 8-AM in SAL

471 showed green fluorescence. Magnification: 400×. ER, endoplasmic reticulum; MSC,
472 mouse Schwann cell; SE, standard error.

473

474

Table 1 Real-time PCR primers

Gene	Forward primer (5'–3')	Reverse primer (5'–3')
PMP22	CTGCCAGCTCTTCACTCTCA	GTTGACATGCCACTCACTGT
GAPDH	GGCCTCCAAGGAGTAAGAAA	GCCCCTCCTGTTATTATGG

MSC

pEGFP-C3-trasfected × 25000

17073-16.tif
Print Mag: 27200x @ 7.0 in
10:12:58 3/8/2017

800 nm
HV=80.0kV
Direct Mag: 25000x
AMT Camera System

Untransfected ×30000

17032-18.tif
Print Mag: 32500x @ 7.0 in
16:32:45 3/2/2017

600 nm
HV=80.0kV
Direct Mag: 30000x
AMT Camera System

×25000

17073-25.tif
Print Mag: 27200x @ 7.0 in
10:41:38 3/8/2017

800 nm
HV=80.0kV
Direct Mag: 25000x
AMT Camera System

×30000

17032-33.tif
Print Mag: 32900x @ 7.0 in
08:24:53 3/3/2017

600 nm
HV=80.0kV
Direct Mag: 30000x
AMT Camera System

pEGFP-C3-VP1

A **B**

The gray level of anti-LC3B western blotting

The autophagy rate by detecting the gray level of anti-LC3B western blotting

siPMP22

×20000

17032-6.tif
Print Mag: 21700x @ 7.0 in
16:13:56 3/2/2017

1 µm
HV=80.0kV
Direct Mag: 20000x
AMT Camera System

×30000

17032-17.tif
Print Mag: 32900x @ 7.0 in
16:31:23 3/2/2017

500 nm
HV=80.0kV
Direct Mag: 30000x
AMT Camera System

A

B