1 Nurtured by nature: Considering the role of environmental and parental legacies in
2 coral ecological performance

3

4 Hollie M. Putnam¹, Raphael Ritson-Williams², Jolly Ann Cruz³, Jennifer M. Davidson²,
 5 Ruth D. Gates²

6

7 ¹University of Rhode Island, Department of Biological Sciences, Kingston, RI USA

8 ²Hawai‘i Institute of Marine Biology, University of Hawai‘i, Mānoa, HI, USA

9 ³Micronesia Islands Nature Alliance, Garapan, Saipan 96950, CNMI

10 * Corresponding author: hputnam@uri.edu, 401-874-9510

11

12 Keywords: trans-generational plasticity, epigenetics, hormesis, *Pocillopora damicornis*,
 13 *Pocillopora acuta*

14

15

16

17

18

19

20

21

22

23

Positive carryover effects in corals

Abstract

The persistence of reef building corals is threatened by human-induced environmental change. Maintaining coral reefs into the future requires not only the survival of adults, but also the influx of recruits to promote genetic diversity and retain cover following adult mortality. Few studies examine the linkages among multiple life stages of corals, despite a growing knowledge of carryover effects in other systems. We provide a novel test of coral parental preconditioning to ocean acidification (OA) to better understand impacts on the processes of offspring recruitment and growth. Coral planulation was tracked for three months following adult exposure to high pCO₂ and offspring from the second month were reciprocally exposed to ambient and high pCO₂. Offspring of parents exposed to high pCO₂ had greater settlement and survivorship immediately following release, retained survivorship benefits during one and six months of continued exposure, and further displayed growth benefits to at least one month post release. Enhanced performance of offspring from parents exposed to high conditions was maintained despite the survivorship in both treatments declining in continued exposure to OA. Preconditioning of the adults while they brood their larvae may provide a form of hormetic conditioning, or environmental priming that elicits stimulatory effects. Defining mechanisms of positive carryover effects, or positive trans-generational plasticity, is critical to better understanding ecological and evolutionary dynamics of corals under regimes of increasing environmental disturbance. Considering parental and environmental legacies in ecological and evolutionary projections may better account for coral reef response to the chronic stress regimes characteristic of climate change.

Introduction

Reef-building corals are the foundation for one of the most diverse and economically important ecosystems (Bishop et al., 2011; Costanza et al., 2014). Corals are sensitive to changes in their environment that destabilize the coral-algal symbiosis and disrupt reef primary production and net ecosystem accretion. Even slight perturbations in temperature, nutrients, light, salinity, and pollution, can negatively affect coral symbiosis and performance (Lesser, 2011). The steady increase in CO₂ emissions (Pachauri et al., 2014) is heightening the magnitude and frequency of global scale environmental perturbations, exacerbating reef decline already occurring due to local pressures.

The emission of CO₂ is driving elevated atmospheric levels, which is in turn changing the chemistry of the oceans. Ocean acidification (OA) is occurring due to the absorption of atmospheric CO₂ by the ocean and the resulting increase and decrease in the concentrations of H⁺ and CO₃⁻² ion concentrations, respectively (Doney, Fabry, Feely, & Kleypas, 2009). OA energetically challenges calcifying marine organisms because they expend more energy for acid-base regulation to both maintain cellular homeostasis (Kaniewska et al., 2012; Stumpp et al., 2012) and calcify in a medium with lower CO₃⁻² ion concentrations. The physiological and ecological consequences of OA range from changes in intracellular pH (Gibbin, Putnam, Davy, & Gates, 2014; Gibbin, Putnam, Gates, Nitschke, & Davy, 2015), to direct reduction of calcification of coral skeletons (Erez, Reynaud, Silverman, Schneider, & Allemand, 2011), to shifts in community composition in low pH areas (Fabricius et al., 2011). Further, the interaction of multiple stressors such as temperature and OA is likely to result in synergistic and antagonistic

Positive carryover effects in corals

responses in comparison to the responses driven by individual stressors (Harvey, Gwynn-Jones, & Moore, 2013; Pandolfi, Connolly, Marshall, & Cohen, 2011).

Given that local coral reef decline is now exacerbated by global issues, a greater effort is being made to better understand the sub-lethal physiological effects of stressors on critical coral reef processes such as coral reproduction and early life stages (Albright, 2011; Byrne & Przeslawski, 2013). Corals are long-lived organisms with complex reproductive life history characteristics (Baird, Guest, & Willis, 2009). Early life history research has focused heavily on the effects of anthropogenic factors (e.g., increased temperature and ocean acidification) on fertilization and development, metamorphosis, settlement, and survivorship in spawning corals (Albright, 2011; Chua, Leggat, Moya, & Baird, 2013; Chua, Leggat, Moya, & Baird, 2013; Foster, Gilmour, Chua, Falter, & McCulloch, 2015; Negri, Marshall, & Heyward, 2007; Randall & Szmant, 2009), and a variety of physiological metrics in brooding corals (Cumbo, Edmunds, Wall, & Fan, 2013; Dufault, Cumbo, Fan, & Edmunds, 2012; Putnam, Edmunds, & Fan, 2010; Rivest & Hofmann, 2014, 2015). Few studies have, however, tracked corals through multiple life history stages including larval supply, settlement and post-settlement survival and growth (Albright, 2011; Ritson-Williams, Arnold, & Paul, 2016; Ross, Ritson-Williams, Olsen, & Paul, 2013), fewer still at the cross-generational scale (Putnam & Gates, 2015), and none at the multigenerational scale. Thus the forecasts for population persistence and replenishment have largely ignored the potentially substantial plasticity driven by environmental and parental legacies (Donelson, Salinas, Munday, & Shama, 2017; Torda et al., 2017).

Positive carryover effects in corals

Phenotypic plasticity, or the generation of multiple performance outcomes within an individual as a function of environment, has the potential to act as a buffer to rapid environmental change and modulate evolutionary response (Chevin, Collins, & Lefèvre, 2013; Gibbin et al., 2017; Putnam & Gates, 2015). Multiple avenues can generate this plasticity. While intra-generational studies of phenotypic plasticity are more common, the study of trans-generational plasticity (TGP) is still in its relative infancy for marine invertebrates (Donelson et al., 2017; Foo & Byrne, 2016; Ross, Parker, & Byrne, 2016; Torda et al., 2017). TGP occurs when environmental legacies manifest through linked life stages (i.e., the norms of reaction are shaped by the environmental conditions of prior generations). Due to the difficulty of disentangling TGP *sensu stricto* (Donelson et al., 2017; Torda et al., 2017), the plasticity exhibited between life stages can also be more generically discussed as carryover effects, which encompass the potential for TGP and developmental acclimation. The lack of data on these life stage linkages in corals is now particularly critical in light of severe ecosystem declines and the growing acknowledgement that adaptive responses are necessary for persistence under rapid climate change (Gaylord et al., 2015; Munday, Warner, Monro, Pandolfi, & Marshall, 2013; Sunday et al., 2014; Sunday, Crim, Harley, & Hart, 2011).

The ecological and evolutionary ramifications of carryover effects from parental or early life stage exposure in longer-lived, ecosystem engineering, calcifying marine invertebrates in response to OA remain to be determined. To date, early life stage exposure often appears to have negative carryover effects. Assessment of Olympia oyster performance following exposure to high pCO₂ at the larval stage by Hettinger and

Positive carryover effects in corals

coauthors revealed slower growth under high conditions. Even after the juveniles were placed in ambient grow out conditions, negative carryover effects remained at 45 days post exposure (Hettinger et al., 2012). Further assessment of the ecological performance of Olympia oyster growth in ambient field conditions four months post exposure to high pCO₂, revealed negative carryover effects on growth rate (Hettinger et al., 2013). Some corals also appear to display negative carryover effects from stressors at the larval stage. In larvae of *Orbicella faveolata*, low salinity treatments caused decreased survival and growth after settlement (Vermeji et al., 2006). Similarly, in larvae of *Porites astreoides* early life exposure to elevated seawater temperatures caused significantly higher mortality in settled coral spat ~3 weeks post exposure in comparison to larvae raised in ambient temperatures (Ross et al., 2013). While studies in both oysters and corals identify carryover effects of early life stage exposure to stressors, this does not account for plasticity that may be induced by parental environment (i.e., TGP).

Depending on the timing of exposure, environmental conditioning can set organisms on different physiological and ecological trajectories (Foo & Byrne, 2016; Ross et al., 2016; Torda et al., 2017). Studies considering the ontogenetic connections in organisms with shorter life spans such as plankton, worms, oysters, clams, and fishes, have documented substantial, and often positive TGP (Chakravarti et al., 2016; Foo & Byrne, 2016; Gibbin et al., 2017; Lane, Campanati, Dupont, & Thiyagarajan, 2015; Miller, Watson, Donelson, McCormick, & Munday, 2012; Parker et al., 2012, 2017; Ross et al., 2016; Thor & Dupont, 2015; Zhao et al., 2018). For example, in the Sydney rock oyster, parental exposure to high pCO₂ enhanced growth, development, and metabolism in F1 offspring exposed to the same high pCO₂ conditions (Parker et al.,

Positive carryover effects in corals

2012). Further, positive TGP was again observed in the F2 generation, in terms of development, growth, and juvenile heart rate in the trans-generational exposure line, relative to the control line (Parker, O'Connor, Raftos, Pörtner, & Ross, 2015). To date, only one study of reef building corals has demonstrated cross-generational acclimatization at the swimming planulae stage (Putnam & Gates, 2015), but the longer-term implications of this plasticity across subsequent life stages are not yet known. These responses demonstrate that understanding the potential carryover effects from parents through ontogeny (i.e., ecological ramifications of TGP) is critical to our view of organism performance on annual to decadal time scales.

The ecological implications of carryover effects have not been considered in projections of coral performance in a future of climate change, yet acclimatization is known to play a key role in intra-generational stress response (Brown & Cossins, 2011; Brown, Dunne, Goodson, & Douglas, 2000; Coles & Brown, 2003), and our initial work in corals suggests positive TGP is possible in corals (Putnam & Gates, 2015). Our current study investigates the effects of adult preconditioning on offspring ecological performance in response to OA conditions. Here, we tested the hypothesis that parental exposure of corals to high pCO₂ during gametogenesis and/or brooding results in beneficial acclimation of the offspring. Further, we tested the hypothesis that these carryover effects are maintained beyond the larval stage, through settlement into the juvenile stage, by assessing settlement and survivorship post release and tracking survivorship and growth for six months in ambient and elevated pCO₂ treatments.

Materials and Methods

Experimental Overview

The experiment consisted of four phases (Fig. 1). First, adult coral colonies were acclimated to ambient conditions in common garden tanks for 34d (Fig. 1a). Second, adults were exposed to two different pCO₂ treatments that fluctuated with a diurnal frequency for the period that contained gametogenesis and brooding, over 3 cycles of planulation (~3 months, Fig 1b.). Third, larvae were collected from the pre-conditioned adults and exposed to the two different pCO₂ conditions in a reciprocal fashion (Fig. 1c-e) in acrylic and mesh chambers containing a settlement tile: Survivorship and settlement of offspring during this phase were assessed after 96h. Fourth, the settled spat were tracked for survivorship and growth (Fig. 1f) in continued reciprocal exposure after one and six months post release.

Positive carryover effects in corals

Figure 1. Experimental Design (a) Coral acclimation and (b) experimental timeline to test the effects of adult preconditioning on larval release larval performance at (c, d) ambient and high pCO₂ conditions in terms of (e) settlement, survivorship and (f) growth and survivorship were examined.

172

173 *Coral Collection and Acclimation*

174 Adult *Pocillopora damicornis* colonies (now considered *P. acuta* based on mitochondrial
175 open reading frame, mtORF, sequencing (Johnston, Forsman, & Toonen, 2018)) were
176 collected from the fringing reefs of southern Kāneohe Bay, O’ahu, Hawai’i
177 (21.429845, -157.793604) under Special Activities Permit SAP2014 from Hawaii’s
178 Division of Aquatic Resources (DAR). Corals were removed from the reef at the dead
179 skeleton base to minimize direct tissue damage and were placed immediately adjacent to
180 each other in a common garden in the field for a 3-day acclimation period (e.g., Putnam,
181 Davidson, & Gates, 2016). Subsequently, corals (n=16) were moved to large, outdoor,

Positive carryover effects in corals

sand-filtered, flow-through seawater, mesocosm tanks (~1,300L) at the Hawai'i Institute of Marine Biology, HIMB (described in Putnam et al., 2016). Irradiance in the mesocosm tanks was reduced to ~60% full irradiance using shade cloth to more closely mimic collection site *in situ* light conditions, and light readings were taken every 15 min with underwater loggers (Odyssey PAR loggers standardized to Li-Cor 192SA cosine sensor; Long et al. 2012; Fig. S1). Seawater temperature was also recorded every 15 min (Hobo Water Temp Pro v2, accuracy = 0.21°C, resolution = 0.02°C, Onset Computer Corporation, Fig. S1). Light, temperature and pH fluctuated within the tanks on natural cycles (Fig. S1). Corals acclimated to these tank conditions for 34d prior to initiation of treatments.

Experimental Exposure

The treatment (high pCO₂) and control conditions (ambient seawater) were maintained using a pH-stat CO₂ injection system (Putnam et al., 2016). Ambient pH and pCO₂ fluctuated daily (e.g., (Drupp, De Carlo, Mackenzie, Bienfang, & Sabine, 2011), driven by ambient conditions and feedbacks from photosynthesis, calcification, and respiration of the organisms on the fringing reef directly off shore of HIMB, where the seawater was obtained. The high CO₂ treatment conditions retained natural diel variability while decreasing the pH (Fig. S1). pH probes from the pH-stat CO₂ injection system were calibrated weekly on the NBS scale and pH and temperature were logged every 15 min in each tank throughout the duration of the experiment. The carbonate chemistry of the seawater was assessed with discrete measurements of pH (total scale), total alkalinity, temperature and salinity according to the Guide to Best Practices (Riebesell, Fabry,

Positive carryover effects in corals

Hansson, Gattuso, & others, 2010). Discrete measurements were made ~daily and water samples were collected ~2x per week for each adult treatment. Given the stability of the total alkalinity and low biomass in the 1,300L tanks, sampling frequency was reduced in the 6 months of offspring exposure to ~ every 1-2 weeks, while pH, temperature, and light were tracked continuously (Fig S1). Total alkalinity samples were analyzed using open cell potentiometric titrations (Dickson, Sabine, & Christian, 2007) and assessed against certified reference materials (CRMs; A. Dickson Laboratory, UCSD; values on average <1% different from TA CRMs); all samples were corrected for any offset from the CRMs. From these measurements, the full suite of carbonate parameters was calculated with the seacarb package (v3.0.11, Gattuso et al., 2016), using the average corrected TA and salinity measured in each treatment tank (Table 1).

Adult Exposure, Planulation, and Settlement

Eight adult corals were exposed to each treatment beginning 06 May 2014. Two colonies died very early in the high pCO₂ treatment prior to June planulation, leaving n=6 in high pCO₂ and n=8 in ambient pCO₂. A common garden approach (n = 2 tanks) was chosen to maximize the similarity of experimental conditions. During the 7-8 days of larval collection each month, adult corals were removed from the tanks and separated into individual ~4.5L bowls (one colony per bowl) with flowing treatment water (i.e., the same experimental conditions as the tanks) from ~5pm to 9am to isolate the larvae released from each colony (e.g., (Putnam & Gates, 2015). The flowing seawater flushed the buoyant larvae into 800 ml tripour beakers with 150 µm mesh bottoms and the

Positive carryover effects in corals

number of larvae released per colony were counted during the months of June, July and August 2014.

July 2014 was the expected seasonal peak of larval release (Fig. 2), and the larvae collected during this time were used for the offspring reciprocal exposure experiments (Fig. 1c, d). Due to the variation in timing of gametogenesis, brooding, and the sexual or asexual origin of the planulae in *Pocillopora damicornis/acuta* (Permata, Kinzie, & Hidaka, 2000; Stimson, 1978; Stoddart, 1983; Stoddart & Black, 1985), it is not possible to definitively state the exact timing of exposure of the offspring gametes or planulae within the adults. Clarifying the timing and mechanisms of carryover effects (i.e. TGP or developmental acclimation) would require, for example, exposing the adults only prior to gametogenesis, only during gametogenesis, or only during brooding. Currently the reproductive biology of *Pocillopora damicornis/acuta*, complicates this assessment. In this case, we are testing the ecological context of TGP (e.g., (Torda et al., 2017) box 2), or carryover effects.

Groups of 16-20 larvae were tracked from each parent colony and placed in a 200ml transparent acrylic chamber with 150µm mesh at each end. To increase replication and account for variability across release date, experiments were conducted on each day of larval release from 12-18 July 2014 on newly released larvae. Each chamber contained a 4.5 x 4.5 x 1cm terracotta brick tile that had been preconditioned on the reef for 1-2 months. Conditions of pH on the fringing reef of Coconut Island in Kāneohe Bay can range from ~7.6 - 8.1 (Guadayol, Silbiger, Donahue, & Thomas, 2014; Silbiger, Guadayol, Thomas, & Donahue, 2014), so these tiles are not naïve to high pCO₂. Further the major constituent was not crustose coralline algae and as such there should not be

Positive carryover effects in corals

strong impacts of pCO₂ on the tiles themselves, thereby providing relatively equal competition dynamics between the newly settled spat and tile communities. The indirect effects of pCO₂ on substrate community, and therefore coral offspring settlement and growth, are important considerations in experimental design for interpretation of results (Albright and Langdon 2011). Chambers were placed into either ambient or high pCO₂ treatments and settlement and survivorship were assessed after 96h. Survivorship was calculated as the total number of living larvae (both swimming and settled) in the chamber divided by the initial number of larvae added. Settlement was counted as those larvae that had settled and metamorphosed to the chamber walls, mesh, or tile divided by the initial number of larvae added to the chamber.

Spat exposure on Tiles

After settlement and survivorship was assessed, new recruits were mapped on each tile and the tiles with settled spat were returned to the mesocosms of their settlement treatment (Figure 1, Figs. S1-S2). After both an additional month and then 6 months of total exposure, tiles were re-assessed under a dissecting scope to count survivorship (number of spat remaining alive relative to the initial amount added to the chamber). Growth of each spat was also measured by counting the number of polyps. Growth rate was calculated as: the (# of polyps - 1 primary polyp) / (# of days post settlement at month 1) for the first month's growth rate, and as the (# of polyps at month 6 - # of polyps at month 1) / (# of days between month 1 and month 6 measurements). Spat that were fused were counted as survivors, but were not used in the growth data analysis. Settlement tile was used as the level of replication to avoid non-independence of multiple spat on a tile.

Positive carryover effects in corals

273

274 *Statistical Analysis*

275 To test if the timing of larval release differed between the control and treatment
 276 conditions at each sampling time-point, a two-sample Kolmogorov-Smirnov test was
 277 used and release by day was treated as a continuous variable (ks.test; stats package;(R
 278 Core Team, 2016). A generalized linear mixed effects model with binomial errors (lme4
 279 package; R Core Team, 2016) was used to test for differences in proportional settlement
 280 and survivorship between the treatment and control using a binomial distribution.
 281 Settlement data were analyzed with parental (Origin) and offspring (Secondary)
 282 treatments and their interaction as fixed effects and settlement tile as a random intercept.
 283 Survivorship and growth data were both measured at multiple time points and, thus, were
 284 modeled with the same fixed effects and interaction, but with a random intercept of
 285 settlement tile nested in time point. A model selection approach was applied and the final
 286 models were selected as those with the lowest delta AIC. Growth data were log
 287 normalized to meet model assumptions (i.e., for normal distribution) and growth data are
 288 plotted as back-transformed means and asymmetrical standard error. Full analytical
 289 details are available on GitHub
 290 (https://github.com/hputnam/HI_Pdam_Parental/releases/tag/Version_20180508) and
 291 Dryad (doi will be added upon acceptance).

292

293 **Results**

294 *Acclimation Conditions*

Positive carryover effects in corals

Corals were acclimated in tanks to fluctuating conditions (Fig. S1) similar to that on the fringing reefs of Kāneohe Bay, O'ahu, Hawai'i (Putnam et al., 2016). During the acclimation period, daily temperature ranged on average from 23.7 - 25.9°C (Fig. S1a). Between 06:00 - 19:00, Photosynthetically Active Radiation (PAR) ranged on average from 62 - 246 $\mu\text{mol s}^{-1} \text{m}^{-2}$ (Fig. S1e).

Experimental Exposure

Light and temperature varied over time within the experiment due to seasonal shifts (see supplemental Figure S1 for detailed description of tank conditions). During the adult pCO₂ exposure, mean temperature ranged from 26.3 - 27.9°C and 26.4 - 28.0°C in ambient and treatment tanks, respectively (Fig. S1), which was on average higher than the acclimation period due to seasonal warming. pH (NBS) ranged from 7.81 - 8.06 in ambient conditions and 7.51 - 7.74 in high pCO₂ conditions for the adult exposures. While this condition may be higher than IPCC pCO₂ predictions for open ocean conditions (Pachauri et al., 2014), lower and more variable pH is common for coastal and reef locations (Price, Martz, Brainard, & Smith, 2012; Rivest, Comeau, & Cornwall, 2017). For example, pH conditions on the fringing reefs adjacent to Coconut Island can range from ~7.6 - 8.1 (Guadayol et al., 2014; Silbiger et al., 2014). Further modeling of the pH change in reef locations under future scenarios results in a 2.5 fold increase in reef pH variation projected with an offshore increase to 900 μatm pCO₂ (Jury, Thomas, Atkinson, & Toonen, 2013). As such, our chosen pH conditions are ecologically relevant in terms of fluctuation and magnitude of potential pH change in the future in this dynamic reef location and do not represent extreme conditions. During the first month of larval

Positive carryover effects in corals

318 exposure, pH ranged from 7.79 - 8.04 in ambient conditions and 7.49 - 7.74 in the high
 319 pCO₂ treatment, and temperature, on average, ranged from 27.0 - 28.5°C and 27.0 -
 320 28.4 °C, respectively (Fig. S1). Lastly, across the six months of juvenile exposure, pH
 321 ranged from 7.77 - 8.04 in ambient conditions and 7.52 - 7.75 in high pCO₂ conditions
 322 and daily temperature, on average, ranged from 25.7 - 27.1°C and 25.8 - 27.0°C,
 323 respectively (Fig. S1). Discrete measurements of carbonate chemistry reveal stable TA
 324 across the 10 months and strong differences in pH and pCO₂ between treatments (Table
 325 1), but as these discrete measurements reflect daylight sampling only, they underestimate
 326 the treatment differences in pH and temperature clearly portrayed in continuous
 327 measurements described above (Fig. S1).

Table 1. Carbonate chemistry parameters for the different phases of the experiment (Figure 1). (a) Adult exposure (05 May 14 - 17 August 14), (b) 1 Month of offspring exposure (12 July 14 - 19 August 14), and (c) 6 Months of offspring exposure (12 July 14 - 12 January 2015). Temperature, salinity, total alkalinity, and pH were measured (N=sample size in each treatment and time point), while the remaining parameters of the carbonate system were calculated using seacarb as described in the methods.

(a) Adult Exposures

Treatment	N	Temp. °C	Salinity psu	pH Total	A _T μmol kg ⁻¹	pCO ₂ μmol kg ⁻¹	CO ₂ μmol kg ⁻¹	HCO ₃ ⁻ μmol kg ⁻¹	CO ₃ ²⁻ μmol kg ⁻¹	DIC μmol kg ⁻¹	Ω _A
Ambient	29	26.81 ± 0.12	33.9 ± 0.1	7.96 ± 0.02	2146 ± 5	482 ± 20	13 ± 1	1705 ± 14	178 ± 6	1896 ± 10	2.9 ± 0.1
High	28	26.78 ± 0.12	33.9 ± 0.1	7.71 ± 0.02	2151 ± 5	940 ± 38	26 ± 1	1878 ± 10	111 ± 4	2014 ± 7	1.8 ± 0.1

(b) Offspring 1 Month

Treatment	N	Temp. °C	Salinity psu	pH Total	A _T μmol kg ⁻¹	pCO ₂ μmol kg ⁻¹	CO ₂ μmol kg ⁻¹	HCO ₃ ⁻ μmol kg ⁻¹	CO ₃ ²⁻ μmol kg ⁻¹	DIC μmol kg ⁻¹	Ω _A
Ambient	8	27.16 ± 0.1	33.8 ± 0.2	7.9 ± 0.01	2138 ± 10	557 ± 21	15 ± 1	1747 ± 13	158 ± 4	1920 ± 12	2.6 ± 0.1
High	8	27.11 ± 0.11	33.8 ± 0.1	7.71 ± 0.03	2142 ± 12	945 ± 78	26 ± 2	1869 ± 23	111 ± 8	2005 ± 18	1.8 ± 0.1

(c) Offspring 6 Months

Treatment	N	Temp. °C	Salinity psu	pH Total	A _T μmol kg ⁻¹	pCO ₂ μmol kg ⁻¹	CO ₂ μmol kg ⁻¹	HCO ₃ ⁻ μmol kg ⁻¹	CO ₃ ²⁻ μmol kg ⁻¹	DIC μmol kg ⁻¹	Ω _A
Ambient	15	26.90 ± 0.35	33.8 ± 0.1	7.97 ± 0.04	2135 ± 7	493 ± 38	13 ± 1	1688 ± 35	180 ± 13	1882 ± 23	2.9 ± 0.2
High	15	26.92 ± 0.35	33.8 ± 0.1	7.73 ± 0.03	2135 ± 7	923 ± 70	25 ± 2	1853 ± 20	114 ± 8	1992 ± 15	1.8 ± 0.1

Positive carryover effects in corals

Planulation and Settlement

Planulae release was monitored on lunar days ~16-24 for the months of June, July, and August, as planulation has been reported to occur following the full moon (lunar day ~15) for *P. damicornis* in Hawai'i (Jokiel, 1985; Richmond & Jokiel, 1984). A clear peak in planulation was observed in July, with lowest release in August (Fig. 2). There was no significant difference in the timing of planulation between treatments in either June (Fig. 2a; $P > 0.05$) or August (Fig. 2c; $P > 0.05$). The general pattern suggested a shift in timing of planulation between treatments (Fig. 2), with a delay in release more prominent in the high $p\text{CO}_2$ condition in July (Fig. 2b; $D = 0.625$, $P = 0.087$).

Figure 2. Average larval release per coral colony. Planulae release as a function of lunar day in (a) June, (b) July, and (c) August in from adult colonies exposed to ambient $p\text{CO}_2$ ($n=8$) and adult colonies exposed to high $p\text{CO}_2$ ($n=6$).

Offspring from high CO_2 -exposed parents displayed significantly higher survivorship following 96h in the settlement chambers ($P < 0.016$ Table S1a), supporting positive

Positive carryover effects in corals

carryover effects. This increase in offspring survival from parents exposed to elevated pCO₂ was approximately equal in both offspring treatments at time 0, with 14.5% and 15.1% greater survivorship in the ambient and high offspring treatments, respectively (Fig. 3a). The settled spat from parents preconditioned to high pCO₂ also showed greater survivorship in both offspring treatments after one month in the reciprocal exposures (Table S1a), with preconditioned offspring having 27.1% greater survivorship at ambient offspring exposure conditions and 16.6% at high pCO₂ exposure (Fig. 3b). After 6 months of reciprocal exposure, parental exposure to high pCO₂ enhanced offspring survivorship by 13.6% in ambient offspring exposures, relative to spat from adults that underwent ambient preconditioning. Offspring from high pCO₂ parents displayed reduced survivorship (7.7%) in high offspring exposure relative to spat from adults preconditioned to ambient (Fig. 3c, Table S1a). Offspring exposure to high pCO₂ also led to lower survivorship regardless of parental preconditioning treatment (Table S1a). Averaged across parental exposures, survivorship decreased with offspring exposure to elevated pCO₂ by 26.1% at time 0, 24.5% at month 1 and 59.9% at month 6. Survivorship declined significantly over time ($P < 0.001$, Table S1a), with survivorship as low as 4.6% of the initial planulae by the final time point.

Planulae settlement was highest when parents were exposed to elevated pCO₂ ($P = 0.021$; Table S1b; Fig. 3d); settlement was enhanced by 17.6% and 11% for planulae settling in ambient and high pCO₂ offspring treatments, respectively. Despite this overall enhanced settlement of planulae from high pCO₂ parents, mean settlement was significantly lower overall (26.1%) in the high pCO₂ offspring treatments ($P < 0.001$; Table S1b), regardless of parental pCO₂ exposure (Fig. 3d).

Positive carryover effects in corals

Figure 3. Parental preconditioning to high pCO₂ enhances offspring performance. Reaction norm plots for (a) survivorship in the chambers, (b) survivorship at month 1, (c) survivorship at month 6, (d) settlement at time 0, (e) growth rate at month 1, and (f) growth rate at month 6. Points display data (mean±sem), while lines indicate only the direction of response. Tile sample sizes for survivorship and settlement time 0, month 1 and month 6 and growth at month 1 ranged from 7-36 (see Table S1). Growth rates were log transformed for analyses and figure displays back-transformed growth rate data.

Lastly, a trend for an Origin*Time interaction ($P=0.056$) was observed in spat growth, where differences in growth with parental origin were apparent at month 1 (growth was on average 1.3-fold higher in offspring from parents preconditioned to high pCO₂; Fig. 3e), but not at month 6 (Fig. 3f). Time also had a significant effect on growth, where by the end of the experiment, at 6 months post release, growth rates were significantly lower than at month 1 ($P<0.0001$, Fig. 3f).

Discussion

Here we present the first ecological assessment of trans-generational or carryover effects in reef-building corals. In our study, the ecological and fitness-related response of coral offspring was enhanced dependent on response and life-stage when their parents were

Positive carryover effects in corals

exposed to a high pCO₂ environment, with carryover effects lasting into the juvenile stage. Projections for future reef persistence are dire (Veron et al., 2009), but many do not incorporate adaptation or acclimatization (Van Hooideonk, Maynard, & Planes, 2013), likely over or under-estimating climate change effects for some stressors. When adaptation and acclimatization are considered, it is clear the trajectories differ from the worst-case scenarios (Ainsworth et al., 2016; Donner, Skirving, Little, Oppenheimer, & Hoegh-Guldberg, 2005; Logan, Dunne, Eakin, & Donner, 2014). Our work provides evidence to support the importance of the role of acclimatory processes in evolutionary thinking in an era of climate change and encourage the examination of mechanisms such as hormesis and epigenetics (e.g., Costantini, 2014; Torda et al., 2017).

Potential for tuning of reproductive timing

Our multi-life stage perspective identifies carryover effects of adult stressor exposure, with reproductive and offspring consequences. Shifts in reproductive timing can have significant consequences for offspring settlement conditions. It is possible that “bet-hedging” or environmental tuning by the parents may result in release of larvae timed to favorable conditions. For example, a shift in reproductive timing of coral planulation with change in temperature across season was recently documented (Fan et al., 2017). In this case, seasonal acclimatization in the parents resulted in change in summer planulae release timing, which minimizes planulae release during strong, likely stressful, upwelling-induced temperature fluctuations of ~10°C (Lee, Chao, Fan, Wang, & Liang, 1997). Adult tuning of larval release of offspring is also possible in our experiment. Kāneohe Bay is a semi-enclosed embayment that has fluctuations in physical

Positive carryover effects in corals

conditions as a function of tidal cycle. Specifically, diurnal pCO₂ fluctuations are greatest when tidal fluctuation is the lowest (Guadayol et al., 2014; Putnam, 2012; Silbiger et al., 2014) in the shallow fringing reefs of Kāneohe Bay. A delay in the peak of planulation from the adults preconditioned to high pCO₂, would then correspond to the timing of lower daily tidal ranges (<https://tidesandcurrents.noaa.gov/stationhome.html?id=1612480>) and thus higher pCO₂ fluctuations that are more similar to the high pCO₂ adult conditions. Another hypothesis that could contribute to a shift in larval release under high pCO₂ condition is adult or offspring energetic constraints. Delay in release could indicate energetic costs to maintaining adult calcification and homeostasis (Stumpp et al., 2012), potentially resulting in decreased parental investment, or increased development time necessary in offspring (Stumpp, Wren, Melzner, Thorndyke, & Dupont, 2011). Further, low pH can influence development processes such as sperm performance, fertilization success, and developmental normalcy and timing (Albright, 2011; Byrne & Przeslawski, 2013).

Given the trend for a shift in the timing of planulation during July when our offspring experiment was completed, it could be hypothesized that parental effects in the reciprocal exposure are due to slight differences in the larval cohorts by day of release (Cumbo et al., 2013; Cumbo, Fan, & Edmunds, 2012; Putnam et al., 2010; Rivest & Hofmann, 2015). For example, peaks in *Symbiodinium* density and photophysiology, and larval size, are positively correlated to peak larval release in *Pocillopora damicornis* in Taiwan (Putnam et al., 2010). These differences in physiology by day of release also translate to variation in susceptibility to changing temperature and pH in *P. damicornis* (Cumbo et al., 2013; Rivest & Hofmann, 2015). The impact of day of release in our case

Positive carryover effects in corals

is likely to be minimal, given the experiment was not conducted on a single day's larval pool, but over 7 consecutive days (Fig. 2), better representing the full range of larval phenotypes from *P. damicornis/acuta*.

Importance of carryover effects for corals

Our work provides further evidence that parental environment matters to offspring performance in this brooding coral species, *Pocillopora acuta*. While ~1 month of exposure to increased temperature and low pH resulted in TGP in this same species (Putnam & Gates, 2015), in our current study with exposure to only OA, reaction norms were primarily parallel, with enhanced performance in those preconditioned. This may suggest that exposure to increased temperature (or the combination of temperature and OA) has more profound, or mechanistically different impacts than OA on processes involved in parental contributions or developmental acclimation (e.g., Byrne, 2012). The enhanced growth of *P. acuta* juveniles under low pH is unexpected given the commonly detrimental effect of ocean acidification on coral calcification (Kroeker, Kordas, Crim, & Singh, 2010). *Pocillopora damicornis* has, however displayed variability in sensitivity of calcification in response to OA, from negative (Putnam et al., 2016) to no effects on growth (Comeau et al., 2014; Comeau, Edmunds, Spindel, & Carpenter, 2013). For instance, ocean acidification did not impact recruitment of *P. damicornis* larvae to the sides of treatment tanks in a mesocosm study also in Kāneohe Bay (Jokiel et al., 2008). Furthermore, physiology at the larvae stage was not strongly affected by OA (Putnam et al 2013), but was affected more so by increased temperature. Enhanced calcification in a trans-generational context has been measured in another marine invertebrate, where

Positive carryover effects in corals

higher shell growth rates in offspring of the Manila clam following exposure of the parents to low pH have been observed (Zhao et al., 2018; Zhao, Schöne, Mertz-Kraus, & Yang, 2017). This positive carryover effect for growth in another marine calcifier supports our findings here with *Pocillopora acuta*, with implications for the presence and importance of TGP beyond a single coral species.

Potential mechanisms underlying parental effects

Several hypotheses may account for the parental contribution to enhanced settlement, survivorship, and growth we documented. It is possible that adults manipulate the investment in their offspring in the form of *Symbiodinium* communities (Padilla-Gamiño, Pochon, Bird, Concepcion, & Gates, 2012), microbiome (Sharp, Distel, & Paul, 2012), size, protein, lipids, or carbohydrates (Hartmann, Marhaver, Chamberland, Sandin, & Vermeij, 2013). Further the role of mitochondrial performance has been posited as a mechanism of rapid adaptation in coral larvae (Dixon et al., 2015) and mitochondrial performance has been linked to parental environment in the trans-generational acclimatization of marine worms (Gibbin et al., 2017). These mechanisms could provide metabolic boosts or conversely detriments during this energetically demanding life stage (Edmunds, Cumbo, & Fan, 2013); e.g., the presence of clade D symbionts reduces growth in coral juveniles (Little, Van Oppen, & Willis, 2004). DNA methylation and other epigenetic mechanisms linked to gene expression regulation (Putnam et al., 2016; Putnam & Gates, 2015; Torda et al., 2017) could also provide a mechanism of heritable cross-generational priming. Identification of differential DNA methylation patterns between the offspring of exposed and unexposed oysters is an initial line of evidence for an epigenetic role in TGP for marine calcifiers (Rondon et al., 2017). The minute

Positive carryover effects in corals

number of methylation changes and the lack of direct impact on RNA transcript abundance or differential splicing (Rondon et al., 2017), however, illustrates how complex these mechanisms can be. While differential provisioning of planulae symbiotically, energetically, and epigenetically is possible, the goal of examining ecological effects on the process of recruitment precluded destructive sampling for such hypotheses in our study, but they remain important considerations in future work.

Brooded embryos may experience developmental acclimation while growing within the parental polyp (Putnam & Gates, 2015). It is possible that low pH in the gastrovascular cavity (GVC) at the site of development could condition planulae for low pH when released. This hypothesis would account for the differences seen in our study between ambient and low pH preconditioned parents if the GVCs were modified differently between treatments to retain a treatment offset. To date, limited data are available for GVC pH under a variety of conditions (Agostini et al., 2011; Cai et al., 2016). The data available suggest both a high pH and low pH within the GVC and associated mesenteries (Cai et al., 2016), with no uniform picture of what planulae within the polyp are exposed to relative to external conditions. Further work with multiple coral species is necessary to disentangle the role of developmental acclimation from TGP, including experiments focused on the magnitude and timing of signals that will induce carryover effects and exposures through the F1 and F2 generation (Torda et al., 2017).

Environmental hardening through hormetic priming

Acclimatization occurs through short-term compensatory processes including modulation of biochemical activity and gene expression in response to external stimuli, occurring on

Positive carryover effects in corals

daily, seasonal, and annual scales, within a generation, and across a generation. Often implicit in the usage of the term acclimatization is the inference that acclimatory processes are beneficial and fully compensatory, *sensu* the Beneficial Acclimation Hypothesis (BAH; Huey, Berrigan, Gilchrist, & Herron, 1999), but acclimatization is not always adaptive (Edmunds, 2014; Edmunds & Gates, 2008; Huey et al., 1999). One conceptual explanation for inconsistency of BAH is the framework of hormetic priming (Costantini, 2014; Costantini, Metcalfe, & Monaghan, 2010), which deserves consideration with regards to explaining patterns of TGP. Hormesis is defined as the stimulation of function through mild exposure(s) (Southam & Ehrlich, 1943). Hormetic priming thus occurs when exposure to a sub-lethal stressor results in stimulatory response to increasing levels of a condition (e.g., Fig. 4), for example, as we have documented here in the enhanced performance of larvae of *Pocillopora acuta* following parental preconditioning. Hormesis is a biphasic response and therefore does not necessitate positive acclimation to all future exposures (Fig. 4).

Hormetic priming as a mechanism of acclimatization could explain both the beneficial and detrimental effects of increased temperature that have been documented in certain types of repeated exposures (Ainsworth et al., 2016; Brown et al., 2000; Grottoli et al., 2014), as well as variability in performance. For example, Ainsworth et al., (2016) identified a “protective trajectory” of sea surface temperatures (SST) that corals experienced, which lead to acquired thermal tolerance in *Acropora aspera*. Conversely a “repetitive bleaching trajectory”, with high frequency bleaching events that lack recovery time resulted in substantial symbiotic cell loss, bleaching, and mortality (Ainsworth et al., 2016). Here it is likely these responses are on differing sides of the zero equivalence

Positive carryover effects in corals

527 point in terms of the biphasic nature of hormetic processes (Y axis in Fig. 4A; (Costantini
528 et al., 2010; Lushchak, 2014). An example of hormetic priming in corals would be that of
529 mild ROS exposure resulting in protein expression priming, thereby reducing the ROS
530 damage in subsequent events (Brown, Downs, Dunne, & Gibb, 2002). Beyond
531 antioxidant genes, frontloading, or the constitutive up-regulation of expression of
532 canonical heat stress genes in *Acropora hyacinthus* samples from the high thermal
533 variability pool in American Samoa, could be another outcome of hormetic priming, with
534 implications for the thermal tolerance of those primed individuals (Barshis et al., 2013).

Positive carryover effects in corals

535

Figure 4. Hormetic priming may contribute to magnitude, pattern, and temporal variation in the benefits of TGP. The dashed line indicates the result of no hormetic priming, the gray dashed line is no change, the dotted line (black area) indicates a hormetic response with no conditioning, while the solid line (hashed area) indicates the hormetic response with preconditioning (Constantini et al., 2010; Lushchak 2014). Shaded areas indicate the hormetic zones, where the stimulation of performance from the increasing stressor is above the line of no change. Variation in the intensity, duration, and life stage of prior exposure may shift the A) magnitude, or B) shape of the hormetic zone. For example, in sessile benthic organisms, carryover effects from parental exposure may result in an enhanced hormetic zone relative to no conditioning. Further there may be C) temporal constraints on hormetic conditioning. For example, the hormetic zone may differ from parental exposure versus developmental exposure, or as in the case of our study, display a decline in positive carryover effects with increasing time post conditioning (as visualized by the temporal shift in hormetic zone).

Positive carryover effects in corals

Corals in our study displayed life stage-dependent TGP. Despite the enhancement in performance seen in early stages, the parental effects of preconditioning to high pCO₂ are absent in the offspring by 6 months post release. While maintenance of maternal effects may be expected in short lived organisms, as generation time increases there is a greater potential mismatch between parental and offspring conditions (e.g., Fig. S1). This change in parental effects over time is not unexpected due to potential maladaptive tradeoffs (Burgess & Marshall, 2014), especially the case in long-lived corals (Torda et al., 2017). Seasonal, annual, and decadal environmental changes likely elicit tradeoffs between the early life stage performance benefits of TGP and the costs of TGP the organism may incur later in life. For example, maternal exposure of soil arthropods to high temperatures resulted in increased thermal resistance at the juvenile stage, but later drove reduced F1 fecundity (Zizzari & Ellers, 2014). Further the expectation of positive TGP is linked to the predictability of the stressor (e.g., anticipatory parental effects; APE's; Burgess & Marshall, 2014; Donelson et al., 2017). Specifically, TGP through hormetic priming would likely be optimized by high environmental autocorrelation (i.e., a strong match between parent and offspring conditions would result in extended enhancement). The greater the temporal shift from the parental environment (e.g., 6 months here Fig. S1) the less likely there is to be a benefit from parental enhancement and the more likely within generation priming would become important (e.g., Fig. 4C). Seasonal changes in physical parameters would then be expected to result in a loss of benefit from TGP as pH continues to change, in a biphasic fashion (Fig. 4C). Additionally, with respect to overall growth rates, the growth from months 2-6 would be expected to decline due to seasonal effects associated with decreases in temperature and

Positive carryover effects in corals

light in the winter months (Fig. S1; Fitt, McFarland, Warner, & Chilcoat, 2000; Thornhill et al., 2011). The temporal transience of parental effects documented in our study argues that for parental effects to translate into long term “memory” more constant environmental and biological feedbacks are necessary (Ptashne, 2013).

Implications for reef-building corals

Experiments designed to examine TGP and carryover effects provide a glimmer of hope for coral reef organisms that acclimatization may act as a buffer against a rapidly changing climate (Putnam & Gates, 2015; Torda et al., 2017). Further experiments are necessary to distinguish between true TGP, parental effects, and developmental acclimation, and their underlying mechanisms. This could be achieved in spawning coral systems, or by manipulating the timing of exposure in the brooding coral system to target isolated stages (i.e., adult, gametogenesis, brooding, and larval). Additionally, tests of the stability and later-life tradeoffs of TGP, as well as mechanisms of environmental “memory” through aspects such as mitochondrial function (Gibbin et al., 2017), DNA methylation (Putnam et al., 2016), or microbiome inheritance (Putnam, Barott, Ainsworth, & Gates, 2017; Webster & Reusch, 2017) will unveil the complex contributions of the holobiont partners to meta-organism function and acclimatization potential.

Our work challenges the paradigm of inevitable coral decline due to rapid climate change by identifying ecologically-relevant beneficial parental effects in offspring, in response to adult preconditioning to ocean acidification. We suggest hormesis, or environmental priming may conceptually explain enhanced tolerance and performance seen in acclimatization studies, as well as explaining the lack of a ubiquitous beneficial

Positive carryover effects in corals

acclimation due to the biphasic nature of hormesis. With regards to a conservation and management actions (i.e. assisted evolution; van Oppen et al., 2017; van Oppen, Oliver, Putnam, & Gates, 2015), environmental priming is not a one-size-fits all phenomenon (i.e., does not always confer beneficial acclimation). Instead conditioning methods would necessitate a Goldilocks approach, as variation in cellular conditions and physiology between species requires a variety of exposures for optimal performance outcomes (Putnam et al., 2017). It is not clear, however, if the duration of the benefits would extend far beyond the exposures, as phenotype-environment mismatches increase with season and anthropogenic effects. The performance and fitness tradeoffs of acclimatization, hormetic triggers, and heritability of potential epigenetic feedbacks present promising areas of further study with respect to carryover effects and the ecological and evolutionary trajectories of reef-building corals.

Acknowledgements

This work was supported by funding from NSF OCE-PRF 1323822 to HMP, and NSF EPS-0903833, and NSF URM 0829272. We would like to thank Rashim Khadka and the HIMB facilities staff for their assistance in this work and N. Silbiger, M. Iacchei for constructive comments on the manuscript. This is HIMB contribution number xxx and SOEST contribution number XXX.

References

- Agostini, S., Suzuki, Y., Higuchi, T., Casareto, B. E., Yoshinaga, K., Nakano, Y., & Fujimura, H. (2011). Biological and chemical characteristics of the coral gastric cavity. *Coral Reefs*, 31(1), 147–156. <https://doi.org/10.1007/s00338-011-0831-6>
- Ainsworth, T. D., Heron, S. F., Ortiz, J. C., Mumby, P. J., Grech, A., Ogawa, D., Eakin, C.M., Leggat, W. (2016). Climate change disables coral bleaching protection on the Great Barrier Reef. *Science*, 352(6283), 338–342.
- Albright, R. (2011). Reviewing the effects of ocean acidification on sexual reproduction and early life history stages of reef-building corals. *Journal of Marine Biology*, 2011.
- Baird, A. H., Guest, J. R., & Willis, B. L. (2009). Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. *Annual Review of Ecology, Evolution, and Systematics*, 40, 551–571.
- Barshis, D. J., Ladner, J. T., Oliver, T. A., Seneca, F. O., Traylor-Knowles, N., & Palumbi, S. R. (2013). Genomic basis for coral resilience to climate change. *Proceedings of the National Academy of Sciences*, 110(4), 1387–1392.
- Bishop, R. C., Chapman, D. J., Kanninen, B. J., Krosnick, J. A., Leeworthy, B., & Meade, N. F. (2011). *Total Economic Value for Protecting and Restoring Hawaiian Coral Reef Ecosystems: Final Report*. Silver Spring, MD: NOAA Office of National Marine Sanctuaries, Office of Response and Restoration, and Coral Reef Conservation Program. NOAA Technical Memorandum CRCP 16.
- Brown, B. E., & Cossins, A. R. (2011). The potential for temperature acclimatisation of reef corals in the face of climate change. In *Coral Reefs: An Ecosystem in Transition* (pp. 421–433). Springer.

Positive carryover effects in corals

- 629 Brown, B. E., Downs, C. A., Dunne, R. P., & Gibb, S. W. (2002). Exploring the basis of
630 thermotolerance in the reef coral *Goniastrea aspera*. *Marine Ecology Progress*
631 *Series*, 242, 119–129.
- 632 Brown, B. E., Dunne, R. P., Goodson, M. S., & Douglas, A. E. (2000). Bleaching
633 patterns in reef corals. *Nature*, 404(6774), 142.
- 634 Burgess, S. C., & Marshall, D. J. (2014). Adaptive parental effects: the importance of
635 estimating environmental predictability and offspring fitness appropriately. *Oikos*,
636 123(7), 769–776.
- 637 Byrne, M. (2012). Global change ecotoxicology: identification of early life history
638 bottlenecks in marine invertebrates, variable species responses and variable
639 experimental approaches. *Marine Environmental Research*, 76, 3–15.
- 640 Byrne, M., & Przeslawski, R. (2013). Multistressor impacts of warming and acidification
641 of the ocean on marine invertebrates' life histories. *Integrative and Comparative*
642 *Biology*, 53(4), 582–596.
- 643 Cai, W.-J., Ma, Y., Hopkinson, B. M., Grottoli, A. G., Warner, M. E., Ding, Q., Hu, X.,
644 Yuan, X., Schoepf, V., Xu, H., Han, C., Melman, T.F., Hoadley, K.D., Pettay, D.
645 T., Matsui Y., Baumann, J.H., Levas, J.H., Ying Y., Wang, Y. (2016).
646 Microelectrode characterization of coral daytime interior pH and carbonate
647 chemistry. *Nature Communications*, 7, 11144.
- 648 Chakravarti, L. J., Jarrold, M. D., Gibbin, E. M., Christen, F., Massamba-N'Siala, G.,
649 Blier, P. U., & Calosi, P. (2016). Can trans-generational experiments be used to
650 enhance species resilience to ocean warming and acidification? *Evolutionary*
651 *Applications*, 9(9), 1133–1146. <https://doi.org/10.1111/eva.12391>

Positive carryover effects in corals

- 652 Chevin, L.-M., Collins, S., & Lefèvre, F. (2013). Phenotypic plasticity and evolutionary
653 demographic responses to climate change: taking theory out to the field.
654 *Functional Ecology*, 27(4), 967–979.
- 655 Chua, C. M., Leggat, W., Moya, A., & Baird, A. H. (2013). Temperature affects the early
656 life history stages of corals more than near future ocean acidification. *Marine*
657 *Ecology Progress Series*, 475, 85–92.
- 658 Chua, C.-M., Leggat, W., Moya, A., & Baird, A. H. (2013). Near-future reductions in pH
659 will have no consistent ecological effects on the early life-history stages of reef
660 corals. *Marine Ecology Progress Series*, 486, 143–151.
- 661 Coles, S. L., & Brown, B. E. (2003). Coral bleaching—capacity for acclimatization and
662 adaptation. *Advances in Marine Biology*, 46, 183–223.
- 663 Comeau, S., Carpenter, R. C., Nojiri, Y., Putnam, H. M., Sakai, K., & Edmunds, P. J.
664 (2014). Pacific-wide contrast highlights resistance of reef calcifiers to ocean
665 acidification. *Proceedings of the Royal Society of London B: Biological Sciences*,
666 281(1790), 20141339.
- 667 Comeau, S., Edmunds, P. J., Spindel, N. B., & Carpenter, R. C. (2013). The responses of
668 eight coral reef calcifiers to increasing partial pressure of CO₂ do not exhibit a
669 tipping point. *Limnology and Oceanography*, 58(1), 388–398.
- 670 Costantini, D. (2014). Does hormesis foster organism resistance to extreme events. *Front*
671 *Ecol Environ*, 12, 209–210.
- 672 Costantini, D., Metcalfe, N. B., & Monaghan, P. (2010). Ecological processes in a
673 hormetic framework. *Ecology Letters*, 13(11), 1435–1447.
674 <https://doi.org/10.1111/j.1461-0248.2010.01531.x>

Positive carryover effects in corals

- 675 Costanza, R., de Groot, R., Sutton, P., van der Ploeg, S., Anderson, S. J., Kubiszewski, I.,
676 ... Turner, R. K. (2014). Changes in the global value of ecosystem services.
677 *Global Environmental Change*, 26, 152–158.
- 678 Cumbo, V. R., Edmunds, P. J., Wall, C. B., & Fan, T.-Y. (2013). Brooded coral larvae
679 differ in their response to high temperature and elevated pCO₂ depending on the
680 day of release. *Marine Biology*, 160(11), 2903–2917.
- 681 Cumbo, V. R., Fan, T.-Y., & Edmunds, P. J. (2012). Physiological development of
682 brooded larvae from two pocilloporid corals in Taiwan. *Marine Biology*, 159(12),
683 2853–2866.
- 684 Dickson, A. G., Sabine, C. L., & Christian, J. R. (2007). Guide to best practices for ocean
685 CO₂ measurements. *PICES Special Publication 3*.
- 686 Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V.
687 (2015). Genomic determinants of coral heat tolerance across latitudes. *Science*,
688 348(6242), 1460–1462.
- 689 Donelson, J. M., Salinas, S., Munday, P. L., & Shama, L. N. S. (2017). Transgenerational
690 plasticity and climate change experiments: Where do we go from here? *Global*
691 *Change Biology*. <https://doi.org/DOI: 10.1111/gcb.13903>
- 692 Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean Acidification:
693 The Other CO₂ Problem. *Annual Review of Marine Science*, 1(1), 169–192.
694 <https://doi.org/10.1146/annurev.marine.010908.163834>
- 695 Donner, S. D., Skirving, W. J., Little, C. M., Oppenheimer, M., & Hoegh-Guldberg, O. V.
696 E. (2005). Global assessment of coral bleaching and required rates of adaptation
697 under climate change. *Global Change Biology*, 11(12), 2251–2265.

Positive carryover effects in corals

- 698 Drupp, P., De Carlo, E. H., Mackenzie, F. T., Bienfang, P., & Sabine, C. L. (2011).
699 Nutrient inputs, phytoplankton response, and CO₂ variations in a semi-enclosed
700 subtropical embayment, Kaneohe Bay, Hawaii. *Aquatic Geochemistry*, 17(4–5),
701 473–498.
- 702 Dufault, A. M., Cumbo, V. R., Fan, T.-Y., & Edmunds, P. J. (2012). Effects of diurnally
703 oscillating pCO₂ on the calcification and survival of coral recruits. *Proceedings of*
704 *the Royal Society of London B: Biological Sciences*, rspb20112545.
- 705 Edmunds, P. J. (2014). Is acclimation beneficial to scleractinian corals, *Porites spp.*?
706 *Marine Biology*, 161(7), 1531–1542. <https://doi.org/10.1007/s00227-014-2438-2>
- 707 Edmunds, P. J., Cumbo, V. R., & Fan, T.-Y. (2013). Metabolic costs of larval settlement
708 and metamorphosis in the coral *Seriatopora caliendrum* under ambient and
709 elevated pCO₂. *Journal of Experimental Marine Biology and Ecology*, 443, 33–38.
- 710 Edmunds, P. J., & Gates, R. D. (2008). Acclimatization in tropical reef corals. *Marine*
711 *Ecology Progress Series*, 361, 307–310.
- 712 Erez, J., Reynaud, S., Silverman, J., Schneider, K., & Allemand, D. (2011). Coral
713 calcification under ocean acidification and global change. In *Coral reefs: an*
714 *ecosystem in transition* (pp. 151–176). Springer.
- 715 Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., ...
716 Lough, J. M. (2011). Losers and winners in coral reefs acclimatized to elevated
717 carbon dioxide concentrations. *Nature Climate Change*, 1(3), 165–169.
- 718 Fan, T.-Y., Hsieh, Y.-C., Lin, K.-H., Kuo, F.-W., Soong, K., McRae, C. J., Edmunds, P.J.,
719 Fang, L.-S. (2017). Plasticity in lunar timing of larval release of two brooding

Positive carryover effects in corals

- 720 pocilloporid corals in an internal tide-induced upwelling reef. *Marine Ecology*
721 *Progress Series*, 569, 117–127.
- 722 Fitt, W. K., McFarland, F. K., Warner, M. E., & Chilcoat, G. C. (2000). Seasonal patterns
723 of tissue biomass and densities of symbiotic dinoflagellates in reef corals and
724 relation to coral bleaching. *Limnology and Oceanography*, 45(3), 677–685.
- 725 Foo, S. A., & Byrne, M. (2016). Chapter Two - Acclimatization and Adaptive Capacity
726 of Marine Species in a Changing Ocean. In B. E. Curry (Ed.), *Advances in Marine*
727 *Biology* (Vol. 74, pp. 69–116). Academic Press.
- 728 Foster, T., Gilmour, J. P., Chua, C. M., Falter, J. L., & McCulloch, M. T. (2015). Effect
729 of ocean warming and acidification on the early life stages of subtropical
730 *Acropora spicifera*. *Coral Reefs*, 34(4), 1217–1226.
- 731 Gattuso, J. P., Epitalon, J. M., Lavigne, H., Orr, J., Gentili, B., Hagens, M., Hofmann, A.,
732 Proye, A., Soetaert, K., Rae, J. (2016). Seacarb: seawater carbonate chemistry
733 with R. *R Package Version*, 3.1.1(8).
- 734 Gaylord, B., Kroeker, K. J., Sunday, J. M., Anderson, K. M., Barry, J. P., Brown, N. E.,
735 Connell, S.D., Dupont, S., Fabricius, K.E., Hall-Spencer, J.M., Klinger, T.,
736 Milazzo, M., Munday, P.L., Russell, B.D., Sanford, E., Schreiber, S.J.,
737 Thiyagarajan, V., Vaughan M.L.H., Widdecombe, S., Harley, C. D. G. (2015).
738 Ocean acidification through the lens of ecological theory. *Ecology*, 96(1), 3–15.
739 <https://doi.org/10.1890/14-0802.1>
- 740 Gibbin, E. M., Chakravarti, L. J., Jarrold, M. D., Christen, F., Turpin, V., N’Siala, G. M.,
741 Blier, P.U., Calosi, P. (2017). Can multi-generational exposure to ocean warming

Positive carryover effects in corals

742 and acidification lead to the adaptation of life history and physiology in a marine
 743 metazoan? *Journal of Experimental Biology*, 220(4), 551–563.

744 Gibbin, E. M., Putnam, H. M., Davy, S. K., & Gates, R. D. (2014). Intracellular pH and
 745 its response to CO₂-driven seawater acidification in symbiotic versus non-
 746 symbiotic coral cells. *Journal of Experimental Biology*, 217(11), 1963–1969.

747 Gibbin, E. M., Putnam, H. M., Gates, R. D., Nitschke, M. R., & Davy, S. K. (2015).
 748 Species-specific differences in thermal tolerance may define susceptibility to
 749 intracellular acidosis in reef corals. *Marine Biology*, 162(3), 717–723.

750 Grottoli, A. G., Warner, M. E., Levas, S. J., Aschaffenburg, M. D., Schoepf, V.,
 751 McGinley, M., Baumann, J., Matsui, Y. (2014). The cumulative impact of annual
 752 coral bleaching can turn some coral species winners into losers. *Global Change*
 753 *Biology*, 20(12), 3823–3833.

754 Guadayol, Ò., Silbiger, N. J., Donahue, M. J., & Thomas, F. I. (2014). Patterns in
 755 temporal variability of temperature, oxygen and pH along an environmental
 756 gradient in a coral reef. *PloS One*, 9(1), e85213.

757 Hartmann, A. C., Marhaver, K. L., Chamberland, V. F., Sandin, S. A., & Vermeij, M. J.
 758 (2013). Large birth size does not reduce negative latent effects of harsh
 759 environments across life stages in two coral species. *Ecology*, 94(9), 1966–1976.

760 Harvey, B. P., Gwynn-Jones, D., & Moore, P. J. (2013). Meta-analysis reveals complex
 761 marine biological responses to the interactive effects of ocean acidification and
 762 warming. *Ecology and Evolution*, 3(4), 1016–1030.

763 <https://doi.org/10.1002/ece3.516>

Positive carryover effects in corals

- 764 Hettinger, A., Sanford, E., Hill, T. M., Lenz, E. A., Russell, A. D., & Gaylord, B. (2013).
765 Larval carry-over effects from ocean acidification persist in the natural
766 environment. *Global Change Biology*, 19(11), 3317–3326.
- 767 Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N., Hoey, J., Forsch, M.,
768 Page, H.N., Gaylord, B. (2012). Persistent carry-over effects of planktonic
769 exposure to ocean acidification in the Olympia oyster. *Ecology*, 93(12), 2758–
770 2768.
- 771 Huey, R. B., Berrigan, D., Gilchrist, G. W., & Herron, J. C. (1999). Testing the Adaptive
772 Significance of Acclimation: A Strong Inference Approach. *Integrative and*
773 *Comparative Biology*, 39(2), 323–336. <https://doi.org/10.1093/icb/39.2.323>
- 774 Johnston, E. C., Forsman, Z. H., & Toonen, R. J. (2018). A simple molecular technique
775 for distinguishing species reveals frequent misidentification of Hawaiian corals in
776 the genus *Pocillopora*. *PeerJ*, 6, e4355. <https://doi.org/10.7717/peerj.4355>
- 777 Jokiel, P. L. (1985). Lunar periodicity of planula release in the reef coral *Pocillopora*
778 *damicornis* in relation to various environmental factors. In *Proc 5th Int Coral*
779 *Reef Symp* (Vol. 4, pp. 307–312).
- 780 Jokiel, P. L., Rodgers, K. S., Kuffner, I. B., Andersson, A. J., Cox, E. F., & Mackenzie, F.
781 T. (2008). Ocean acidification and calcifying reef organisms: a mesocosm
782 investigation. *Coral Reefs*, 27(3), 473–483. [https://doi.org/10.1007/s00338-008-](https://doi.org/10.1007/s00338-008-0380-9)
783 0380-9
- 784 Jury, C. P., Thomas, F. I., Atkinson, M. J., & Toonen, R. J. (2013). Buffer capacity,
785 ecosystem feedbacks, and seawater chemistry under global change. *Water*, 5(3),
786 1303–1325.

Positive carryover effects in corals

- 787 Kaniewska, P., Campbell, P. R., Kline, D. I., Rodriguez-Lanetty, M., Miller, D. J., Dove,
788 S., & Hoegh-Guldberg, O. (2012). Major Cellular and Physiological Impacts of
789 Ocean Acidification on a Reef Building Coral. *PLOS ONE*, 7(4), e34659.
790 <https://doi.org/10.1371/journal.pone.0034659>
- 791 Kroeker, K. J., Kordas, R. L., Crim, R. N., & Singh, G. G. (2010). Meta-analysis reveals
792 negative yet variable effects of ocean acidification on marine organisms. *Ecology*
793 *Letters*, 13(11), 1419–1434.
- 794 Lane, A., Campanati, C., Dupont, S., & Thiagarajan, V. (2015). Trans-generational
795 responses to low pH depend on parental gender in a calcifying tubeworm.
796 *Scientific Reports*, 5.
- 797 Lee, H.-J., Chao, S.-Y., Fan, K.-L., Wang, Y.-H., & Liang, N.-K. (1997). Tidally induced
798 upwelling in a semi-enclosed basin: Nan Wan Bay. *Journal of Oceanography*, 53,
799 467–480.
- 800 Lesser, M. P. (2011). Coral Bleaching: Causes and Mechanisms. In Z. Dubinsky & N.
801 Stambler (Eds.), *Coral Reefs: An Ecosystem in Transition* (pp. 405–419). Springer
802 Netherlands. https://doi.org/10.1007/978-94-007-0114-4_23
- 803 Little, A. F., Van Oppen, M. J., & Willis, B. L. (2004). Flexibility in algal endosymbioses
804 shapes growth in reef corals. *Science*, 304(5676), 1492–1494.
- 805 Logan, C. A., Dunne, J. P., Eakin, C. M., & Donner, S. D. (2014). Incorporating adaptive
806 responses into future projections of coral bleaching. *Global Change Biology*,
807 20(1), 125–139.
- 808 Lushchak, V. I. (2014). Dissection of the hormetic curve: analysis of components and
809 mechanisms. *Dose-Response*, 12(3), dose–response.

Positive carryover effects in corals

- 810 Miller, G. M., Watson, S.-A., Donelson, J. M., McCormick, M. I., & Munday, P. L.
811 (2012). Parental environment mediates impacts of increased carbon dioxide on a
812 coral reef fish. *Nature Climate Change*, 2(12), 858–861.
- 813 Munday, P. L., Warner, R. R., Monro, K., Pandolfi, J. M., & Marshall, D. J. (2013).
814 Predicting evolutionary responses to climate change in the sea. *Ecology Letters*,
815 16(12), 1488–1500.
- 816 Negri, A. P., Marshall, P. A., & Heyward, A. J. (2007). Differing effects of thermal stress
817 on coral fertilization and early embryogenesis in four Indo Pacific species. *Coral*
818 *Reefs*, 26(4), 759–763.
- 819 Pachauri, R. K., Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., ...
820 others. (2014). *Climate change 2014: synthesis Report. Contribution of working*
821 *groups I, II and III to the fifth assessment report of the intergovernmental panel*
822 *on climate change*. IPCC.
- 823 Padilla-Gamiño, J. L., Pochon, X., Bird, C., Concepcion, G. T., & Gates, R. D. (2012).
824 From parent to gamete: vertical transmission of *Symbiodinium* (Dinophyceae)
825 ITS2 sequence assemblages in the reef building coral *Montipora capitata*. *PLoS*
826 *One*, 7(6), e38440.
- 827 Pandolfi, J. M., Connolly, S. R., Marshall, D. J., & Cohen, A. L. (2011). Projecting coral
828 reef futures under global warming and ocean acidification. *Science*, 333(6041),
829 418–422.
- 830 Parker, L. M., O'Connor, W. A., Byrne, M., Coleman, R. A., Virtue, P., Dove, M., Gibbs,
831 M., Spohr, L., Scanes, E., Ross, P. M. (2017). Adult exposure to ocean
832 acidification is maladaptive for larvae of the Sydney rock oyster *Saccostrea*

Positive carryover effects in corals

- 833 *glomerata* in the presence of multiple stressors. *Biology Letters*, 13(2), 20160798.
- 834 <https://doi.org/10.1098/rsbl.2016.0798>
- 835 Parker, L. M., O'Connor, W. A., Raftos, D. A., Pörtner, H.-O., & Ross, P. M. (2015).
- 836 Persistence of Positive Carryover Effects in the Oyster, *Saccostrea glomerata*,
- 837 following Transgenerational Exposure to Ocean Acidification. *PLOS ONE*, 10(7),
- 838 e0132276. <https://doi.org/10.1371/journal.pone.0132276>
- 839 Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A., & Pörtner, H.-
- 840 O. (2012). Adult exposure influences offspring response to ocean acidification in
- 841 oysters. *Global Change Biology*, 18(1), 82–92.
- 842 Permata, W. D., Kinzie, R. A., & Hidaka, M. (2000). Histological studies on the origin of
- 843 planulae of the coral *Pocillopora damicornis*. *Marine Ecology Progress Series*,
- 844 191–200.
- 845 Price, N. N., Martz, T. R., Brainard, R. E., & Smith, J. E. (2012). Diel variability in
- 846 seawater pH relates to calcification and benthic community structure on coral
- 847 reefs. *PloS One*, 7(8), e43843.
- 848 Ptashne, M. (2013). Epigenetics: core misconception. *Proceedings of the National Academy*
- 849 *of Sciences*, 110(18), 7101–7103.
- 850 Putnam, H. M. (2012). *Resilience and acclimatization potential of reef corals under*
- 851 *predicted climate change stressors*. UNIVERSITY OF HAWAI 'I MĀNOA.
- 852 Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The vulnerability
- 853 and resilience of reef-building corals. *Current Biology*, 27(11), R528–R540.

Positive carryover effects in corals

- 854 Putnam, H. M., Davidson, J. M., & Gates, R. D. (2016). Ocean acidification influences
855 host DNA methylation and phenotypic plasticity in environmentally susceptible
856 corals. *Evolutionary Applications*, 9(9), 1165–1178.
- 857 Putnam, H. M., Edmunds, P. J., & Fan, T.-Y. (2010). Effect of a fluctuating thermal
858 regime on adult and larval reef corals. *Invertebrate Biology*, 129(3), 199–209.
- 859 Putnam, H. M., & Gates, R. D. (2015). Preconditioning in the reef-building coral
860 *Pocillopora damicornis* and the potential for trans-generational acclimatization in
861 coral larvae under future climate change conditions. *Journal of Experimental*
862 *Biology*, 218(15), 2365–2372.
- 863 R Core Team. (2016). *R: A language and environment for statistical computing*. Vienna,
864 Austria: R Foundation for Statistical Computing.
- 865 Randall, C. J., & Szmant, A. M. (2009). Elevated temperature affects development,
866 survivorship, and settlement of the elkhorn coral, *Acropora palmata* (Lamarck
867 1816). *The Biological Bulletin*, 217(3), 269–282.
- 868 Richmond, R. H., & Jokiel, P. L. (1984). Lunar periodicity in larva release in the reef
869 coral *Pocillopora damicornis* at Enewetak and Hawaii. *Bulletin of Marine Science*,
870 34(2), 280–287.
- 871 Riebesell, U., Fabry, V. J., Hansson, L., Gattuso, J.-P., & others. (2010). *Guide to best*
872 *practices for ocean acidification research and data reporting* (Vol. 260).
873 Publications Office of the European Union Luxembourg.
- 874 Ritson-Williams, R., Arnold, S. N., & Paul, V. J. (2016). Patterns of larval settlement
875 preferences and post-settlement survival for seven Caribbean corals. *Marine*
876 *Ecology Progress Series*, 548, 127–138.

Positive carryover effects in corals

- 877 Rivest, E. B., Comeau, S., & Cornwall, C. E. (2017). The Role of Natural Variability in
878 Shaping the Response of Coral Reef Organisms to Climate Change. *Current*
879 *Climate Change Reports*, 3(4), 271–281.
- 880 Rivest, E. B., & Hofmann, G. E. (2014). Responses of the metabolism of the larvae of
881 *Pocillopora damicornis* to ocean acidification and warming. *PLoS One*, 9(4),
882 e96172.
- 883 Rivest, E. B., & Hofmann, G. E. (2015). Effects of temperature and pCO₂ on lipid use
884 and biological parameters of planulae of *Pocillopora damicornis*. *Journal of*
885 *Experimental Marine Biology and Ecology*, 473, 43–52.
- 886 Rondon, R., Grunau, C., Fallet, M., Charlemagne, N., Sussarellu, R., Chaparro, C.,
887 Montagnani, C., Mitta, G., Bachere, E., Akcha, F., Cosseau, C. (2017). Effects of
888 a parental exposure to diuron on Pacific oyster spat methylome. *Environmental*
889 *Epigenetics*, 3(1). <https://doi.org/10.1093/eep/dvx004>
- 890 Ross, C., Ritson-Williams, R., Olsen, K., & Paul, V. J. (2013). Short-term and latent
891 post-settlement effects associated with elevated temperature and oxidative stress
892 on larvae from the coral *Porites astreoides*. *Coral Reefs*, 32(1), 71–79.
- 893 Ross, P., Parker, L., & Byrne, M. (2016). Transgenerational responses of molluscs and
894 echinoderms to changing ocean conditions. *ICES Journal of Marine Science:*
895 *Journal Du Conseil*, fsv254.
- 896 Sharp, K. H., Distel, D., & Paul, V. J. (2012). Diversity and dynamics of bacterial
897 communities in early life stages of the Caribbean coral *Porites astreoides*. *The*
898 *ISME Journal*, 6(4), 790–801. <https://doi.org/10.1038/ismej.2011.144>

Positive carryover effects in corals

- 899 Silbiger, N. J., Guadayol, Ò., Thomas, F. I., & Donahue, M. J. (2014). Reefs shift from
900 net accretion to net erosion along a natural environmental gradient. *Marine*
901 *Ecology Progress Series*, 515, 33–44.
- 902 Southam, C. M., & Ehrlich, J. (1943). Effects of extract of western red-cedar heartwood
903 on certain wood-decaying fungi in culture. *Phytopathology*, 33, 517–524.
- 904 Stimson, J. S. (1978). Mode and timing of reproduction in some common hermatypic
905 corals of Hawaii and Enewetak. *Marine Biology*, 48(2), 173–184.
- 906 Stoddart, J. A. (1983). Asexual production of planulae in the coral *Pocillopora*
907 *damicornis*. *Marine Biology*, 76(3), 279–284.
- 908 Stoddart, J. A., & Black, R. (1985). Cycles of gametogenesis and planulation in the coral
909 *Pocillopora damicornis*. *Marine Ecology Progress Series*, 153–164.
- 910 Stumpp, M., Hu, M. Y., Melzner, F., Gutowska, M. A., Dorey, N., Himmerkus, N.,
911 Holtmann, W.C., Dupont, S.T., Thorndyke, M.C., Bleich, M. (2012). Acidified
912 seawater impacts sea urchin larvae pH regulatory systems relevant for
913 calcification. *Proceedings of the National Academy of Sciences*, 109(44), 18192–
914 18197. <https://doi.org/10.1073/pnas.1209174109>
- 915 Stumpp, M., Wren, J., Melzner, F., Thorndyke, M. C., & Dupont, S. T. (2011). CO₂
916 induced seawater acidification impacts sea urchin larval development I: elevated
917 metabolic rates decrease scope for growth and induce developmental delay.
918 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*
919 *Physiology*, 160(3), 331–340.

Positive carryover effects in corals

- 920 Sunday, J. M., Calosi, P., Dupont, S., Munday, P. L., Stillman, J. H., & Reusch, T. B.
- 921 (2014). Evolution in an acidifying ocean. *Trends in Ecology & Evolution*, 29(2),
- 922 117–125.
- 923 Sunday, J. M., Crim, R. N., Harley, C. D. G., & Hart, M. W. (2011). Quantifying Rates of
- 924 Evolutionary Adaptation in Response to Ocean Acidification. *PLOS ONE*, 6(8),
- 925 e22881. <https://doi.org/10.1371/journal.pone.0022881>
- 926 Thor, P., & Dupont, S. (2015). Transgenerational effects alleviate severe fecundity loss
- 927 during ocean acidification in a ubiquitous planktonic copepod. *Global Change*
- 928 *Biology*, 21(6), 2261–2271.
- 929 Thornhill, D. J., Rotjan, R. D., Todd, B. D., Chilcoat, G. C., Iglesias-Prieto, R., Kemp, D.
- 930 W., LaJeunesse, T.C., McCabe Reynolds, J., Schmidt, G.W., Shannon, T., Warner,
- 931 M.E., Fitt, W. K. (2011). A connection between colony biomass and death in
- 932 Caribbean reef-building corals. *PLoS ONE*, 6(12), e29535.
- 933 <https://doi.org/10.1371/journal.pone.0029535>
- 934 Torda, G., Donelson, J. M., Aranda, M., Barshis, D. J., Bay, L., Berumen, M. L., ...
- 935 others. (2017). Rapid adaptive responses to climate change in corals. *Nature*
- 936 *Climate Change*, 7(9), 627–636.
- 937 Van Hooidonk, R., Maynard, J. A., & Planes, S. (2013). Temporary refugia for coral
- 938 reefs in a warming world. *Nature Climate Change*, 3(5), 508–511.
- 939 van Oppen, M. J. H., Gates, R. D., Blackall, L. L., Cantin, N., Chakravarti, L. J., Chan, W.
- 940 Y., Cormick, C., Crean, A., Damjanovic, K., Epstein, H., Peplow, L.M., Jones,
- 941 T.A., Miller, M., Pears, R.J., Torda, G., Wachenfeld, D., Weeks, A.R., Putnam,

Positive carryover effects in corals

- 942 H.M. (2017). Shifting paradigms in restoration of the world's coral reefs. *Global*
- 943 *Change Biology*. <https://doi.org/10.1111/gcb.13647>
- 944 van Oppen, M. J., Oliver, J. K., Putnam, H. M., & Gates, R. D. (2015). Building coral
- 945 reef resilience through assisted evolution. *Proceedings of the National Academy*
- 946 *of Sciences*, 112(8), 2307–2313.
- 947 Veron, J. E. N., Hoegh-Guldberg, O., Lenton, T. M., Lough, J. M., Obura, D. O., Pearce-
- 948 Kelly, P., Sheppard, C.R.C., Spalding, M., Stafford-Smith, M.G., Rogers, A. D.
- 949 (2009). The coral reef crisis: The critical importance of < 350ppm CO₂. *Marine*
- 950 *Pollution Bulletin*, 58(10), 1428–1436.
- 951 Webster, N. S., & Reusch, T. B. (2017). Microbial contributions to the persistence of
- 952 coral reefs. *The ISME Journal*.
- 953 Zhao, L., Schöne, B. R., Mertz-Kraus, R., & Yang, F. (2017). Sodium provides unique
- 954 insights into transgenerational effects of ocean acidification on bivalve shell
- 955 formation. *Science of The Total Environment*, 577, 360–366.
- 956 Zhao, L., Yang, F., Milano, S., Han, T., Walliser, E. O., & Schöne, B. R. (2018).
- 957 Transgenerational acclimation to seawater acidification in the Manila clam
- 958 *Ruditapes philippinarum*: Preferential uptake of metabolic carbon. *Science of The*
- 959 *Total Environment*, 627, 95–103.
- 960 Zizzari, Z. V., & Ellers, J. (2014). Rapid shift in thermal resistance between generations
- 961 through maternal heat exposure. *Oikos*, 123(11), 1365–1370.
- 962 <https://doi.org/10.1111/oik.01496>
- 963