

1 Title: Countries' geographic latitude and their human populations' cholesterol and blood
2 pressure

3 Yuhao Liu, BS¹; Robert D. Brook, MD²; Xuefeng Liu, PhD³; and James Brian Byrd, MD, MS⁴

4

5 ¹Department of Statistics, University of Michigan, Ann Arbor, MI, USA

6 ²Department of Medicine, University of Michigan Medical School, Ann Arbor MI, USA

7 ³Department of Systems, Populations and Leadership, University of Michigan School of Nursing,
8 Ann Arbor, MI, USA

9 ⁴Department of Medicine, University of Michigan Medical School, Ann Arbor, MI, USA

10

11 Corresponding Author: James Brian Byrd, MD, MS, University of Michigan, 5570C MSRB II,
12 Ann Arbor, MI 48109; Tel: (734) 998-7991, Email: jbyrd@med.umich.edu

13

14 Short running title: Latitude, cholesterol, and blood pressure

15

16 **Acknowledgements**

17 JBB is funded by the National Institutes of Health award K23HL128909. We thank
18 computational biologist Kasper Hansen for suggesting latitude might have a relationship with
19 blood pressure after examining a map JBB had made. We thank Edward Roccella and
20 Brahmajee Nallamothu for their thoughtful critical review of the manuscript.

21 **Competing Financial Interests Delcaration:** The authors have no conflicts of interest.

22 **Abstract**

23 **Background** Sunlight has been hypothesized to play a role in variation in cardiovascular
24 disease according to geographic latitude. **Objectives** To evaluate the plausibility of sunlight as
25 a factor in populations' average cholesterol and blood pressure **Methods** We analyzed World
26 Health Organization data including 180 or more countries' age-standardized average
27 cholesterol, age-standardized mean systolic blood pressure (BP), and age-standardized
28 prevalence of raised BP, by geographic latitude, over decades. We also performed analysis by
29 ultraviolet B light (UVB) exposure. **Results** Mean cholesterol increases with the distance of a
30 country from the Equator. This relationship has changed very little since 1980. Similarly, in
31 1975, mean systolic BP and prevalence of raised BP were higher in countries farther from the
32 Equator. However, the relationship between latitude and BP has changed dramatically; by 2015,
33 the opposite pattern was observed in women. Countries' average UVB exposure has a stable
34 relationship with cholesterol over recent decades, but a changing relationship with BP.

35 **Conclusions** Since sunlight exposure in a country is relatively fixed and its relationship with BP
36 has changed dramatically in recent decades, countries' average sunlight exposure is an unlikely
37 explanation for contemporary country-level variation in BP. However, our findings are consistent
38 with a putative effect of sunlight on countries' average cholesterol, as well as a no longer
39 detectable effect on BP decades ago. A parsimonious potential explanation for the relationship
40 between light and cholesterol is that 7-dehydrocholesterol can be converted to cholesterol, or in
41 the presence of ultraviolet light, can instead be converted to vitamin D.

42

43 **Introduction**

44 In the 1980s, the incidence of cardiovascular disease was noted to be higher in countries farther
45 from the Equator.(Fleck 1989) A popular hypothesis is that sunlight exerts a protective effect by
46 reducing cholesterol and/or blood pressure.(Grimes et al. 1996; Patwardhan et al. 2017;
47 Rostand 1997) A classic study in rabbits(Altschul 1953) and a more recent study in
48 mice,(Geldenhuis et al. 2014) a cross-sectional study of humans,(Prodam et al. 2016) and a
49 small clinical trial(Patwardhan et al. 2017) provide evidence of an effect of ultraviolet radiation
50 on cholesterol. A parsimonious potential explanation is that 7-dehydrocholesterol can be
51 converted to cholesterol, or in the presence of ultraviolet light, it can instead be converted to
52 vitamin D.(Geldenhuis et al. 2014; Patwardhan et al. 2017) Sunlight's effect on blood pressure
53 is hypothesized to have a more complex biological basis, which might involve vitamin D, nitric
54 oxide, and melanin.(Feelisch et al. 2010) There is experimental evidence to support an effect of
55 ultraviolet B light on blood pressure.(Krause et al. 1998) Sunlight might also change
56 cardiovascular risk factors by influencing countries' agriculture, for example the availability of
57 fresh fruits and vegetables. Whether effects of sunlight on cholesterol or blood pressure
58 manifest at the level of global health is not well understood since most ecological studies
59 relating sunlight exposure and/or latitude to cardiovascular risk factors are old and cross-
60 sectional and involve relatively few countries.

61 New treatments for elevated cholesterol and blood pressure have been developed and
62 disseminated since the 1970s, even as changes in health habits have swept the globe. It is
63 possible that the relationship between latitude and cholesterol or blood pressure has changed
64 accordingly, suggesting that sunlight is at most a small factor compared to other aspects of
65 lifestyle. If instead, latitude or ultraviolet radiation has an unchanged relationship with a
66 cardiovascular risk factor over decades despite lifestyle changes, a role for sunlight is more
67 likely. Using decades of longitudinal data from over 180 countries, we examined the relationship

68 between latitude and total cholesterol, and systolic blood pressure, with attention to changes
69 over time. We also examined the relationship between ultraviolet B light exposure and countries'
70 cholesterol or blood pressure. In addition, we examined whether sex differences exist in these
71 relationships.

72 **Methods**

73 We obtained the country latitudes used in Google's Public Data Explorer project.(Google) From
74 the World Health Organization's (WHO) Global Health Observatory,(World Health Organization)
75 we obtained country-level age-standardized estimates of: mean total cholesterol (1980-2009),
76 mean systolic blood pressure (1975-2015), and the prevalence of raised blood pressure (1975-
77 2015). Raised blood pressure was defined by WHO as systolic blood pressure \geq 140 mm Hg or
78 diastolic blood pressure \geq 90 mm Hg). Additional methods underlying the blood pressure
79 measurements have been published.(NCD Risk Factor Collaboration 2017)

80 In addition, we obtained country-level ultraviolet B light exposure (averaged between 1997-
81 2003) from the WHO Global Health Observatory. Some countries' names were designated
82 differently in the Google Public Data Explorer latitudes dataset and the WHO datasets. After
83 harmonizing the country names, we used local regression (LOESS) plots to visualize the
84 relationship between latitude and these endpoints. The R package '*ggplot2*' and its command
85 '*geom_smooth*' command were used to generate the LOESS curves and corresponding 95%
86 confidence intervals. We also calculated Spearman correlation coefficients for distance from the
87 Equator and countries' mean cholesterol, mean systolic blood pressure, or the prevalence of
88 raised blood pressure. We further evaluated whether our findings were stable over recent
89 decades. Analyses were performed using R 3.4.3.(R Core Team (2017))

90

91

92 Results

93 Latitude-based analyses included 188 countries' mean cholesterol values, 180 countries' mean
94 systolic blood pressures, and 189 countries' prevalences of raised blood pressure. Ultraviolet B
95 radiation exposure-based analyses included 185 countries' mean cholesterol values and 186
96 countries' raised blood pressure prevalences.

97 Since at least the 1980s, average total cholesterol has been lowest in women (**Figure 1A**) and
98 men (**Figure 1B**) living in countries at or near the Equator and has increased symmetrically with
99 distance north or south of the Equator. Despite a downward shift in countries' average
100 cholesterol over recent decades, this symmetry around the Equator has been preserved.
101 Absolute distance of a country from the Equator has correlated with total cholesterol in women
102 (**Figure 1C**) and men (**Figure 1D**) since at least the 1980s ($r_s = 0.66$, $P < 2.2e^{-16}$ [females]; $r_s =$
103 0.67 , $P < 2.2e^{-16}$ [males] in 1980; $r_s = 0.51$, $P = 1.2e^{-13}$ [females]; $r_s = 0.57$, $P < 2.2e^{-16}$ [males] in
104 2009).

105 In 1975, countries' mean systolic blood pressure followed a pattern similar to mean cholesterol,
106 with systolic blood pressure increasing with distance north or south of the Equator in females
107 (**Figure 2A**) and to a lesser extent in males (**Figure 2B**). As with cholesterol, countries' mean
108 systolic blood pressure exhibited symmetry north and south of the Equator. In contrast to
109 cholesterol, mean systolic blood pressure's relationship with latitude has been unstable over
110 time. In fact, by 2015, the pattern had reversed itself in females (**Figure 2A**). In males,
111 countries' mean systolic blood pressure had a complex, bimodal relationship with latitude in
112 2015 (**Figure 2B**). Particularly in females, the changes in latitude's relationship with age-
113 standardized mean systolic blood pressure have been similar in the Northern and Southern
114 hemispheres, preserving the north-south symmetry seen in the 1970s, with the Equator
115 appearing as an inflection point. The slope of the relationship between absolute distance from
116 the Equator and mean systolic blood pressure has been decreasing over time in females

117 **(Figure 2C)**--in whom it has reversed direction in recent years to a negative relationship--and in
118 males **(Figure 2D)**, $r_s = 0.52$, $P=4.0e^{-14}$ [females]; $r_s = 0.49$, $P=2.9e^{-12}$ [males] in 1975; $r_s = -0.28$,
119 $P=0.0002$ [females]; $r_s = 0.07$, $P=0.32$ [males] in 2015). Compared to mean systolic blood
120 pressure, countries' prevalences of raised blood pressure had similar LOESS plots **(Figures 3A**
121 **and 3B)**, but stronger symmetry around the Equator and marginally stronger correlation in the
122 1970s ($r_s = 0.57$, $P<2.2e^{-16}$ [females]; $r_s = 0.60$, $P<2.2e^{-16}$ [males] in 1975; $r_s = -0.29$, $P=4.9e^{-5}$
123 [females], $r_s = 0.13$, $P=0.09$ [males] in 2015, **Figure 3**).

124 The relationship between ultraviolet B radiation and cholesterol or blood pressure mirrored their
125 relationships with latitude. For decades, countries with higher average ultraviolet B radiation
126 levels have had lower mean cholesterol in females **(Figure 4A)** and males **(Figure 4B)**. As with
127 latitude, the relationship between ultraviolet B radiation and raised blood pressure has been
128 changing over time, with clear reversal of this relationship in females in 2015 compared to 1975
129 **(Figure 4C)** and dramatic changes in males **(Figure 4D)**.

130 **Discussion**

131 The major new findings of this study are that for at least four decades, males' and females'
132 average cholesterol levels have been higher in countries farther from the Equator, whereas
133 mean systolic blood pressure and the prevalences of raised blood pressure showed a similar
134 pattern decades ago, but no longer. In women, blood pressure's relationship with latitude has
135 reversed over time. Countries' average ultraviolet B radiation has had a relatively constant
136 relationship for decades with their populations' mean cholesterol. The relationship between
137 countries' average ultraviolet B radiation exposure and blood pressure has changed
138 dramatically over time.

139 In a small study (n=338) conducted at three sites in British Columbia, plasma cholesterol
140 decreased with increasing latitude, a finding the authors attributed to differences in Rhesus

141 Blood Group system.(Alfred et al. 1974) Conversely, hypertension prevalence or blood pressure
142 have been reported to increase with distance north or south of the Equator or with increased
143 solar radiation exposure within single countries(Cabrera et al. 2016; Rostand et al. 2016) and in
144 studies of smaller groups of countries(Duranton et al. 2018; Rostand 1997) than we have
145 analyzed. However, adjusting for vitamin D levels had no effect on the solar radiation-blood
146 pressure relationship in a recent study, suggesting blood pressure's variation by solar radiation
147 levels is not mediated by vitamin D.(Rostand et al. 2016) Public availability of high-quality data
148 facilitated our more comprehensive analysis of latitude's and ultraviolet B radiation's relationship
149 with males' and females' cardiovascular risk factors estimated at the national level, around the
150 globe and over decades. The focus of our study was to understand the relationship between
151 latitude and cholesterol or blood pressure at the whole-globe level. Our findings contrast with
152 the prior single-country analysis of latitude and cholesterol, suggesting a possible Simpson's
153 paradox, in which findings at a smaller scale of analysis are at odds with findings at a larger
154 scale of analysis. Countries' unique social history might explain such a paradox. Our latitude-
155 blood pressure findings using data from decades ago are consistent with results from decades
156 ago, but since then, the relationship not only has changed in men, it has reversed in women.
157 The factors explaining the stable relationship between countries' latitude and their populations'
158 cholesterol are likely different from those explaining the relationship between latitude and blood
159 pressure, which has changed dramatically over time. Exposure to sunlight has been proposed
160 to explain the relationship between latitude and blood pressure,(Feelisch et al. 2010) but the
161 marked changes in the relationship indicate it is no longer a key determinant of country-level
162 differences in blood pressure, if it once was. Increased sunlight lowers cholesterol in
163 experimental settings in rabbits and humans.(Altschul 1953; Patwardhan et al. 2017) Sunlight
164 exposure could plausibly explain or contribute to the more stable relationship between latitude
165 and cholesterol. An indirect effect of sunlight on cardiovascular risk factors through differences

166 in agriculture is an alternate, plausible explanation of the relationship between UV light and
167 cholesterol. There are likely other plausible explanations, as well. For example, countries' gross
168 domestic product varies by latitude. Interestingly, these differences in economic productivity
169 might also be related to climate,(Masters and McMillan 2001) and thus related to sunlight.

170 Strengths of the current study include longitudinal analysis of high-quality data from more than
171 180 countries and the bringing together of multiple publicly available datasets to throw new light
172 on an old question. The study is truly global in scale. The principal limitation of the study is
173 directly related to this strength: ecological studies can show us what is true around the globe,
174 but they permit only relatively weak inferences about how to explain what is seen. A second
175 limitation is that our analysis compared countries' UVB light exposure averaged between 1997-
176 2003 to risk factors collected over a broader time span. Our analysis assumes stability of UVB
177 light exposure, whereas there have been some regional changes in UVB light exposure due to
178 the hole in the ozone layer. This issue affects only the UVB light analyses.


179 Ischemic heart disease (IHD) is the leading cause of death globally. A long-term 10% reduction
180 in total cholesterol lowers risk of ischemic heart disease by 50% at age 40 and 20% at age
181 70,(Law et al. 1994) and a 20 mm Hg lower usual systolic blood pressure is associated with a
182 50% decrease in death from IHD and 50% decrease in death from stroke.(Lewington et al.
183 2002) As we seek new means of understanding cardiovascular risk reduction, additional studies
184 to better understand the effect of sunlight on cardiovascular risk are needed.

185 **Conclusions**

186 Since sunlight exposure in a country is relatively fixed and its relationship with BP has changed
187 dramatically in recent decades, countries' average sunlight exposure is an unlikely explanation
188 for contemporary country-level variation in BP. However, our findings are consistent with a
189 putative effect of sunlight on countries' average cholesterol, as well as a no longer detectable

190 effect on BP decades ago. A parsimonious potential explanation for the relationship between
191 light and cholesterol is that 7-dehydrocholesterol can be converted to cholesterol, or in the
192 presence of ultraviolet light, can instead be converted to vitamin D.


193 **Figure 1.**



195 **Figure 2.**


A.

1975 – 2015, Females




B.

1975 – 2015, Males




C.

1975 – 2015, Females




D.

1975 – 2015, Males



196
197

198 **Figure 3.**




199

200

Figure 4.


A.

1980 – 2009, Females




B.

1980 – 2009, Males




C.

1975 – 2015, Females



D.

1975 – 2015, Males



References

Alfred BM, Lee M, Desai ID. 1974. A relationship between plasma cholesterol level, latitude, and the rhesus blood group system. *Hum Biol* 46:641-646.

Altschul R. 1953. Inhibition of experimental cholesterol arteriosclerosis by ultraviolet irradiation. *N Engl J Med* 249:96-99.

Cabrera SE, Mindell JS, Toledo M, Alvo M, Ferro CJ. 2016. Associations of blood pressure with geographical latitude, solar radiation, and ambient temperature: Results from the Chilean health survey, 2009-2010. *Am J Epidemiol* 183:1071-1073.

Duranton F, Kramer A, Szwarc I, Bieber B, Gayraud N, Jover B, et al. 2018. Geographical variations in blood pressure level and seasonality in hemodialysis patients. *Hypertension* 71:289-296.

Feelisch M, Kolb-Bachofen V, Liu D, Lundberg JO, Revelo LP, Suschek CV, et al. 2010. Is sunlight good for our heart? *Eur Heart J* 31:1041-1045.

Fleck A. 1989. Latitude and ischaemic heart disease. *Lancet* 1:613.

Geldenhuis S, Hart PH, Endersby R, Jacoby P, Feelisch M, Weller RB, et al. 2014. Ultraviolet radiation suppresses obesity and symptoms of metabolic syndrome independently of vitamin D in mice fed a high-fat diet. *Diabetes* 63:3759-3769.

Google. Canonical latitudes for public data explorer project. Available: Available from https://developers.google.com/public-data/docs/canonical/countries_csv.

Grimes DS, Hindle E, Dyer T. 1996. Sunlight, cholesterol and coronary heart disease. *QJM* 89:579-589.

Krause R, Buhning M, Hopfenmuller W, Holick MF, Sharma AM. 1998. Ultraviolet B and blood pressure. *Lancet* 352:709-710.

Law MR, Wald NJ, Thompson SG. 1994. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 308:367-372.

Lewington S, Clarke R, Qizilbash N, Peto R, Collins R. 2002. Age-specific relevance of usual blood pressure to vascular mortality: A meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 360:1903-1913.

Masters WA, McMillan MS. 2001. Climate and scale in economic growth. *Journal of Economic Growth* 6:167-186.

NCD Risk Factor Collaboration. 2017. Worldwide trends in blood pressure from 1975 to 2015: A pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet* 389:37-55.

Patwardhan VG, Mughal ZM, Padidela R, Chiplonkar SA, Khadilkar VV, Khadilkar AV. 2017. Randomized control trial assessing impact of increased sunlight exposure versus vitamin d supplementation on lipid profile in indian vitamin d deficient men. *Indian J Endocrinol Metab* 21:393-398.

Prodam F, Zanetta S, Ricotti R, Marolda A, Giglione E, Monzani A, et al. 2016. Influence of ultraviolet radiation on the association between 25-hydroxy vitamin d levels and cardiovascular risk factors in obesity. *J Pediatr* 171:83-89.e81.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rostand SG. 1997. Ultraviolet light may contribute to geographic and racial blood pressure differences. *Hypertension* 30:150-156.

Rostand SG, McClure LA, Kent ST, Judd SE, Gutierrez OM. 2016. Associations of blood pressure, sunlight, and vitamin d in community-dwelling adults. *J Hypertens* 34:1704-1710.

World Health Organization. Global health observatory data repository. Available: Available from <http://apps.who.int/gho/data/node.imr>.

Figure Legends.

Figure 1. LOESS plots of latitude or distance from the Equator and countries' mean total cholesterol (age-standardized). The top two panels show countries' latitude and the mean cholesterol in females (**Panel A**) and males (**Panel B**). The vertical dotted lines represent 23.5 degrees North and South, the Tropics of Cancer and Capricorn, which define the tropics. The lower two panels show LOESS plots of absolute distance from the Equator in degrees and countries' mean total cholesterol (age-standardized) in females (**Panel C**) and males (**Panel D**). To provide confidence intervals while maintaining visual clarity, the oldest and most recent years' LOESS curves show the 95% confidence interval in pink and grey, respectively.

Figure 2. LOESS plots of latitude or distance from the Equator and countries' mean systolic blood pressure (age-standardized estimate). The top two panels show mean systolic blood pressure according to latitude in females (**Panel A**) and males (**Panel B**), using data from every 10 years between 1975 and 2015. The lower two panels are LOESS plots of absolute distance from the Equator in degrees and countries' mean systolic blood pressure (age-standardized estimate) in females (**Panel C**) and males (**Panel D**), respectively. To provide confidence intervals while maintaining visual clarity, the oldest and most recent years' LOESS curves show the 95% confidence interval in pink and grey, respectively.

Figure 3. LOESS plots of latitude or distance from the Equator and countries' prevalence of raised blood pressure (SBP \geq 140 or DBP \geq 90 [age-standardized estimate]). The upper two panels show the prevalence of raised blood pressure according to latitude in females (**Panel A**) and males (**Panel B**), using data from every 10 years between 1975 and 2015. The vertical dotted lines represent 23.5 degrees North and South, the Tropics of Cancer and Capricorn, which define the tropics. The lower two panels are LOESS plots of absolute distance from the Equator in degrees and countries' prevalence of raised blood pressure in females (**Panel C**) and males (**Panel D**), respectively. To provide confidence intervals while maintaining visual clarity,

the oldest and most recent years' LOESS curves show the 95% confidence interval in pink and grey, respectively.

Figure 4.

LOESS plots of countries' log-transformed ultraviolet B radiation exposure (averaged between 1997-2003) and their populations' mean cholesterol or prevalence of raised blood pressure (SBP \geq 140 or DBP \geq 90 [age-standardized estimate]). The upper two panels show mean cholesterol in females (**Panel A**) and males (**Panel B**) according to ultraviolet B radiation exposure. The lower two panels show LOESS plots of the relationship between log-transformed ultraviolet B radiation exposure and prevalence of raised blood pressure in females (**Panel C**) and males (**Panel D**). To provide confidence intervals while maintaining visual clarity, the oldest and most recent years' LOESS curves show the 95% confidence interval in pink and grey, respectively.