Autolysin-Independent DNA Release in *Streptococcus pneumoniae* *in vitro* Biofilms

Mirian Domenech,^{a,b} Ernesto García^{a,b#}

^aDepartamento de Biotecnología Microbiana y de Plantas, Centro de Investigaciones Biológicas (CSIC), Ramiro de Maeztu 9, 28040 Madrid, Spain.

^bCIBER de Enfermedades Respiratorias (CIBERES), Madrid, Spain

Running Head: Extracellular DNA in Pneumococcal Biofilms

#Address correspondence to Ernesto García, e.garcia@cib.csic.es.

Word counts: Abstract, 232; Importance, 147; Text, 5,251.

ABSTRACT Biofilms are defined as layers of cells of microorganisms adhered to the surface of a substrate and embedded in an extracellular matrix and provide an appropriate environment for increased genetic exchange. Extracellular DNA (eDNA) is an essential component of the extracellular matrix of microbial biofilms, but the pathway(s) responsible for DNA release are largely unknown. Autolysis (either spontaneous or phage-induced) has been proposed the major event leading to the appearance of eDNA. The 'suicidal tendency' of *Streptococcus pneumoniae* is well-known, with lysis mainly caused by the triggering of LytA, the major autolytic amidase. However, the LytC lysozyme and CbpD (a possible murein hydrolase) have also been shown involved. The present work examines the relationship between eDNA, autolysins, and the formation and maintenance of *in vitro* pneumococcal biofilms, via fluorescent labelling combined with confocal laser scanning microscopy, plus genetic transformation experiments. Bacterial DNA release mechanisms other than those entailing lytic enzymes were shown to be involved by demonstrating that horizontal gene transfer in biofilms takes place even in the absence of detectable autolytic activity. It had been previously suggested that the quorum sensing systems ComABCDE and LuxS/AI-2 are involved in the production of eDNA as a response to the accumulation of quorum sensing signals, although our immunofluorescence results do not support this hypothesis. Evidence that the release of DNA is somehow linked to the production of extracellular vesicles by *S. pneumoniae* is provided.

IMPORTANCE Most human bacterial infections are caused by microorganisms growing as biofilms. Bacteria in biofilms are less susceptible to antimicrobials and to killing by the host immune system, are very difficult to eliminate and cause recalcitrant and persistent diseases. Extracellular DNA is one of the major components of the bacterial biofilm matrix. In the present study, we provide direct evidence of the existence of biologically active (transforming), extracellular DNA in *Streptococcus pneumoniae* biofilms. In

46 previous studies, the involvement of three pneumococcal choline-binding
 47 proteins with autolytic activity (LytA, LytC and CbpD) in DNA release had been
 48 reported. In contrast, we demonstrate here that pneumococcal *in vitro* biofilms
 49 do contain eDNA, even in the absence of these enzymes. Moreover, our results
 50 suggest that DNA release in *S. pneumoniae* biofilms is connected with the
 51 production of extracellular vesicles and that this DNA is associated to the outer
 52 part of the vesicles.

53

The human pathogen *Streptococcus pneumoniae* is a leading cause of pneumonia, meningitis and bloodstream infections in the elderly, and one of the main pathogens responsible for middle ear infections in children. It is carried asymptomatically in the nasopharynx of many healthy adults, and in as many as 20–40% of healthy children (colonization begins shortly after birth) (1). Pneumococcal biofilms appear on adenoid and mucosal epithelium in children with recurrent middle-ear infections and otitis media with effusion, and on the sinus mucosa of patients with chronic rhinosinusitis, and they can be also formed *in vitro* (2, 3). Biofilm formation in *S. pneumoniae* is an efficient way of evading both the classical and the PspC-dependent alternative complement pathways (4).

Over 60% of all human bacterial infections, and up to 80% of those that become chronic, are thought to involve growth in biofilms. A biofilm is defined as an accumulation of microorganisms embedded in a self-produced extracellular matrix (ECM) adhered to an abiotic or living surface (5). The ECM is composed of different polymers, mainly polysaccharides, proteins, and nucleic acids. The requirement of extracellular deoxyribonucleic acid (eDNA) in ECM formation and maintenance has been documented in a variety of Gram-positive and Gram-negative bacteria (6). eDNA binds to polysaccharides and/or proteins, protecting bacterial cells from physical and/or chemical challenges, as well as providing biofilms with structural integrity. It is a major component of the *S. pneumoniae* ECM (7-11). Various pneumococcal surface proteins, *e.g.*, several members of the choline-binding family of proteins (CBPs) (11, 12) and the pneumococcal serine-rich repeat protein (PsrP) (13), form tight complexes with eDNA via electrostatic interactions, a mechanism proposed widespread among microorganisms (14). It should be noted that PsrP appears in $\approx 60\%$ of clinical pneumococcal isolates, whereas the main CBPs — LytA (SPD_1737/SPD_RS09190), LytB (SPD_0853/SPD_RS04550), LytC

(SPD_1403/SPD_RS07385) or CbpD (SPD_2028/SPD_RS10645) among others — are present in all *S. pneumoniae* strains.

The source of eDNA may vary across microorganisms and in part appears because of autolysis, phage-induced lysis, and/or active secretion systems, as well as through association with extracellular vesicles (EV). In *S. pneumoniae*, the release of DNA during limited lysis of the culture, *i.e.*, by controlled autolysis directed by the main CBP autolysins (LytA *N*-acetylmuramoyl- γ -alanine amidase [EC 3.5.1.28; NAM-amidase] and LytC lysozyme [EC 3.2.1.17; muramidase]), as well as prophage-mediated lysis, have been proposed as biofilm-promoting in part of the bacterial population (9, 11). The NAM-amidase LytA, the main autolytic enzyme of *S. pneumoniae*, is kept under control by lipoteichoic acid — a membrane-bound teichoic acid that contains choline — during exponential growth (15), and regulated at the level of substrate recognition (16). LytC also acts as an autolysin when pneumococci are incubated at about 30°C, a temperature close to that of the upper respiratory tract (17); this lysozyme might be post-transcriptionally inhibited by CbpF (SPD_0345/SPD_RS01835) (18).

Studies in different microorganisms suggest that the appearance of eDNA in biofilms may also be a response to the accumulation of quorum sensing (QS - a cell density-dependent communication system that regulates cooperative behavior) signals (for a recent review, see reference 19). Two early studies on genetic transformation in planktonic cultures showed *S. pneumoniae* to release measurable amounts of DNA in the absence of detectable autolysis (20). Although these pioneering studies were forgotten for years, more recent investigations have shown that LytA, LytC and CbpD (a putative cell wall-degrading enzyme) are directly responsible for autolytic DNA release from only a subset of cells in competent pneumococcal planktonic cultures (21, 22). In fact, the killing of non-competent sister cells by competent pneumococci — a phenomenon named fratricide — promotes autolysis and DNA release (23).

It is well known that competence induction in *S. pneumoniae* depends on a QS mechanism (24). LytA, LytC and CbpD are all required for autolysis (and for the concomitant DNA release) when pneumococci grow under planktonic conditions (25). It is currently believed that the limited damage caused by CbpD activates LytA and LytC, resulting in the more extensive lysis of target cells than that achieved by CbpD alone. LytA and LytC are constitutively synthesized by non-competent cells. However, while the expression of LytA increases during competence, LytC is not part of the competence regulon in *S. pneumoniae* (26). A slightly different situation has been proposed to occur in biofilms. The lysis of target (non-competent) cells in biofilms requires CbpD to act in conjunction with LytC, whereas LytA is not required for efficient fratricide-mediated gene exchange (26). In these experiments, however, a direct visualization of eDNA in the ECM was not reported. Interestingly, the transcription of both *lytA* and *cbpD* also appears to be regulated by the LusX/autoinducer-2 (AI-2) QS (27).

Via the use of an *in vitro* biofilm model system, the present work provides evidence that a small (but significant) proportion of biologically active, eDNA in pneumococcal biofilms is released into the medium by an alternative (or complementary) pathway to cell autolysis. It would appear that this occurs independent of the activity of LytA, LytC and CbpD and the QS systems (ComABCDE and LuxS/AI).

RESULTS

Visualization of eDNA in the pneumococcal biofilm. *S. pneumoniae* R6 biofilms grown for 5 h at 34°C in C+Y medium were stained with a combination of SYTO 59, DDAO (7-hydroxy-9H-[1,3-dichloro-9,9-dimethylacridin-2-one]) and anti-double-stranded (ds) DNA monoclonal antibodies (α -dsDNA). When examined under the confocal laser scanning microscope (CLSM), abundant, mostly cell-associated eDNA was observed (Fig. 1). However, when scanned at 488 nm (green), immunostained eDNA appeared as a lattice-like array

consisting of long DNA fibers, mainly at the top of the biofilm (Fig. 1C). Only seldomly they were associated with the cocci (marked with yellow arrows in Fig. 1I). At the bottom of the biofilm, small areas of what appeared to be compacted eDNA (but no fibers) were seen (data not shown).

Notably, DNA fibers were not observed when DDAO was employed. Many reports have used DDAO for staining eDNA in biofilms (28). However, a recent evaluation of eDNA stains in biofilms of various species has shown that this compound was neither completely cell impermeant nor capable to reveal DNA-containing fibrillar structures (29). As our results were in agreement with these data, DDAO was not used in additional experiments.

To study the dynamics of eDNA release, strain R6 was incubated under biofilm-forming conditions for up to 5 h at 34°C. Immunostaining with α -dsDNA of sessile (adherent) cells revealed the existence of eDNA even at early incubation times (Fig. 2). At 3 h, eDNA filaments were visible, although the majority of eDNA appeared as dots and patches, which may represent different stages of condensation of DNA–protein aggregates, as previously suggested (12). In 5 h-old biofilms, eDNA threads were infrequent and mainly located at the top of the biofilm (see above). Previous studies have revealed the existence of a mature ECM (consisting of DNA, proteins and polysaccharides) at this time point (11). Immunostaining planktonic cultures (*i.e.*, non-adherent cells) revealed the existence of eDNA filaments that were more abundant in younger than in older cultures (Fig. 2).

Evidence of the presence of eDNA in biofilms formed by different pneumococcal mutants. The involvement of eDNA in biofilm formation and maintenance was ascertained using strain P046, which lacks the two main autolytic CBPs. Strain P234 was employed as a control; this has a point mutation in the *pspC* (= *cbpA*) gene (SPD_2017/SPD_RS10590), which encodes a CBP important in virulence (30). PspC, which lacks any autolytic activity and is partly involved in biofilm formation *in vitro* (on polystyrene

microtiter plates) (7) and *in vivo* (in a murine nasopharynx colonization model) (3), also forms complexes with the eDNA (12). Compared to the parental R6 strain, both mutants showed reduced biofilm-forming capacity, in agreement with previous findings (7). Interestingly, biofilm formation, but not culture growth, was greatly impaired when pneumococci were grown in the presence of DNase I (Fig. 3). Moreover, the incubation of preformed biofilms with DNase I drastically diminished the number of biofilm-associated sessile cells. As a whole, however, preformed biofilms were less reduced by DNase I treatment than growing biofilms, strongly suggesting that eDNA is more important and/or more exposed during the early stages of biofilm formation. Alternatively, eDNA may become resistant to DNase enzymes during biofilm maturation by forming complexes with other macromolecules such as CBPs.

The presence of eDNA in the biofilms of $\Delta comC$ (SPD_2066/SPD_RS10845) or $\Delta luxS$ (SPD_0309/SPD_RS1650) mutants was also analyzed. These genes are essential for the functioning of two QS systems documented as being involved in biofilm formation (see above). The *comC* gene codes for the pre-CSP (competence-stimulating peptide), which is matured and exported by the ComA–ComB complex as an unmodified 17-residue-long peptide pheromone. LuxS is an S-ribosylhomocysteine lyase, and is responsible for the production of the QS molecule homoserine lactone autoinducer 2 (AI-2). It has been reported that transcription of competence genes (including *lytA* and *cbpD*) is reduced in a $\Delta luxS$ strain (27). As shown above for two CBP mutants, the $\Delta comC$ and $\Delta luxS$ mutants produced only $\approx 50\%$ of the biofilm formed by the R6 strain (Fig. 3). Positive evidence for the involvement of eDNA in formation and maintenance of these biofilms was also obtained by treatment with DNase I (Fig. 3).

Contribution of lytic CBPs to eDNA release. The presence of eDNA in the ECM of biofilms formed by strain P046 — a double *LytA*[−] *LytC*[−] mutant — was unexpected since it is generally believed that at least one of the autolysins is required for DNA release (see above). To gain further insight, the existence of

eDNA in the biofilm formed by a mutant deficient in CbpD (or its combination with the *lytA* and *lytC* mutations) was analyzed. A *pspC* mutant (strain P234) was included as a control. Notably, eDNA was present even in the ECM of strain P204, a mutant deficient in the three CBPs, *i.e.*, LytA, LytC and CbpD (Fig. 4). The morphology of the biofilms formed by strain P204 differed from those of R6, with the former biofilms containing fewer microcolonies and larger eDNA patches than the wild type.

It is well known that when pneumococcal cells are incubated with 2% choline chloride, CBPs are released into the medium. Those with enzymatic activity are completely inhibited (31), but transformability is not altered (32). Under these conditions, however, the biofilm-forming capacity is severely reduced (7). This was confirmed in the present work, and might be attributed (at least in part) to the drastically diminished eDNA content of the biofilm. Quite unexpectedly, R6 biofilms formed in the presence of 2% choline chloride still showed the presence of eDNA (Fig. 4).

The competence QS system is not involved in eDNA release. The presence of eDNA in biofilms formed by additional *S. pneumoniae* strains possessing combinations of mutations affecting the *comA* gene and various lytic genes was studied by CLSM. These strains were R391, P203, P204, and P213. The P147 strain (*ciaH*_{Tupelo_VT}; SPD_0702/SPD_RS0372) was also included since in previous work our group has shown that CiaR/H, a two-component signal transduction system that mediates the stress response, is in some way: a) implicated in the triggering of the LytA autolysin (33), b) required for efficient *in vitro* biofilm formation and nasopharyngeal colonization in a mouse model (34), and c) involved in the control of competence for genetic transformation (35). CLSM observations of *in vitro* biofilms showed eDNA to be also present in the biofilms of every mutant tested (Fig. 5).

Transforming capacity of eDNA in pneumococcal biofilms. The above results indicate that *in vitro* pneumococcal biofilms are capable of releasing

eDNA in a manner apparently independent of the activity of the three pneumococcal autolysins LytA, LytC and CbpD — even though these were generally believed necessary for DNA release in planktonic cultures. To investigate whether the eDNA in the biofilms formed by pneumococci simultaneously deficient in these three lytic enzymes has transforming activity, the efficacy of gene transfer in mixed biofilms was determined. For this, the reciprocal (donor and recipient) transforming capacity of two strain pairs was measured: in addition to being autolysin proficient (or not), one pair harbored the well-known low efficiency (LE) *nov1* marker (a C:G to T:A transition mutation conferring novobiocin resistance) (36), and the other the high efficiency (HE) marker *str41* (an A:T to C:G transversion), which bestows streptomycin resistance (37). To allow for further characterization of the direction of DNA transfer, the latter pair of strains used was also resistant to optochin. In pneumococcal transformation, the LE markers return 5–10% as many transformed cells as do HE markers (which are little degraded, or not at all). Table 1 shows that, in the biofilms, the spontaneous transformation of autolysin-proficient strains (P233 and P273) took place at levels typically observed in planktonic cultures when using naked chromosomal DNA (between 0.1 and 1% of total viable cells). Moreover, the heteroduplex DNA base mismatch repair system (Hex), which is responsible for marker-specific variations in transforming efficiency in planktonic cultures (38), was functional in the pneumococcal biofilms, as deduced from the relative transfer efficiency of the LE *nov1* and HE *str41* mutations (Table 1). In agreement with that reported by other authors (26), the transformation frequency was reduced by more than 100-fold in mixed biofilms formed by the mutants deficient in the three lytic enzymes. On average, however, one among 10^5 pneumococci showed transformation to streptomycin resistance, demonstrating that a small amount of biologically active eDNA was still present in the ECM of the biofilms formed by the triple deficient mutant.

Previous studies performed with different bacteria have suggested that, in addition to autolytic phenomena, eDNA might be associated with EV. Recent evidence indicates that *S. pneumoniae*, actively sheds extracellular nano-sized EV, as do many other microorganisms (39-41). Whether pneumococcal EV contains DNA, however, was never examined. Here, high-speed sediments of biofilm filtrates — a crude preparation of EV with associated DNA (see Fig. S2 in the supplemental material)— were used to perform additional transformation experiments. The results confirmed a measurable amount of transforming DNA to be present in these crude preparations, and that this DNA could be destroyed by treatment with DNase I (Table 2). Comparable results were obtained when strain P271 was incubated under planktonic conditions.

DISCUSSION

It has been known for decades that biofilm ECM contains eDNA, but its active role in biofilm appearance and maintenance was not recognized until Whitchurch *et al.* added DNase I to a *Pseudomonas aeruginosa* biofilm and watched the biofilm disappear (42). Since then, many reports have confirmed that a plethora of bacteria require eDNA to establish and maintain biofilms (6). The destruction of this eDNA provides a way of fighting biofilm-producing pathogens. Indeed, numerous clinical studies have shown that aerosolized DNase I (dornase alpha, a recombinant DNase I from the human pancreas) is highly effective in this respect, improving the lung function of patients with cystic fibrosis (43).

In vivo studies have shown that eDNA is a major element of biofilms. However, whether it is of bacterial or environmental (including host) origin (or both) is controversial; certainly it is difficult to determine which is the case under *in vivo* conditions. Studies using *in vitro* biofilm models (grown on plastic, glass, or other abiotic surfaces) provide a convenient way to screen large sets of strains, treatments, or growing conditions (44).

Among the possible origins of eDNA in pneumococcal biofilms (14), autolysins (either from bacterial or phage origin) have been proposed to have critical role. Previous studies performed at our own and other laboratories have employed indirect (e.g., DNase I treatment and/or intrabiofilm transformation experiments) and direct methods (e.g., staining with a variety of DNA-specific fluorophores) to disclose the presence of eDNA in pneumococcal biofilms (7-9, 11, 26, 27). Immunostaining with α -dsDNA combined with CLSM also revealed the existence of long filaments of eDNA in biofilms formed either by *Enterococcus faecalis* (45) or *Haemophilus influenzae* ECM (46). Moreover, the presence of eDNA-containing fibrous structures in the ECM of *Staphylococcus aureus* and *Propionibacterium acnes* biofilms has also been reported using α -dsDNA and atmospheric scanning electron microscopy (47, 48). The use of different technical approaches reinforces the idea that the fibrous assemblies of eDNA observed in some bacterial biofilms actually exist and also offers a novel perspective of the ECM structure of pneumococcal biofilms. Since most of the DNA filaments were found at the top of the mature biofilm (where actively growing cells reside) within 3 h of growth (Fig. 1), it is unlikely that these eDNA fibers were formed exclusively via autolysis. The same conclusion might be drawn from the CLSM images of immunostained planktonic cultures (Fig. 2).

In agreement with previous results, pneumococcal mutants either lacking the major autolysins or deficient in Com or LuxS/AI2 QSs showed limited biofilm-forming capacity. However, the results obtained following DNase I treatment of growing or pre-formed biofilms strongly suggest that those biofilms still contain eDNA. Direct CLSM visualization of the corresponding immunostained biofilms fully confirmed the presence of eDNA, even when the pneumococci were incubated in the presence of 2% choline chloride; this is known to induce the complete non-competitive inhibition of CBPs with enzyme activity, and to release all CBPs, whether enzymatic or not, from their bacterial surface attachments. It should be noted, however, that even at this high concentration,

choline does not separate the CBP–DNA complexes that form part of the ECM of *S. pneumoniae* biofilms (11, 12). Nevertheless, since the complete inhibition of the enzymatic activity of autolysins takes place under these conditions, an exclusive, direct role for such enzymes in eDNA release appears to be unlikely. More direct evidence was obtained using biofilms formed by strain P204, a triple $\Delta lytA \Delta lytC \Delta cbpD$ mutant, in which the presence of eDNA was also verified (Fig. 4). The release of eDNA was also observed in biofilms formed by $\Delta comA$ mutants or combined $\Delta comA$ /autolysin-deficient mutations. Most notably, the existence of biologically active eDNA in pneumococcal biofilms — even when autolysin-deficient strains were used — was fully confirmed by *in situ* reciprocal transformation experiments (Table 1). Deletion of either *lytA*, *lytC* and/or *cbpD* does not alter the transformability of the mutants compared to their parental strains when chromosomal DNA or plasmid(s) is used as donor material (17, 49). Wei and Håvarstein studied the impact of LytA, LytC and CbpD on transformation efficiency in biofilm-grown pneumococci using an approach with some similarity to that employed here (26). The authors made use of *in vitro* mixed biofilms containing spectinomycin-resistant (Spc^r) $\Delta comA$ ‘attacker cells’ — harboring concomitant deletions in the three autolysin genes (strain SPH149), or not (strain RH1) — and Nov^r non-competent ($\Delta comA \Delta comE$) ‘target cells’ (strain RH401). Upon the addition of CSP, the attackers (but not the targets, due to their being $\Delta comE$) acquired competence and were transformed by the DNA released from the target cells through the fratricidal killing caused by the lytic enzymes induced by the attackers. A *ca.* 40-fold reduction in transformation efficiency was observed in biofilms composed of SPH149 attackers (autolysin-deficient) and RH401 cells (0.009%) with respect to that seen in biofilms containing the same target cells but involving a lysis-proficient strain (RH1) as the attacker (0.34%). Moreover, a further near 40-fold reduction in gene transfer frequency (0.00026%) was seen when SPH149 attackers were incubated with $\Delta lytA \Delta lytC$ targets cells (strain SPH148) (26).

Target cells containing a $\Delta cbpD$ mutation were not tested, possibly because, as previously reported for mixed planktonic cultures, CbpD-proficient attacker cells (RH1) were 1000-fold more efficient in transformations involving *cbpD*⁺ target cells than were CbpD-deficient attacker cells (49).

Pneumococcal biofilms that form during nasopharyngeal colonization may provide an optimal environment for increased genetic exchange with enhanced natural transformation *in vivo* (3). The presence of eDNA in the biofilm matrix has generally been attributed to the autolysis of a subpopulation of cells via fratricidal killing, suicidal killing, and/or the controlled release of DNA via signal transduction (19). Autolysis-independent eDNA release has been documented in some Gram-positive bacteria including *Bacillus subtilis* (50), enterococci (45, 51), and staphylococci (52, 53). Taking the present results together, it is clear that mechanisms involved in active eDNA release (perhaps associated with the production of EV), other than those directly dependent on autolysins, are at work in *S. pneumoniae*. This is important since an important feature of biofilms is the development of chemical gradients (i.e., pH, redox potential, and ions) (reviewed in reference 28). For example, in *P. aeruginosa* biofilms the pH value towards the center of a microcolony (≈ 6.0) is lower than that at the edge of the biofilm or in the growth medium (≈ 6.8) (54). Although not directly tested, a similar situation might be relevant in pneumococcal biofilms because it is well known that *S. pneumoniae* autolysis is inhibited at low pH values (≤ 6.0) (55). In this case, autolysin-independent DNA release would allow the maintenance of horizontal gene transfer events within the biofilm. Although the amount of biologically active DNA released in the absence of detectable autolytic activity is limited, it appears to be sufficient to partially promote a biofilm lifestyle and, importantly, to allow horizontal gene transfer. It is also possible that, as suggested by the CLSM observations, a substantial amount of eDNA may be present in biofilms, although lacking most transforming activity. It is tempting to speculate that the DNA–protein complexes present in the ECM may hinder the

biological activity of eDNA as a transforming agent while contributing to the formation and preservation of the biofilm.

As reported for *Streptococcus mutans*, lysis-independent EV can transport DNA, contributing to horizontal gene transfer (56). Recent results also suggest that signal transduction mechanisms may be involved in the regulation of EV production in some Gram-positive bacteria (57, 58), but evidence for this is lacking in *S. pneumoniae*. As strongly suggested by the inhibitory effect of DNase I treatment in transforming efficiency, the biologically active eDNA released independent of detectable autolysis in *S. pneumoniae* appears to be located outside the EV. Although the loading of DNA into EV is thought to be widespread, experimental evidence showing that most EV-associated genomic DNA is present externally has been recently shown in *P. aeruginosa* (59, 60). However, other possibilities are also conceivable. For example, the stability of EV under various conditions may vary and a vesicle that loses membrane integrity may 'leak' its constituents into the supernatant (vesicle destabilization), rather than break up, as reported in *Bacillus anthracis*, *S. aureus* and other microorganisms (61). Interestingly, a very recent study has reported that the LytA NAM-amidase is not needed for production of EV, although the presence of EV-associated DNA was not analyzed (41).

An alternative (or complementary) mechanism for eDNA release may be bacterial type IV secretion systems (T4SS) that selectively deliver macromolecules to other cells or to the extracellular medium. An outstanding feature of these secretion systems is their ability to secrete both proteins and DNA molecules, a particularity that distinguishes them from other types of secretion system. The existence of a type IV secretion-like system involved in eDNA secretion was first described in *Neisseria gonorrhoeae* (62). More recently, it has been found that the release of eDNA from the cytoplasm of *H. influenzae* into the ECM requires the expression of an inner-membrane complex with homology to type IV secretion-like systems, plus the ComE outer-

membrane pore through which the type IV pilus is extruded (63). Type IV pili are surface-exposed fibers that mediate many functions in bacteria, including locomotion, adherence to eukaryotic cells, biofilm formation, DNA uptake (competence), and protein secretion. Although initially considered to be exclusive to Gram-negative bacteria, they are also present in Gram-positive, although their role(s) is just beginning to emerge (64). Recently, several studies have revealed the existence of type IV competence-induced pili — predominantly composed of the ComGC pilin (SPD_1861; SPD_RS09815) — on the surface of *S. pneumoniae* cells (65). Since these pili bind DNA, it has been proposed that the transformation pilus is the primary DNA receptor on the bacterial cell during transformation in *S. pneumoniae*. It is tempting to speculate that, during competence, intracellular DNA may also be secreted into the ECM with the participation of the type IV pili, perhaps involving the aqueous pore formed by ComEC (SPD_0844; SPD_RS0450) in the cytoplasmic membrane (66). Further studies are required to test this hypothesis.

MATERIALS AND METHODS

Strains, media and growth conditions. Table 3 lists the pneumococcal strains used in this study; all were grown in pH 8-adjusted C medium (CpH8) supplemented with 0.08% yeast extract (C+Y) medium, or not, as required (7). Cells were incubated at 37°C without shaking, and growth monitored by measuring absorbance at 550 nm (A_{550}). When used, antibiotics were added at the following concentrations: erythromycin 0.5 $\mu\text{g ml}^{-1}$, kanamycin 250 $\mu\text{g ml}^{-1}$, novobiocin 10 $\mu\text{g ml}^{-1}$, optochin 5 $\mu\text{g ml}^{-1}$, tetracycline 1 $\mu\text{g ml}^{-1}$, spectinomycin 100 $\mu\text{g ml}^{-1}$, and streptomycin 100 $\mu\text{g ml}^{-1}$. DNase I (from bovine pancreas, DN25) was purchased from Sigma-Aldrich. For the construction of mutants (Table 3), the appropriate *S. pneumoniae* strains were transformed with chromosomal or plasmid DNA in C medium supplemented with 0.08% bovine serum albumin after treating cells with 250 ng ml^{-1} synthetic CSP-1 at 37°C for

10 min to induce competence, followed by incubation at 30°C during DNA uptake.

Biofilm formation was determined by the ability of cells to adhere to the walls and base of 96-well (flat-bottomed) polystyrene microtiter dishes (Costar 3595; Corning Incorporated), using a modification of a previously reported protocol (67). Unless stated otherwise, cells grown in C+Y medium to an A_{550} of ≈ 0.5 – 0.6 , sedimented by centrifugation, resuspended in an equal volume of the indicated pre-warmed medium, diluted 1/10 or 1/100, and then dispensed at a concentration of 200 μ l per well. Plates were incubated at 34°C for 3, 4.5 and 5 h and bacterial growth determined by measuring the A_{595} using a VersaMax microplate absorbance reader (Molecular Devices). The biofilm formed was stained with 1% crystal violet (67).

Intrabiofilm gene transfer. Exponentially growing cultures of donor strains (see above) were seeded together in a 1:1 ratio in polystyrene microtiter plates and incubated in C+Y medium at 34°C. Previous results have indicated that biofilm-grown pneumococcal cells must be actively growing to become competent (26). Hence, biofilms formation was allowed for only 4.5 h; non-adherent cells were removed and adherent cells were washed with PBS, disaggregated by gentle pipetting and slow vortexing (4). The latter were then serially diluted and plated on blood agar plates containing Nov plus Str. Nov^r Str^r transformants were then picked from plates containing Opt to ascertain their parental strain. For each donor strain the total number of viable cells was determined using blood agar plates containing either Nov or Opt plus Str.

Preparation of extracellular vesicles and microscopical examination. EV-enriched centrifugation fractions were prepared from *S. pneumoniae* following standard procedures (61). Briefly, for biofilms, one liter of C+Y medium was inoculated with 10 ml of an exponentially growing culture of strain P271, distributed in 50 Petri dishes (10 cm diameter), and incubated at 34°C for 4.5 h. The non-adherent cells in the dishes were pipetted off and the biofilm-growing

cells ($6-7 \times 10^9$ CFU) suspended in 75 ml of fresh C+Y medium. After centrifugation ($9500 \times g$, 20 min, 4°C) the supernatant was filtered through a $0.2 \mu\text{m}$ pore-size filter (Millipore). The filtrate was centrifuged at $100,000 \times g$ for 1 h at 4°C to sediment the vesicular fraction into a pellet. The supernatant was then discarded and the pellet suspended in a small volume of C+Y medium (EV fraction) and stored in aliquots at -20°C . Aliquots of the EV fraction were also treated with DNase I ($10 \mu\text{g ml}^{-1}$) for 1 h at 37°C . EDTA (50 mM), SDS (1%) and proteinase K ($100 \mu\text{g ml}^{-1}$) were then added and the mixtures incubated for 2 h at 37°C . Extraction was performed with phenol, and precipitation with ethanol, following standard procedures. Finally, the pellet was dissolved in a small volume of 10 mM Tris-HCl, pH 8.0 containing 1 mM EDTA. Cultures of the same pneumococcal strain were also grown under planktonic conditions until late exponential phase ($\approx 3 \times 10^8$ CFU ml^{-1}) at 37°C without agitation, and processed as biofilm-grown pneumococci for EV preparation.

EV preparations (7 μl) were spotted on a glass slide, air dried, stained with the red fluorescence, membrane dye FM 4-64 (Molecular Probes), and incubated with α -dsDNA (ab27156, Abcam) followed by incubation with Alexa fluor 488-labeled goat anti-mouse IgG (A-11001, Invitrogen) (see below). Specimens were observed under a Leica DM4000B fluorescence microscope and viewed under a Leica HC PL APO63 $\times$ /1.40–0.60 oil objective. *S. pneumoniae* P271 biofilms formed on glass surfaces were prepared for low-temperature scanning electron microscopy (LTSEM) as previously described (7), and the samples observed at -135°C using a DSM 960 Zeiss scanning electron microscope. For transmission electron microscopy (TEM) analysis, 5 μl of EV were placed for 5 min at room temperature on carbon-reinforced, Formvar-coated copper grids (300 mesh), which had been rendered hydrophilic by glow-discharge using a Quorum GloQube apparatus (Quorum Technologies). The sample was quickly washed with ultrapure water and negative staining was performed using 1% sodium phosphotungstate for 5 min.

The excess stain was removed, and the sample was allowed to dry.

Micrographs were taken on a JEOL JEM 1230 working at 80 kV.

Quantification of eDNA. DNA quantification was performed by spectrophotometry (with concentrated samples) and/or using genetic transformation experiments (68). It is well known that a first-order concentration dependence (near unity) exists in chromosomal DNA transformation in pneumococci and other bacteria. This was confirmed in this study (see Fig. S1 in the supplemental material). It should be underlined that, since the size of the *S. pneumoniae* R6 genome is about 2 Mb, the DNA content of a single CFU, *i.e.*, a diplococcus, equals approximately 4 fg.

Microscopic observation of biofilms. For the observation of *S. pneumoniae* biofilms by CLSM, pneumococcal strains were grown on glass-bottomed dishes (WillCo-dish) for 3–5 h at 34°C as previously described (11). Following incubation, the culture medium was removed and the biofilm rinsed with phosphate-buffered saline (PBS) to remove non-adherent bacteria. The biofilms were then stained with DDAO (2 µM) (H6482, Invitrogen), α -dsDNA (at 2–25 µg ml⁻¹ each) and/or SYTO 59 (10 µM) (S11341, Invitrogen). All staining procedures involved incubation for 10–20 min at room temperature in the dark, except when biofilms were incubated with mouse α -dsDNA (2 µg ml⁻¹); this involved a fixation step at room temperature with 3% paraformaldehyde for 10 min. The biofilms were then rinsed with 0.5 ml PBS and incubated for 1 h at 4°C followed by 30 min incubation at room temperature in the dark with Alexa fluor 488-labelled goat anti-mouse IgG (1:500). After staining, the biofilms were gently rinsed with 0.5 ml PBS. Observations were made at a magnification of 63× using a Leica TCS-SP2-AOBS-UV or TCS-SP5-AOBS-UV CLSM equipped with an argon ion laser. Laser intensity and gain were kept the same for all images. Images were analyzed using LCS software from LEICA. Projections were obtained in the x–y (individual scans at 0.5–1 µm intervals) and x–z (images at 5–6 µm intervals) planes.

Statistical analysis. Data comparisons were performed using the two-tailed Student *t*-test.

ACKNOWLEDGMENTS

We thank J.-P. Claverys and J. Vidal for kindly providing pneumococcal strains, M. Moscoso, P. García and J. Yuste for helpful comments and critical reading of the manuscript, A. Burton for revising the English version, M. T. Seisdedos and G. Elvira for their help with CLSM, and E. Cano and S. Ruiz for skillful technical assistance.

This research was supported by grant SAF2017-88664 from Ministerio de Economía, Industria y Competitividad (MEICOM). Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES) is an initiative of the Instituto de Salud Carlos III (ISCIII).

REFERENCES

1. Weiser JN. 2010. The pneumococcus: why a commensal misbehaves. *J Mol Med* 88:97-102. <https://doi.org/10.1007/s00109-009-0557-x>.
2. Domenech M, García E, Moscoso M. 2012. Biofilm formation in *Streptococcus pneumoniae*. *Microb Biotechnol* 5:455-465. <https://doi.org/10.1111/j.1751-7915.2011.00294.x>.
3. Chao Y, Marks LR, Pettigrew MM, Hakansson AP. 2015. *Streptococcus pneumoniae* biofilm formation and dispersion during colonization and disease. *Front Cell Infect Microbiol* 4:194. <https://doi.org/10.3389/fcimb.2014.00194>.
4. Domenech M, Ramos-Sevillano E, García E, Moscoso M, Yuste J. 2013. Biofilm formation avoids complement immunity and phagocytosis of *Streptococcus pneumoniae*. *Infect Immun* 81:2606-2615. <https://doi.org/10.1128/IAI.00491-13>.

5. Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappin-Scott HM. 1995. Microbial biofilms. *Annu Rev Microbiol* 49:711-745. <https://doi.org/10.1146/annurev.mi.49.100195.003431>.
6. Okshevsky M, Meyer RL. 2015. The role of extracellular DNA in the establishment, maintenance and perpetuation of bacterial biofilms. *Crit Rev Microbiol* 41:341-352. <https://doi.org/10.3109/1040841X.2013.841639>.
7. Moscoso M, García E, López R. 2006. Biofilm formation by *Streptococcus pneumoniae*: role of choline, extracellular DNA, and capsular polysaccharide in microbial accretion. *J Bacteriol* 188:7785-7795. <https://doi.org/10.1128/JB.00673-06>.
8. Hall-Stoodley L, Nistico L, Sambanthamoorthy K, Dice B, Nguyen D, Mershon WJ, Johnson C, Hu FZ, Stoodley P, Ehrlich GD, Post JC. 2008. Characterization of biofilm matrix, degradation by DNase treatment and evidence of capsule downregulation in *Streptococcus pneumoniae* clinical isolates. *BMC Microbiol* 8:173. <https://doi.org/10.1186/1471-2180-8-173>.
9. Carrolo M, Frias MJ, Pinto FR, Melo-Cristino J, Ramirez M. 2010. Prophage spontaneous activation promotes DNA release enhancing biofilm formation in *Streptococcus pneumoniae*. *PLoS One* 5:e15678. <https://doi.org/10.1371/journal.pone.0015678>.
10. Trappetti C, Gualdi L, Di Meola L, Jain P, Korir C, Edmonds P, Iannelli F, Ricci S, Pozzi G, Oggioni M. 2011. The impact of the competence quorum sensing system on *Streptococcus pneumoniae* biofilms varies depending on the experimental model. *BMC Microbiol* 11:75. <https://doi.org/10.1186/1471-2180-11-75>.
11. Domenech M, García E, Prieto A, Moscoso M. 2013. Insight into the composition of the intercellular matrix of *Streptococcus pneumoniae* biofilms. *Environ Microbiol* 15:502-516. <https://doi.org/10.1111/j.1462-2920.2012.02853.x>.

12. Domenech M, Ruiz S, Moscoso M, García E. 2015. *In vitro* biofilm development of *Streptococcus pneumoniae* and formation of choline-binding protein–DNA complexes. *Environ Microbiol Rep* 7:715-727. <https://doi.org/10.1111/1758-2229.12295>.
13. Schulte T, Mikaelsson C, Beaussart A, Kikhney A, Deshmukh M, Wolniak S, Pathak A, Ebel C, Löfling J, Fogolari F, Henriques-Normark B, Dufrêne YF, Svergun D, Nygren P-Å, Achour A. 2016. The BR domain of PsrP interacts with extracellular DNA to promote bacterial aggregation; structural insights into pneumococcal biofilm formation. *Sci Rep* 6:32371. <https://doi.org/10.1038/srep32371>.
14. Arenas J, Tommassen J. 2017. Meningococcal biofilm formation: let's stick together. *Trends Microbiol* 25:113-124. <https://doi.org/10.1016/j.tim.2016.09.005>.
15. Höltje JV, Tomasz A. 1975. Biological effects of lipoteichoic acids. *J Bacteriol* 124:1023-1027.
16. Sandalova T, Lee M, Henriques-Normark B, Hesek D, Mobashery S, Mellroth P, Achour A. 2016. The crystal structure of the major pneumococcal autolysin LytA in complex with a large peptidoglycan fragment reveals the pivotal role of glycans for lytic activity. *Mol Microbiol* 101:954-967. <https://doi.org/10.1111/mmi.13435>.
17. García P, González MP, García E, García JL, López R. 1999. The molecular characterization of the first autolytic lysozyme of *Streptococcus pneumoniae* reveals evolutionary mobile domains. *Mol Microbiol* 33:128-138. <https://doi.org/10.1046/j.1365-2958.1999.01455.x>.
18. Molina R, González A, Stelter M, Pérez-Dorado I, Kahn R, Morales M, Moscoso M, Campuzano S, Campillo NE, Mobashery S, García JL, García P, Hermoso JA. 2009. Crystal structure of CbpF, a bifunctional choline-binding protein and autolysis regulator from *Streptococcus pneumoniae*. *EMBO Rep* 10:246-251. <https://doi.org/10.1038/embo.2008.245>.

19. Ibáñez de Aldecoa AL, Zafra O, González-Pastor JE. 2017. Mechanisms and regulation of extracellular DNA release and its biological roles in microbial communities. *Front Microbiol* 8:1390. <https://doi.org/10.3389/fmicb.2017.01390>.
20. Ottolenghi E, Hotchkiss RD. 1962. Release of genetic transforming agent from pneumococcal cultures during growth and disintegration. *J Exp Med* 116:491-519. <https://doi.org/10.1084/jem.116.4.491>.
21. Moscoso M, Claverys J-P. 2004. Release of DNA into the medium by competent *Streptococcus pneumoniae*: kinetics, mechanism and stability of the liberated DNA. *Mol Microbiol* 54:783-794. <https://doi.org/10.1111/j.1365-2958.2004.04305.x>.
22. Kausmally L, Johnsborg O, Lunde M, Knutsen E, Håvarstein LS. 2005. Choline-binding protein D (CbpD) in *Streptococcus pneumoniae* is essential for competence-induced cell lysis. *J Bacteriol* 187:4338-4345. <https://doi.org/10.1128/JB.187.13.4338-4345.2005>.
23. Claverys J-P, Håvarstein LS. 2007. Cannibalism and fratricide: mechanisms and raisons d'être. *Nat Rev Microbiol* 5:219-229. <https://doi.org/10.1038/nrmicro1613>.
24. Shanker E, Federle JM. 2017. Quorum sensing regulation of competence and bacteriocins in *Streptococcus pneumoniae* and *mutans*. *Genes* 8:15. <https://doi.org/10.3390/genes8010015>.
25. Guiral S, Mitchell TJ, Martin B, Claverys J-P. 2005. Competence-programmed predation of noncompetent cells in the human pathogen *Streptococcus pneumoniae*: genetic requirements. *Proc Natl Acad Sci U S A* 102:8710-8715. <https://doi.org/10.1073/pnas.0500879102>.
26. Wei H, Håvarstein LS. 2012. Fratricide is essential for efficient gene transfer between pneumococci in biofilms. *Appl Environ Microbiol* 78:5897-5905. <https://doi.org/10.1128/aem.01343-12>.

27. Trappetti C, Potter AJ, Paton AW, Oggioni MR, Paton JC. 2011. LuxS mediates iron-dependent biofilm formation, competence and fratricide in *Streptococcus pneumoniae*. Infect Immun 79:4550-4558. <https://doi.org/10.1128/iai.05644-11>.
28. Schlafer S, Meyer RL. 2017. Confocal microscopy imaging of the biofilm matrix. J Microbiol Methods 138:50-59. <https://doi.org/10.1016/j.mimet.2016.03.002>.
29. Okshevsky M, Meyer RL. 2014. Evaluation of fluorescent stains for visualizing extracellular DNA in biofilms. J Microbiol Methods 105:102-104. <https://doi.org/10.1016/j.mimet.2014.07.010>.
30. Andre GO, Converso TR, Politano WR, Ferraz LFC, Ribeiro ML, Leite LCC, Darrieux M. 2017. Role of *Streptococcus pneumoniae* proteins in evasion of complement-mediated immunity. Front Microbiol 8:224. <https://doi.org/10.3389/fmicb.2017.00224>.
31. López R, García E. 2004. Recent trends on the molecular biology of pneumococcal capsules, lytic enzymes, and bacteriophage. FEMS Microbiol Rev 28:553-580. <https://doi.org/10.1016/j.femsre.2004.05.002>.
32. Steinmoen H, Knutsen E, Håvarstein LS. 2002. Induction of natural competence in *Streptococcus pneumoniae* triggers lysis and DNA release from a subfraction of the cell population. Proc Natl Acad Sci U S A 99:7681-7686. <https://doi.org/10.1073/pnas.112464599>.
33. Moscoso M, Domenech M, García E. 2010. Vancomycin tolerance in clinical and laboratory *Streptococcus pneumoniae* isolates depends on reduced enzyme activity of the major LytA autolysin or cooperation between CiaH histidine kinase and capsular polysaccharide. Mol Microbiol 77:1052-1064. <https://doi.org/10.1111/j.1365-2958.2010.07271.x>.
34. Johnston C, Hauser C, Hermans PWM, Martin B, Polard P, Bootsma HJ, Claverys J-P. 2016. Fine-tuning of choline metabolism is important for

pneumococcal colonization. Mol Microbiol 100:972-988.

<https://doi.org/10.1111/mmi.13360>.

35. Laux A, Sexauer A, Sivaselvarajah D, Kaysen A, Brückner R. 2015. Control of competence by related non-coding csRNAs in *Streptococcus pneumoniae* R6. Front Genet 6:246.

<https://doi.org/10.3389/fgene.2015.00246>.

36. Muñoz R, Bustamante M, de la Campa AG. 1995. Ser-127-to-Leu substitution in the DNA gyrase B subunit of *Streptococcus pneumoniae* is implicated in novobiocin resistance. J Bacteriol 177:4166-4170.

<https://doi.org/10.1128/jb.177.14.4166-4170.1995>.

37. Salles C, Créancier L, Claverys J-P, Méjean V. 1992. The high level streptomycin resistance gene from *Streptococcus pneumoniae* is a homologue of the ribosomal protein S12 gene from *Escherichia coli*.

Nucleic Acids Res 20:6103. <https://doi.org/10.1093/nar/20.22.6103>.

38. Claverys J-P, Lacks SA. 1986. Heteroduplex deoxyribonucleic acid base mismatch repair in bacteria. Microbiol Rev 50:133-165.

39. Olaya-Abril A, Prados-Rosales R, McConnell MJ, Martín-Peña R, González-Reyes JA, Jiménez-Munguía I, Gómez-Gascón L, Fernández J, Luque-García JL, García-Lidón C, Estévez H, Pachón J, Obando I, Casadevall A, Pirofski L-a, Rodríguez-Ortega MJ. 2014. Characterization of protective extracellular membrane-derived vesicles produced by *Streptococcus pneumoniae*. J Proteomics 106:46-60.

<https://doi.org/10.1016/j.jprot.2014.04.023>.

40. Volgers C, Benedikter BJ, Grauls GE, Hellebrand PHM, Savelkoul PHM, Stassen FRM. 2017. Effects of N-acetyl-L-cysteine on the membrane vesicle release and growth of respiratory pathogens. FEMS Microbiol Lett 364:fnx087. <https://doi.org/10.1093/femsle/fnx087>.

41. Codemo M, Muschiol S, Iovino F, Nannapaneni P, Plant L, Wai SN, Henriques-Normark B. 2018. Immunomodulatory effects of pneumococcal

- extracellular vesicles on cellular and humoral host defenses. mBio
9:e00559-00518. <https://doi.org/10.1128/mBio.00559-18>.
42. Whitchurch CB, Tolker-Nielsen T, Ragas PC, Mattick JS. 2002.
Extracellular DNA required for bacterial biofilm formation. Science
295:1487. <https://doi.org/10.1126/science.295.5559.1487>.
43. Yang CL, Chilvers M, Montgomery M, Nolan SJ. 2017. Dornase alfa for
cystic fibrosis. Paediatr Respir Rev 21:65-67.
<https://doi.org/10.1016/j.prrv.2016.09.001>.
44. Magana M, Sereti C, Ioannidis A, Mitchell CA, Ball AR, Magiorkinis E,
Chatzipanagiotou S, Hamblin MR, Hadjifrangiskou M, Tegos GP. 2018.
Options and limitations in clinical investigation of bacterial biofilms. Clin
Microbiol Rev 31:e00084-00016. <https://doi.org/10.1128/cmr.00084-16>.
45. Barnes AMT, Ballering KS, Leibman RS, Wells CL, Dunne GM. 2012.
Enterococcus faecalis produces abundant extracellular structures
containing DNA in the absence of cell lysis during early biofilm formation.
mBio 3:e00193-00112. <https://doi.org/10.1128/mBio.00193-12>.
46. Domenech M, Pedrero-Vega E, Prieto A, García E. 2016. Evidence of the
presence of nucleic acids and β -glucan in the matrix of non-typeable
Haemophilus influenzae in vitro biofilms. Sci Rep 6:36424.
<https://doi.org/10.1038/srep36424>.
47. Sugimoto S, Okuda K-i, Miyakawa R, Sato M, Arita-Morioka K-i, Chiba A,
Yamanaka K, Ogura T, Mizunoe Y, Sato C. 2016. Imaging of bacterial
multicellular behaviour in biofilms in liquid by atmospheric scanning electron
microscopy. Sci Rep 6:25889. <https://doi.org/10.1038/srep25889>.
48. Okuda K-i, Nagahori R, Yamada S, Sugimoto S, Sato C, Sato M, Iwase T,
Hashimoto K, Mizunoe Y. 2018. The composition and structure of biofilms
developed by *Propionibacterium acnes* isolated from cardiac pacemaker
devices. Front Microbiol 9:182. <https://doi.org/10.3389/fmicb.2018.00182>.

49. Johnsborg O, Eldholm V, Bjørnstad ML, Håvarstein LS. 2008. A predatory mechanism dramatically increases the efficiency of lateral gene transfer in *Streptococcus pneumoniae* and related commensal species. Mol Microbiol 69:245-253. <https://doi.org/10.1111/j.1365-2958.2008.06288.x>.
50. Zafra O, Lamprecht-Grandío M, González de Figueras C, González-Pastor JE. 2012. Extracellular DNA release by undomesticated *Bacillus subtilis* is regulated by early competence. PLoS One 7:e48716. <https://doi.org/10.1371/journal.pone.0048716>.
51. Paganelli FL, Willems RJL, Jansen P, Hendrickx A, Zhang X, Bonten MJM, Leavis HL. 2013. *Enterococcus faecium* biofilm formation: identification of major autolysin AtlA_{Efm}, associated Acm surface localization, and AtlA_{Efm}-independent extracellular DNA release. mBio 4:e00154. <https://doi.org/10.1128/mBio.00154-13>.
52. Rajendran NB, Eikmeier J, Becker K, Hussain M, Peters G, Heilmann C. 2015. Important contribution of the novel locus *comEB* to extracellular DNA-dependent *Staphylococcus lugdunensis* biofilm formation. Infect Immun 83:4682-4692. <https://doi.org/10.1128/iai.00775-15>.
53. DeFrancesco AS, Masloboeva N, Syed AK, DeLoughery A, Bradshaw N, Li G-W, Gilmore MS, Walker S, Losick R. 2017. Genome-wide screen for genes involved in eDNA release during biofilm formation by *Staphylococcus aureus*. Proc Natl Acad Sci U S A 114:E5969-E5978. <https://doi.org/10.1073/pnas.1704544114>.
54. Hunter RC, Beveridge TJ. 2005. Application of a pH-sensitive fluoroprobe (C-SNARF-4) for pH microenvironment analysis in *Pseudomonas aeruginosa* biofilms. Appl Environ Microbiol 71:2501-2510. <https://doi.org/10.1128/aem.71.5.2501-2510.2005>.
55. Ronda-Lain C, Lopez R, Tapia A, Tomasz A. 1977. Role of the pneumococcal autolysin (murein hydrolase) in the release of phage

- progeny bacteriophage and in the bacteriophage-induced lysis of the host cells. J Virol 21:366-374.
56. Liao S, Klein MI, Heim KP, Fan Y, Bitoun JP, Ahn S-J, Burne RA, Koo H, Brady LJ, Wen ZT. 2014. *Streptococcus mutans* extracellular DNA is upregulated during growth in biofilms, actively released via membrane vesicles, and influenced by components of the protein secretion machinery. J Bacteriol 196:2355-2366. <https://doi.org/10.1128/jb.01493-14>.
57. Resch U, Tsatsaronis JA, Le Rhun A, Stübiger G, Rohde M, Kasvandik S, Holzmeister S, Tinnefeld P, Wai SN, Charpentier E. 2016. A two-component regulatory system impacts extracellular membrane-derived vesicle production in group A *Streptococcus*. mBio 7:e00207-00216. <https://doi.org/10.1128/mBio.00207-16>.
58. Obana N, Nakao R, Nagayama K, Nakamura K, Senpuku H, Nomura N, Bäumlér AJ. 2017. Immunoactive clostridial membrane vesicle production is regulated by a sporulation factor. Infect Immun 85:e00096-00017. <https://doi.org/10.1128/iai.00096-17>.
59. Renelli M, Matias V, Lo RY, Beveridge TJ. 2004. DNA-containing membrane vesicles of *Pseudomonas aeruginosa* PAO1 and their genetic transformation potential. Microbiology 150:2161-2169. <https://doi.org/10.1099/mic.0.26841-0>.
60. Bitto NJ, Chapman R, Pidot S, Costin A, Lo C, Choi J, D'Cruze T, Reynolds EC, Dashper SG, Turnbull L, Whitchurch CB, Stinear TP, Stacey KJ, Ferrero RL. 2017. Bacterial membrane vesicles transport their DNA cargo into host cells. Sci Rep 7:7072. <https://doi.org/10.1038/s41598-017-07288-4>.
61. Brown L, Wolf JM, Prados-Rosales R, Casadevall A. 2015. Through the wall: extracellular vesicles in Gram-positive bacteria, mycobacteria and fungi. Nat Rev Microbiol 13:620-630. <https://doi.org/10.1038/nrmicro3480>.

62. Hamilton HL, Domínguez NM, Schwartz KJ, Hackett KT, Dillard JP. 2005. *Neisseria gonorrhoeae* secretes chromosomal DNA via a novel type IV secretion system. Mol Microbiol 55:1704-1721. <https://doi.org/10.1111/j.1365-2958.2005.04521.x>.
63. Jurcisek JA, Brockman KL, Novotny LA, Goodman SD, Bakaletz LO. 2017. Nontypeable *Haemophilus influenzae* releases DNA and DNABII proteins via a T4SS-like complex and ComE of the type IV pilus machinery. Proc Natl Acad Sci U S A 114:E6632-E6641. <https://doi.org/10.1073/pnas.1705508114>.
64. Melville S, Craig L. 2013. Type IV pili in Gram-positive bacteria. Microbiol Mol Biol Rev 77:323-341. <https://doi.org/10.1128/mmb.00063-12>.
65. Laurenceau R, Péhau-Arnaudet G, Baconnais S, Gault J, Malosse C, Dujeancourt A, Campo N, Chamot-Rooke J, Le Cam E, Claverys J-P, Fronzes R. 2013. A type IV pilus mediates DNA binding during natural transformation in *Streptococcus pneumoniae*. PLoS Pathog 9:e1003473. <https://doi.org/10.1371/journal.ppat.1003473>.
66. Bergé M, Moscoso M, Prudhomme M, Martin B, Claverys J-P. 2002. Uptake of transforming DNA in Gram-positive bacteria: a view from *Streptococcus pneumoniae*. Mol Microbiol 45:411-421. <https://doi.org/10.1046/j.1365-2958.2002.03013.x>.
67. Domenech M, Araújo-Bazán L, García E, Moscoso M. 2014. *In vitro* biofilm formation by *Streptococcus pneumoniae* as a predictor of post-vaccination emerging serotypes colonizing the human nasopharynx. Environ Microbiol 16:1193-1201. <https://doi.org/10.1111/1462-2920.12370>.
68. Hotchkiss RD. 1957. Methods for characterization of nucleic acid. Methods Enzymol 3:708-715. [https://doi.org/10.1016/S0076-6879\(57\)03448-5](https://doi.org/10.1016/S0076-6879(57)03448-5).
69. Hoskins J, Alborn WE, Jr., Arnold J, Blaszcak LC, Burgett S, DeHoff BS, Estrem ST, Fritz L, Fu D-J, Fuller W, Geringer C, Gilmour R, Glass JS, Khoje H, Kraft AR, Lagace RE, LeBlanc DJ, Lee LN, Lefkowitz EJ, Lu J,

- Matsushima P, McAhren SM, McHenney M, McLeaster K, Mundy CW, Nicas TI, Norris FH, O'gara M, Peery RB, Robertson GT, Rockey P, Sun P-M, Winkler ME, Yang Y, Young-Bellido M, Zhao G, Zook CA, Baltz RH, Jaskunas R, Rosteck PRJ, Skatrud PL, Glass JI. 2001. Genome of the bacterium *Streptococcus pneumoniae* strain R6. J Bacteriol 183:5709-5717. <https://doi.org/10.1128/JB.183.19.5709-5717.2001>.
70. Martin B, Prats H, Claverys J-P. 1985. Cloning of the *hexA* mismatch-repair gene of *Streptococcus pneumoniae* and identification of the product. Gene 34:293-303. [https://doi.org/10.1016/0378-1119\(85\)90138-6](https://doi.org/10.1016/0378-1119(85)90138-6).
71. Ronda C, García JL, López R. 1988. Characterization of genetic transformation in *Streptococcus oralis* NCTC 11427: expression of the pneumococcal amidase in *S. oralis* using a new shuttle vector. Mol Gen Genet 215:53-57. <https://doi.org/10.1007/BF00331302>.
72. Fenoll A, Muñoz R, García E, de la Campa AG. 1994. Molecular basis of the optochin-sensitive phenotype of pneumococcus: characterization of the genes encoding the F₀ complex of the *Streptococcus pneumoniae* and *Streptococcus oralis* H⁺-ATPases. Mol Microbiol 12:587-598. <https://doi.org/10.1111/j.1365-2958.1994.tb01045.x>.
73. Vidal JE, Ludewick HP, Kunkel RM, Zähler D, Klugman KP. 2011. The LuxS-dependent quorum-sensing system regulates early biofilm formation by *Streptococcus pneumoniae* strain D39. Infect Immun 79:4050-4060. <https://doi.org/10.1128/iai.05186-11>.
74. Vidal JE, Howery KE, Ludewick HP, Nava P, Klugman KP. 2013. Quorum-sensing systems LuxS/autoinducer 2 and Com regulate *Streptococcus pneumoniae* biofilms in a bioreactor with living cultures of human respiratory cells. Infect Immun 81:1341-1353. <https://doi.org/10.1128/iai.01096-12>.

TABLE 1 Transformation efficiency in *S. pneumoniae* biofilms^a

Strain		Transformation frequency ^b	
		Str ^r	Nov ^r
P233 (Nov ^r)	P273 (Opt ^r Str ^r)	0.45 ± 0.10	0.04 ± 0.02
P272 (<i>lytA lytC</i> <i>cbpD</i> Nov ^r)	P271 (<i>lytA lytC</i> <i>cbpD</i> Opt ^r Str ^r)	$0.27 \times 10^{-2} \pm 0.05 \times 10^{-2}$	$0.30 \times 10^{-3} \pm 0.19 \times 10^{-3}$

^aExponentially growing cultures of the indicated donor (and recipient) strains were mixed in polystyrene microtiter plates and incubated in C+Y medium at 34°C. After 4.5 h incubation, non-adherent cells were removed and biofilm-grown cells were washed and resuspended with PBS, diluted, and plated on blood agar plates containing Nov plus Str. Nov^r Str^r transformants were then picked from plates containing Opt to ascertain their parental strain. Total viable cells of each strain were determined using plates containing either Nov or Opt plus Str, and ranged from 1.5×10^8 to 1.8×10^8 for P233 + P273 biofilms, and from 3.2×10^8 to 3.4×10^8 CFU ml⁻¹ for P272 + P271 biofilms.

^bTransformation frequency is defined as the number of transformants (CFU ml⁻¹) multiplied by 100, divided by the total number of bacteria (CFU ml⁻¹). Values correspond to the means ± standard errors for four independent experiments.

TABLE 2 Transforming eDNA in pneumococcal biofilms and planktonic cultures^a

DNA origin	Total DNA content (μg)	Cell equivalents (%)
Biofilm		
Intracellular	65	1.6×10^{10} (100)
EV-associated	7.5×10^{-5}	1.9×10^4 (0.00012)
EV-associated + DNase I	ND	–
Planktonic culture		
Intracellular	3×10^3	7.5×10^{11} (100)
EV-associated	1.2×10^{-2}	3×10^6 (0.0004)
EV-associated + DNase I	ND	–

^aEV were prepared from biofilms or planktonic cultures of strain P271 (*lytA lytC cbpD* Opt^f; Str^r) and DNA was purified as described under Materials and Methods. Total chromosomal DNA (intracellular) was also purified. Competent R6 cells were used as recipients in transformation experiments. Transformants were selected using streptomycin-containing plates. Results are the means for three independent determinations. ND, not detected.

848 **TABLE 3** *S. pneumoniae* strains used

Strain	Description ^a	Source or reference
R6	Nonencapsulated D39 derivative; <i>lytA</i> ⁺	69
R6BC	R6 <i>lytB::ermC lytC::tet</i> ; Ery ^r Tet ^r	7
R800	R6 derivative	70
M22	A laboratory multi-resistant strain; <i>nov1 str41</i> ; Nov ^r ; Str ^r	71
M222	M22 derivative; M22 transformed with <i>Streptococcus oralis</i> NCTC 11427 chromosomal DNA; Opt ^r ; Str ^r	72
SPJV05	D39 derivative <i>luxS</i> null mutant, Ery ^r	73
SPJV10	D39 derivative <i>comC</i> null mutant, Ery ^r	74
P042	R800 but <i>lytA::aphIII lytC::tet</i> ; Kan ^r Tet ^r	7
P046	R800 but <i>lytA::aphIII lytC::ermC</i> ; Kan ^r Ery ^r	7
P104	R6 but <i>lytA::cat</i> ; Cm ^r	This study
P147	R6 <i>ciaH</i> _{Tupelo_VT}	33
P203	R391 but <i>cbpD::spc</i> ; R391 transformed with R1582 chromosomal DNA; Spc ^r	This study
P204	P042 but <i>cbpD::spc</i> ; P042 transformed with R1582 chromosomal DNA; Spc ^r	This study
P211	P104 but <i>comA::kan</i> ; P104 transformed with R391 chromosomal DNA; Kan ^r	This study
P213	P211 but <i>lytC</i> ; P211 transformed with R6BC chromosomal DNA; Tet ^r	This study
P233	R6 transformed with M22 chromosomal DNA; Nov ^r	This study
P234	R6 <i>pspC</i> ; spontaneous mutant ^b	This study
P235	R6 but <i>luxS</i> null mutant; R6 transformed with SPJV05 chromosomal DNA; Ery ^r	This study

P236	R6 but <i>comC</i> null mutant; R6 transformed with SPJV10 chromosomal DNA; Ery ^r	This study
P271	P204 transformed with M222 chromosomal DNA; Opt ^r ; Str ^r	This study
P272	P204 transformed with M222 chromosomal DNA; Nov ^r	This study
P273	R6 transformed with M222 chromosomal DNA; Opt ^r ; Str ^r	This study
R391	R800 but <i>comA::kan</i> ; Kan ^r	25
R1582	R800 but <i>cbpD::spc</i> ; Spc ^r	25

849 ^aCm, chloramphenicol; Ery, erythromycin; Kan, kanamycin; Nov, novobiocin;
850 Opt, optochin; Tet, tetracycline; Spc, spectinomycin; Str, streptomycin; VT,
851 vancomycin tolerance. ^r, resistant.

852 ^bThe *pspC* mutation corresponds to a single base substitution (G:C to A:T)
853 converting Trp (TGG) at position 512 to a stop codon (TAG). The wild type
854 *pspC* gene encodes a 701 amino acid residue-long protein.

855

856

FIGURE LEGENDS

FIG 1 Evidence of the existence of eDNA in pneumococcal biofilms using CLSM. A biofilm of *S. pneumoniae* R6 grown for 5 h at 34°C in C+Y medium was stained with a combination of SYTO 59 (A, red), DDAO (B, blue), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (C, green). Image D is a merge of channels A and B. Image E is a merge of channels A and C. Image F is a merge of channels B and C. Image G is a merge of the three channels. Images H, I and J correspond, respectively, to an enlarged vision of the area marked with squares in D, E and F. Yellow arrows point to eDNA stained only with DDAO or α -dsDNA–Alexa Fluor-488. White arrows indicate the location of SYTO 59-stained bacteria together with eDNA labelled either with DDAO (H, magenta) or with α -dsDNA–Alexa Fluor-488 (I, yellow). The red arrow in image J points to doubly labelled (DDAO plus α -dsDNA–Alexa Fluor-488) eDNA (J, light blue). Scale bars = 25 μ m.

FIG 2 Dynamics of eDNA release in pneumococcal biofilms. *S. pneumoniae* R6 was grown under biofilm-forming conditions. After 3 and 5 h incubation at 34°C, adherent (biofilm) and non-adherent (planktonic) cells were independently stained with a combination of SYTO 59 (red) and α -dsDNA–Alexa Fluor-488 (green). Scale bars = 25 μ m.

FIG 3 Inhibition and dispersal of pneumococcal biofilms with DNase I. (A) The indicated *S. pneumoniae* strains were grown overnight at 37°C to an A_{550} value of 0.5 (corresponding to the late exponential phase of growth) in C+Y medium, centrifuged, and adjusted to an A_{550} of 0.6 with fresh medium. The cell suspensions were then diluted 100-fold, and 200 μ l aliquots distributed into the wells of microtiter plates, which were then incubated for 5 h at 34°C (open bars). Other samples received DNase I (100 μ g ml⁻¹) (red bars) and were incubated as above (inhibition assay). In other cases, and after biofilm development (4 h at 34°C), DNase I (green bars) was added at 100 μ g ml⁻¹,

and incubation allowed to proceed for an additional 1 h at 34°C before staining with CV to quantify biofilm formation (dispersal assay). In all assays, black bars indicate growth (adherent plus non-adherent cells). *, $P < 0.001$ compared with the corresponding, untreated control.

FIG 4 Influence of autolysins on biofilm formation and eDNA release revealed by CLSM. Biofilms were stained with a combination of SYTO 59 (red), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (green). The R6 strain was also incubated in C+Y medium containing 2% choline chloride (R6 + 2% cho). Horizontal and vertical three-dimensional reconstructions of 55 (x - y plane) or 65 scans (x - z plane) are shown. Scale bars = 25 μ m.

FIG 5 Influence of competence induction on biofilm formation and eDNA release revealed by CLSM. Biofilms were stained with a combination of SYTO 59 (red), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (green). Scale bars = 25 μ m.

Supplemental Material

Additional information may be found in the online version of this article:

FIG S1 Calibration curve for *S. pneumoniae*-transforming DNA. Competent R6 cells were used as recipient bacteria. Values represent the means $\pm$ standard errors for three independent transformation experiments. The dotted line corresponds to a slope of 1.

FIG S2 Observation of EV produced by *S. pneumoniae* P271. (A–C) Fluorescent labeling of an EV preparation with FM 4-64 (red; A) and α -dsDNA, followed by incubation with Alexa fluor 488-labeled goat anti-mouse IgG (green; B). Image C is a merger of the two channels. Scale bars = 10 μ m. (D–E) Electron micrographs of EVs. (D) LTSEM image of a pneumococcal biofilm. Arrows point to spherical blebs protruding from cells. The arrowhead indicates a putatively cell extruded EV. (E, F) TEM micrographs showing negatively stained EV. In some cases, EV appear to coalesce and fuse (F).

FIG 1 Evidence of the existence of eDNA in pneumococcal biofilms using CLSM. A biofilm of *S. pneumoniae* R6 grown for 5 h at 34°C in C+Y medium was stained with a combination of SYTO 59 (A, red), DDAO (B, blue), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (C, green). Image D is a merge of channels A and B. Image E is a merge of channels A and C. Image F is a merge of channels B and C. Image G is a merge of the three channels. Images H, I and J correspond, respectively, to an enlarged vision of the area marked with squares in D, E and F. Yellow arrows point to eDNA stained only with DDAO or α -dsDNA–Alexa Fluor-488. White arrows indicate the location of SYTO 59-stained bacteria together with eDNA labelled either with DDAO (H, magenta) or with α -dsDNA–Alexa Fluor-488 (I, yellow). The red arrow in image J points to doubly labelled (DDAO plus α -dsDNA–Alexa Fluor-488) eDNA (J, light blue). Scale bars = 25 μ m.

FIG 2 Dynamics of eDNA release in pneumococcal biofilms. *S. pneumoniae* R6 was grown under biofilm-forming conditions. After 3 and 5 h incubation at 34°C, adherent (biofilm) and non-adherent (planktonic) cells were independently stained with a combination of SYTO 59 (red) and α -dsDNA–Alexa Fluor-488 (green). Scale bars = 25 μ m.

FIG 3 Inhibition and dispersal of pneumococcal biofilms with DNase I. The indicated *S. pneumoniae* strains were grown overnight at 37°C to an A_{550} value of 0.5 (corresponding to the late exponential phase of growth) in C+Y medium, centrifuged, and adjusted to an A_{550} of 0.6 with fresh medium. The cell suspensions were then diluted 100-fold, and 200 μ l aliquots distributed into the wells of microtiter plates, which were then incubated for 5 h at 34°C (open bars). Other samples received DNase I (100 μ g ml⁻¹) (red bars) and were incubated as above (inhibition assay). In other cases, and after biofilm development (4 h at 34°C), DNase I (green bars) was added at 100 μ g ml⁻¹, and incubation allowed to proceed for an additional 1 h at 34°C before staining with CV to quantify biofilm formation (dispersal assay). In all assays, black bars indicate growth (adherent plus non-adherent cells). *, $P < 0.001$ compared with the corresponding, untreated control.

FIG 4 Influence of autolysins on biofilm formation and eDNA release revealed by CLSM. Biofilms were stained with a combination of SYTO 59 (red), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (green). The R6 strain was also incubated in C+Y medium containing 2% choline chloride (R6 + 2% cho). Horizontal and vertical three-dimensional reconstructions of 55 (x - y plane) or 65 scans (x - z plane) are shown. Scale bars = 25 μ m.

FIG 5 Influence of competence induction on biofilm formation and eDNA release revealed by CLSM. Biofilms were stained with a combination of SYTO 59 (red), and α -dsDNA, followed by Alexa Fluor-488 goat anti-mouse IgG (green). Scale bars = 25 μ m.