Deficiency for Piwi results in transmission of a heritable stress that promotes longevity via DAF-16/Foxo

Bree Heestand^{1,2,3*}, Matt Simon^{1,2*}, Stephen Frenk^{1,2}, and Shawn Ahmed^{1,2,3}

¹Department of Genetics, University of North Carolina, Chapel Hill, NC, USA

²Department of Biology, University of North Carolina, Chapel Hill, NC, USA

³Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill,
NC, USA

To whom correspondence should be addressed. E-mail: shawn@med.unc.edu

* These authors contributed equally to this manuscript

Abstract

The *C. elegans* Piwi Argonaute protein PRG-1 and associated piRNAs protect the genomes of germ cells by suppressing the expression of transposons and potentially deleterious foreign nucleic acids. Deficiency for *prg-1* compromises germ cell immortality, resulting in normal fertility for many generations followed by progressively reduced fertility and ultimately sterility. The sterility phenotype of *prg-1* mutants was recently shown to be a form of reproductive arrest, which implies that *prg-1* mutants may become sterile in response to a form of heritable stress. The DAF-16 stress resistance and longevity factor can promote germ cell immortality of *prg-1* mutants by activating a systemic RNAi pathway. We found that this RNAi pathway was not required for the somatic longevity function of DAF-16. Given that *prg-1* mutant germ cells may transmit a form of heritable stress, we studied the somatic longevity of *prg-1* mutant adults. We found that early generation *prg-1* mutants had normal lifespans, but that late-generation adults that displayed reduced fertility or sterility were long lived. Germ cells of long-lived late-generation *prg-1* mutants gave rise to F1 cross progeny that were heterozygous for *prg-1*, fertile and also long lived. However, in the absence of DAF-16, the heritable stress transmitted by *prg-1* mutant germ cells was deleterious and caused lifespan to shorten. We conclude that deficiency for the genomic surveillance factor PRG-1/Piwi results in germ cells that transmit a heritable stress that promotes somatic longevity via DAF-16/Foxo, which could be relevant transgenerational regulation of aging.

Introduction

In principle, aging can be divided into two categories: the aging of post-mitotic cells, such as neurons or muscle cells, and the aging of cells that proliferate, such as hematopoietic stem cells. Primary human fibroblasts are deficient for telomerase and can divide for a limited number of cell divisions (40-60) due to telomere erosion, termed the Hayflick limit (1). Telomere erosion induces senescence, a state of permanent cell cycle arrest, but senescence can also be induced by other stresses that include aberrant cellular proliferation conditions (2). In many organisms, somatic aging occurs as a consequence of aging of proliferating cells and of cells that are post-mitotic, but the relationship between these two forms of aging is not well understood (3). The somatic cells of *C. elegans* adults are post-mitotic, whereas mitotic germ cells of *C. elegans* adults proliferate for several days. It has been shown that sterility induced by either physical or genetic ablation of the germline can elicit an increase in adult longevity (4), which implies that there may be a trade-off between reproduction and aging (5).

Germ cells are an effectively immortal cell lineage that can be maintained indefinitely. We study germ cells of the nematode *C. elegans* as a transgenerational model of aging of cells that proliferate. We identified *mortal germline* (*mrt*) mutants that initially display normal levels of fertility but after a number of generations these mutants succumb to complete sterility, which can be thought of as transgenerational senescence of the germ line (6). Some *mrt* mutants are deficient for telomerase-mediated telomere maintenance and their germ cells transmit chromosome termini that shorten with each generation, ultimately eliciting end-to-end chromosome fusions and genome

catastrophe (7, 8). Given that germ cells of human families with telomerase deficiency transmit shortened telomeres that primarily elicit the lethal disorders aplastic anemia and pulmonary fibrosis that shorten lifespan (9), we previously examined post-mitotic aging of *C. elegans* adults that were deficient for the telomerase reverse transcriptase, *trt-1*. We found that early-, middle- and late-generation *trt-1* mutants had wildtype lifespans, even though late-generation strains possessed telomere fusions and had lost segments of subtelomeric genomic DNA from some chromosome termini (8). These results indicate that telomere attrition, which is a known cause of proliferative senescence, does not impact the aging of post-mitotic *C. elegans* adult somatic cells.

A second pathway that promotes germ cell immortality is a genomic silencing pathway where small RNAs associate with Argonaute proteins to promote heterochromatin formation (10-12). This pathway initiates with the perinuclear Argonaute protein PRG-1, the *C. elegans* ortholog of Piwi, which localizes to P granules of germ cells and interacts with thousands of 21 nucleotide piRNAs that scan the genome for viruses, transposons or foreign nucleic acids such as man-made transgenes (13-19). When a genomic intruder is identified, possibly via a process that can detect genomic incongruities that occur when homologous chromosomes are paired during meiosis (20, 21), PRG-1 and associated piRNAs interact with RNA made from the foreign genetic element and recruit RNA-dependent RNA polymerases that act in P-granule associated Mutator bodies to promote biogenesis of 22 nucleotide effector siRNAs named 22G RNAs (22). 22G RNAs generated in response to PRG-1/piRNAs can interact with the P granule Argonaute protein WAGO-1 or the nuclear Argonaute protein HRDE-1, which promote genomic silencing via nuclear RNAi factors and histone

modifying proteins (10, 13, 17). The PRG-1/Piwi pathway therefore represents a genomic immune system for germ cells that silences foreign genetic elements (23, 24).

Freshly outcrossed *prg-1*/piRNA mutants display normal levels of fertility for several generations (12). However, in later generations *prg-1* mutants show a decline in fertility such that very few progeny are produced prior to the onset of reproductive delay and sterility (25). Some late-generation animals display pronounced effects on reproduction, such that sterility occurs on day 1 of adulthood, followed by fertility on days 2-4 adulthood. However, some *prg-1* mutants that are sterile on day 1 of adulthood remain in an indefinite state of reproductive quiescence termed Adult Reproductive Diapause (ARD) that can be reversed if environmental conditions change (25). States of developmental arrest, termed diapause, occur in many species in response to environmental stresses such as nutritional or seasonal cues, and ARD has been shown to occur in *C. elegans* in response to starvation (26-28). We therefore hypothesized that *prg-1* mutant germ cells may transmit a form of Heritable Stress that eventually becomes severe enough to induce a state of reproductive arrest (25).

We previously found that the number of generations that *prg-1* mutant strains can grow, termed transgenerational lifespan, can be significantly extended by weekly bouts of starvation (12). This effect was mediated by the DAF-16/Foxo transcription factor that responds to a variety of stresses and can promote adult longevity (3, 29). Constitutive activation of DAF-16 (by mutation of the upstream *daf-2* insulin/IGF-1 receptor) restores germ cell immortality to *prg-1* mutants (12). This effect requires 22G RNAs created by the Mutator body protein MUT-7 as well as the H3K4 demethylase SPR-5, both of which promote genomic silencing, implying that DAF-16 activity represses the

transgenerational sterility phenotype of *prg-1* mutants by activating a compensatory small RNA-mediated silencing pathway (12). In the accompanying paper, by Simon, Spichal, Heestand and colleagues, we demonstrate that DAF-16 acts in somatic cells to upregulate a systemic small RNA pathway that requires Dicer and the double-stranded RNA (dsRNA) binding protein RDE-4, which function together to promote primary siRNA biogenesis in response to exogenous or endogenous dsRNA triggers (22).

microRNAs are produced from hairpin precursor molecules that processed by the ribonucleases Dicer and Drosha in conjunction with DGCR8/*pash-1* (30). A conditional allele of *pash-1* has revealed that loss of miRNAs shortens lifespan, although miRNAs were not required for the long life of *daf-2* mutants whose longevity depends on the DAF-16 stress response transcription factor (31). Specific microRNAs have also been demonstrated to regulate aging in *C. elegans* (32-35), including some microRNAs whose levels change with age (36-38).

Here we address the relationship between aging of post-mitotic cells with transgenerational aging of germ cells by studying the adult longevity of *prg-1* mutants. Our results suggest that *prg-1* mutants transmit an epigenetic factor, termed heritable stress, that is intimately related to the function of DAF-16/Foxo and that can promote adult longevity.

Results

The systemic small RNA pathway activated by DAF-16 does not affect adult longevity

daf-2 mutants have been reported to be hypersensitive to RNA interference (39), and *daf-2* mutations suppress the transgenerational sterility phenotype of *prg-1* mutants by activating a systemic RNAi pathway that requires Dicer and the dsRNA binding protein RDE-4 (see related manuscript by Simon, Spichal, Heestand and colleagues). Because RDE-4 promotes the biogenesis of endogenous small RNAs and *rde-4* mutants have relatively minor overt phenotypes (11, 40, 41), we asked whether deficiency for *rde-4* affects the lifespan of *daf-2* mutants. We found that *daf-2; rde-4* double mutant lifespan was not significantly different than that of *daf-2* single mutant controls (Fig 1A, S1 Table, $p=1$), indicating that the small RNA pathway that can be activated by DAF-16 in *prg-1* mutants is dispensable for its function in adult longevity.

Heritable Stress of *prg-1* mutants induces longevity but not dauer formation

In *C. elegans*, various environmental stresses including temperature, crowding and starvation can induce an alternative developmental stage termed the Dauer Larva, which is a state of development arrest that is long-lived and highly stress resistant (42). Late-generation *prg-1* mutants can enter a reversible state of reproductive arrest, suggesting that *prg-1* mutant germ cells may transmit a heritable stress (25). We therefore asked whether late-generation *prg-1* mutants that were close to sterility gave rise to dauer larvae. We scored late-generation *prg-1* mutants that were poised to become sterile at the stressful temperature of 25°C, but did not observe any dauer larvae, similar to wild-type controls, and to early and fertile late-generation *prg-1* mutant controls (Fig 1C). To verify that *prg-1* mutants were not defective for inducing dauer larvae, we crossed a *daf-2* mutation, which leads to high levels of dauers at 25°C, to

create *prg-1*; *daf-2* double mutants and found that there was no defect in entry into dauer arrest (Fig 1C). This indicates that the heritable stress that is transmitted by *prg-1* mutant germ cells can specifically induce developmental arrest of the adult germ line and does not perturb larval development, even though dauer larvae are induced by a variety of stresses like heat, crowding and starvation (42).

C. elegans worms that are subjected to dietary restriction, acute starvation or other stresses live much longer than worms fed ad libitum (26, 43-46). These observations are consistent with the phenomenon of hormesis, in which adult longevity can be promoted by stresses that are strong but not debilitating (47, 48). We therefore asked whether *prg-1* mutant lifespan is affected by the heritable stress transmitted by *prg-1* mutant germ cells, by measuring longevity of early-generation, late-generation *prg-1* mutants that were sterile and late-generation *prg-1* mutants that were fertile. We found that early-generation *prg-1* mutants, with low amounts of heritable stress, were not significantly different than wild-type animals (Fig 1B, S1 Table, $p=1$). However, fertile but stressed late-generation and sterile *prg-1* mutant adults displayed significant 46.5% and 81.8% increases in median lifespan in comparison to wild-type controls, respectively (Fig 1B, S1 Table, $p=0.0008$ and $p=1.55E-16$) suggesting that increased levels of heritable stress in germ cells is sufficient to confer an increase in somatic longevity.

***prg-1* mutant germ cells transmit longevity to heterozygous *prg-1/+* progeny**

We tested the hypothesis that *prg-1* mutant germ cells transmit a form of heritable stress that affects longevity by crossing late-generation *prg-1* mutant

hermaphrodites with wildtype males and measuring longevity of their F1 cross-progeny. We performed crosses of three independently derived lines of late-generation *prg-1* hermaphrodites with males that were heterozygous for an *unc-13* mutation and compared the lifespans of their *prg-1* +/- F1 progeny with those of early- and late-generation *prg-1* mutant controls. To ensure that the presence of the *unc-13* balancer mutation did not confer an unexpected dominant effect on longevity, we also measured the lifespan of +/-*unc-13* heterozygous controls and found that these were not significantly different from early-generation *prg-1* mutant controls (Fig 2A, S1 Table, p=1). During the initial days of the longevity experiment (reproductive period), we transferred adults to new plates every day until they stopped laying progeny. Although late-generation *prg-1* mutant control adults give rise to very few progeny, we observed that their *prg-1* +/- F1 progeny had a marked increase in brood size, demonstrating that the strong fertility defect of late-generation *prg-1* mutants is not transmitted to their heterozygous progeny. However, all three independent lines of *prg-1* +/- F1 progeny displayed significant increases in lifespan compared to +/-*unc-13*, early-generation *prg-1*, and early-generation *prg-1* +/- controls (Fig 2A,B, S1 Table, p>0.05 in all comparisons). Notably, the lifespan of these *prg-1* +/- F1 cross progeny was not statistically different than late-generation *prg-1* mutant adults (Fig 2 A,B, S1 Table, p>0.72 in all cases).

Heritable stress is detrimental to longevity in the absence of *daf-16*

The DAF-16 transcription factor is well known to promote longevity (3, 29, 49, 50) and has been previously shown to interact in three distinct ways with deficiency for *prg-1*. First, when *daf-2* mutations that activate DAF-16 are introduced into early-generation

prg-1 mutants, DAF-16 promotes germ cell immortality and suppresses the transgenerational sterility defect of *prg-1* mutants (12). However, knockdown of *daf-2* by mutation or RNAi does not suppress the sterility phenotype of late-generation *prg-1* mutants (25), indicating that DAF-16 may lose its ability to promote germ cell immortality in late-generation *prg-1* mutants. A second function of DAF-16 occurs when *prg-1* mutants become sterile. Most of these animals display a pronounced germ cell atrophy phenotype as L4 larvae mature in to 1 day old adults, but deficiency for *daf-16* significantly alters the germline profiles of sterile *prg-1* mutant adults, leading to reduced levels of germ cell atrophy (25). This implies that *daf-2* signaling promotes germ cell atrophy as sterile *prg-1* mutant L4 larvae mature into 1 day old adults. A third function of DAF-16 occurs for 5 day old sterile *prg-1* mutants, which are in a state of prolonged reproductive arrest, a small fraction of which can become fertile if moved to an alternative food source (25). This exit from reproductive arrest requires DAF-16. We therefore asked whether the longevity of late-generation *prg-1* mutants depends on DAF-16. Fertile and sterile late-generation *prg-1 daf-16* animals lived significantly shorter than either early-generation *prg-1 daf-16* double mutants (late $p=3.53E-12$ and sterile $p=1.55E-35$) or *daf-16* single mutant controls (late $p=0.028$ and sterile $p=2.71E-19$), (Fig 2C, S1 Table), indicating that DAF-16 is responsible for the longevity observed in late-generation *prg-1* mutants. Furthermore, sterile late-generation *prg-1 daf-16* double mutants were shorter lived than their fertile late-generation siblings (Fig 2C, S1 Table, $p=5.41E-13$) implying that a detrimental form of heritable stress is transmitted by late-generation *prg-1* mutants.

Given that DAF-16 promotes the longevity of late-generation *prg-1* mutants, we used a *daf-16::GFP* transgene that rescues the adult longevity defect of *daf-16*; *daf-2* double mutants (51) to create *prg-1*; *daf-16::GFP* mutant lines. We found that early- and late-generation *prg-1* mutant lines displayed DAF-16::GFP expression that was primarily cytoplasmic, similar to that of *daf-16::GFP* transgenic controls (Fig 3A,B). While DAF-16::GFP was only mildly nuclear in fertile late-generation *prg-1* mutants (Fig 3C,E), we found that sterile late-generation *prg-1* mutant strains displayed strong nuclear localization of DAF-16::GFP in somatic tissue, similar to that observed with *daf-16::GFP*; *daf-2* mutant controls, where DAF-16 is constitutively activated (Fig 3D,E) (51).

Transgenerational activation of P bodies in *prg-1* mutants

Processing bodies (P bodies) and stress granules are typically cytoplasmic aggregates that are composed of proteins and RNA that function to protect RNAs under stressful conditions. P bodies appear under conditions of stress, including many types of environmental stress such as heat shock, radiation, osmotic stress, oxidative stress and ER stress (52). Given that *prg-1* mutant germ cells may transmit a form of heritable stress that promotes somatic longevity, we crossed GFP-tagged component of P bodies and stress granules DCAP-1 (53) into *prg-1* mutant backgrounds and monitored GFP expression in early- and late-generation strains. We observed that DCAP-1::GFP aggregates were rarely present in wild-type and early-generation *prg-1* mutant strains but were significantly induced in late-generation sterile *prg-1* mutants (Fig 3 F-I), similar to what has been previously observed under stressful conditions (53). These results are

consistent the possibility that late-generation *prg-1* mutants may experience a form of stress.

Discussion

daf-2 mutants have been previously reported to be hypersensitive to the exogenous RNAi pathway that normally targets foreign viruses and transposons (48). The PRG-1/Piwi silencing pathway also represents a form of immunity that protects the germ line from foreign genetic elements (54, 55). We found that deficiency for the *prg-1*-mediated genomic silencing pathway leads to a transgenerational sterility phenotype that can be suppressed by mutation of *daf-2*, via an endogenous systemic small RNA pathway that is activated by DAF-16 (see accompanying manuscript by Simon, Spichal, Heestand and colleagues). Given that DAF-16 can act cell-non-autonomously in distinct somatic tissues to promote longevity of *daf-2* adults (56), we asked whether this systemic RNAi pathway was relevant to somatic longevity of *daf-2* mutants and found it was not. This indicates that DAF-16 does not simply respond to stress by upregulating pathways that promote somatic stress resistance and longevity, but that it possesses a distinct function that promotes germ cell immortality when PRG-1/Piwi is deficient (12). Consistently, DAF-16 has been previously linked to the innate immune response to bacterial or fungal pathogens (57-60), an activity that is separable from its function in somatic longevity (57).

The disposable soma theory of aging posits that trade-offs occur when deciding on activating processes that benefit either somatic longevity or reproduction (5, 61). However, our study implies that these choices are not mutually exclusive and that

stresses that compromise germ cell function can be counteracted by a pathway that promotes somatic longevity in times of stress. Our data suggests that a trade-off may occur for the DAF-16 pathway in the context of deficiency for *prg-1*/piRNAs and lead us to propose the following model (Fig 4): In early generation *prg-1* mutants, DAF-16 activity is wildtype from the perspective of somatic longevity and DAF-16::GFP fluorescence. In this context, DAF-16 can be activated transiently or constitutively to suppress the sterility phenotype of *prg-1* mutants (12). In a population of wild animals that experience a virus- or transposon-based attack that compromises the PRG-1/Piwi silencing system (54, 55), germlines of those animals that experience strong bouts of environmental stress may be better able to withstand a genomic parasite. Indeed, the feast or famine lifestyle of *C. elegans* populations that rely on ephemeral food sources such as rotten fruits or vegetables may experience significant levels DAF-16 activity on a consistent basis, which may suppress dysfunction of *prg-1*/Piwi (62, 63). If, however, *prg-1* deficient strains are transferred regularly under laboratory conditions of replete food, these strains transmit a form of heritable stress that builds up to the point where it engages DAF-16 to promote somatic longevity (Fig 1). Fertile late-generation *prg-1* mutant strains were long-lived and showed a mild and transient DAF-16::GFP nuclear localization pattern. Given that genetic or RNAi knockdown of *daf-2* in late-generation *prg-1* mutant strains no longer ameliorates their sterility defect (25), we propose that if DAF-16 is already engaged in responses such as heritable stress-induced adult longevity that may preclude DAF-16 from activating the systemic RNAi pathway that suppresses the sterility of *prg-1* mutants (12). An alternative explanation is that the

heterochromatin defect of late-generation *prg-1* mutants is so severe that it can no longer be suppressed by activation of DAF-16.

Support for the concept that heritable stress engages specific facets of the DAF-16 pathway comes from our observation that late-generation *prg-1* mutants do not display dauer larvae that are commonly produced under circumstances of stress (Fig 1C) (42). Instead, DAF-16 specifically promotes longevity and reproductive arrest in late-generation *prg-1* mutant adults (25), both of which may represent adaptive responses to the heritable stress transmitted by *prg-1* mutant germ cells. We found that F1 cross progeny of late-generation *prg-1* mutants were also long-lived yet lacked the strong fertility defect of their *prg-1* mutant mothers, which suggests a dilution of the stress, but also partial separation between the stress and the organismal response to it. In the absence of *daf-16*, late-generation *prg-1* mutants lived shorter than *daf-16* single mutants, which are themselves short-lived (64, 65). This is consistent with the longevity of late-generation *prg-1* single mutants being a hormetic stress response (47, 48), where stress is beneficial unless the relevant stress response pathway is inactivated.

Additional support for the concept of heritable stress is provided by the observation that strong increases in P bodies occur in late-generation *prg-1* mutants. As P bodies or stress granules are created in response to many forms of stress (52), their presence implies that the heritable stress transmitted by *prg-1* mutant germ cells induces a stress response in adult somatic tissues. P bodies or stress granules can promote viral resistance and aspects of the innate immune response (52), which are also properties of perinuclear germ granules that house the PRG-1/Piwi/piRNA genome silencing proteins (66). Transgenerational dysfunction of the PRG-1/piRNA silencing

pathway leads to desilencing of transposons (12, 67, 68), and expression of one or more transposon classes could facilitate stress granule formation in a manner that promotes lifespan extension. While the precise identity of the heritable stress transmitted by *prg-1* mutant germ cells remains uncertain, our data provide strong evidence that germ cells can transmit factors other than DNA mutations that can have a significant impact on longevity. We suggest that desilencing of heterochromatin of *prg-1* mutant germ cells over the generations may perturb an essential biochemical pathway that corresponds to the heritable stress that leads to increased longevity and P body formation in late-generation *prg-1* mutants.

Inactivation of members of the ASH-2 trithorax histone H3K4 methylation complex for a single generation promotes an adult longevity phenotype that can be transmitted transgenerationally in *C. elegans* (69). Functionally, ASH-2/trithorax transcriptional activation complex might be predicted to oppose the activity the PRG-1/piRNA genome silencing pathway, and it was demonstrated that the RBR-2 H3K4 demethylase that promotes genome silencing in germ cells is required for the longevity caused by inactivation of ASH-2/trithorax (69). RBR-2 is required for germ cell immortality at 25°C (70), likely because it acts downstream of PRG-1/piRNAs. Moreover, ASH-2 trithorax inactivation was capable of extending the lifespan of *daf-16* mutants, whereas late-generation *prg-1* mutants were very short-lived in the absence of *daf-16* (Fig 2). Together, these results suggest that independent epigenetic defects can promote longevity in a manner that is heritable, such that knockdown of the ASH-2 trithorax transcriptional activation complex induces an immediate somatic longevity

phenotype, whereas inactivation of the PRG-1/piRNA genomic silencing complex for many generations leads to a functionally distinct form of longevity.

Like RBR-2, the SPR-5 H3K4 demethylase is required for germ cell immortality (70). While this study was in progress it was shown that late- but not early-generation *spr-5* mutants are long lived (71). However, *daf-16* is not required for the longevity of late-generation *spr-5* mutants (71), implying that deficiency for *prg-1* and *spr-5* lead to distinct transgenerational epigenetic defects that promote distinct mechanisms of longevity.

The transgenerational sterility of *prg-1* mutants suggests a defect that accumulates as germ cells proliferate over the generations, which is potentially analogous to proliferative aging or senescence of mammalian somatic cells (72). Several links between aging and dysregulation of the epigenome have been documented, such as desilencing of retrotransposons in senescent human cells, and epigenetic silencing of retrotransposons by the anti-aging protein SIRT6 in the mouse (73-75). The induction of reproductive arrest in response to heritable stress therefore suggests a form of inheritance that is potentially relevant to epigenetic modulation of senescence of mammalian somatic cells.

Our study suggests that mutation of *prg-1*/Piwi can induce a form of heritable stress that is transmitted by gametes and can promote longevity, which could represent one facet of the evolutionary biology of aging in metazoans. An analogous form of non-Mendelian stress that is transmitted by germ cells could affect the rate of aging or adult onset disorders in mammals, potentially by modulating aging of post-mitotic somatic

cells in a manner that depends on Foxo or by promoting replicative aging of proliferating cells.

Materials and Methods

Strains

Unless noted otherwise, all strains were cultured at 20°C on Nematode Growth Medium (NGM) plates seeded with *E. coli* OP50. Strains used include Bristol N2 wild type, *unc-13(e450) I*, *prg-1(n4357) I*, *daf-16(mu86) I*, *kri-1(ok1251) I*, *daf-2(e1368) III*, zls356 [*daf-16p::daf-16a/b::GFP + rol-6(su10060)*, *P_{dcap-1}::dcap-1::gfp::3' UTR^{dcap-1}*

prg-1 mutations were outcrossed versus an outcrossed stock of *unc-13(e450)*, and freshly isolated homozygous F2 lines were established for analysis. To create the linked *prg-1*, *daf-16* double mutant, *prg-1*, *dpy-24* and *unc-13*, *daf-16* double mutants were first created, then progeny of *prg-1*, *dpy-24* / *unc-13*, *daf-16* heterozygotes that lost *unc-13* were identified, and the resulting putative *prg-1*, *daf-16* recombinant chromosomes were made homozygous and PCR genotyped to verify the presence of *prg-1* or *daf-16* deletions.

Lifespan analysis

Animals were singled from non-starved plates and transferred every two days until the end of their reproductive cycles. Animals were kept uncrowded and free of

contamination throughout experiment and transferred to new plates once per week. All animals were maintained at 20°C.

Statistical analysis

Statistical analysis was performed using the R statistical computing environment (76) and the survival R package (77). For lifespan experiments, pairwise log-rank tests were performed to detect significant differences between groups. For the P bodies experiment, a Kruskal-Wallace test was performed to detect the presence of any significant differences within the dataset. Pairwise Mann Whitney U tests were performed as a post-hoc test to detect the presence of significant differences between specific groups. All reported p-values were adjusted using Bonferroni correction to control the familywise error rate.

Acknowledgements

We thank members of the Ahmed lab for critical reading of the manuscript. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). This study was supported by NIH grant RO1 GM083048 (S.A).

Competing interests: Authors declare no competing interests.

Author contributions: B.H. and M.S. performed experiments. S.F. analyzed the data. B.H. and S.A. wrote the manuscript.

References

1. Shay JW, Wright WE. Senescence and immortalization: role of telomeres and telomerase. *Carcinogenesis*. 2005;26(5):867-74.
2. He S, Sharpless NE. Senescence in Health and Disease. *Cell*. 2017;169(6):1000-11.
3. Bitto A, Wang AM, Bennett CF, Kaeberlein M. Biochemical Genetic Pathways that Modulate Aging in Multiple Species. *Cold Spring Harb Perspect Med*. 2015;5(11).
4. Hsin H, Kenyon C. Signals from the reproductive system regulate the lifespan of *C. elegans*. *Nature*. 1999;399(6734):362-6.
5. Kirkwood TB, Holliday R. The evolution of ageing and longevity. *Proc R Soc Lond B Biol Sci*. 1979;205(1161):531-46.
6. Smelick C, Ahmed S. Achieving immortality in the *C. elegans* germline. *Ageing Res Rev*. 2005;4(1):67-82.
7. Ahmed S, Hodgkin J. MRT-2 checkpoint protein is required for germline immortality and telomere replication in *C. elegans*. *Nature*. 2000;403(6766):159-64.
8. Meier B, Clejan I, Liu Y, Lowden M, Gartner A, Hodgkin J, et al. trt-1 is the *Caenorhabditis elegans* catalytic subunit of telomerase. *PLoS Genet*. 2006;2(2):e18.
9. Armanios M, Blackburn EH. The telomere syndromes. *Nat Rev Genet*. 2012;13(10):693-704.
10. Buckley BA, Burkhart KB, Gu SG, Spracklin G, Kershner A, Fritz H, et al. A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature*. 2012;489(7416):447-51.
11. Sakaguchi A, Sarkies P, Simon M, Doebley AL, Goldstein LD, Hedges A, et al. *Caenorhabditis elegans* RSD-2 and RSD-6 promote germ cell immortality by maintaining small interfering RNA populations. *Proc Natl Acad Sci U S A*. 2014;111(41):E4323-31.
12. Simon M, Sarkies P, Ikegami K, Doebley AL, Goldstein LD, Mitchell J, et al. Reduced insulin/IGF-1 signaling restores germ cell immortality to *Caenorhabditis elegans* Piwi mutants. *Cell reports*. 2014;7(3):762-73.
13. Ashe A, Sapetschnig A, Weick EM, Mitchell J, Bagijn MP, Cording AC, et al. piRNAs Can Trigger a Multigenerational Epigenetic Memory in the Germline of *C. elegans*. *Cell*. 2012;150(1):88-99.
14. Bagijn MP, Goldstein LD, Sapetschnig A, Weick EM, Bouasker S, Lehrbach NJ, et al. Function, targets, and evolution of *Caenorhabditis elegans* piRNAs. *Science*. 2012;337(6094):574-8.
15. Batista PJ, Ruby JG, Claycomb JM, Chiang R, Fahlgren N, Kasschau KD, et al. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in *C. elegans*. *Mol Cell*. 2008;31(1):67-78.
16. Das PP, Bagijn MP, Goldstein LD, Woolford JR, Lehrbach NJ, Sapetschnig A, et al. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the *Caenorhabditis elegans* germline. *Mol Cell*. 2008;31(1):79-90.

17. Lee HC, Gu W, Shirayama M, Youngman E, Conte D, Jr., Mello CC. C. elegans piRNAs mediate the genome-wide surveillance of germline transcripts. *Cell*. 2012;150(1):78-87.
18. Shirayama M, Seth M, Lee HC, Gu W, Ishidate T, Conte D, Jr., et al. piRNAs initiate an epigenetic memory of nonself RNA in the C. elegans germline. *Cell*. 2012;150(1):65-77.
19. Wang G, Reinke V. A C. elegans Piwi, PRG-1, regulates 21U-RNAs during spermatogenesis. *Curr Biol*. 2008;18(12):861-7.
20. Kelly WG, Aramayo R. Meiotic silencing and the epigenetics of sex. *Chromosome Res*. 2007;15(5):633-51.
21. Leopold LE, Heestand BN, Seong S, Shtessel L, Ahmed S. Lack of pairing during meiosis triggers multigenerational transgene silencing in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 2015;112(20):E2667-76.
22. Billi AC, Fischer SE, Kim JK. Endogenous RNAi pathways in C. elegans. *WormBook*. 2014:1-49.
23. Juliano C, Wang J, Lin H. Uniting germline and stem cells: the function of Piwi proteins and the piRNA pathway in diverse organisms. *Annu Rev Genet*. 2011;45:447-69.
24. Kasper DM, Gardner KE, Reinke V. Homeland security in the C. elegans germ line: insights into the biogenesis and function of piRNAs. *Epigenetics*. 2014;9(1):62-74.
25. Heestand B, Simon M, Frenk S, Titov D, Ahmed S. Transgenerational Sterility of Piwi Mutants Represents a Dynamic Form of Adult Reproductive Diapause. *Cell reports*. 2018;23(1):156-71.
26. Angelo G, Van Gilst MR. Starvation protects germline stem cells and extends reproductive longevity in C. elegans. *Science*. 2009;326(5955):954-8.
27. Padilla PA, Ladage ML. Suspended animation, diapause and quiescence: arresting the cell cycle in C. elegans. *Cell Cycle*. 2012;11(9):1672-9.
28. Tatar M, Yin C. Slow aging during insect reproductive diapause: why butterflies, grasshoppers and flies are like worms. *Exp Gerontol*. 2001;36(4-6):723-38.
29. DiLoreto R, Murphy CT. The cell biology of aging. *Mol Biol Cell*. 2015;26(25):4524-31.
30. Blakaj A, Lin H. Piecing together the mosaic of early mammalian development through microRNAs. *J Biol Chem*. 2008;283(15):9505-8.
31. Lehrbach NJ, Castro C, Murfitt KJ, Abreu-Goodger C, Griffin JL, Miska EA. Post-developmental microRNA expression is required for normal physiology, and regulates aging in parallel to insulin/IGF-1 signaling in C. elegans. *RNA*. 2012;18(12):2220-35.
32. Bethke A, Fielenbach N, Wang Z, Mangelsdorf DJ, Antebi A. Nuclear hormone receptor regulation of microRNAs controls developmental progression. *Science*. 2009;324(5923):95-8.
33. Boehm M, Slack F. A developmental timing microRNA and its target regulate life span in C. elegans. *Science*. 2005;310(5756):1954-7.
34. Boulias K, Horvitz HR. The C. elegans microRNA mir-71 acts in neurons to promote germline-mediated longevity through regulation of DAF-16/FOXO. *Cell Metab*. 2012;15(4):439-50.
35. Kato M, Slack FJ. Ageing and the small, non-coding RNA world. *Ageing Res Rev*. 2013;12(1):429-35.

36. de Lencastre A, Pincus Z, Zhou K, Kato M, Lee SS, Slack FJ. MicroRNAs both promote and antagonize longevity in *C. elegans*. *Curr Biol*. 2010;20(24):2159-68.
37. Ibanez-Ventoso C, Yang M, Guo S, Robins H, Padgett RW, Driscoll M. Modulated microRNA expression during adult lifespan in *Caenorhabditis elegans*. *Aging Cell*. 2006;5(3):235-46.
38. Pincus Z, Smith-Vikos T, Slack FJ. MicroRNA predictors of longevity in *Caenorhabditis elegans*. *PLoS Genet*. 2011;7(9):e1002306.
39. Samuelson AV, Klimczak RR, Thompson DB, Carr CE, Ruvkun G. Identification of *Caenorhabditis elegans* genes regulating longevity using enhanced RNAi-sensitive strains. *Cold Spring Harb Symp Quant Biol*. 2007;72:489-97.
40. Blanchard D, Parameswaran P, Lopez-Molina J, Gent J, Saynuk JF, Fire A. On the nature of in vivo requirements for *rde-4* in RNAi and developmental pathways in *C. elegans*. *RNA Biol*. 2011;8(3):458-67.
41. Lee RC, Hammell CM, Ambros V. Interacting endogenous and exogenous RNAi pathways in *Caenorhabditis elegans*. *RNA*. 2006;12(4):589-97.
42. Hu PJ. Dauer. *WormBook*. 2007:1-19.
43. Greer EL, Brunet A. Different dietary restriction regimens extend lifespan by both independent and overlapping genetic pathways in *C. elegans*. *Aging Cell*. 2009;8(2):113-27.
44. Houthoofd K, Braeckman BP, Johnson TE, Vanfleteren JR. Life extension via dietary restriction is independent of the Ins/IGF-1 signalling pathway in *Caenorhabditis elegans*. *Exp Gerontol*. 2003;38(9):947-54.
45. Kaeberlein TL, Smith ED, Tsuchiya M, Welton KL, Thomas JH, Fields S, et al. Lifespan extension in *Caenorhabditis elegans* by complete removal of food. *Aging Cell*. 2006;5(6):487-94.
46. Lakowski B, Hekimi S. The genetics of caloric restriction in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 1998;95(22):13091-6.
47. Calabrese EJ, Iavicoli I, Calabrese V. Hormesis: why it is important to biogerontologists. *Biogerontology*. 2012;13(3):215-35.
48. Shore DE, Ruvkun G. A cytoprotective perspective on longevity regulation. *Trends Cell Biol*. 2013;23(9):409-20.
49. Kapahi P, Kaeberlein M, Hansen M. Dietary restriction and lifespan: Lessons from invertebrate models. *Ageing Res Rev*. 2017;39:3-14.
50. Kaplan RE, Baugh LR. L1 arrest, *daf-16*/FoxO and nonautonomous control of post-embryonic development. *Worm*. 2016;5(2):e1175196.
51. Lin K, Hsin H, Libina N, Kenyon C. Regulation of the *Caenorhabditis elegans* longevity protein DAF-16 by insulin/IGF-1 and germline signaling. *Nat Genet*. 2001;28(2):139-45.
52. Protter DS, Parker R. Principles and Properties of Stress Granules. *Trends Cell Biol*. 2016;26(9):668-79.
53. Rousakis A, Vlanti A, Borbolis F, Roumelioti F, Kapetanou M, Syntichaki P. Diverse functions of mRNA metabolism factors in stress defense and aging of *Caenorhabditis elegans*. *PLoS One*. 2014;9(7):e103365.
54. Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science*. 2007;318(5851):761-4.

55. Ernst C, Odom DT, Kutter C. The emergence of piRNAs against transposon invasion to preserve mammalian genome integrity. *Nat Commun.* 2017;8(1):1411.
56. Libina N, Berman JR, Kenyon C. Tissue-specific activities of *C. elegans* DAF-16 in the regulation of lifespan. *Cell.* 2003;115(4):489-502.
57. Evans EA, Chen WC, Tan MW. The DAF-2 insulin-like signaling pathway independently regulates aging and immunity in *C. elegans*. *Aging Cell.* 2008;7(6):879-93.
58. Jensen VL, Simonsen KT, Lee YH, Park D, Riddle DL. RNAi screen of DAF-16/FOXO target genes in *C. elegans* links pathogenesis and dauer formation. *PLoS One.* 2010;5(12):e15902.
59. Kurz CL, Tan MW. Regulation of aging and innate immunity in *C. elegans*. *Aging Cell.* 2004;3(4):185-93.
60. Troemel ER, Chu SW, Reinke V, Lee SS, Ausubel FM, Kim DH. p38 MAPK regulates expression of immune response genes and contributes to longevity in *C. elegans*. *PLoS Genet.* 2006;2(11):e183.
61. Nystrom T. Aging in bacteria. *Curr Opin Microbiol.* 2002;5(6):596-601.
62. Felix MA, Braendle C. The natural history of *Caenorhabditis elegans*. *Curr Biol.* 2010;20(22):R965-9.
63. Jobson MA, Jordan JM, Sandrof MA, Hibshman JD, Lennox AL, Baugh LR. Transgenerational Effects of Early Life Starvation on Growth, Reproduction, and Stress Resistance in *Caenorhabditis elegans*. *Genetics.* 2015;201(1):201-12.
64. Lin K, Dorman JB, Rodan A, Kenyon C. *daf-16*: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science.* 1997;278(5341):1319-22.
65. Ogg S, Paradis S, Gottlieb S, Patterson GL, Lee L, Tissenbaum HA, et al. The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature.* 1997;389(6654):994-9.
66. Strome S, Updike D. Specifying and protecting germ cell fate. *Nat Rev Mol Cell Biol.* 2015;16(7):406-16.
67. McMurchy AN, Stempor P, Gaarenstroom T, Wysolmerski B, Dong Y, Aussanikava D, et al. A team of heterochromatin factors collaborates with small RNA pathways to combat repetitive elements and germline stress. *Elife.* 2017;6.
68. Ni JZ, Chen E, Gu SG. Complex coding of endogenous siRNA, transcriptional silencing and H3K9 methylation on native targets of germline nuclear RNAi in *C. elegans*. *BMC Genomics.* 2014;15:1157.
69. Greer EL, Maures TJ, Hauswirth AG, Green EM, Leeman DS, Maro GS, et al. Members of the H3K4 trimethylation complex regulate lifespan in a germline-dependent manner in *C. elegans*. *Nature.* 2010;466(7304):383-7.
70. Alvares SM, Mayberry GA, Joyner EY, Lakowski B, Ahmed S. H3K4 demethylase activities repress proliferative and postmitotic aging. *Aging Cell.* 2014;13(2):245-53.
71. Greer EL, Becker B, Latza C, Antebi A, Shi Y. Mutation of *C. elegans* demethylase *spr-5* extends transgenerational longevity. *Cell Res.* 2016;26(2):229-38.
72. Sharpless NE, DePinho RA. How stem cells age and why this makes us grow old. *Nat Rev Mol Cell Biol.* 2007;8(9):703-13.

73. De Cecco M, Criscione SW, Peckham EJ, Hillenmeyer S, Hamm EA, Manivannan J, et al. Genomes of replicatively senescent cells undergo global epigenetic changes leading to gene silencing and activation of transposable elements. *Aging Cell*. 2013;12(2):247-56.
74. Gorbunova V, Boeke JD, Helfand SL, Sedivy JM. Human Genomics. Sleeping dogs of the genome. *Science*. 2014;346(6214):1187-8.
75. Van Meter M, Kashyap M, Rezazadeh S, Geneva AJ, Morello TD, Seluanov A, et al. SIRT6 represses LINE1 retrotransposons by ribosylating KAP1 but this repression fails with stress and age. *Nat Commun*. 2014;5:5011.
76. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. 2017.
77. T T. A Package for Survival Analysis in S. version 2.38. <https://CRAN.R-project.org/package=survival>. 2015.

Figure 1. *prg-1* mutants show a transgenerational increase in lifespan.

Lifespan curves for (A) *daf-2 (e1368) rde-4 (ne301)* and (B) early, late, and sterile generation *prg-1* mutants. (C) Percent worms at each developmental stage after 72 hours at 25°C. n.s. is not significant, *** $p < 0.001$, p values determined using log rank analysis and reported in S1 Table.

Figure 2. Epigenetic inheritance of heritable stress.

(A) Lifespan curves for *+/unc-13*, early or late (stressed) generation *prg-1*, and early or late *prg-1 +/-* F1 progeny. (B) independent lines of heterozygous (*prg-1 +/-*) late generation F1 progeny compared to homozygous non-crossed sisters. $p=1$ (C) *daf-16 (mu86)* is required for somatic lifespan extension in late and sterile generation *prg-1 (n4357)* mutants. n.s. is not significant, *** $p<0.001$, p values determined using log rank analysis and reported in S1 Table.

Figure 3. Induction of somatic stress response with heritable stress.

(A) DAF-16::GFP remains primarily cytoplasmic in adult wild-type animals. (B) Mutant *prg-1(n4357)* animals display no nuclear localization of *daf-16::GFP* in early generation animals, but demonstrates partial nuclear localization in fertile late generation (C) and strong (D) nuclear localization in sterile generation adults. White arrows indicate nuclear localization. Scales bars = 50 μ m. (E) P body aggregates visualized using *dcap-1::gfp* in wild-type, (F) early *prg-1(n4357)*, (G) and stressed late generation *prg-1(n4357)* mutants. Scale bars = 50 μ m. (H) Percent of wild-type, early generation *prg-1*, fertile late (stressed) generation *prg-1*, and sterile late generation *prg-1* worms with no, partial, or full *daf-16::GFP* nuclear localization phenotypes. (n=25, 22, 22, 20 respectively). (I) Number of P body aggregates from (F-H). * is p=0.002 Kruskal Wallace test.

Figure 4. Model for heritable stress-induced longevity.

Loss of *prg-1* leads to a transgenerational increase in heritable stress which promotes somatic longevity through DAF-16/Foxo.

	Median Lifespan	Mean Lifespan	Maximum Lifespan	Change Median	Change Mean	Change Maximum	Total Worms	Total Worms Omitted	Comparison	Log-Rank p Value
Wild-type (N2)	17.06	17.69	27				120	29		
<i>daf-2(e1368)</i>	32.07	28.93	53	88.00%	63.51%	96.30%	120	35	N2 vs <i>daf-2(e1368)</i>	8.10E-23
<i>rde-4(ne301)</i>	18.56	18.79	34	8.77%	6.22%	25.93%	120	28	<i>rde-4(ne301)</i> vs <i>daf-2(e1368)</i>	1.93E-18
<i>daf-2(e1368) rde-4(ne301)</i>	28.70	29.59	43	54.67%	57.46%	26.47%	120	34	<i>rde-4(ne301)</i> vs <i>daf-2 rde-4</i>	7.10E-23
Wild-type (N2)	17.30	18.37	29				120	33		
<i>daf-2(e1368)</i>	29.57	30.40	46	70.93%	66.48%	58.62%	120	34	N2 vs <i>daf-2(e1368)</i>	1.85E-25
<i>rde-4(ne301)</i>	17.17	18.24	34	-0.77%	-0.71%	17.24%	120	23	<i>rde-4(ne301)</i> vs <i>daf-2(e1368)</i>	1.25E-25
<i>daf-2(e1368) rde-4(ne301)</i>	29.78	28.96	48	71.46%	58.82%	41.18%	120	35	<i>rde-4(ne301)</i> vs <i>daf-2 rde-4</i>	8.03E-20
Wild-type (N2)	14.58	15.44	30				115	26		
<i>prg-1(n4357) Early</i>	14.00	16.38	34	-3.98%	6.09%	13.33%	125	31	N2 vs <i>prg-1(n4357) Early</i>	1
<i>prg-1(n4357) Stressed</i>	21.00	19.08	34	44.03%	23.58%	13.33%	110	4		
<i>prg-1(n4357) Stressed</i>	21.36	20.38	54	46.50%	31.99%	80.00%	125	23	N2 vs <i>prg-1(n4357) Stressed</i>	7.76E-04
<i>prg-1(n4357) Sterile</i>	26.50	26.95	54	81.76%	74.55%	80.00%	125	14	N2 vs <i>prg-1(n4357) Sterile</i>	1.58E-16
Wild-type (N2)	14.50	16.41	29				100	56		
<i>unc-13/+</i>	18.75	19.00	34	29.31%	15.79%	17.24%	100	47		
<i>prg-1(n4357) Early</i>	19.67	19.76	32	4.91%	4.00%	-5.88%	100	36	<i>prg-1(n4357) Early</i> vs <i>unc-13 -/+</i>	1
<i>prg-1(n4357) Stressed</i>	25.57	25.67	46	30.00%	29.89%	43.75%	100	40	<i>prg-1(n4357) Early</i> vs <i>prg-1(n4357) Stressed</i>	7.41E-05
<i>prg-1(n4357) Stressed</i>	26.30	25.13	45	33.71%	27.16%	40.63%	100	37	<i>prg-1(n4357) Early</i> vs <i>prg-1(n4357) Stressed</i>	4.30E-05
<i>prg-1(n4357)/unc-13 Early</i>	18.00	18.62	34	-4.00%	-2.02%	0.00%	100	40	<i>prg-1(n4357) -/+ Early</i> vs <i>unc-13 -/+</i>	1
<i>prg-1(n4357) -/+ Stressed</i>	24.36	25.32	43	35.32%	35.98%	26.47%	100	43	<i>prg-1(n4357) -/+ Early</i> vs <i>prg-1(n4357) -/+ Stressed</i>	1.50E-06
<i>prg-1(n4357) -/+ Stressed</i>	24.00	24.26	41	33.33%	30.13%	20.59%	100	62	<i>prg-1(n4357) -/+ Early</i> vs <i>prg-1(n4357) -/+ Stressed</i>	1
<i>prg-1(n4357) -/+ Stressed</i>	26.75	27.10	43	48.61%	45.56%	26.47%	100	59	<i>prg-1(n4357) -/+ Early</i> vs <i>prg-1(n4357) -/+ Stressed</i>	1
Wild-type (N2)	16.59	18.00	34				120	13		
<i>daf-16(mu86)</i>	13.06	14.67	26	-21.28%	-18.50%	-23.53%	120	6		
<i>daf-16(mu86)</i>	13.16	14.82	24	-20.68%	-17.67%	-29.41%	117	3		
<i>prg-1(n4357) daf-16(mu86) St</i>	17.26	17.61	24	32.16%	20.04%	-7.69%	120	3	<i>daf-16</i> vs <i>prg-1 daf-16 Early</i>	1.43E-03
<i>prg-1(n4357) daf-16(mu86) St</i>	11.31	13.14	22	-34.47%	-25.38%	-8.33%	120	1	<i>prg-1 daf-16 Early</i> vs <i>prg-1 daf-16 Stressed</i>	3.53E-12
<i>prg-1(n4357) daf-16(mu86) St</i>	8.59	9.73	18	-50.23%	-44.75%	-25.00%	115	2	<i>prg-1 daf-16 Stressed</i> vs <i>prg-1 daf-16 Sterile</i>	5.41E-13

Supplemental S1 Table. Analysis of longevity assays. Log rank analysis with Bonferroni correction comparing lifespan assays.