Haplopappus platylepis resin for pest control

1

2

3 **“*Haplopappus platylepis* (Asteraceae) resin: an adhesive trap for pest control of**
 4 **crawling arthropods, with antimicrobial potential”**

5

6 Cristian A. Villagra^{1*}¶, Verónica Macías-Marabolí^{1&}, Constanza Schapheer^{2&}, Jorge
 7 Bórquez ^{3&}, Mario J. Simirgiotis^{4&}, Javier Echeverría⁵, Marcia González-Teuber⁵,
 8 Alejandro Urzúa^{5*}¶.

9 ¹Instituto de Entomología, Universidad Metropolitana de Ciencias de la Educación,
 10 Santiago, Chile.

11 ² Laboratorio de Sistemática y Evolución de Plantas, Departamento de Silvicultura y
 12 Conservación de la Naturaleza, Universidad de Chile. Av. Santa Rosa 11315, La Pintana,
 13 Santiago, Chile

14 ³Laboratorio de Productos Naturales, Departamento de Química, Facultad de Ciencias
 15 Básicas, Universidad de Antofagasta, Chile.

16 ⁴Instituto de Farmacia, Facultad de Ciencias, Universidad Austral de Chile, Chile.

17 ⁵Laboratorio de Química Ecológica, Facultad de Química y Biología, Universidad de
 18 Santiago de Chile, Chile.

19

20 *Corresponding authors

21 cristian.villagra@umce.cl, alejandro.urzua@usach.cl

22 ¶ These authors are Joint Senior Authors

23 & These authors contributed equally to this work

Haplopappus platylepis resin for pest control

Abstract

The use of plant secondary metabolites has been incorporated as key part of integrated pest management and as an alternative to the use of pesticides. This may even be more relevant regarding domiciliary pest insects, capable of vectoring pathogens to humans. In these environments control is more difficult due to its possible effect on non-target organisms and human health. Here we evaluated the use of the resinous exudate of Chile's endemic bush *Haplopappus platylepis* (Asteraceae) as a sticky trap for crawling pest insects. We used *Blatta orientalis* Linneus (oriental cockroach), a cosmopolitan synanthropic pest, as test organism. We compared effectiveness on cockroach-trapping of *H. platylepis*' resin versus a commercially available sticky trap, and analyzed these two sticky substances using UHPLC-DAD-MS and GC-MS. We found that *H. platylepis* resin was as effective as the commercial adhesive on trapping *B. orientalis*. Plant resinous exudate was composed by a mixture of flavonoids, labdane diterpenoids and unsaturated fatty acids oxylipins, which are known for their antimicrobial and antioxidant properties. In contrast, the commercial sticky trap was rich in 1-bromohexadecane and 2-chlorocyclohexanol, which have been described as allergens and as potentially toxic to humans. Considering these findings, we suggest the use of the resinous extract of *H. platylepis* as an effective adhesive trapping method against pest cockroaches and possibly other crawling synanthropic arthropods cohabiting with humans. We highlight the importance of novel, non-toxic and eco-friendly products as strategies to be applied in the management of insect pests.

Keywords: synanthropic pest, integrated pest management, labdane terpenoids antimicrobial properties.

Introduction

Synthetic insecticides are controversial as they may represent a potential risk for human health and non-target organisms, beside its contribution to air and soil pollution [1–3]. Furthermore, controlling effects on pests can be rapidly ameliorated due to the evolution of resistance on target organisms [4–6]. This is especially concerning in the case of synanthropic arthropods related to vector-borne and zoonotic diseases inhabiting household, food storage facilities and hospitals [7,8]. These pests are hard to control due to their proximity to human-used spaces, restricting even more the use of various chemical control methods [9,10].

This is the case of several crawling pest arthropods including arachnids such as ticks [11,12], and insects belonging to: Hemiptera, like bedbugs [13] and triatomines [14] and Blattodea: such as pest cockroaches [15,16]. Synanthropic cockroaches [17] such as *Periplaneta americana* (Blattidae), *Blattella germanica* (Ectobiidae) and *Blatta orientalis* (Blattidae) have evolved associated to human-modified environments and usually act as vectors of allergens and diverse pathogenious microorganisms responsible for human diseases [18–21]. Thus, these insects represent a serious threat for human health [22].

The use of insecticides for the control of these insects has been extremely difficult, as cockroaches may become resistant to commonly-used chemical compounds [6]. Moreover, many insecticides at sublethal doses, are repellent to cockroaches and they are capable to avoid its contact [23]. In addition, some studies have shown that the use of pesticide against cockroach infestation paradoxically increases the level of the cockroach allergens Bla g 1 and Bla g 2, and possibly other allergens [24,25]. For example, adults of

B. germanica exposed to sub-lethal doses of the pesticide boric acid increase the production of the major allergen of Bla g 2 [25], which can lead to significant health problems, including asthma, eczemas skin reactions and allergic rhinitis [26]. Furthermore, it has been demonstrated the evolution of antibiotic resistance in pathogenic strains carried by *P. americana* and *B. germanica* collected from domiciliary and intensive care hospital facilities [27–30].

Therefore, in order to avoid the development of resistances either in the animal or their microbial counterparts, control strategies must combine the suppression of both crawling arthropod vectors and its associated pathogens. This approach must also consider current concerns on the safe use of pesticides for controlling difficult insect pests, especially regarding inhabited and food storing places [31,32]. In this work we studied the chemical composition of the resinous exudate of a Chilean endemic shrub *Haplopappus platylepis* Phil. (Asteraceae), focusing with particular interest on the presence of antimicrobial potential compounds. Coupled with this, we studied if adhesive extracts of this secretion can be used for the control of pest crawling arthropods, testing its adhesive function against the cosmopolitan pest cockroach *Blatta orientalis* Linnaeus, 1758 (Blattodea: Blattidae).

The use of plant-derived substances, capable of repelling and/or killing synanthropic pests, has been shown in several studies as an effective alternative to insecticides [33–35]. Among these, plant resins have demonstrated to be effective not only against several arthropods [36], but also in the combat against pathogenic microorganisms [37,38]. Moreover, the use of sticky traps could represent a more

Haplopappus platylepis resin for pest control

restrictible pesticide format in comparison with air-borne product, where spray drift unwanted consequences on human health have been reported [39].

In addition, adhesive traps can be displayed in refuge areas where airborne products can not easily reach [40], and reduce pest insects mechanically by catching them [41].

Moreover, these collected insects allow pest density monitoring [42]. This latter is a guide during decision-making for the most appropriate control measurement [43].

Considering the above-mentioned information, adhesive plant secretions such as resinous extractions may arise as suitable candidate for safe pest control of house pest and zoonotic vector insect [44].

Haplopappus platylepis, also known as “Devil’s Lollipop”, produces an adhesive resinous secretion covering its leaves and forming a natural sticky trap over floral buds [45]. This plant belongs to an asteraceous lineage presenting copious resin production with known antibacterial and antifungal properties, widely distributed in north and central Chile [38,46–48]. Previously, under field conditions, we showed that *H. platylepis*’ sticky exudate was capable of trapping several groups of insects that were fatally adhered during its blooming season [45]. In this study, we evaluated the potential use of *H. platylepis* inflorescence’s sticky exudate as an alternative adhesive trap for pest crawling insects. For these propose we tested it, in laboratory bioassays, on a common global household pest: the oriental cockroach *B. orientalis*. We compared its effectiveness on adhering pest cockroaches in relation to a commercial adhesive trap (Eco-opción®). In addition, we analyzed and compared the chemical composition of the sticky exudate of *H. platylepis* and the commercial adhesive trap using UHPLC-DAD-MS (ultra-high-performance liquid chromatography-diode array detector- mass spectrometry) and GC-

Haplopappus platylepis resin for pest control

MS (gas chromatography-mass spectrometry). Finally, we reviewed for bioactivity of compounds detected in both natural and commercial adhesives, in order to assess both their potential toxicity and harmful effects for humans, as well as any additional biological properties, especially focusing against pathogenic microorganisms.

Materials and methods

Plant material and trap extractions

Plant specimens of *Haplopappus platylepis* Phil. (Asteraceae) were determined following Klingenberg's monography for *Haplopappus* genus [49]. Floral buds of devil's lollypop were collected during March 2016 at Los Molles, Provincia de Petorca, V Region de Valparaíso, Chile (32°14'07.0"S 71°31'24"W) and at Punta Hueso, Pichidanguí, Provincia de Choapa, IV Region de Coquimbo, Chile (32°10'27"S 71°31'21"W). Samples were preserved until analysis at -10° C. Voucher specimens (SGO 166498) were deposited in the Herbarium of the "Museo Nacional de Historia Natural" (MCCN), Santiago, Chile.

The sticky exudate of *H. platylepis* was obtained by dipping fresh plant material (300 g) in cold CH₂Cl₂ (8 L) for 48 h, following Urzúa 2004's method[50]. The resulting extract was filtered through a cotton layer and concentrated to a sticky residue (36 g, 12%) Commercial adhesive trap used was Eco-Opción® (Anasac Corporation, Santiago, Chile), sticky trap offered for the control of cursorial domiciliary pest such as ants, cockroaches and spiders. Each unit brings four 29.6x23.3 cm cardboard sticky traps with a total adhesive surface of 11x13 cm. The adhesive mixture from the cardboard was removed with a spatula and followed above-mentioned procedure for extraction. Extracts

of both natural (*H. platylepis* inflorescence's resin) and commercial sticky traps were kept under 4°C for further chemical analyses (see below).

Insects

Oriental cockroaches used in this work were obtained from a population maintained in our laboratory since year 2014. Further specimens used for this study were collected from locations in San Miguel, Santiago, Metropolitan Region, Chile (33°29'54"S 70°38'42"W). For taxonomic identification a general key for cosmopolitan and pest cockroaches present in Chile was used [51]. Insects were kept in captivity under laboratory conditions (20°-25°C and 40%-50% humidity) in 120x50x15 cm plastic rearing boxes, fed with dog food (MasterDog Adult ®) and water *ad libitum*, at Instituto de Entomología, UMCE. *Blatta orientalis* from both sexes were used for sticky-trapping bioassays (with body lengths among 5 to 25 mm, measured dorsally from head to last abdominal segments).

Trapping bioassays

Two treatments and one control were defined for the experiment. Treatments corresponded to cardboard surfaces (40x13cm) painted either with *H. platylepis* resinous exudate or with the commercial trap's adhesive. For control, a cardboard surface (40x13cm) with no adhesive mixture added was used. Each of these options was presented individually in the experimental arena. For this, the cardboard section was placed in the center of the horizontal space inside the arena, fixing its position with double-contact tape (Fig. 1). For each replicate 10 individuals from different sizes

(measured as explained above) were placed in the experimental arena habituation area (Fig. 1), a subdivision of the box from where insect were released without contact them directly. For each trial we lifted the opening section of the habituation area and gave light pulses (10s) during three instances of the experiment: 0, 180 and 360s. At each of these pulses cockroaches tended to leave the habituation area and run to the other extreme of the box crossing the cardboard section. Total time of each test was 6min. After this period, for each treatment and control the number of individuals found attached to the cardboard was counted. Trapped insects were ultimately sacrificed by applying cold temperature (-10 °C). For each of these alternatives we repeated this test 10 times. Before using the experimental arena for each trial, this was cleaned with ethanol (95%), distilled water and dried in order to remove any chemical cue. The response variable was the proportion of insects trapped in each trial for each treatment. As data did not meet the criterion of normality distribution (Hammer, 1999), it was analyzed with a non-parametric analysis of variance Kruskal-Wallis followed by *post hoc* Mann Whitney test. In order to determine if *H. platylepis* inflorescence's resin and the commercial sticky trap are equally efficient trapping cockroaches of different sizes (seven ranges: from 5 to 7; 8 to 10; 11 to 13; 14 to 16; 17 a 19; 20 to 22 and 23 to 25mm), insect proportion per range, captured in both traps, was compared. This was analyzed by using a Chi square test for two proportions [53]. All analyses were done with the PAST Paleontological Statistic, version 3.15.

Fig. 1. Bioassay setup A. Experimental arena: a. Background pattern. b. Treatment Area, c. Darkened walls, d: Hatch e: Habituation cubicle. f: Arena's door. **B.** Sticky trap made

Haplopappus platylepis resin for pest control

- 186 with *H. platylepis* resin (upper picture) and Eco-opción® adhesive (lower picture).
- 187 Trapped roaches are highlighted with arrows.

Chemicals

UHPLC-MS solvents, LC-MS formic acid and reagent grade chloroform were from Merck (Santiago, Chile). Ultrapure water was obtained from a Millipore water purification system (Milli-Q Merck Millipore, Chile). HPLC standards, (kaempferol, quercetin, isorhamnetin, eriodictyol, luteolin, apigenin, naringenin, all standards with purity higher than 95 % by HPLC) were purchased either from Sigma Aldrich (Saint Louis, Mo, USA), ChromaDex (Santa Ana, CA, USA), or Extrasynthèse (Genay, France).

UHPLC-DAD-MS analyses

Chemical resinous components were analyzed by using ultra-high-performance liquid chromatography-diode array detector-tandem mass spectrometry (UHPLC-DAD-MS). UHPLC-DAD-MS analysis was performed using a Thermo Scientific Dionex Ultimate 3000 UHPLC system hyphenated with a Thermo Q exactive focus machine as it was reported by Simirgiotis et al. (2016). 5 mg of the resinous exudate were dissolved in 2 mL of methanol and filtered with a PTFE filter for a final injection of 10 µL into the instrument. Measurements were done as previously reported by Simirgiotis et al. (2016). The generation of molecular formulas was performed using high resolution accurate mass analysis (HRAM) and matching with the isotopic pattern. Lastly, analyses were confirmed using MS/MS data and comparing the fragments found with the literature.

LC and MS parameters

Haplopappus platylepis resin for pest control

Liquid chromatography was performed using an UHPLC C18 column (Acclaim, 150 mm × 4.6 mm ID, 2.5 µm, Thermo Fisher Scientific, Bremen, Germany) operated at 25 °C. The detection wavelengths were 254, 280, 330 and 354 nm, and DAD was recorded from 200 to 800 nm for peak characterization. Mobile phases were 1 % formic aqueous solution (A) and acetonitrile (B). The gradient program time (min, % B) was: (0.00, 5); (5.00, 5); (10.00, 30); (15.00, 30); (20.00, 70); (25.00, 70); (35.00, 5) and 12 minutes for column equilibration before each injection. The flow rate was 1.00 mL min⁻¹, and the injection volume was 10 µL. Standards and the resin extract dissolved in methanol were kept at 10°C during storage in the autosampler. The HESI II and Orbitrap spectrometer parameters were optimized as previously reported [54].

GC-MS analyses

Chemical composition of the commercial adhesive trap was analyzed by gas chromatography-mass spectrometry (GC-MS). GC-MS analysis was performed using a Thermo Scientific Trace GC Ultra linked to an ISQ quadrupole mass spectrometric detector with an integrated data system (Xcalibur 2.0, Thermo Fisher Scientific Inc., Waltham, MA, USA), equipped with a capillary column (Rtx-5 MS, film thickness 0.25 µm, 60 x 0.25 mm, Restek Corporation, Bellefonte, PA, USA) The operating conditions were as follows: on-column injection; injector temperature, 250 °C; detector temperature, 280 °C; carrier gas, He at 1.25 mL/min; oven temperature program: 40 °C increase to 260 °C at 4 °C/min, and then 260 °C for 5 min. The mass spectra were obtained at an ionization voltage of 70 eV. Recording conditions employed a scan time of 1.5 s and a mass range of 40 to 400 amu. The identification of compounds in the chromatographic

profiles was achieved by comparison of their mass spectra with a library database (NIST08, NIST, Gaithersburg, MD, USA) and by comparison of their calculated retention indices with those reported in the literature [55] for the same type of column.

Results

Trapping bioassays

The proportion of insects found over the cardboards was statistically different among treatments ($H(X^2) = 19.43$, $p < 0.001$, Kruskal-Wallis, Fig. 2A). *H. platylepis* inflorescence's sticky exudate and the commercial sticky trap differed with statistical significance from control clean cardboard (in both cases: U Mann-Whitney pairwise, $p < 0.001$). However, no differences were found in post hoc test for the total number of insects attached on cardboards between the *H. platylepis*' resin and the commercial sticky trap (U Mann-Whitney pairwise, $p = 0.691$). When the proportion of cockroaches trapped by *H. platylepis*' sticky exudate and by the commercial sticky trap for each size range was compared, no statistical differences were found between natural and commercial sticky traps ($X^2 = 1.57$, $p = 0.211$) (Fig. 2B).

Fig. 2. Cockroach adhesion results. A. Mean and 1SE for the proportion of *B. orientalis* found over the cardboard (Y axis) painted with: *H. platylepis* resin (green), Eco-opción® commercial adhesive (red) and control (clean cardboard, black) obtained from 10 replicates each (X axis). Different letters correspond to statistical differences after post hoc test at $p < 0,05$. B. Proportion of cockroaches trapped (Y axis) by either *H. platylepis* resin (light grey) or Eco-opción® commercial adhesive (dark grey) for each insect size

range (X axis). No statistical differences were found for each pair compared.

Chemical analyses

The data-dependent scan experiment was very useful for the identification of unknown compounds since it provides high resolution and accurate mass product ion spectra from precursor ions that are unknown beforehand within a single run. Combining data-dependent scans and MSⁿ experiments, phytochemicals were tentatively identified in *H. platylepis* including simple phenolic acids flavones, flavanones, fatty acids, and labdane diterpenoids. UHPLC Q-orbitrap mass spectrometry analysis of *H. platylepis* sticky exudate showed the presence of twenty seven metabolites in the chromatograms (Fig. 3) including: 7 flavonoids (peaks **5**, **6**, **8-10**, **15** and **16**), 3 phenolic acids (peaks **1-3**), 8 fatty acids (Peaks **4**, **7**, **13**, **14**, **18**, **21**, **22** and **25**), and 9 labdane terpenoids (peaks **11**, **12**, **17**, **19,20**, **23**, **24**, **26**, and **27**). The detailed identification is explained below (Table 1, Figs. 4 and 1S).

Fig. 3: UHPLC chromatograms A. TIC (total ion current, negative mode) and B. UV at 280 nm, of *H. platylepis* resin.

Fig. 4: Proposed biogenetic relationships between labdane diterpenoids.

Haplopappus platylepis resin for pest control

Table 1: High resolution UHPLC PDA-Q-orbitrap identification of metabolites in *Haplopappus platylepis* resin.

Peak #	Retention time (min)	UV max (nm)	Tentative identification	Elemental composition [M-H]	Theoretical mass (m/z)	Measured mass (m/z)	Accuracy (δppm)	MS ⁿ ions (δppm)
1	11.43	-	12-Hydroxyjasmonate	C ₁₂ H ₁₇ O ₄ ⁻	225.11276	225.11313	4.27	
2	12.93	-	Dihydroxyphaseic acid	C ₁₅ H ₂₁ O ₅ ⁻	281.13953	281.13945	-0.28	
3	13.71	325	Ferulic acid	C ₁₀ H ₉ O ₄ ⁻	193.05063	193.05040	-1.19	
4	18.76	285	Trihydroxyoctadecaenoic acid	C ₁₈ H ₃₃ O ₅ ⁻	329.23335	329.23367	0.97	
5	19.05	255, 354	7,3'-dimethoxyquercetin	C ₁₇ H ₁₃ O ₇ ⁻	329.06668	329.06702	1.03	
6	19.26	287	Hesperetin	C ₁₆ H ₁₃ O ₆ ⁻	301.07176	301.07199	0.76	160.84154, 135.04446
7	19.38	285	Trihydroxyoctadecadienoic acid	C ₁₈ H ₃₁ O ₅ ⁻	327.21770	327.21799	0.89	
8	19.56	287	5,3',5'-trihydroxy-3,7,4'-trimethoxyflavanone	C ₁₇ H ₁₅ O ₇ ⁻	331.08261	331.08233	1.22	
9	20.02	255-354	5,3'-dihydroxy-3,7,4'-trimethoxyflavone	C ₁₈ H ₁₅ O ₈ ⁻	343.08233	343.08267	1.25	313.03580 (C ₁₆ H ₉ O ₇ ⁻ , [M-OCH ₃ -CH ₃])
10	20.04	255-354	7, 3', 5'-trimethoxymyricetin	C ₁₈ H ₁₅ O ₈ ⁻	359.07724	359.07748	0.58	285.04031 (C ₁₅ H ₉ O ₆ ⁻ , kaempferol)
11	20.07	289	Dehydropinifolic acid	C ₂₀ H ₃₃ O ₄ ⁻	337.23843	337.23886	1.28	
12	20.10	289	Pinifolic acid (labd-8(20)-en-15,18-dioic acid)	C ₂₀ H ₃₁ O ₄ ⁻	335.22278	335.22287	0.98	
13	21.13	305	Trihydroxyheneicosahexaenoic acid	C ₂₁ H ₂₉ O ₅ ⁻	361.20205	361.20242	1.02	
14	21.36	303	dihydroxyeicosapentaenoic acid	C ₂₀ H ₂₉ O ₄ ⁻	333.20713	333.20740	0.90	273.18622 (C ₁₈ H ₂₅ O ₂ ⁻); [M ⁻ - (CO ₂ - CH ₃ - H)]
15	21.71	255-354	3,7-dimethoxyquercetin	C ₁₇ H ₁₃ O ₇ ⁻	329.06668	329.06705	1.12	
16	21.96	255-354	3,5, dihydroxy-3',4',7-trimethoxyflavone	C ₁₈ H ₁₅ O ₈ ⁻	343.08233	343.08273	1.17	313.03467 (C ₁₆ H ₉ O ₇ ⁻ , (M ⁻ -OCH ₃ - CH ₃))

Haplopappus platylepis resin for pest control

17	22.12	-	(epi) Pinifolic acid	C ₂₀ H ₃₁ O ₄ ⁻	335.22278	335.22287	0.54	317.21219 (C ₂₀ H ₂₉ O ₃ ⁻ ; [M ⁻ - H ₂ O]; 273.18652 (C ₁₈ H ₂₅ O ₂ ⁻)
18	22.87	302	Tetrahydroxytetrahexaenoic acid	C ₂₄ H ₃₅ O ₆ ⁻	419.24423	419.24391	3.37	319.22806
19	22.92	289	18-hydroxy-8(17)en-15-labdanoic acid	C ₂₀ H ₃₃ O ₃ ⁻	321.24377	321.24377	0.00	
20	23.94	289	Dehydropinifolic acid isomer	C ₂₀ H ₃₃ O ₄ ⁻	337.23843	337.23886	1.28	
21	24.25	308	Hydroxyeicosapentaenoic acid	C ₂₀ H ₂₉ O ₃ ⁻	317.21222	317.21255	1.04	
22	22.87	303	Hydroxyeicosatetraenoic acid	C ₂₀ H ₃₁ O ₃ ⁻	319.22787	319.22821	1.07	
23	25.40	289	13-en-Pinifolic acid methyl ester	C ₂₁ H ₃₁ O ₄ ⁻	347.22278	347.22311	1.04	273.18616 (C ₁₈ H ₂₅ O ₂ ⁻); 239.26134 (C ₁₆ H ₃₁ O ⁻)
24	25.78	289	Pinifolic acid methyl ester	C ₂₁ H ₃₃ O ₄ ⁻	349.23843	349.23880	1.06	
25	25.88	306	Trihydroxydocosaheptaenoic acid	C ₂₂ H ₃₁ O ₅ ⁻	375.21770	375.21823	1.41	
26	25.99	289	18-acetyl-13,8 (17)dien-15-labdanoic acid	C ₂₂ H ₃₃ O ₄ ⁻	361.23843	361.23877	0.94	
27	26.56	289	18-acetyl-8(17)en-15-labdanoic acid	C ₂₂ H ₃₅ O ₄ ⁻	363.25408	363.25443	1.13	321.24319 (C ₁₈ H ₂₅ O ₂ ⁻ ; M ⁻ - H ₂ O)

280

281 Flavonoids

282 Peak **15** with a [M-H]⁻ ion at *m/z* 329.06705 was identified 3,7-dimethoxyquercetin
 283 (C₁₇H₁₃O₇⁻) and peak **5** with an ion at *m/z* 329.06702 as its isomer: 7,3'-
 284 dimethoxyquercetin (Table 1). Peak **9** with a [M-H]⁻ ion at *m/z* 343.08276 was identified
 285 as the trimethoxylated flavonoid 5,3'-dihydroxy-3,7,4'-trimethoxyflavone (C₁₈H₁₅O₈⁻),
 286 while peak **10** with a [M-H]⁻ ion at *m/z* 359.07745 as 7,3',5'-trimethoxymyricetin
 287 (C₁₈H₁₅O₈⁻). Peak **16** with a pseudomolecular ion at *m/z* 343.08273 was identified as 3,5-
 288 dihydroxy-3',4',7-trimethoxyflavone (C₁₈H₁₅O₈⁻). The flavanone hesperetin, peak **6**, have

Haplopappus platylepis resin for pest control

been previously reported as main component in extracts of several *Nolana* species by some of us (Simirgiotis, et al., 2015) and its HR-MS ($C_{16}H_{13}O_6^-$) and UV data matched the one obtained in our chromatograms (m/z : 301.07176). Another flavanone, peak **8** with a $[M-H]^-$ ion at m/z 331.08261 was identified as 5,3',5'-trihydroxy-3,7,4'-trimethoxyflavanone ($C_{17}H_{15}O_7^-$).

Phenolic acids

The examination of the chromatograms revealed the presence of 3 phenolic acids: dihydroxyphaseic acid (peak **2**, ion at m/z 281.13945, $C_{15}H_{21}O_5^-$) [56], ferulic acid (peak **1**, m/z 193.05040) and 12-hydroxy jasmonate (peak **3**, m/z 225.11313) [57].

Fatty acids

Several peaks were tentatively identified as the dietary antioxidant polyhydroxylated unsaturated fatty acids known as oxylipins [58,59], antioxidant fatty acids. Peak **4** with a $[M-H]^-$ ion at m/z 329.23367 was identified as trihydroxy-octadecenoic acid ($C_{18}H_{33}O_5^-$), and peak **7** as its diene derivative ($C_{18}H_{31}O_5^-$), as previously reported by some of us from Keule fruits [59]. Peak **13** with a pseudomolecular ion at m/z 361.20242 was identified as trihydroxyheneicosahexaenoic acid ($C_{21}H_{29}O_5^-$). Peak **14** with a $[M-H]^-$ ion at m/z 333.20743 was identified as a dihydroxyeicosapentaenoic acid ($C_{20}H_{29}O_4^-$) while peak **18** with a $[M-H]^-$ ion at m/z 419.24391 was identified as dihydroxytetracosatrienoic acid ($C_{24}H_{35}O_6^-$) [58]. Peak **21** and **22** were identified as hydroxyeicosapentaenoic acid and hydroxyeicosatetraenoic acid ($C_{20}H_{29}O_3^-$) and ($C_{20}H_{31}O_3^-$), respectively. Finally, peak **25**

Haplopappus platylepis resin for pest control

with a $[M-H]^-$ ion at m/z 375.21823 was identified as trihydroxydocosaheanoic acid ($C_{22}H_{31}O_5^-$).

Labdane terpenoids

Labdane terpenoids corresponded to derivatives of pinifolic acid (labd-8(20)-en-15,18-dioic acid, peak **12**, $C_{20}H_{36}O_3$) [60] most of them reported for the first time in this species. Thus, peak **11** with a $[M-H]^-$ ion at m/z 337.23886 was identified as its hydrogenated derivative of dehydropinifolic acid ($C_{20}H_{33}O_4^-$) and peak **17** with a $[M-H]^-$ ion at m/z 335.22296 as an isomer of pinifolic acid ($C_{20}H_{31}O_4^-$), probably the epimer at C-4 of the latter. Peak **24** was identified as pinifolic acid methyl ester ($C_{21}H_{33}O_4^-$) and peak **23** as its derivative 13-en-pinifolic acid methyl ester ($C_{21}H_{31}O_4^-$). Peak **20** with a $[M-H]^-$ ion at m/z 337,23886 was identified as pinifolic acid derivative ($C_{20}H_{33}O_4^-$). Three compounds were identified as labdanoic acid derivatives [61]. Thus, peak **19** with a $[M-H]^-$ ion at m/z 321.24377 was identified as 18-hydroxy-8(17)en-15-labdanoic acid ($C_{20}H_{33}O_3^-$), Peak **27** with a $[M-H]^-$ ion at m/z 363.25449 was identified as 18-acetyl-8(17)en-15-labdanoic acid ($C_{22}H_{35}O_4^-$) and peak **26** as its diene derivative ($C_{22}H_{33}O_4^-$) (Fig. 4).

Components identified in the commercial sticky trap

GC-MS identified only two compound in the commercial sticky trap as: 1-bromohexadecane and 2-chlorocyclohexanol.

Discussion

The aim of this study was to compare the effectiveness of a natural sticky trap against a commercial one in capturing cockroaches by adhesion. In addition, the chemical composition of both traps was analyzed in order to estimate potential harmful effects for humans as well as potential antimicrobial chemical compounds. Our results provide evidence that the natural sticky trap of *H. platylepis* was as effective as the commercial one on trapping pest cockroaches. Considerable differences, however, were found in the chemical composition between the natural and the commercial trap. Whereas the former was rich in plant-derived antimicrobial compounds, the latter was rich in halogenated compounds, whose potential toxic effects for humans have been previously reported.

The *H. platylepis* sticky exudate seems to offer multiple benefits in relation to its use for controlling synanthropic pest crawling insect, such as cockroaches. First, because of its stickiness, it resulted as effective as the commercial trap for capturing cursorial insects, and second, due to its chemical composition rich in antibacterial compounds [62], it shows a further potential for controlling pest arthropod-borne transmitted pathogens. As far as we know, most of the compounds identified for *H. platylepis* resin are reported for the first time in this species. Antibacterial properties of *H. platylepis* sticky exudate can be associated with the phytochemical families detected in the mixture [62]. For instance, flavonoids have shown a wide-spectrum of inhibitory activity against a variety of human pathogens, including antibiotic-resistant Gram-positive and Gram-negative bacteria, viruses and fungus [62–66]. Labdane diterpenoids are also well known as antimicrobials [67,68]. It has been proved that the presence of a carboxylic acid in the C-15 position, which acted as a hydrogen-bond donor (HBD), is essential for the antibacterial activity of *ent*-labdanes [64]. Furthermore, derivatives of pinifolic acid,

which were characterized in the *H. platylepis* sticky exudate, showed this main structural characteristic of labdanes. In addition, pinifolic acid has been previously reported as an effective compound in the treatment of leishmaniasis [69], a global insect-borne disease related to trypanosomes [70]. Long-chain polyunsaturated fatty acids, which were also abundant in *H. platylepis* resin, including oxylipins, have been widely tested for its antimicrobial activity [71–75]. Therefore, further functions of chemical compounds found in *H. platylepis*' resinous exudate expand the potential value of this plant-derived adhesive to act as a control against various vectoring-disease scenarios.

Synanthropic crawling arthropods are usual carriers of several human pathogens [76]. In the case of *B. orientalis*, it has been described to bear several human pathogenic bacteria genera such as *Mycobacteria*, *Klebsiella*, *Staphylococcus*, *Escherichia* and *Enterobacter* [77,78]. Therefore, the occurrence of compounds with anti-microbial functions in the sticky exudate of *H. platylepis* may synergistically contribute as an integrative pest control method, not only directly affecting the insect pests but also its associated pathogenic microorganisms. The commercial sticky trap, in contrast, is poor in its chemical composition and lacks antimicrobial compounds. 1-Bromohexadecane (**1**) and 2-chlorocyclohexanol (**2**) were the only two compounds identified on the commercial trap. Both are known as halogenated compounds. Based on Globally Harmonized System of Classification and Labeling of Chemicals (GHS), both are characterized as irritant for humans, due to the fact that these compounds induce skin corrosion (category 2), respiratory tract irritation (category 3) as well as severe eye irritation (category 2A) (European Chemical Agency- ECHA, 2017). This chemical profile suggests that this commercial trap would not be innocuous for human health; nevertheless, it is

commercially offered as an eco-friendly option. Our results highly suggest that *H. platylepis* sticky exudate may be a suitable alternative for controlling synanthropic crawling insects, including cockroaches, at low cost and with additional benefits such as potential antimicrobial properties. These virtues of *H. platylepis* sticky exudate trap fit the current needs and trends in pest control, where several methodologies must be integrated in order to generate novel alternatives in consideration of human and environmental health [79]. Further research is needed in order to test this adhesive resin in other formats for insect trapping as well as to evaluate its effectiveness against other pest insects. For instance, resinous materials have been considered among the updated alternatives for controlling domiciliary termites [44].

Conclusions

Results here demonstrated that devil's lollypop resin is a natural source of terpenoids and flavonoids with potential applications as insecticide and antibacterial. Using UHPLC-DAD-MS we have identified 27 secondary metabolites in *H. platylepis*' resin. Most of which, as far as we know, are reported here for the first time. Many of these compounds are flavones, flavanones, phenolic acids, fatty acids, and labdane terpenoids. This chemical knowledge may be helpful for further research on *H. platylepis* and its applications in biomedicine and pest and pathogens control industry. In conclusion, this plant is a rich source of phenolic and clerodane compounds with insecticide and antibacterial activity that may be used as an effective biocontrol agent against zoonotic crawling insects and their associate microorganisms .

Supporting Information

Fig. A.1: Full HR-MS spectra and structures of compounds 3 (a), 9 (b), 10 (c), 12 (d), 14 (e), 22 (f), 23 (g), 26 (h) and 27 (i).

Acknowledgments

We thank Catherine Cabello and Angel Olguín for help during fieldwork and laboratory work.

Funding

This research was funded by FONDECYT Iniciación No. 11100109 and CONICYT Inserción No. 79100013 granted to Cristian Villagra, RSG N° 21286-2 to Constanza Schapheer, Proyecto Fortalecimiento USACH USA1799-UA253010, Universidad de Santiago de Chile granted to Alejandro Urzúa, Javier Echeverria and Marcia Gonzalez, and CONICYT PAI/ACADEMIA No. 79160109 to Javier Echeverria.

References

1. Nicolopoulou-Stamati P, Maipas S, Kotampasi C, Stamatis P, Hens L. Chemical Pesticides and Human Health: The Urgent Need for a New Concept in Agriculture. Front Public Heal. Frontiers; 2016;4: 148. doi:10.3389/fpubh.2016.00148
2. Bernardes MFF, Pazin M, Pereira LC, Dorta DJ. Impact of Pesticides on

- 424 Environmental and Human Health. Toxicology Studies - Cells, Drugs and
425 Environment. InTech; 2015. doi:10.5772/59710
- 426 3. Finizio A, Villa S. Environmental risk assessment for pesticides. Environ Impact
427 Assess Rev. InTech; 2002;22: 235–248. doi:10.1016/S0195-9255(02)00002-1
- 428 4. Metcalf RL. Insect resistance to insecticides. Pestic Sci. 1989;26: 333–358.
429 doi:10.1002/ps.2780260403
- 430 5. Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease.
431 Annu Rev Entomol. 2000;45: 371–391. doi:10.1146/annurev.ento.45.1.371
- 432 6. Naqqash MN, Gökçe A, Bakhsh A, Salim M. Insecticide resistance and its
433 molecular basis in urban insect pests. Parasitology Research. 2016. pp. 1363–1373.
434 doi:10.1007/s00436-015-4898-9
- 435 7. Bonnefoy X, Kampen H, Sweeney K. Public Health Significance of Urban Pests.
436 Public health significance of urban pests. 2008. doi:10.1016/S1473-
437 3099(09)70222-1
- 438 8. Chavasse DC, Yap HH. Chemical Methods for the Control of Vectors and Pests of
439 Public Health Importance. 1997. pp. 16–21.
- 440 9. Menasria T, Moussa F, El-Hamza S, Tine S, Megri R, Chenchouni H. Bacterial
441 load of German cockroach (*Blattella germanica*) found in hospital environment.
442 Pathog Glob Health. 2014;108: 141–7. doi:10.1179/2047773214Y.0000000136
- 443 10. Rivault C, Cloarec A, Le Guyader A. Bacterial load of cockroaches in relation to
444 urban environment. Epidemiol Infect. 1993;110: 317–325.
445 doi:10.1017/S0950268800068254
- 446 11. Buckingham SC. Tick-borne disease. Clinical Infectious Disease, Second Edition.

2015. pp. 797–799. doi:10.1017/CBO9781139855952.136
12. LaSala PR, Holbrook M. Tick-borne flaviviruses. Clinics in Laboratory Medicine. 2010. pp. 221–235. doi:10.1016/j.cll.2010.01.002
13. Delaunay P, Blanc V, Del Giudice P, Levy-Bencheton A, Chosidow O, Marty P, et al. Bedbugs and infectious diseases. Clinical Infectious Diseases. 2011. pp. 200–210. doi:10.1093/cid/ciq102
14. Rassi A, Rassi A, Marin-Neto JA. Chagas disease. The Lancet. 2010. pp. 1388–1402. doi:10.1016/S0140-6736(10)60061-X
15. Cloarec A, Rivault C, Fontaine F, Le Guyader A. Cockroaches as carriers of bacteria in multi-family dwellings. Epidemiol Infect. 1992;109: 483–490. doi:10.1017/S0950268800050470
16. Schapheer C, Sandoval G, Villagra C. Pest Cockroaches May Overcome Environmental Restriction Due to Anthropization. J Med Entomol. 2018;In press.
17. Nasirian H. Infestation of cockroaches (Insecta: Blattaria) in the human dwelling environments: A systematic review and meta-analysis. Acta Tropica. 2017. pp. 86–98. doi:10.1016/j.actatropica.2016.12.019
18. Kassiri H, Kassiri A, Kazemi S. Investigation on American cockroaches medically important bacteria in Khorramshahr hospital, Iran. Asian Pacific J Trop Dis. 2014;4: 201–203. doi:10.1016/S2222-1808(14)60505-3
19. Fakoorziba MR, Shahriari-Namadi M, Moemenbellah-Fard MD, Hatam GR, Azizi K, Amin M, et al. Antibiotics susceptibility patterns of bacteria isolated from American and German cockroaches as potential vectors of microbial pathogens in hospitals. Asian Pacific J Trop Dis. Elsevier; 2014;4: S790–S794.

- 470 20. Arruda LK, Ferriani VPL, Vailes LD, Pomés A, Chapman MD. Cockroach
471 allergens: environmental distribution and relationship to disease. Curr Allergy
472 Asthma Rep. Springer; 2001;1: 466–473.
- 473 21. Burgess NR, McDermott SN, Whiting J. Aerobic bacteria occurring in the hind-gut
474 of the cockroach, *Blatta orientalis*. J Hyg (Lond). 1973;71: 1–7.
475 doi:10.1017/S0022172400046155
- 476 22. Arruda LK, Vailes LD, Ferriani VPL, Santos ABR, Pomés A, Chapman MD.
477 Cockroach allergens and asthma. Journal of Allergy and Clinical Immunology.
478 2001. pp. 419–428. doi:10.1067/mai.2001.112854
- 479 23. Wooster MT, Ross MH. Sublethal responses of the German cockroach to vapors of
480 commercial pesticide formulations. Entomol Exp Appl. Wiley Online Library;
481 1989;52: 49–55.
- 482 24. Chew GL, Burge HA, Dockery DW, Muilenberg ML, Weiss ST, Gold DR.
483 Limitations of a home characteristics questionnaire as a predictor of indoor
484 allergen levels. Am J Respir Crit Care Med. Am Thoracic Soc; 1998;157: 1536–
485 1541.
- 486 25. Zhang YC, Perzanowski MS, Chew GL. Sub-lethal exposure of cockroaches to
487 boric acid pesticide contributes to increased Bla g 2 excretion. Allergy. Wiley
488 Online Library; 2005;60: 965–968.
- 489 26. Pomés A, Arruda LK. Investigating cockroach allergens: aiming to improve
490 diagnosis and treatment of cockroach allergic patients. Methods. Elsevier;
491 2014;66: 75–85.
- 492 27. Pai HH, Chen WC, Peng CF. Isolation of bacteria with antibiotic resistance from

493 household cockroaches (*Periplaneta americana* and *Blattella germanica*). *Acta*
494 *Trop.* 2005;93: 259–265. doi:10.1016/j.actatropica.2004.11.006

495 28. Bouamama L, Sorlozano A, Laglaoui A, Lebbadi M, Aarab A, Gutierrez J.
496 Antibiotic resistance patterns of bacterial strains isolated from *Periplaneta*
497 *americana* and *Musca domestica* in Tangier, Morocco. *J Infect Dev Ctries.* 2010;4:
498 194–201.

499 29. Moges F, Eshetie S, Endris M, Huruy K, Muluye D, Feleke T, et al. Cockroaches
500 as a Source of High Bacterial Pathogens with Multidrug Resistant Strains in
501 Gondar Town, Ethiopia. *Biomed Res Int.* 2016;2016. doi:10.1155/2016/2825056

502 30. Tilahun B, Worku B, Tachbele E, Terefe S, Kloos H, Legesse W. High load of
503 multi-drug resistant nosocomial neonatal pathogens carried by cockroaches in a
504 neonatal intensive care unit at Tikur Anbessa specialized hospital, Addis Ababa,
505 Ethiopia. *Antimicrob Resist Infect Control.* 2012;1. doi:10.1186/2047-2994-1-12

506 31. Quarles W. IPM reduces pesticides, cockroaches, and asthma. *IPM Pract.* 2009;31:
507 8.

508 32. Schal C, Hamilton R. Integrated suppression of synanthropic cockroaches. *Annu*
509 *Rev Entomol.* 1990;35: 521–551.

510 33. Ferrero A, Sánchez Chopa C, Werdin González J, Alzogaray R. Repellence and
511 toxicity of *Schinus molle* extracts on *Blattella germanica*. *Fitoterapia.* 2007;78:
512 311–314. doi:10.1016/j.fitote.2006.11.021

513 34. Sánchez Chopa C, Alzogaray R, Ferrero A. Repellency Assays with *Schinus molle*
514 var. *areira* (L.) (Anacardiaceae) Essential Oils against *Blattella germanica* L.
515 (Blattodea : Blattellidae). *BioAssay.* 2006;1: 1–3.

Haplopappus platylepis resin for pest control

- 516 35. Yoon C, Kang S-H, Yang J-O, Noh D-J, Indiragandhi P, Kim G-H. Repellent
517 activity of citrus oils against the cockroaches *Blattella germanica*, *Periplaneta*
518 *americana* and *P. fuliginosa*. *J Pestic Sci.* 2009;34: 77–88.
519 doi:10.1584/jpestics.G07-30
- 520 36. Castella G, Chapuisat M, Moret Y, Christe P. The presence of conifer resin
521 decreases the use of the immune system in wood ants. *Ecol Entomol.* 2008;33:
522 408–412. doi:10.1111/j.1365-2311.2007.00983.x
- 523 37. Shuaib M, Ali A, Ali M, Panda B, Ahmad M. Antibacterial activity of resin rich
524 plant extracts. *J Pharm Bioallied Sci.* 2013;5: 265. doi:10.4103/0975-7406.120073
- 525 38. Urzúa A, Jara F, Tojo E, Wilkens M, Mendoza L, Rezende MC. A new
526 antibacterial clerodane diterpenoid from the resinous exudate of *Haplopappus*
527 *uncinatus*. *J Ethnopharmacol.* 2006;103: 297–301.
- 528 39. Lam J, Sutton P, Kalkbrenner A, Windham G, Halladay A, Koustas E, et al. A
529 systematic review and meta-analysis of multiple airborne pollutants and autism
530 spectrum disorder. *PLoS One.* 2016;11. doi:10.1371/journal.pone.0161851
- 531 40. Hewitt AJ. Drift filtration by natural and artificial collectors: A literature review.
532 Macon, MO, USA.; 2001.
- 533 41. Smith LM, Appel AG. Comparison of several traps for catching German
534 cockroaches (Dictyoptera: Blattellidae) under laboratory conditions. *J Econ*
535 *Entomol. BioOne;* 2008;101: 151–158.
- 536 42. Nalyanya G, Gore JC, Linker HM, Schal C. German cockroach allergen levels in
537 North Carolina schools: comparison of integrated pest management and
538 conventional cockroach control. *J Med Entomol.* 2014;46: 420–427.

- 539 43. Barzman M, Bàrberi P, Birch ANE, Boonekamp P, Dachbrodt-Saaydeh S, Graf B,
540 et al. Eight principles of integrated pest management. *Agronomy for Sustainable*
541 *Development*. 2015. pp. 1199–1215. doi:10.1007/s13593-015-0327-9
- 542 44. Verma M, Sharma S, Prasad R. Biological alternatives for termite control: A
543 review. *International Biodeterioration and Biodegradation*. 2009. pp. 959–972.
544 doi:10.1016/j.ibiod.2009.05.009
- 545 45. Villagra CA, Meza AA, Urzúa A. Differences in arthropods found in flowers
546 versus trapped in plant resins on *Haplopappus platylepis* Phil. (Asteraceae): Can
547 the plant discriminate between pollinators and herbivores? *Arthropod Plant*
548 *Interact*. 2014;8: 411–419. doi:10.1007/s11829-014-9328-x
- 549 46. Urzúa A. Secondary Metabolites in the Epicuticle of *Haplopappus Foliosus* D.C.
550 (Asteraceae). *J Chil Chem Soc*. 2004;49: 137–141. doi:10.4067/S0717-
551 97072004000200006
- 552 47. Urzúa A, Andrade L. Comparative chemical composition of the resinous exudates
553 from *Haplopappus foliosus* and *H. uncinatus*. *Biochem Syst Ecol*. 2000;28: 491–
554 493.
- 555 48. Vargas HA, Rasmann S, Ramirez-Verdugo P, Villagra CA. *Lioptilodes friasi*
556 (Lepidoptera: Pterophoridae) Niche Breadth in the Chilean Mediterranean
557 Matorral Biome: Trophic and Altitudinal Dimensions. *Neotrop Entomol*. 2017;47.
558 doi:10.1007/s13744-017-0514-2
- 559 49. Klingenberg L. Monographie der südamerikanischen Gattungen *Haplopappus*
560 *Cass.* Und *Notopappus* L. Klingenberg (Asteraceae-Astereae). Berlin: Series
561 *Bibliotheca Botanica*, Heft 157 0067- 7892, Schweizerbartsche

- Verlagsbuchhandlung; 2007. p. 331.
50. Urzúa A. Secondary metabolites in the epicuticle of *Haplopappus foliosus* DC. (Asteraceae). J Chil Chem Soc. 2004;49: 137–141. doi:10.4067/S0717-97072004000200006
51. Camousseight A. Baratas o cucarachas. In: Canals M, Cattán P, editors. Zoología Médica II: Invertebrados. Primera. Editorial Universitaria; 2008. p. 392.
52. Hammer O. PAST Reference manual. Natural History Museum. 1999.
53. Richardson J. The analysis of 2 x 2 contingency tables - Yet again. Stat Med. 2011;30: 890.
54. Simirgiotis MJ, Quispe C, Bórquez J, Schmeda-Hirschmann G, Avendaño M, Sepúlveda B, et al. Fast high resolution Orbitrap MS fingerprinting of the resin of *Heliotropium taltalense* Phil. from the Atacama Desert. Ind Crops Prod. Elsevier; 2016;85: 159–166.
55. Adams RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. 4th ed. Stream C, editor. Allured Publishing Corporation; 2007.
56. Korovetska H, Novák O, Turečková V, Hájíčková M, Gloser V. Signalling mechanisms involved in the response of two varieties of *Humulus lupulus* L. to soil drying: II. changes in the concentration of abscisic acid catabolites and stress-induced phytohormones. Plant Growth Regul. Springer; 2016;78: 13–20.
57. Kapp K, Hakala E, Orav A, Pohjala L, Vuorela P, Püssa T, et al. Commercial peppermint (*Mentha× piperita* L.) teas: Antichlamydial effect and polyphenolic composition. Food Res Int. Elsevier; 2013;53: 758–766.

Haplopappus platylepis resin for pest control

- 585 58. Jiménez-Sánchez C, Lozano-Sánchez J, Rodríguez-Pérez C, Segura-Carretero A,
586 Fernández-Gutiérrez A. Comprehensive, untargeted, and qualitative RP-HPLC-
587 ESI-QTOF/MS2 metabolite profiling of green asparagus (*Asparagus officinalis*). J
588 Food Compos Anal. Elsevier; 2016;46: 78–87.
- 589 59. Simirgiotis MJ, Ramirez JE, Hirschmann GS, Kennelly EJ. Bioactive coumarins
590 and HPLC-PDA-ESI-ToF-MS metabolic profiling of edible queule fruits
591 (*Gomortega keule*), an endangered endemic Chilean species. Food Res Int.
592 Elsevier; 2013;54: 532–543.
- 593 60. Doménech-Carbó MT, de La Cruz-Cañizares J, Osete-Cortina L, Doménech-Carbó
594 A, David H. Ageing behaviour and analytical characterization of the Jatobá resin
595 collected from *Hymenaea stigonocarpa* Mart. Int J Mass Spectrom. Elsevier;
596 2009;284: 81–92.
- 597 61. De Gutierrez AN, Catalan CAN, Díaz JG, Herz W. Sesquiterpene lactones and
598 other constituents of *Stevia jujuyensis*. Phytochemistry. Elsevier; 1992;31: 1818–
599 1820.
- 600 62. Urzúa A, Echeverría J, Espinoza J. Lipophilicity and antibacterial activity of
601 flavonols: Antibacterial activity of resinous exudates of *haplopappus litoralis*, *H.*
602 *chrysantemifolius* and *H. scrobiculatus* | Lipofilia y actividad antibacteriana de
603 flavonoles: Actividad antibacteriana de los ex. Bol Latinoam y del Caribe Plantas
604 Med y Aromat. 2012;
- 605 63. Cushnie TPT, Lamb AJ. Recent advances in understanding the antibacterial
606 properties of flavonoids. Int J Antimicrob Agents. Elsevier; 2011;38: 99–107.
- 607 64. Echeverría J, Urzúa A, Sanhueza L, Wilkens M. Enhanced Antibacterial Activity

Haplopappus platylepis resin for pest control

- 608 of Ent-Labdane Derivatives of Salvic Acid (7 α -Hydroxy-8(17)-ent-Labden-15-Oic
- 609 Acid): Effect of Lipophilicity and the Hydrogen Bonding Role in Bacterial
- 610 Membrane Interaction. *Molecules*. 2017; doi:10.3390/molecules22071039
- 611 65. Echeverría J, Opazo J, Mendoza L, Urzúa A, Wilkens M. Structure-Activity and
- 612 Lipophilicity Relationships of Selected Antibacterial Natural Flavones and
- 613 Flavanones of Chilean Flora. *Molecules*. 2017; doi:10.3390/molecules22040608
- 614 66. Echeverría J, González-Teuber M, Urzúa A. Antifungal activity against Botrytis
- 615 cinerea of labdane-type diterpenoids isolated from the resinous exudate of
- 616 *Haplopappus velutinus* Remy (Asteraceae). *Nat Prod Res*. 2018; 1–5.
- 617 doi:10.1080/14786419.2018.1443093
- 618 67. Chinou I. Labdanes of natural origin-biological activities (1981-2004). *Curr Med*
- 619 *Chem*. Bentham Science Publishers; 2005;12: 1295–1317.
- 620 68. Singh M, Pal M, Sharma RP. Biological activity of the labdane diterpenes. *Planta*
- 621 *Med*. Georg Thieme Verlag Stuttgart· New York; 1999;65: 2–8.
- 622 69. Santos AO dos, Izumi E, Ueda-Nakamura T, Dias-Filho BP, Veiga-Júnior VF da,
- 623 Nakamura CV. Antileishmanial activity of diterpene acids in copaiba oil. *Mem*
- 624 *Inst Oswaldo Cruz*. SciELO Brasil; 2013;108: 59–64.
- 625 70. Dostálová A, Volf P. Leishmania development in sand flies: parasite-vector
- 626 interactions overview. *Parasit Vectors*. 2012;5: 276. doi:10.1186/1756-3305-5-276
- 627 71. Prost I, Dhondt S, Rothe G, Vicente J, Rodriguez MJ, Kift N, et al. Evaluation of
- 628 the antimicrobial activities of plant oxylipins supports their involvement in defense
- 629 against pathogens. *Plant Physiol*. Am Soc Plant Biol; 2005;139: 1902–1913.
- 630 72. Desbois AP, Smith VJ. Antibacterial free fatty acids: activities, mechanisms of

631 action and biotechnological potential. Appl Microbiol Biotechnol. Springer;
632 2010;85: 1629–1642.

633 73. Martin-Arjol I, Bassas-Galia M, Bermudo E, Garcia F, Manresa A. Identification
634 of oxylipins with antifungal activity by LC–MS/MS from the supernatant of
635 *Pseudomonas* 42A2. Chem Phys Lipids. Elsevier; 2010;163: 341–346.

636 74. Desbois AP, Lawlor KC. Antibacterial activity of long-chain polyunsaturated fatty
637 acids against *Propionibacterium acnes* and *Staphylococcus aureus*. Mar Drugs.
638 Multidisciplinary Digital Publishing Institute; 2013;11: 4544–4557.

639 75. Trapp MA, Kai M, Mithöfer A, Rodrigues-Filho E. Antibiotic oxylipins from
640 *Alternanthera brasiliana* and its endophytic bacteria. Phytochemistry. Elsevier;
641 2015;110: 72–82.

642 76. Clem RJ, Passarelli AL. Baculoviruses: Sophisticated Pathogens of Insects. PLoS
643 Pathog. 2013;9. doi:10.1371/journal.ppat.1003729

644 77. Kazda J. The chronology of mycobacteria and the development of mycobacterial
645 ecology. The ecology of Mycobacteria: impact on animal's and human's health.
646 Springer; 2009. pp. 1–11.

647 78. Frishman AM, Alcamo IE. Domestic cockroaches and human bacterial disease.
648 Pest Control. 1977;45: 16–46.

649 79. Roni M, Murugan K, Panneerselvam C, Subramaniam J, Nicoletti M,
650 Madhiyazhagan P, et al. Characterization and biotoxicity of *Hypnea musciformis*-
651 synthesized silver nanoparticles as potential eco-friendly control tool against
652 *Aedes aegypti* and *Plutella xylostella*. Ecotoxicol Environ Saf. 2015;121: 31–38.
653 doi:10.1016/j.ecoenv.2015.07.005

Haplopappus platylepis resin for pest control

654

655

656

657

658

Haplopappus platylepis resin for pest control

660

Haplopappus platylepis resin for pest control

662

A

B

A

B