1 **UDP-glucose:anthocyanidin 3-*O*-glucoside-2''-*O*-glucosyltransferase catalyzes**
2 **further glycosylation of anthocyanins in purple *Ipomoea batatas***
3 **Running Title: Further anthocyanin glycosylation in sweetpotato**

4 Hongxia Wang¹, Chengyuan Wang², Weijuan Fan^{1,3}, JunYang³, Ingo Appelhagen⁴,
5 Yinliang Wu^{1,5} and Peng Zhang^{1,5}

6 ¹National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in
7 Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Shanghai
8 Institutes for Biological Sciences, Chinese Academy of Science, Shanghai 200032,
9 China

10 ²Center for Computational Medicine and Bioinformatics, University of Michigan,
11 Ann Arbor, MI 48109-2218, USA

12 ³Shanghai Key Laboratory of Plant Functional Genomics and Resources, Shanghai
13 Chenshan Plant Science Research Center, Chinese Academy of Science, Shanghai
14 Chenshan Botanical Garden, Shanghai 201602, China

15 ⁴John Innes Centre, Norwich Research Park, Colney, Norwich NR4 7UH, United
16 Kingdom

17 ⁵University of Chinese Academy of Sciences, Beijing 100049, China

18 Authors for correspondence:

19 Hongxia Wang, Tel: +86 2154924097, Email: hxwang@sibs.ac.cn

20 Peng Zhang, Tel: +86 2154924096, Email: zhangpeng@sibs.ac.cn

21 **Abstract**

22 Glycosylation contributes to the diversity and stability of anthocyanins in plants. The
23 process is catalyzed by various glucosyltransferases using different anthocyanidin
24 aglycones and glycosyl donors. An anthocyanidin
25 3-*O*-glucoside-2''-*O*-glucosyltransferase (3GGT) from purple sweetpotato (cv.
26 Ayamurasaki) served for the catalytic conversion of anthocyanidin 3-*O*-glucoside into

anthocyanidin 3-*O*-sophoroside, which is functionally different from the 3GGT ortholog of *Arabidopsis*. The phylogenetic analysis indicates regioselectivity of 3GGT using UDP-xylose or UDP-glucose as the glycosyl is divergent between *Convolvulaceae* and *Arabidopsis*. Homology-based protein modeling and site-directed mutagenesis of Ib3GGT and At3GGT suggested that the Thr-138 of Ib3GGT is a key amino acid residue for UDP-glucose recognition and plays a major role in sugar donor selectivity. The wild type and *ugt79b1* mutants of *Arabidopsis* plants overexpressing *Ib3GGT* produced the new component cyanidin 3-*O*-sophoroside. Moreover, *Ib3GGT* expression was associated with anthocyanin accumulation in different tissues during Ayamurasaki plant development and was regulated by the transcription factor IbMYB1. The localization assay of Ib3GGT showed that further glycosylation occurs in the cytosol and not endoplasmic reticulum. The present study revealed the function of Ib3GGT in further glycosylation of anthocyanins and its Thr-138 is the key amino acid residue for UDP-glucose recognition.

Key words: *Ipomoea batatas*, anthocyanins, further glycosylation, UDP-glucose, glucosyltransferase, regioselectivity.

Introduction

Anthocyanins are major secondary metabolites responsible for color variation in plants, exhibiting health-promoting properties (de Pascual-Teresa and Sanchez-Ballesta, 2008; He and Giusti, 2010). The basic structures of anthocyanins are mono- and di-glycosylated forms in common anthocyanidins, which include cyanidin, delphinidin, malvidin, pelargonidin, peonidin, and petunidin (Moglia et al., 2014). Different sugar moieties, i.e., glucose, galactose, xylose, arabinose, or fructose can be linked to hydroxyl groups at 3, 5, 7, 3', and 5' positions, with the glycosylation at the 3rd position on the C-ring ubiquitously (Andersen and Jordheim, 2010). Glycosylation of 3-OH is catalyzed by a series of UDP carbohydrate-dependent glycosyltransferases (UGTs), which utilize the nucleotide-activated sugars as donor substrates and anthocyanidin aglycones or anthocyanins as acceptors. These activities increase the structural diversity of anthocyanins by adding different types and/or numbers of sugar moieties on various positions (Gachon et al., 2005). The glycosylation of anthocyanin is speculated to occur on the cytoplasmic surface of the

58 endoplasmic reticulum (ER), and may serve as a signal for the transport of
59 anthocyanins to vacuoles via multiple pathways; this transport is essential for the
60 stable storage of anthocyanins in vacuoles (Ono et al., 2006; Matsuba et al., 2010; Sun
61 et al., 2012; Zhao et al., 2011; Zhao, 2015). Glycosylation also participates in the fine
62 adjustment and stabilization of flower pigmentation in ornamental plants
63 (Yonekura-Sakakibara et al., 2012).

64 Monoglycosylation of anthocyanidins produces anthocyanidin 3-*O*-glucosides,
65 the first major stable colored pigments in the anthocyanin biosynthesis pathway
66 (Griesser et al., 2008a; Montefiori et al., 2011). Deficiency of the activity of the
67 corresponding UDP-glucose:flavonoid 3-*O*-glycosyltransferase (UF3GT), in maize
68 *bronze1* and *Arabidopsis anl1*, results in a significantly suppressed accumulation of
69 anthocyanin (Fedoroff et al., 1984; Kubo et al., 2007). Until now, UF3GT is one of
70 the well-characterized UGTs related to anthocyanin biosynthesis (Gachon et al., 2005;
71 Yonekura-Sakakibara and Hanada, 2011). Further glycosylation of anthocyanidin
72 3-*O*-glucosides involves diverse sugars in different species, such as UDP-rhamnose,
73 UDP-glucose, UDP-xylose, and UDP-arabinose, as donor substrates to be added at a
74 species-specific position to the glycosides of mono 3-*O*-glycosylated anthocyanins
75 (Yonekura-Sakakibara et al., 2012). The mutants affected in this further glycosylation
76 function may be impacted in the anthocyanin accumulation, as reported in petunia and
77 Japanese morning glory (Kroon et al., 1994; Morita et al., 2005). Since all the UGT
78 proteins are highly similar in their secondary and tertiary structures, with a defined
79 fold structure and highly conserved putative secondary product glycosyltransferase
80 (PSPG) motif (Breton et al., 2006; Lairson et al., 2008; Osmani et al., 2009),
81 structure-based modeling have identified the key residues of UF3GT responsible for
82 sugar donor specificity (Kubo et al., 2004) in *Arabidopsis* (Kim et al., 2013), *Freesia*
83 *hybrid* (Sun et al., 2016), grapes (Offen et al., 2006; Ono et al., 2010), lamiales
84 (Noguchi et al., 2009), perilla (Noguchi et al., 2009), and red daisy (Osmani et al.,
85 2009). Nevertheless, the residues involved in UDP-sugar selectivity in 3GGT are yet
86 unknown.

87 Although the anthocyanidin decoration by glycosylation is progressive, it
88 commonly begins with 3-*O*-glycosylation to ensure the stability of the aglycon.
89 Additional glycosylation leads to the compound and functional diversity, thereby

contributing to several varieties of anthocyanins in the plant (Gachon et al., 2005; Caputi et al., 2012). To date, more than 600 anthocyanins or their derivatives have been identified in nature (Glover and Martin, 2012); however, only a limited number of genes encoding UFGTs in different species have been well characterized. Several flavonoid 3-*O*-glycosyltransferases have been characterized in *Arabidopsis* (Kubo et al., 2007; Saito et al., 2013), strawberry (Griesser et al., 2008a, b), grapes (Offen et al., 2006), and maize (Fedoroff et al., 1984). In addition, for further flavonoid glycosylation multiple UGTs were also characterized, including anthocyanidin 3-*O*-glucoside 6''-*O*-rhamnosyltransferase in *Petunia hybrida* (Kroon et al., 1994), anthocyanidin 3-*O*-glucoside 2''-*O*-glucuronosyltransferase in red daisy flowers (Sawada et al., 2005), and flavonol 3-*O*-glucoside 2''-*O*-glucosyltransferase in *Arabidopsis* (Yonekura-Sakakibara et al., 2014). Apparently, the divergence towards different glycosylation types occurs at this step. At the same 2'' position, different glycosylation types, i.e., glycosylation or xylosylation, are found in various plant species. In morning glory, anthocyanidin 3-*O*-glucoside 2''-*O*-glucosyltransferase catalyzes the addition of a glucose molecule to anthocyanidin 3-*O*-glucosides on the 2'' position to form anthocyanidin 3-*O*-sophorosides (Morita et al., 2005). In *Arabidopsis*, further glycosylation of the 3-*O*-glucoside is catalyzed by anthocyanidin 3-*O*-glucoside 2''-*O*-xylosyltransferase (AtA3G2XylT, i.e., At3GGT) to add one xylose molecule specifically to the first glucose residue (Yonekura-Sakakibara et al., 2012). Further decorations, for example, to the diversity or functionality, malonylation and aromatic acylation rely on the glycosylation of anthocyanidins (Sasaki et al., 2014).

Purple sweetpotato (*Ipomoea batatas*) accumulates a lot of anthocyanins in storage roots. Anthocyanidin 3-*O*-glucoside-2''-*O*-glucoside (anthocyanin 3-*O*-sophoroside) and derivatives are the major anthocyanin compounds (Tian et al., 2005). So far, at least 26 components, mostly caffeoylated, coumarylated or feruloylated anthocyanidin glucosides have been identified (Truong et al., 2009; Lee et al., 2013). In contrast, 11 anthocyanins have been identified in *Arabidopsis*; all of them derived from cyanidin 3-*O*-glucoside-2''-*O*-xyloside (Tohge et al., 2005; Yonekura-Sakakibara et al., 2012; Kavinich et al., 2014). Therefore, unlike *Arabidopsis*, purple sweetpotato uses UDP-glucose as sugar donor for further

glycosylation of anthocyanidin 3-*O*-glucosides to form anthocyanidin 3-*O*-sophorosides. In the present study, we characterized a UFGT, termed as UDP-glucose:anthocyanidin 3-*O*-glucoside-2''-*O*-glucosyltransferase (IbA3G2GluT, i.e., Ib3GGT) that catalyzes the anthocyanin glycosylation in purple sweetpotato and its key amino acid for sugar donor selectivity.

Materials and methods

Plant materials

The purple-fleshed sweetpotato (*Ipomoea batatas* Lam.) cultivar Ayamurasaki was used in this study. The *in vitro* shoot cultures were subcultured on SBM medium (MS salts including vitamins + 0.3 mg/L VB1 + 30 g/L sucrose, pH 5.8) in plant growth chambers under a 16 h photoperiod provided by cool-white fluorescent tubes (~50 $\mu\text{mol}/\text{m}^2/\text{s}$), at 25°C and 50% relative humidity. One-month-old plantlets were transplanted into plastic pots containing well-mixed soil (soil:peat:perlite, 1:1:1) and grown in the greenhouse (16 h/8 h light/dark cycle, 25°C day/night). Various tissues including leaves, stems, fibrous roots, and storage roots of sweetpotato plants at different developmental stages were harvested from the pot- or field-grown plants for multiple analyses. The *Arabidopsis* plants were grown under a 16 h/8 h light/dark cycle, at 22°C in the growth chamber.

Plasmid construction and production of transgenic sweetpotato

The open reading frame of *Ib3GGT* (1380 bp) was amplified from the cDNA of sweetpotato Ayamurasaki using the primers Ib3GGTF (5'-CGGGGTACCATGGGTCTCAAGCAACAAC-3', KpnI site underlined) and Ib3GGTR (5'-AATGTCGACTCATCCAAGGAGATCCTGCA-3', SalI site underlined). This fragment was inserted into the KpnI/SalI sites of the pCAMBIA1301-based plant expression vector to generate the binary vector pOE-Ib3GGT containing the expression cassette of *Ib3GGT* driven by the CaMV 35S promoter. The pRNAi-Ib3GGT binary vector was manipulated to express double-stranded hairpin RNA of the 252 bp *Ib3GGT* fragment (382–633 bp) based on the pRNAi-DFR vector (Wang et al., 2013). Then, pOE-Ib3GGT and pRNAi-Ib3GGT were introduced into *Agrobacterium tumefaciens* strain LBA4404 for sweetpotato transformation, as described previously (Yang et al., 2011). Transgenic plants were produced and verified for *Ib3GGT* expression by real-time RT-PCR. For total *Ib3GGT*

154 expression, an internal primer pair of *Ib3GGT* was designed for detecting the *Ib3GGT*
155 expression in WT, OE-*Ib3GGT* and RNAi-*Ib3GGT* plants by real-time RT-PCR
156 (Table S2). The *Actin* gene of sweetpotato was used as an internal control for gene
157 amplification.

158 **Transformation and analysis of *Ib3GGT*-overexpressing *Arabidopsis***

159 Two independent UGT79B1 *Arabidopsis* transposon mutants, *ugt79b1-1* and
160 *ugt79b1-2* (Kuromori et al., 2004; Ito et al., 2005), along with the WT Nossen and
161 ecotype Col-0 were transformed with *A. tumefaciens* LB4404 harboring pOE-*Ib3GGT*,
162 using the floral dip method (Clough and Bent, 1998). The transformants were selected
163 on 1/2 MS medium containing 50 mg/L hygromycin for Nossen and mutants or 25
164 mg/L hygromycin for Col-0 plants. The RNA extracted from T3 homozygous
165 *Arabidopsis* seedlings was used for RT-PCR analysis. The primer pairs used to detect
166 the expression of *At3GGT* and *Ib3GGT* in WT and transgenic *Arabidopsis* plants were
167 designed using software Primer 3.0 and listed in Table S2. *At3GGT* were amplified a
168 223-bp fragment from position +369 to +591 bp and *Ib3GGT* were amplified a 189-bp
169 fragment from +1009 to +1197 bp. The *Actin* gene of *Arabidopsis* was used as a
170 reference gene.

171 **Phylogenetic Analysis**

172 To construct a phylogenetic tree, 16 UGT protein sequences obtained from NCBI
173 GenBank were aligned by ClustalW and implemented in MEGA6 (Tamura et al.,
174 2013). Ten closely related UGTs were used to illustrate the relationship. The
175 maximum likelihood method was used to obtain the alignment results (Stamatakis,
176 2014). Bootstrap values were obtained with 1000 replications.

177 **Site-directed mutagenesis and *in vitro* enzymatic assay of recombinant *Ib3GGT* 178 and *At3GGT***

179 The full-length sequence of the *Ib3GGT* gene was amplified by PCR using the
180 primers IbGGT-FP (5'-CCCAAGCTTATGGGTTCTCAAGCAACAAC-3', HindIII
181 site underlined) and IbGGT-RP
182 (5'-CGCGGATCCTCACATCACCATCACCATCACTCCAAGGAGATCCTGCA-3',
183 BamHI site and 6 His sites underlined). The full-length *At3GGT* was amplified by
184 PCR using the primers AtGGT-FP

185 (5'-GGGGTACCATGGGTGTTTTTGGATCGAA-3', KpnI site underlined) and
 186 AtGGT-RP
 187 (5'-CGGAATTCCTCACATCACCATCACCATCACTGACTTCACAAGTTCAATTA
 188 AATT-3', EcoRI site and 6 His sites underlined). Site-directed mutations were
 189 generated by changing the Thr-138 nucleotide ACC into ATT in *Ib3GGT* and Ile-142
 190 ATC into ACT in *At3GGT* using PCR-based amplification with a Phusion
 191 Site-Directed Mutagenesis Kit (Thermo Scientific). The sequence fragments, with or
 192 without the mutation of the 3GGTs, were cloned into the pYES2 vector and
 193 introduced in *Saccharomyces cerevisiae* BY4742 according to the manufacturer's
 194 instructions (Cat# V825-20, Invitrogen). The recombinant 3GGT proteins were
 195 induced by replacing the carbon source from 2% glucose to 2% galactose in the SC-U
 196 medium. The reaction mixture for the 3GGT enzymatic assay consisted of 100 mM
 197 phosphate buffer (pH 7.0), 0.6 mM flavonoid aglycones (cyanidin,
 198 cyanidin 3-*O*-glucoside, cyanidin 3,5-*O*-diglucoside, or flavonol 3-*O*-glucoside), 1
 199 mM UDP-glucose, and 20 µL of crude yeast extract as the enzymatic solution in a
 200 reaction volume of 100 µL. After incubation for 2 h at 37°C, the reaction was
 201 terminated by centrifugation. The enzymatic activity of mutant 3GGT was assessed
 202 by cyanidin 3-*O*-glucoside as the acceptor substrate and different UDP-sugars
 203 (UDP-glucose, UDP-xylose, UDP-galactose or UDP-arabinose) as a sugar donor.

204 **LC-MS analyses of metabolites obtained by enzymatic reaction**

205 Ten µL of filtered supernatants were analyzed on an Agilent
 206 HPLC1200-MSD/Q-TOF 6520 system (Agilent, Waldbronn, Germany) as described
 207 previously (Wang et al., 2013). Briefly, the mobile phase consisted of 0.5% (v/v)
 208 acetic acid in water (eluent A) and 100% acetonitrile (eluent B). The samples eluted at
 209 a flow rate of 0.2 mL/min passed through a reverse-phase C18 column (Agilent
 210 ZORBAX Eclipse XDB, 4.6 × 50 mm ID, 1.8 µm), and a DAD detector at 530 nm
 211 monitored the anthocyanin. Subsequently, an ESI interfaced Q-TOF mass detector
 212 (m/z 40–1500) collected the mass m/z data that were processed by Agilent Mass
 213 Hunter Qualitative Analysis (version 3.0) for the estimation of accurate molecular
 214 mass as well as spectrum evaluation. Cyanidin 3-*O*-sophoroside (Tongtian, Shanghai,
 215 China) was used as a standard.

216 **Subcellular localization of *Ib3GGT* in plant cells**

217 The *Ib3GGT* gene was amplified by PCR using Pfu polymerase (Takara, Shanghai,
218 China) to obtain a non-stop coding sequence using the primers FPGGT_L
219 (5'-AATGTCGACATGGGTTCTCAAGCAACAAC-3', SalI site underlined) and
220 RPGGT_L (5'-GGACTAGTCCAAGGAGATCCTGCAGTT-3', SpeI site underlined).
221 Ib3GGT-eGFP was constructed by inserting the *Ib3GGT* fragment into the
222 corresponding sites of a modified pCambia1300 to fuse with the eGFP coding
223 sequence. The construction of the ER-marker (Nelson et al., 2007) and the expression
224 construct for mRFP (Claudia et al., 2017) has been described elsewhere. The ER
225 marker, ER-mCherry, contains a signal peptide of AtWAK2 at the N-terminal and a
226 synthetic HDEL at the C-terminal (He et al., 1999; Nelson et al., 2007). All constructs
227 were introduced into *A. tumefaciens* GV3101 (pMP90). The growth conditions for *N.*
228 *benthamiana* and *A. tumefaciens*, as well as the agro-infiltration procedure, were
229 described previously (Leuzinger et al., 2013). The images were acquired 36 h
230 post-infiltration with a Leica SP8X confocal microscope equipped with a Leica HC
231 PL APO CS2 63x/1.20 water immersion objective. The GFP fluorescence was
232 detected by hybrid detector HyD1 in the range of 500–540 nm and excited using the
233 488-nm line of an argon ion laser. mCherry and mRFP fluorescence were detected in
234 the range of 580–630 nm by HyD2 after excitation at 561 nm with a diode-pumped
235 solid-state laser. Both fluorophores were recorded line-by-line sequentially at a 3- to
236 4-fold average in a background noise-dependent manner. The Leica Application Suite
237 X software was used for image acquisition and intensity estimations.

238 **Anthocyanin measurement and detection**

239 Total anthocyanins in the WT and transgenic lines were extracted using previously
240 described methods with slight modifications (Wang et al., 2013). The total content of
241 anthocyanin in the WT and transgenic lines was quantified as cyanidin
242 3-*O*-sophoroside equivalent. The anthocyanin autofluorescence in epidermal cells of
243 sweetpotato leaves was examined using a PCM-2000/Nikon Eclipse 600
244 laser-scanning microscope (Nikon, Japan) equipped with an argon and helium-neon
245 laser (excitation 488 nm, emission 544 nm).

246 **Luciferase assay**

247 The *Ib3GGT* promoter (2000 bp) was amplified by the primers Ib3GGTprFP
248 (3'-AACTGCAGTTCAGTCAGGCAATCACAGG-5', PstI site underlined) and

249 Ib3GGTprRP (3'-CGCGGATCCAATAATACCTAGCTAGCT-5', BamHI site
250 underlined) and cloned into the pLL00R vector to generate the luciferase reporter
251 vector. The *IbMYB1* gene was amplified by the primers IbMYB1FP
252 (3'-GGGGTACCATGGTTATTTTCATCTGTATG, KpnI site underlined) and
253 IbMYB1RP (3'-AACTGCAGTTAGCTTAACAGTTCTGAC-5', PstI site underlined)
254 and subcloned into pCambia1300 to generate the CaMV35S-IbMYB1 effector
255 plasmid. *A. tumefaciens* strain GV3101 harboring the *Ib3GGT* promoter-LUC reporter
256 and CaMV 35S-IbMYB1 effector was infiltrated into the 5-week-old *N. benthamiana*
257 leaves using a needleless syringe for assessing the luciferase activity. The plants were
258 grown for 48 h (16 h/8 h light/dark cycle, 25°C day/night), followed by injecting the
259 leaves with 0.94 mM luciferin as substrate. The leaves were collected in the dark after
260 3 min and luciferase signals detected on a Tanon-5200 image system. The LUC
261 reporter empty vector with 35S-IbMYB1 or *Ib3GGT* promoter-LUC reporter with
262 empty effector vector was also co-infiltrated as a negative control. These experiments
263 were repeated at least three times, and similar results were obtained.

264 **Molecular modeling of Ib3GGT and At3GGT active sites**

265 The 3D models of Ib3GGT and At3GGT were generated using SWISS-MODEL
266 workspace (Biasini et al., 2014; Wetterhorn et al., 2016) and I-TASSER server
267 (Hiromoto et al., 2006) based on the structure of *N-/O*-glucosyltransferase of *A.*
268 *thaliana* that served as a template (UGT72B1 PDB ID: 2VCE, Brazier-Hicks et al.,
269 2007). The substrate binding sites were predicted by superposing both models to
270 UGT72B1 using the COOT program (Emsley et al., 2010).

271 **Statistical analyses**

272 All data were represented as mean $\pm$ SD from at least three biological replicates.
273 One-way ANOVA analyses were performed by using SPSS Statistics 17.0 to Duncan's
274 multiple comparison tests. A value of $P < 0.05$ was considered as statistically
275 significant difference.

276 **Results**

277 **Comparison of anthocyanins indicates different further glycosylation patterns in** 278 **sweetpotato and *Arabidopsis***

279 In purple sweetpotato cv. Ayamurasaki, anthocyanins include aromatically acylated

anthocyanidin 3-*O*-sophoroside and derivatives, whereas in *Arabidopsis* Col-0, anthocyanin components are anthocyanidin 3-*O*-glucoside-2''-*O*-xylosyl derivatives (Table S1). This phenomenon implies that, although the first glycosylation step of anthocyanins is similar for the production of anthocyanidin 3-*O*-glucosides and is catalyzed by UDP-glucose:flavonoid 3-*O*-glucosyltransferases, further modifications of anthocyanidin 3-*O*-glucosides diverge based on the utilization of different sugar donors, i.e. glucose in sweetpotato and xylose in *Arabidopsis*. In *Arabidopsis*, glucosyltransferase UGT79B1 (At3GGT) catalyzes the conversion of UDP-xylose and cyanidin 3-*O*-glucoside into cyanidin 3-*O*-glucoside-2''-*O*-xyloside (Tohge et al., 2005; Saito et al., 2013). In sweetpotato, a novel glucosyltransferase, UDP-glucose:anthocyanidin 3-*O*-glucoside-2-*O*-glucosyltransferase (Ib3GGT), was predicted to participate in further glycosylation.

292 **Cloning and phylogenetic characterization of Ib3GGT**

The full-length *Ib3GGT* CDS sequence (GenBank accession number EF108571) was identified from a sweetpotato cDNA library by comparison with the *At3GGT* sequence. The 1380-bp *Ib3GGT* gene harbors an open-reading frame encoding 459 amino acids (aa) with a calculated molecular mass of 50.87 kDa and an isoelectric point 6.537. Further sequence analysis of Ib3GGT showed that its amino acid sequence shared the common domain of PSPG box (334–377 aa, Fig. 1A) with other UF3GGTs in the C-terminal region (Osmani et al., 2009). In addition, although the sugar donor specificity was reported to be partially determined by the last amino acid residue of the PSPG box, i.e., glutamine (Gln) for UDP-glucose and histidine (His) for UDP-galactose (Kubo et al., 2004), the last residue of PSPG in Ib3GGT (at 377 aa), At3GGT (UDP-glucose:flavonol 3-*O*-glucoside-2''-*O*-glucosyltransferase), At3GGT, and Ip3GGT are Gln that is conserved among these glucosyltransferases. This phenomenon indicated that other amino acid residues in their sequences might contribute towards sugar donor specificity, which necessitates further elucidation.

Phylogenetic analysis showed that Ib3GGT belongs to a cluster of typical further glucosyltransferases, and is most closely related to Ip3GGT of *Ipomoea purpurea* (Morita et al., 2005), showing 94.3% identity (Fig. 1B). Ib3GGT is also homologous to At3GGT and At3GGTf with 45.7% and 45.6% identity, respectively.

311 **Ib3GGT is an enzyme that catalyzes the glycosylation of anthocyanidin** 312 **3-*O*-glucoside into anthocyanidin 3-*O*-sophoroside**

313 To further examine the function of Ib3GGT *in vitro*, recombinant His-tag fusion
314 Ib3GGT and At3GGT proteins expressed in the yeast expression vector pYES2
315 (Invitrogen, USA), were used for assessing the enzymatic activity. The specificity of
316 Ib3GGT was examined using different sugar acceptors and donors. The recombinant
317 Ib3GGT protein only catalyzed the conversion of cyanidin 3-*O*-glucoside into
318 cyanidin 3-*O*-sophoroside using UDP-glucose as a sugar donor (Fig. 2A). Other
319 glucosyl acceptors such as cyanidin, cyanidin 3,5-*O*-diglucoside, and flavonol
320 3-*O*-glucoside could not serve as substrates, and hence, no product was detected (Fig.
321 2B, 2C, 2D), similar to the negative control (empty vector) (Fig. 2E). In addition, the
322 Ib3GGT protein could use peonidin 3-*O*-glucoside as the glucosyl acceptor to form
323 peonidin 3-*O*-sophoroside (Fig. 2F). These findings indicated that Ib3GGT used
324 anthocyanidin 3-*O*-glucoside as the glucosyl acceptor.

325 Ib3GGT specificity was also confirmed using four different UDP-sugars (Table
326 1). No detectable or predominant UGT activity was detected by UDP-sugars except
327 UDP-glucose, indicating that Ib3GGT is highly specific to UDP-glucose. The weak
328 utilization of UDP-xylose to produce cyanidin 3-*O*-glucoside-2''-*O*-xyloside
329 indicated the low affinity to this substrate. In contrast, the At3GGT protein was
330 capable of using only UDP-xylose (not for UDP-glucose) as the sugar donor to
331 catalyze cyanidin 3-*O*-glucoside into cyanidin 3-*O*-glucoside-2''-*O*-xyloside (Table 1,
332 Yonekura-Sakakibara et al., 2012), showing the divergence in the specificity of sugar
333 donors by the two UF3GGTs from different species.

334 **Thr-138 of Ib3GGT contributes to sugar donor preference**

335 To further identify the key amino acid residue of Ib3GGT responsible for sugar donor
336 recognition, docking experiments were performed based on the 3D structures of over
337 10 different glycosyltransferases enzymes from various plants (Brazier-Hicks et al.,
338 2007; Hiromoto et al., 2006; Hiromoto et al., 2013; Modolo et al., 2009; Offen et al.,
339 2006; Shao et al., 2005; Wetterhorn et al., 2016). The overall structures of these
340 glycosyltransferases share a similar folding topology: two Rossmann-like domains
341 formed a cleft which contains two substrates binding sites and one functional
342 conserved histidine residues located between these sites (Supplementary Fig.S1). By

343 using DALI services 8 (Holm and Laakso, 2016), more UGT homologous structures
344 were analyzed and the root mean square deviations (RMSDs) between each other
345 ranged from 1.1 to 2.5 Å over the core structure region (Supplementary Fig. S2).
346 Although the sugar acceptor ligands in these structures are diverse, the sugar donors
347 (most of them are UDP-glucose) are similar and share a group of conserved residues
348 in their binding pockets (Supplementary Fig. S2).

349 Two different methods, SWISS-MODEL (Biasini et al., 2014) and I-TASSER
350 Suite (Yang et al., 2015), were used for Ib3GGT structure modeling building. To
351 qualify the modeling results, the overall structures of Ib3GGT with the two service
352 models were compared with proteins 2VG8 and 2VCH; then the corresponding
353 residues in the modeling structures were also checked. Both sets of results were
354 similar, especially in the ligand binding sites (Supplementary Fig. S3). The RMSD
355 between two Ib3GGT modeling structures is 1.72Å and two At3GGT modeling
356 structures is 1.92Å, showing similar structure folding of the two services. Thus, the
357 results of SWISS-MODEL service were used to compare the Ib3GGT/At3GGT
358 modeling structures with an *Arabidopsis thaliana* O-glucosyltransferase (PDB ID:
359 2VCE, Brazier-Hicks et al., 2007) by superposing all three structures together. The
360 UDP-glucose in the template structure (2VCE) was also well bound in
361 Ib3GGT/At3GGT modeling structures. In Ib3GGT modeling structure,
362 Thr22/Ser276/Glu360/Gln337/Gln338/Trp334 form a binding pocket and interact
363 with the uridine group of UDP-glucose (Fig. 3A, Supplementary Fig.S1B);
364 Glu277/His352/Ser357 show tightly interactions with diphosphate group
365 (Supplementary Fig.S1C). All these residues are extremely conserved in UGTs (Hsu
366 et al., 2018; Thompson et al., 2017).

367 The two modeling structures shared a common group of residues in binding the
368 sugar donors (Glu277/Asp376/Gln377 in Ib3GGT and Gln285/Glu385/Gln386 in
369 At3GGT). The only difference in sugar binding pocket is the Thr-138 in Ib3GGT
370 which is equivalent to Ile-142 in At3GGT. The distance between O-6-glucose and
371 Thr-138 is 2.7Å (Supplementary Fig. S4), which can form a tight interaction.

372 However, the distance in At3GGT is 1.7Å (Supplementary Fig.S4B), which is too
373 short to bind the UDP-glucose. While replacing UDP-glucose with UDP-xylose in the
374 same position may lead to weak interaction between the xylose group and Thr-138 in
375 Ib3GGT (Supplementary Fig.S4C) and forms a 3.9Å hydrophobic interaction in
376 At3GGT (Supplementary Fig. S4D). These modeling observations are consistent with
377 our enzymatic activity assays (Table 1). Therefore, we hypothesized that the residue
378 Thr-138 in Ib3GGT and its equivalence Ile-142 in At3GGT are the key residues for
379 sugar donor specificity in purple sweetpotato and Arabidopsis, respectively.

380 To further attest to our hypothesis, firstly the corresponding site-directed mutants,
381 namely Ib3GGT^{T138I} (Thr-138 changed to Ile-138) and At3GGT^{I142T} (Ile-142 changed
382 to Thr-142), were constructed. Their enzyme activity showed that both the intact
383 protein and Ib3GGT^{T138I} could catalyze UDP-xylose to cyanidin
384 3-*O*-glucoside-2''-*O*-xyloside. However, the Ib3GGT^{T138I} mutant failed to use
385 UDP-glucose (Fig. 3B). On the other hand, At3GGT^{I142T} could not only primarily
386 catalyze UDP-xylose to produce cyanidin 3-*O*-glucoside-2''-*O*-xyloside but also use
387 UDP-glucose to synthesize cyanidin 3-*O*-sophorosides. These findings confirmed that
388 T138 is a key residue for sugar (glucose/xylose) recognition in Ib3GGT.

389 To check whether the Thr-138 is a key residue for other sugar recognition, 3D
390 models generated for UDP-galactose and UDP-arabinose were compared in the same
391 position with UDP-glucose and UDP-xylose. The only difference between
392 UDP-galactose and UDP-glucose is the direction of O₄H moiety which changes the
393 distance between O₄H moiety with the main chain N from 3.1Å to 4.93Å, thus
394 UDP-galactose should have less binding affinity than UDP-glucose (Supplementary
395 Fig. S5A, S5B). The UDP-arabinose also has less binding affinity than UDP-xylose as
396 the disappearance of the interaction between O5 with His-20 (Supplementary Fig.
397 S5C, S5D). As expected, no enzymatic activities were detected for Ib3GGT or
398 Ib3GGT^{T138I} using UDP-galactose and UDP-arabinose as sugar donors (Table 1).

399 To verify whether other species have the same mechanism of sugar donor
400 selectivity, two 3GGT proteins containing Thr-138 residue from *Prunus persica*
401 (Pp3GGT, XP_007213494) and *Lupinus angustifolius* (La3GGT, XP_019424989)

and one containing Ile-138 from *Camelina sativa* (Cs3GGT, XP_018450414) were cloned (Supplementary Fig. S6). Both Pp3GGT and La3GGT prefer UDP-glucose rather than UDP-xylose as sugar donor (Table 1). Their weak utilization of UDP-xylose indicated the low affinity to this substrate. On the other hand, the Cs3GGT protein was capable of using only UDP-xylose as the sugar donor (Table 1). Therefore, Thr-138 residue plays a key role in specificity of UDP-glucose donors by the two kinds of UF3GGTs. These results indicated that plant UGTs may share a same mechanism in sugar donor selectivity.

***Ib3GGT* expression in *Arabidopsis* produces new anthocyanin molecules**

To further validate the activity of *Ib3GGT* in planta, the *Ib3GGT* gene driven by the CaMV 35S promoter was overexpressed in *Arabidopsis* Col-0 and the UGT79B1 transposon insertion mutants, *ugt79b1-1* and *ugt79b1-2* (Kuromori et al., 2004; Ito et al., 2005). More than 10 independent transgenic plant lines were produced for each transformation event and their T3 homozygous lines (Fig. 4A). Further RT-PCR analysis confirmed the overexpression of *Ib3GGT* in these transgenic lines of Col-0, *ugt79b1-1* and *ugt79b1-2* (Fig. 4B, 4C). In the T3 homozygous *Ib3GGT*-OE transgenic lines, a new peak with an *m/z* value corresponding to cyanidin 3-*O*-sophoroside was detected in comparison to WT Col-0 by HPLC-electrospray ionization (ESI)-tandem mass spectrometry (MS/MS) analysis (Fig. 4D), although the purple-color phenotype and anthocyanin content at the cotyledon-stage seedling was indistinguishable (Fig. 4A, 4E). Moreover, the seedlings of *ugt79b1-1* and *ugt79b1-2* lines, which lacked the purple coloration as compared to the WT Nossen on 4.5% sucrose containing media, showed recovered anthocyanin accumulation when overexpressing the *Ib3GGT* gene (Fig. 4A, bottom panel, Supplementary Fig. S7). The comparison of anthocyanin profiles among WT, *ugt79b1-1*, and *ugt79b1-1* overexpressing *Ib3GGT* showed the production of cyanidin 3-*O*-sophoroside in transgenic lines (Fig. 4D; Supplementary Fig. S7). These results confirmed that *Ib3GGT* could specifically catalyze the conversion of cyanidin 3-*O*-glucoside to cyanidin 3-*O*-sophoroside in *Arabidopsis*, a biological process absent in this plant.

***Ib3GGT* expression is associated with anthocyanin accumulation and organ development and is regulated by IbMYB1**

433 Anthocyanin accumulation in Ayamurasaki plants showed an organ-dependent pattern.
 434 The immature leaves and mature storage roots contained maximum levels of
 435 anthocyanins (Fig. 5A); while mature leaves and fibrous roots had the least amounts.
 436 Among leaves, Lf1 reached a concentration of 0.6324 mg/g, approximately 7-fold that
 437 of Lf5. The *Ib3GGT* expression analyzed by real-time PCR in different organs also
 438 showed a similar pattern — high expression was found in immature leaves as well as
 439 developing and mature storage roots (Fig. 5A). *Ib3GGT* was expressed abundantly in
 440 developing storage roots (Dt, Fig. 5A) as compared to mature roots (Mt), which
 441 accumulated 30% more anthocyanins (0.4276 mg/g, Fig. 5A). Overall, *Ib3GGT*
 442 expression was associated with anthocyanin accumulation in different organs of
 443 Ayamurasaki plants.

444 The transcription factor IbMYB1 predominantly regulates the anthocyanin
 445 biosynthesis in purple sweetpotato (Mano et al., 2007), and its expression in
 446 Ayamurasaki plants was associated with anthocyanin accumulation. Therefore,
 447 *Ib3GGT* could potentially be a target gene of IbMYB1 that regulates the expression of
 448 downstream genes by binding the G-box element (CACGTG) in their promoters
 449 (Mano et al., 2007). Nonetheless, the promoter region of *Ib3GGT* showed a G-box
 450 element at position 992 as analyzed by the PlantCARE software (Supplementary Fig.
 451 S8). To confirm that *Ib3GGT* expression was stimulated by IbMYB1, a luciferase
 452 gene reporter driven by a 2000 bp promoter of *Ib3GGT* was assayed for luciferase
 453 activity in tobacco leaves after co-agroinfiltration with the effector, which harbors the
 454 *CaMV 35S::IbMYB1* expression cassette. Interestingly, a strong luciferase activity
 455 was detected. Reporter only or effector with a corresponding empty vector failed to
 456 detect the luminescent signals (Fig. 5B). These findings indicated that IbMYB1
 457 regulates the *Ib3GGT* expression in sweetpotato Ayamurasaki plants.

458 **Regulation of *Ib3GGT* expression in sweetpotato Ayamurasaki alters the** 459 **anthocyanin content but not the overall component profile**

460 To further elucidate the role of *Ib3GGT* in sweetpotato, *Ib3GGT*-overexpressing
 461 (OE-*Ib3GGT*) or -RNAi (RNAi-*Ib3GGT*) transgenic Ayamurasaki plants were
 462 analyzed. Multiple independent transgenic plant lines were produced and propagated
 463 in the greenhouse. Compared to the WT, RNAi-*Ib3GGT* lines showed reduced
 464 anthocyanin levels in the leaves of pot-grown plants, whereas OE-*Ib3GGT* plants

showed an increased anthocyanin accumulation in the top leaves (Fig. 6A, 6B). The expression of *Ib3GGT* was down-regulated in the RNAi-*Ib3GGT* lines and up-regulated in the OE-*Ib3GGT* lines by real-time PCR analyses (Fig. 6C). Anthocyanin level in the third leaf was reduced to 28.5% of WT in the RNAi-*Ib3GGT*-2 and increased up to 112% of WT in the OE-*Ib3GGT*-2 (Fig. 6A, 6B, 6D). The changes in the anthocyanin levels were correlated to *Ib3GGT* expression in these plants (Fig. 6C, 6D). Nevertheless, the overall profile of anthocyanins did not alter in these plants (Fig. 6E), indicating that *Ib3GGT* is involved in an early stage of anthocyanin modifications. Furthermore, the auto-fluorescence assayed in leaf epidermal cells showed a dramatic reduction in the fluorescent intensity in RNAi-*Ib3GGT* lines, while WT and OE-*Ib3GGT* transgenic plants displayed strong signals (Fig. 6F). Similar trends of altered anthocyanin accumulation were also observed in the field-grown corresponding plants (Supplementary Fig. S9).

***Ib3GGT* functions in the cytosol**

Anthocyanins have been suggested to be synthesized on the outer surface of the ER. To determine the location of the glycosylation of anthocyanins by *Ib3GGT*, the putative transit peptides were predicted *in silico* by Signal IP3.0, but none was found in the full *Ib3GGT* protein sequence. To test whether *Ib3GGT* is associated with ER, the N- and C-terminal fusions of *Ib3GGT* to eGFP regulated by the CaMV 35S promoter, together with an ER-marker or a soluble mRFP, were transiently expressed in *Nicotiana benthamiana* leaves (Fig. 7). Both, the N- and C-terminal *Ib3GGT* fusion proteins were found to localize in the cytosol, similar to the soluble mRFP (Fig. 7C, 7D). However, some signal of the eGFP-*Ib3GGT* fusions was also observed in the nucleus. Additionally, when *Ib3GGT*-eGFP was expressed together with ER-mCherry, no co-localization was found (Fig. 7A, 7B). Thus, *Ib3GGT* is a soluble protein in the cytosol and not associated with the ER (Poustka et al., 2007).

Discussion

Further glycosylation modifications of anthocyanins in the cytosol

Further glycosylation of anthocyanins is an essential step in their biosynthesis, accumulation, and stability (Yonekura-Sakakibara et al., 2008; Zhang et al., 2014). In purple sweetpotato, the major anthocyanins include cyanidin and peonidin

3-sophorosides as well as their acylated derivatives (Truong et al., 2009; Lee et al., 2013), indicating that further glycosylation is required for the conversion from anthocyanidin 3-*O*-glucosides into anthocyanidin 3-*O*-sophorosides. In this study, we found that Ib3GGT was responsible for the reaction using UDP-glucose as the sugar donor. Unlike sweetpotato, further glycosylation of anthocyanins is xylosylation catalyzed by UGT79B1 in *Arabidopsis* (Saito et al., 2013). Apparently, *Arabidopsis* lacks the enzymes to form cyanidin 3-*O*-sophoroside using cyanidin 3-*O*-glucosides as the substrate for further modification, as the overexpressing Ib3GGT in *Arabidopsis* only showed a peak of cyanidin 3-*O*-sophoroside without a change in other anthocyanin components. Therefore, Ib3GGT is a key player of anthocyanin for further glycosylation modifications in sweetpotato.

Further glycosylation is a critical step determining the subsequent anthocyanin modifications, such as malonylation and acylation (Yonekura-Sakakibara et al., 2008; Andersen and Jordheim, 2010). The Ib3GGT as UDP-glycosyltransferase can add a sugar residue to anthocyanidin 3-*O*-glucosides but not anthocyanidin 3,5-*O*-diglucosides (Fig. 2A, 2C). Thus, we conclude that 5GT catalyzes a glucose molecule into the 5th position of C-ring that can hinder the transfer of a glucose molecule into the 2'' position of anthocyanidin 3-*O*-glucosides. In certain order modification of anthocyanin, this phenomenon might be a crucial factor for providing the condition for subsequent modification in anthocyanin. Ib3GGT cannot catalyze flavonol-3-*O*-glucoside as acceptor substrate, demonstrating that Ib3GGT has substrate specificity in purple sweetpotato (Fig. 2D). However, At3GGT can catalyze flavonol-3-*O*-glucoside and anthocyanin-3-*O*-glucoside as a broad substrate in *Arabidopsis* (Saito et al., 2013).

As a primary sedative mechanism that maintains metabolic homeostasis in plants, glycosylation contributes to the diversity in synthesizing various secondary plant metabolites, thereby altering the biological functions of these metabolites (Jones and Vogt, 2001; Gachon et al., 2005). Apparently, divergence occurs among species by adaption of glycosyltransferase substrate specificity. In peach, the PpUGT79B is responsible for glycosylation by adding a rhamnoside molecule to anthocyanidin 3-*O*-glucosides forming the anthocyanidin 3-*O*-rutinoside (Cheng et al., 2014). Ib3GGT was functional in transgenic *Arabidopsis* by producing

anthocyanin-3-sophoroside; while the overexpression of UGT79B1 (At3GGT) in sweetpotato did not catalyze the production of anthocyanin 3-*O*-glucoside-2''-*O*-xylose (Supplementary Fig. S10). These findings indicated that anthocyanin glycosylation in sweetpotato diverges from that of *Arabidopsis* towards the specific sugar acceptor. Interestingly, the further glycosylation of anthocyanin occurs in the cytosol, not like other UGTs that are mainly ER membrane-bound enzymes (Poustka et al., 2007; Zhao, 2015).

535 **Key amino acids in UGTs affect both sugar donor preference and regioselectivity**

536 The phylogenetic comparisons of flavonoid GGTs suggested that the potentially
537 conserved amino acid residues are involved in further substrate-selectivity. Four
538 amino acid residues (Trp-334/Gln-337/Glu-360/His-352 in Ib3GGT) are generally
539 conserved across all known flavonoid 3-*O*-glucoside-2''-*O*-glycosyltransferases. The
540 close relationship between Ib3GGT and UGT79B1 in the phylogenetic tree also
541 indicated that the sugar donor selectivity of flavonoid GGTs was established after
542 species differentiation (Saito et al., 2013). In sweetpotato, Ib3GGT accepts
543 UDP-glucose as sugar donor to conjugate to anthocyanins, such as cyanidin
544 3-*O*-glucoside or peonidin 3-*O*-glucoside. Interestingly, *Arabidopsis* also has a
545 UDP-glucose:flavonoid 3-*O*-glucoside-2''-*O*-glycosyltransferase (At3GGTF), which
546 preferentially uses flavonol 3-*O*-glucoside and UDP-glucose as substrates (Kubo et al.,
547 2007). Thr-138 as the key residue for UDP-glucose recognition is also conserved in
548 glucosyltransferases that uses UDP-glucose as sugar donor in morning glory,
549 *Arabidopsis* (At3GGTF), *Ricinus communis*, and *Glysin max* (Supplementary Fig. S6).
550 Corresponding to the Thr-138 residue, the Ile-142 is the residue for UDP-xylose
551 recognition in *Arabidopsis*. Instead, the corresponding sites of Ile in *Camelina sativa*,
552 Thr in *Tarenaya hassleriana* and *Brassica napus*, and Val in *Eucalyptus grandis* are
553 also responsible for recognizing UDP-xylose (Supplementary Fig. S6).

554 **Anthocyanin glycosyltransferases are regulated by transcription factors (TFs)**

555 Anthocyanin biosynthesis is a finely regulated system involving multiple TFs
556 associated with plant development (Pireyre and Burow, 2015; Xu et al., 2015). For
557 example, the temporal and spatial regulation of anthocyanin production in flowers
558 mediated by TFs, R2R3-MYB, basic Helix-Loop-Helix (bHLH), or WD40 type
559 (reviewed in Davies et al., 2012) brings the colorful variation in the world. Hitherto,

the only well-characterized TF in sweetpotato is R2R3-MYB type IbMYB1, which controls anthocyanin biosynthesis specifically in tuberous roots by inducing all the structural anthocyanin genes (Mano et al., 2007). In this study, the accumulation of anthocyanin in different organs was strongly associated with the *Ib3GGT* expression, which indicates its divergent regulation during plant development. Importantly, the activation of the *Ib3GGT* promoter by IbMYB1 confirmed that *Ib3GGT* is highly regulated by the TF in storage roots. The relatively low level of *IbMYB1* transcript in the leaves might reflect its tissue specificity. In *Arabidopsis*, it is well-documented that the R2R3-MYB TF can induce glycosyltransferases such as UGT79B1 (Tohge et al., 2005; Yonekura-Sakakibara et al., 2008; Stracke et al., 2010).

In summary, sweetpotato Ib3GGT catalyzes the anthocyanidin 3-*O*-glucosides into anthocyanidin 3-*O*-sophorosides using UDP-glucose as a sugar donor. The Thr-138 of Ib3GGT is a key residue for sugar donor selectivity in the further glycosylation that contributes to the stability and diversity of anthocyanins. The Ib3GGT glycosylation occurs in the cytosol and is regulated by IbMYB1 TF. The present study provides further insights regarding the glycosylation enzymes involved in secondary metabolism in divergence that can assist in developing a useful approach to diversifying certain flavonoids in crops.

Supplementary data

Fig. S1. UDP-glucose binding sites of Ib3GGT. (A) Chemical structure of UDP-glucose. (B,C) Close-up views of the interactions of uridine moiety (B), and diphosphate moiety. (D) Overall structure of Ib3GGT modeling structure. UDP-glucose is showed with ball-and-stick model and the protein structure is showed with cartoon; the helix/sheet/loops are showed in cyan/red/magenta, respectively.

Fig. S2. Structure alignment of UGT homologs. (A) All structures are showed with ribbon and superposed by Coot (Emsley et al., 2010). The PDB numbers of these structures are 2ACW, 2C1Z, 2PQ6, 2VG8, 3HBF, 3WC4, 5GL5, 5NLM, 5TMB and 5U6M. (B) Ligands in superposed structures are showed with sticks.

Fig. S3. Sequence alignment of Ib3GGT with 2VG8 and 2VCH proteins using SWISS-MODEL and I-TASSER.

590 **Fig. S4.** Sugar donor binding sites in Ib3GGT and At3GGT. (A, B, C, D) Close-up
591 views of the interactions of UDP-glucose with Ib3GGT (A), UDP-glucose with
592 At3GGT (B), UDP-xylose with Ib3GGT (C), and UDP-xylose with At3GGT (D).
593 UDP-glucose and UDP-xylose are showed with stick-and-ball in magenta and red,
594 respectively; side chain residues are showed with stick; the hydrogen bonds are
595 indicated by dashed lines.

596 **Fig. S5.** Difference in binding affinity of sugar analogies. (A, B, C, D)
597 Ball-and-stick model of UDP-galactose (A), UDP-glucose (B), UDP-arabinose (C),
598 and UDP-xylose (D). They are showed in grey, magenta, pink, and yellow,
599 respectively. Chemical structure of galactose, glucose, arabinose, and xylose are
600 showed in the corresponding positions.

601 **Fig. S6.** Amino acid sequence comparison of GGT analogies with Ib3GGT. The
602 Thr-138 site is boxed.

603 **Fig. S7.** Anthocyanin pigmentation and component profiles in seedlings of the
604 *ugt79b1-2* mutant and *Ib3GGT*-overexpressing *ugt79b1-2* transgenic line
605 (*ugt19b1-2*+*Ib3GGT*-OE).

606 **Fig. S8.** *Ib3GGT* promoter sequence showing the two G-box sites.

607 **Fig. S9.** Leaf and root phenotypes of field-grown wild-type (WT), RNAi-*Ib3GGT*-2
608 and OE-*Ib3GGT*-2 plant lines.

609 **Fig. S10.** Analysis of anthocyanin compounds in wild-type (WT) and
610 *At3GGT*-overexpressing (OE-*At3GGT*) sweetpotato by HPLC-MS.

611 **Table S1:** Anthocyanin compounds in sweetpotato (Tian et al., 2005) and *Arabidopsis*
612 (Tohge et al., 2005)

613 **Table S2:** List of primers for gene expression analysis in *Arabidopsis* and sweetpotato
614 plant lines.

615 **Acknowledgments**

616 This work was supported by the grants from the National Natural Science Foundation
617 of China (31501357, 31771854), the Collaborative Innovation Action – Agricultural

618 Science and Technology Innovation Program of Chinese Academy of Agricultural
619 Sciences (CAAS-XTCX2016009), and XDPB0402 of CAS. We thank Prof. Cathie
620 Martin from JIC for providing helpful suggestions and corrections for the manuscript.
621 We also thank our colleagues Prof. Peng Zhang for protein structure analysis, Miss
622 Yuanhong Shan for HPLC-MS-MS analysis and Mr. XiaoshuGao for confocal
623 microscopy.

624 **Reference**

- 625 **Andersen ØM, Jordheim M.** 2010. Anthocyanins. In: Encyclopedia of Life Sciences
626 (ELS). Chichester: John Wiley and Sons Ltd.
- 627 **Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F,**
628 **Gallo CT, Bertoni M, Bordoli L.** 2014. SWISS-MODEL: modelling protein tertiary
629 and quaternary structure using evolutionary information. *Nucleic Acids Research* **42**,
630 W252-W258.
- 631 **Brazier-Hicks M, Offen WA, Gershater MC, Revett TJ, Lim EK, Bowles DJ,**
632 **Davies G J, Edwards R.** 2007. Characterization and engineering of the bifunctional
633 N- and O-glucosyltransferase involved in xenobiotic metabolism in plants.
634 *Proceedings of the National Academy of Sciences, USA* **104**, 20238-20243.
- 635 **Breton C, Šnajdrová L, Jeanneau C, Koča J, Imberty A.** 2006. Structures and
636 mechanisms of glycosyltransferases. *Glycobiology* **16**, 29R-37R.
- 637 **Caputi L, Malnoy M, Goremykin V, Nikiforova S, Martens S.** 2012. A
638 genome-wide phylogenetic reconstruction of family 1 UDP-glycosyltransferases
639 revealed the expansion of the family during the adaptation of plants to life on land.
640 *Plant Journal* **69**, 1030-1042.
- 641 **Cheng J, Wei G, Zhou H, Gu C, Vimolmangkang S, Liao L, Han Y.** 2014.
642 Unraveling the mechanism underlying the glycosylation and methylation of
643 anthocyanins in peach. *Plant Physiology* **166**, 1044-1058.
- 644 **Claudia C, Philippe R, Edith F, Vincent B, Yumei Z, Sharyn EP, Juan J R,**
645 **Martin Y, Christophe D.** 2017. The class III peroxidase PRX17 is a direct target of
646 the MADS-box transcription factor AGAMOUS-LIKE15 (AGL15) and participates in

647 lignified tissue formation. *New Phytologist* **213**, 250-263.

648 **Clough SJ, Bent AF.** 1998. Floral dip: a simplified method for
649 *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* **16**,
650 735-743.

651 **Davies KM, Albert NW, Schwinn KE.** 2012. From landing lights to mimicry: the
652 molecular regulation of flower colouration and mechanisms for pigmentation
653 patterning. *Funct. Plant Biology* **39**, 619-638.

654 **de Pascual-Teresa S, Sanchez-Ballesta MT.** 2008. Anthocyanins: from plant to
655 health. *Phytochemistry Reviews* **7**, 281-299.

656 **Emsley P, Lohkamp B, Scott WG, Cowtan K.** 2010. Features and development of
657 Coot. *Acta Crystallographica Section D: Structural Biology* **66**, 486-501.

658 **Fedoroff NV, Furtak DB, Nelson OE.** 1984. Cloning of the bronze locus in maize by
659 a simple and generalizable procedure using the transposable controlling element
660 Activator (Ac). *Proceedings of the National Academy of Sciences, USA* **81**,
661 3825-3829.

662 **Gachon CM, Langlois-Meurinne M, Saindrenan P.** 2005. Plant secondary
663 metabolism glycosyltransferases: the emerging functional analysis. *Trends in Plant*
664 *Science* **10**, 542-549.

665 **Glover BJ, Martin C.** 2012. Anthocyanins. *Current Biology* **22**, R147-R150.

666 **Griesser M, Hoffmann T, Bellido ML, Rosati C, Fink B, Kurtzer R, Aharoni A,**
667 **Munoz-Blanco J, Schwab W.** 2008a. Redirection of flavonoid biosynthesis through
668 the down-regulation of an anthocyanidin glucosyltransferase in ripening strawberry
669 fruit. *Plant Physiology* **146**, 1528-1539.

670 **Griesser M, Vitzthum F, Fink B, Bellido ML, Raasch C, Munoz-Blanco J,**
671 **Schwab W.** 2008b. Multi-substrate flavonol O-glucosyltransferases from strawberry
672 (*Fragaria* × *anayasa*) achene and receptacle. *Journal of Experimental Botany* **59**,
673 2611-2625.

674 **He J, Giusti MM.** 2010. Anthocyanins: natural colorants with health-promoting
675 properties. *Annual Review of Food Science and Technology* **1**, 163-187.

676 **He ZH, Cheeseman I, He D, Kohorn BD.** 1999. A cluster of five cell
677 wall-associated receptor kinase genes, Wak1-5, are expressed in specific organs of
678 Arabidopsis. *Plant Molecular Biology* **39**, 1189-1196.

679 **Hiromoto T, Honjo E, Noda N, Tamada T, Kazuma K, Suzuki M, Blaber M,**
680 **Kuroki R.** 2015. Structural basis for acceptor-substrate recognition of
681 UDP-glucose:anthocyanidin 3-O-glucosyltransferase from *Clitoria ternatea*. *Protein*
682 *Science* **24**, 395-407.

683 **Hiromoto T, Honjo E, Tamada T, Noda N, Kazuma K, Suzuki M, Kuroki R.**
684 2013. Crystal structure of UDP-glucose:anthocyanidin 3-O-glucosyltransferase from
685 *Clitoria ternatea*. *Journal of Synchrotron Radiation* **20**, 894-898.

686 **Holm L, Laakso LM.** 2016. Dali server update. *Nucleic Acids Research* **44**, 351-355.

687 **Hsu TM, Welner DH, Russ ZN, Cervantes B, Prathuri RL, Adams PD, Dueber**
688 **JE.** 2018. Employing a biochemical protecting group for a sustainable indigo dyeing
689 strategy. *Nature Chemical Biology* **14**, 256-261.

690 **Ito T, Motohashi R, Kuromori T, Noutoshi Y, Seki M, Kamiya A, Mizukado S,**
691 **Sakurai T, Shinozaki K.** 2005. A resource of 5,814 dissociation transposon-tagged
692 and sequence-indexed lines of Arabidopsis transposed from start loci on chromosome
693 5. *Plant Cell Physiology* **46**, 1149-1153.

694 **Jones P, Vogt T.** 2001. Glycosyltransferases in secondary plant metabolism:
695 tranquilizers and stimulant controllers. *Planta* **213**, 164-174.

696 **Kim HS, Kim B, Sung S, Kim M, Mok H, Chong Y, Ahn J.** 2013. Engineering
697 flavonoid glycosyltransferases for enhanced catalytic efficiency and extended
698 sugar-donor selectivity. *Planta* **238**, 683-693.

699 **Kovinich N, Kanyanjan G, Chanoca A, Riedl K, Otegui MS, Grotewold E.** 2014.
700 Not all anthocyanins are born equal: distinct patterns induced by stress in Arabidopsis.
701 *Planta* **240**, 931-940.

702 **Kroon J, Souer E, de Graaff A, Xue Y, Mol J, Koes R.** 1994. Cloning and structural
703 analysis of the anthocyanin pigmentation locus *Rt* of *Petunia hybrida*: characterization
704 of insertion sequences in two mutant alleles. *Plant Journal* **5**, 69-80.

705 **Kubo A, Arai Y, Nagashima S, Yoshikawa T.** 2004. Alteration of sugar donor
706 specificities of plant glycosyltransferases by a single point mutation. Archives of
707 Biochemistry and Biophysics **429**, 198-203.

708 **Kubo H, Nawa N, Lupsea SA.** 2007. Anthocyaninless1 gene of Arabidopsis thaliana
709 encodes a UDP-glucose:flavonoid-3-O-glucosyltransferase. Journal of Plant Research
710 **120**, 445-449.

711 **Kuromori T, Hirayama T, Kiyosue Y, Takabe H, Mizukado S, Sakurai T,**
712 **Akiyama K, Kamiya A, Ito T, Shinozaki K.** 2004. A collection of 11800 single-copy
713 Ds transposon insertion lines in Arabidopsis. Plant Journal **37**, 897-905.

714 **Lairson LL, Henrissat B, Davies GJ, Withers SG.** 2008. Glycosyltransferases:
715 structures, functions, and mechanisms. Annual Review of Biochemistry **77**, 521-555.

716 **Leuzinger K, Dent M, Hurtado J, Stahnke J, Lai H, Zhou X, Chen Q.** 2013.
717 Efficient agroinfiltration of plants for high-level transient expression of recombinant
718 proteins. Journal of Visualized Experiments **77**, 50521.

719 **Lee MJ, Park JS, Choi DS, Jung MY.** 2013. Characterization and quantitation of
720 anthocyanins in purple-fleshed sweet potatoes cultivated in Korea by HPLC-DAD and
721 HPLC-ESI-QTOF-MS/MS. Journal of Agricultural and Food Chemistry **61**,
722 3148-3158.

723 **Mano H, Ogasawara F, Sato K, Higo H, Minobe Y.** 2007. Isolation of a regulatory
724 gene of anthocyanin biosynthesis in tuberous roots of purple-fleshed sweetpotato.
725 Plant Physiology **143**, 1252-1268.

726 **Matsuba Y, Sasaki N, Tera M, Okamura M, Abe Y, Okamoto E, Nakamura H,**
727 **Funabashi H, Takatsu M, Saito M, Matsuoka H, Nagasawa K, Ozeki Y.** 2010. A
728 novel glucosylation reaction on anthocyanins catalyzed by acyl-glucose-dependent
729 glucosyltransferase in the petals of carnation and delphinium. Plant Cell **22**,
730 3374-3389.

731 **Modolo LV, Li L, Pan HY, Blount JW, Dixon RA, Wang XQ.** 2009. Crystal
732 structures of glycosyltransferase UGT78G1 reveal the molecular basis for
733 glycosylation and deglycosylation of (iso)flavonoids. Journal of Molecular Biology

734 **392**, 1292-1302.

735 **Moglia A, Lanteri S, Comino C, Hill L, Knevitt D, Cagliero C, Rubiolo P,**
736 **Bornemann S, Martin C.** 2014. Dual catalytic activity of
737 hydroxycinnamoyl-Coenzyme A quinate transferase from tomato allows it to
738 moonlight in the synthesis of both mono-and dicaffeoylquinic acids. *Plant Physiology*.
739 **166**, 1777-1787.

740 **Montefiori M, Espley RV, Stevenson D, Cooney J, Datson PM, Saiz A, Atkinson**
741 **RG, Hellens RP, Allan AC.** 2011. Identification and characterisation of F3GT1 and
742 F3GGT1, two glycosyltransferases responsible for anthocyanin biosynthesis in
743 red-fleshed kiwifruit (*Actinidia chinensis*). *Plant Journal* **65**, 106-118.

744 **Morita Y, Hoshino A, Kikuchi Y, Okuhara H, Ono E, Tanaka Y, Fukui Y, Saito N,**
745 **Nitasaka E, Noguchi H.** 2005. Japanese morning glory dusky mutants displaying
746 reddish-brown or purplish-gray flowers are deficient in a novel glycosylation enzyme
747 for anthocyanin biosynthesis, UDP-glucose:anthocyanidin 3-O-glucoside-2
748 "-O-glucosyltransferase, due to 4-bp insertions in the gene. *Plant Journal* **42**, 353-363.

749 **Nelson BK, Cai X, Nebenfuhr A.** 2007. A multicolored set of in vivo organelle
750 markers for co-localization studies in Arabidopsis and other plants. *Plant Journal* **51**,
751 1126-1136.

752 **Noguchi A, Horikawa M, Fukui Y, Fukuchi-Mizutani M, Iuchi-Okada A,**
753 **Ishiguro M, Kiso Y, Nakayama T, Ono E.** 2009. Local differentiation of sugar donor
754 specificity of flavonoid glycosyltransferase in Lamiales. *Plant Cell* **21**, 1556-1572.

755 **Offen W, Martinez-Fleites C, Yang M, Kiat-Lim E, Davis BG, Tarling CA, Ford**
756 **CM, Bowles DJ, Davies GJ.** 2006. Structure of a flavonoid glucosyltransferase
757 reveals the basis for plant natural product modification. *EMBO Journal* **25**,
758 1396-1405.

759 **Ono E, Fukuchi-Mizutani M, Nakamura N, Fukui Y, Yonekura-Sakakibara K,**
760 **Yamaguchi M, Nakayama T, Tanaka T, Kusumi T, Tanaka Y.** 2006. Yellow
761 flowers generated by expression of the aurone biosynthetic pathway. *Proceedings of*
762 *the National Academy of Sciences, USA* **103**, 11075-11080.

763 **Ono E, Homma Y, Horikawa M, Kunikane-Doi S, Imai H, Takahashi S, Kawai Y,**
764 **Ishiguro M, Fukui Y, Nakayama T.** 2010. Functional differentiation of the
765 glycosyltransferases that contribute to the chemical diversity of bioactive flavonol
766 glycosides in grapevines (*Vitis vinifera*). *Plant Cell* **22**, 2856-2871.

767 **Osmani SA, Bak S, Møller BL.** 2009. Substrate specificity of plant UDP-dependent
768 glycosyltransferases predicted from crystal structures and homology modeling.
769 *Phytochemistry* **70**, 325-347.

770 **Pireyre M, Burow M.** 2015. Regulation of MYB and bHLH transcription factors: a
771 glance at the protein level. *Molecular Plant* **8**, 378-388.

772 **Poustka F, Irani N G, Feller A, Lu Y, Pourcel L, Frame K, Grotewold E.** 2007. A
773 trafficking pathway for anthocyanins overlaps with the endoplasmic
774 reticulum-to-vacuole protein-sorting route in *Arabidopsis* and contributes to the
775 formation of vacuolar inclusions. *Plant Physiology* **145**, 1323-1335.

776 **Saito K, Yonekura-Sakakibara K, Nakabayashi R, Higashi Y, Yamazaki M,**
777 **Tohge T, Fernie AR.** 2013. The flavonoid biosynthetic pathway in *Arabidopsis*:
778 structural and genetic diversity. *Plant Physiology and Biochemistry* **72**, 21-34.

779 **Sasaki N, Nishizaki Y, Ozeki Y, Miyahara T.** 2014. The role of acyl-glucose in
780 anthocyanin modifications. *Molecules* **19**, 18747-18766.

781 **Sawada S, Suzuki H, Ichimaida F, Yamaguchi MA, Iwashita T, Fukui Y, Hemmi**
782 **H, Nishino T, Nakayama T.** 2005. UDP-glucuronic acid:anthocyanin
783 glucuronosyltransferase from red daisy (*Bellis perennis*) flowers. *Enzymology and*
784 *phylogenetics of a novel glucuronosyltransferase involved in flower pigment*
785 *biosynthesis. Journal of Biological Chemistry* **280**, 899-906.

786 **Shao H, He XZ, Achnine L, Biount JW, Dixon RA, Wang XQ.** 2005. Crystal
787 structures of a multifunctional triterpene/flavonoid glycosyltransferase from
788 *Medicago truncatula*. *Plant Cell* **17**, 3141-3154.

789 **Shen Y, Chen Y, Wu J, Shaner NC, Robert E, Campbell RE.** 2017. Engineering of
790 mCherry variants with long Stokes shift, red-shifted fluorescence, and low
791 cytotoxicity. *PLoS One* **12(2)**, e0171257.

792 **Stamatakis A.** 2014. RAxML version 8: a tool for phylogenetic analysis and
793 post-analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313.

794 **Stracke R, Jahns O, Keck M, Tohge T, Niehaus K, Fernie AR, Weisshaar B.** 2010.
795 Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol
796 glycoside accumulation in *Arabidopsis thaliana* plants reveals MYB11-, MYB12- and
797 MYB111-independent flavonol glycoside accumulation. *New Phytologist* **188**,
798 985-1000.

799 **Sun Y, Li H, Huang JR.** 2012. *Arabidopsis* TT19 functions as a carrier to transport
800 anthocyanin from the cytosol to tonoplasts. *Molecular Plant* **5**, 387-400.

801 **Sun W, Liang L, Meng X, Li Y, Gao F, Liu X, Wang L.** 2016. Biochemical and
802 molecular characterization of a flavonoid 3-O-glycosyltransferase responsible for
803 anthocyanins and flavonols biosynthesis in *Freesia hybrida*. *Front Plant Science* **7**,
804 410.

805 **Tamura K, Stecher G, Peterson D, Filipski A, Kumar S.** 2013. MEGA6: molecular
806 evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30**,
807 2725-2729.

808 **Thompson AMG, Iancu CV, Neet KE, Dean JV, Choe JY.** 2017. Differences in
809 salicylic acid glucose conjugations by UGT74F1 and UGT74F2 from *Arabidopsis*
810 *thaliana*. *Scientific Reports* **7**, 46629.

811 **Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima J, Awazuhara M, Inoue E,**
812 **Takahashi H, Goodenowe DB, Kitayama M.** 2005. Functional genomics by
813 integrated analysis of metabolome and transcriptome of *Arabidopsis* plants
814 over-expressing an MYB transcription factor. *Plant Journal* **42**, 218-235.

815 **Truong VD, Deighton N, Thompson RT, McFeeters RF, Dean LO, Pecota KV,**
816 **Yencho GC.** 2009. Characterization of anthocyanins and anthocyanidins in
817 purple-fleshed sweetpotatoes by HPLC-DAD/ESI-MS/MS. *Journal of Agricultural*
818 *and Food Chemistry* **58**, 404-410.

819 **Tian Q, Konczak I, Schwartz SJ.** 2005. Probing anthocyanin profiles in purple

820 sweet potato cell line (*Ipomoea batatas* L. Cv. Ayamurasaki) by high-performance
 821 liquid chromatography and electrospray ionization tandem mass spectrometry. *Journal*
 822 *of Agricultural and Food Chemistry* **53**, 6503-6509.

823 **Wang H, Fan W, Li H, Yang J, Huang J, Zhang P.** 2013. Functional
 824 characterization of dihydroflavonol-4-reductase in anthocyanin biosynthesis of purple
 825 sweet potato underlies the direct evidence of anthocyanins function against abiotic
 826 stresses. *PLoS One* **8**, e78484.

827 **Wetterhorn KM, Newmister SA, Caniza RK, Busman K, McCormick SP,**
 828 **Berthiller F, Adam G, Rayment I.** 2016. Crystal structure of Os79 (Os04g0206600)
 829 from *Oryza sativa*: A UDP-glucosyltransferase involved in the detoxification of
 830 deoxynivalenol. *Biochemistry* **55**, 6175-6186.

831 **Xu W, Dubos C, Lepiniec L.** 2015. Transcriptional control of flavonoid biosynthesis
 832 by MYB–bHLH–WDR complexes. *Trends Plant Science* **20**, 176-185.

833 **Yang J, Bi HP, Fan WJ, Zhang M, Wang HX, Zhang P.** 2011. Efficient
 834 embryogenic suspension culturing and rapid transformation of a range of elite
 835 genotypes of sweet potato (*Ipomoea batatas* [L.] Lam.). *Plant Science* **181**, 701-711.

836 **Yang JY, Yan RX, Roy A, Xu D, Poisson J, Zhang Y.** 2015. The I-TASSER Suite:
 837 protein structure and function prediction. *Nature Methods* **12**, 7-8.

838 **Yonekura-Sakakibara K, Nakayama T, Yamazaki M, Saito K.** 2008. Modification
 839 and stabilization of anthocyanins. In: *Anthocyanins*. Springer New York, 169-190.

840 **Yonekura-Sakakibara K, Fukushima AR, Nakabayashi K, Hanada F, Matsuda S,**
 841 **Sugawara E, Inoue T, Kuromori T, Ito K, Shinozaki B.** 2012. Two
 842 glycosyltransferases involved in anthocyanin modification delineated by
 843 transcriptome independent component analysis in *Arabidopsis thaliana*. *Plant Journal*
 844 **69**, 154-167.

845 **Yonekura-Sakakibara K, Hanada K.** 2011. An evolutionary view of functional
 846 diversity in family 1 glycosyltransferases. *Plant Journal* **66**, 182-193.

847 **Yonekura-Sakakibara K, Nakabayashi R, Sugawara S, Tohge T, Ito T, Koyanagi**
 848 **M, Saito K.** 2014. A flavonoid 3-O-glucoside:2''-O-glucosyltransferase responsible

849 for terminal modification of pollen-specific flavonols in *Arabidopsis thaliana*. *Plant*
850 *Journal* **79**, 769-782.

851 **Zhang Y, Butelli E, Martin C.** 2014. Engineering anthocyanin biosynthesis in plants.
852 *Current Opinion in Plant Biology* **19**, 81-90.

853 **Zhao J, Huhman D, Shadle G, He XZ, Sumner LW, Tang Y, Dixon RA.** 2011.
854 MATE2 mediates vacuolar sequestration of flavonoid glycosides and glycoside
855 malonates in *Medicago truncatula*. *Plant Cell* **23**, 1536-1555.

856 **Zhao J.** 2015. Flavonoid transport mechanisms: how to go, and with whom. *Trends*
857 *Plant Science* **20**, 576-585.

858

859

860

861

862

863

864

865

866

867

868

869

870

871 **Tables**

872 **Table 1:** Sugar substrate specificity of different 3GGT proteins

Sugar donor	Relative activity (%)					
	Ib3GGT	At3GGT	Ib3GGT ^{T1} _{38I}	Pp3GGT	La3GG _T	Cs3GGT
UDP-glucose	100 ± 8.5	ND	ND	100 ± 10.5	100 ± 9.3	ND
UDP-xylose	10.4 ± 1.2	100 ± 15.0	6.1 ± 0.79	9.4 ± 1.4	11.4 ± 1.2	100 ± 11.7
UDP-galactose	ND	ND	ND	--	--	--
UDP-arabinose	ND	ND	ND	--	--	--

873 Note: The reactions were performed with cyanidin 3-*O*-glucoside as the sugar
874 acceptor. ND, not detected; --, not tested. The method for calculation of sugar donor
875 specificity is according to Yonekura-Sakakibara et al. (2012). Abbreviations for
876 species: At, *Arabidopsis thaliana*; Ib, *Ipomoea batatas*; Pp, *Prunus persica*; La,
877 *Lupinus angustifolius*; Cs, *Camelina sativa*.

878 **Figure legends**

879 **Figure 1** Alignment of amino acid sequences and phylogenetic tree of flavonoid
880 glycosyltransferases. **(A)** Multiple alignments of amino acid sequences of sweetpotato
881 Ib3GGT, morning glory Ip3GGT, and *Arabidopsis* At3GGT. The underlined
882 nucleotides represent the putative C-terminal UDP binding motif for
883 glycosyltransferases (PSPG). **(B)** Non-rooted molecular phylogenetic tree of
884 flavonoid glycosyltransferases from selected plant UDP-glycosyltransferases. All
885 amino acids were aligned using CLUSTALW. Bootstrap values from 100 retrials are
886 indicated at each branch. The scale shows 0.2 amino acid substitution per site. The
887 GenBank accession numbers or genome sequence codes for the sequences are shown
888 in parentheses: AtA3G2"XylT (NP_200217); AtF3G2"GlcT (NP_200212);
889 BpA3G2"GlcAT (AB190262); CmF7G2"RhaT (AY048882); CrF3G6"GlcT
890 (BAH80312); CsF3G2"XylT(XP_018450414); CsaF3G2"GlcT(CCG85331);
891 CsiF7G6"RhaT (NP_001275829); Ib3GGT (ABL74480); IbF3G2"XylT
892 (XP_019151635); IpA3G2"GlcT (AB192315); InF3G2"GlcT (XP_019194233);
893 ItF3G2"GlcT (itf02g12970.t1); ItF3G2"GlcT (itb02g08330.t1); ItF3G2"XylT
894 (itb03g28310.t1); ItF3G2"XylT (itf03g22690.t2); LaF3G2"GlcT(XP_019424989);
895 PhA3G6"RhaT (CAA81057); PpF3G2"Glc (XP_007213494). Abbreviations for
896 species: Ac, *Actinidia chinensis*; At, *Arabidopsis thaliana*; Bp, *Bellis perennis*; Cm,
897 *Citrus maxima*; Cr, *Catharanthus roseus*; Cs, *Camelina sativa*; Csa, *Crocus sativus*;
898 Csi, *Citrus sinensis*; Ib, *Ipomoea batatas*; In, *Ipomoea nil*; Ip, *Ipomoea purpurea*; Itf,
899 *Ipomoea trifida*; Itl, *Ipomoea triloba*; La, *Lupinus angustifolius*; Ph, *Petunia hybrida*;
900 Pp, *Prunus persica*.

901 **Figure 2** Functional assays of Ib3GGT recombinant protein using UDP-glucose and
902 different acceptor substrates by HPLC. **(A)** Cyanidin 3-*O*-glucoside as acceptor
903 substrate; **(B)** Cyanidin as acceptor substrate. **(C)** Cyanidin 3,5-*O*-diglucoside as
904 acceptor substrate. **(D)** Quercetin 3-*O*-glucoside as acceptor substrate. **(E)** Cyanidin
905 3-*O*-glucoside as acceptor substrate without Ib3GGT protein treatment. **(F)** Peonidin
906 3-*O*-glucoside as acceptor substrate.

907 **Figure 3** Three-dimensional modeling of Ib3GGT and At3GGT interacting with a
908 sugar donor and a glycone acceptor. **(A)** Active center of Ib3GGT showing the key
909 amino acid residues for sugar donor and acceptor positions. **(B)** Docking illustration
910 of sugar donors and a glycone acceptor in the binding pocket of WT and mutant
911 Ib3GGT and At3GGT. The performance of their reactions using cyanidin

912 3-*O*-glucoside with sugar nucleotide UDP-glucose or UDP-xylose is shown in the
913 bottom panel. The percentage of relative enzyme activity is indicated in the
914 parentheses.

915 **Figure 4** Anthocyanin characterization of transgenic *Arabidopsis* plants
916 overexpressing *Ib3GGT* gene. (A) Anthocyanin pigmentation in seedlings of WTs
917 (Col0 and Nossen), *Ib3GGT*-overexpressing Col0 lines (*Ib3GGT*-OE), *ugt79b1*
918 mutant (*ugt79b1-1*), and *Ib3GGT*-overexpressing *ugt79b1* line
919 (*ugt79b1-1*+*Ib3GGT*-OE). (B) The expression of *At3GGT* and *Ib3GGT* in the WT
920 Col0 and *Ib3GGT*-overexpressing Col0 lines (*Ib3GGT*-OE1 and *Ib3GGT*-OE2) by
921 real-time RT-PCR analysis. (C) RT-PCR detection of *At3GGT* and *Ib3GGT*
922 expression in the WTs (Col0 and Nossen), two independent *Ib3GGT*-overexpressing
923 Col0 lines (*Ib3GGT*-OE1 and *Ib3GGT*-OE2), *ugt79b1* mutant, and two
924 *Ib3GGT*-overexpressing *ugt79b1* lines (*ugt79b1-1*+*Ib3GGT*-OE1 and
925 *ugt79b1-1*+*Ib3GGT*-OE2). Different letters indicate significant differences (one-way
926 ANOVA, $P < 0.05$). (D) Anthocyanin component profiles by HPLC/PDA/MS in the
927 seedlings of WTs (Col0 and Nossen), *ugt79b1* mutant, and *Ib3GGT*-overexpressing
928 *ugt79b1* line (*ugt79b1-1*+*Ib3GGT*-OE). Blue arrows indicate the new peaks of
929 cyanidin 3-*O*-sophoroside. (E) Anthocyanin content in the WTs (Col0 and Nossen),
930 two independent *Ib3GGT*-overexpressing Col0 lines (*Ib3GGT*-OE1 and
931 *Ib3GGT*-OE2), *ugt79b1* mutant, and two *Ib3GGT*-overexpressing *ugt79b1* line
932 (*ugt79b1-1*+*Ib3GGT*-OE1 and *ugt79b1-1*+*Ib3GGT*-OE2).

933 **Figure 5** Correlation of anthocyanin accumulation and gene expression in various
934 organs of sweetpotato cv. Ayamurasaki. (A) Profiles of anthocyanin accumulation,
935 and *Ib3GGT* and *IbMYB1* transcript levels as detected by qRT-PCR in different organs.
936 Lf1, Lf2, Lf3, and Lf4 represent leaves of different developmental stages: St, stem;
937 Ft1, white fibrous root; Ft2, red fibrous root; Dt, developing root; Mt, mature root.
938 Values are mean $\pm$ SD (n = 6). (B) Luciferase assay of *Ib3GGT* promoter activity
939 (reporter) regulated by *IbMBY1* (effector) in agroinfiltrated tobacco leaves.

940 **Figure 6** Anthocyanin characterization of wildtype and *Ib3GGT* transgenic
941 sweetpotato plants. (A) Pot-grown plant phenotypes. WT, wild type; OE-*Ib3GGT* line,
942 transgenic plants overexpressing *Ib3GGT*; RNAi-*Ib3GGT* line, *Ib3GGT* RNAi

943 transgenic plants. **(B)** Anthocyanin pigmentation in top leaves of WT,
 944 RNAi-Ib3GGT-2, and OE-Ib3GGT-2 plants. Both the adaxial (left) and abaxial (right)
 945 leaf surfaces are shown. **(C)** Relative transcription levels of native *Ib3GGT* and the
 946 *Ib3GGT* transgene in WT and transgenic lines assessed by qRT-PCR. **(D)**
 947 Anthocyanin content in WT, RNAi-Ib3GGT, and OE-Ib3GGT plant lines. Different
 948 letters indicate significant differences (one-way ANOVA, $P < 0.05$). **(E)** Component
 949 profiles of anthocyanins in WT, RNAi-Ib3GGT-2, and OE-Ib3GGT-2 plants, as
 950 assessed by HPLC. **(F)** Anthocyanin autofluorescence in leaf epidermal cells of
 951 RNAi-Ib3GGT-2 and OE-Ib3GGT-2 plants.

952 **Figure 7** Subcellular localization of Ib3GGT after transient expression in *Nicotiana*
 953 *benthamiana* leaves. **(A)** Optical section through a pavement cell, co-expressing
 954 Ib3GGT-eGFP (left panel) and ER-mCherry (middle panel). GFP and mCherry signals
 955 are distinct in the merged image (right panel). The outlined region is shown at a
 956 higher magnification in (B). **(B)** Relative fluorescence intensity along the axis is
 957 marked by the dotted arrow in the right panel. **(C)** Co-expression of eGFP-Ib3GGT
 958 (left panel) and soluble mRFP (middle panel). White pixels of the merged image show
 959 an overlay of both channels (right panel). The marked region was enlarged in (D). **(D)**
 960 Intensity plot, as presented before. eGFP-Ib3GGT and soluble mRFP were
 961 co-localized in the cytoplasm. Scale bar, 15 μ m.

962

A

SWEET_POTATO_3GGT	MG...SQATTHHWAMYPWFGVGHLLTAFFRLANKLASKGHRISFLIPKNTQSKLASFNLPHLVSFVPI TVPSI PGLPPG	76
MORNING_GLORY_3GGT	MG...SQATTYHWAMYPWFGVGHLLTGFFRLANKLAGKGRISFLIPKNTQSKLESFNLPHLVSFVPI VVPSI PGLPPG	76
ARABIDOPSIS_3GGTF	MGVFGSNESSSVSI VMYPWLAFGHMTFFLLHLSNKLAEKGHKIVFLLPKKALNQLLEPLNLYPNLITFHTISIPQVKGGLPPG	80
ARABIDOPSIS_3GGT	MG.....SKFHAFMFPWFGFGHMTAFLHLANKLAEKDKITFLLPKKARKQLESNLNFFDCIVFQTLTIPSVQGLPPG	73
SWEET_POTATO_3GGT	AETISDVPPFSSTHLLMEAMDKTQTDIEIILKNLEVDVVFDETHWLPGLARKIGIKSVFYSTISPLMHWGFALSPERR...	153
MORNING_GLORY_3GGT	AETISDVPPFSTHLLMEAMDKTQNDIEIILKDLKVDVVFDETHWLPGLARKIGIKSVFYSTISPLMHWGFALSPERR...	153
ARABIDOPSIS_3GGTF	AETISDVPPFLLTHLLAVAMDKTRPEVETIFRTIKPDLVFYDSAHWPEIAKPIGAKTVCFNI VSAASIALSLVPSAEREV	160
ARABIDOPSIS_3GGT	AETISDIPISLGSLASAMDRIRIQVKEAVSVGKPDLIFFDFAHWPEIAREYGVKSVNEITISAACVAISEVP.....	147
SWEET_POTATO_3GGT	VAGKQLTEADNMKAPASFPDPSIKLHAHEARGFTARTVMKFGGDITFFDRI FTAVSESDGLAYSTCREIEGCFCDYIETQ	233
MORNING_GLORY_3GGT	VVGKQLTEADNMKAPASFPDPSIKLHAHEARGFTARTVMKFGGDITFFDRI FTAVSESDGLAYSTCREIEGCFCDYIETQ	233
ARABIDOPSIS_3GGTF	IDGKENSGEELAKTPLGYSSKVVLRPHEAKSLSF.VVRKHEAIGSFFDGKVTAMRNCDAI AIRTCRETEGKFCDYISRO	239
ARABIDOPSIS_3GGT	..CR..SQDDLGSTPGYSSKVVLRGHEITNSLSF.LSYPEGDGTSFYERIWI GLKNCDVSI SIRTCEMEGKFCDEIENQ	222
SWEET_POTATO_3GGT	FKKPVLLAGPALPVP..SKSTMEQKVSDDLKFKKEGSIYCAFGSECTLRKEQFQELLWGLELTGMPFFAALKAPFGTD	310
MORNING_GLORY_3GGT	FQKPVLLAGPALPVP..SKSTMEQKVSDDLKFKKEGSIYCAFGSECTLRKDKFQELLWGLELTGMPFFAALKAPPFETE	310
ARABIDOPSIS_3GGTF	YSKPVYLLTGPVLPQSGQNPQSLDPQAEWLAKFNHGSVVFCAGSQPVVNKI DQFQELCLGLESTGFPFLVAIKPPSGVS	319
ARABIDOPSIS_3GGT	FQRKVVLLTGPMLPEPDNSKP.LEDQVRQWLKFDPGSVIYCALGSDIILEK DQFQELCLGMEITGLPFLVAIKPPKGS	300
SWEET_POTATO_3GGT	SIEAAIPEELREKIHGKGVHGGWVQCQLFLQHPSVGCVFVSHCGWASLSEALVNDCCI VLLPQVGDQIINARI MSVSLKV	390
MORNING_GLORY_3GGT	SVEAAIPEELREKIQGRGVHGGWVQCQLFLQHPSVGCVFVSHCGWASLSEALVNDCCI VLLPQVGDQIINARI MSVSLKV	390
ARABIDOPSIS_3GGTF	TVEEALPEGFKERVQGRGVVFGWVQQPLVLNHPVSGCVFVSHCGFGSNWESLMSDCCI VLVPPHGEQILNARLMTTEENEV	399
ARABIDOPSIS_3GGT	TIQEALPKGFEBRVKARGVWVGWVQCPLLAHPSIGCFVSHCGFGSNWESLMSDCCI VFI PHLGEQILNTRLMTSEELKV	380
SWEET_POTATO_3GGT	GVEVEKGEEDGVFSRESVCKAVKAVVDEKSEIGREVRGNHDKLRGFLLNADLDSKYVDSFNQKIQDILG...	459
MORNING_GLORY_3GGT	GVEVEKGEEDGVFSRESVCKAVKAVVDEKSEIGREVRGNHDKLRGFLLNADLDSKYVDSFNQKIQDILG...	459
ARABIDOPSIS_3GGTF	AVEVER.EKKWFSRCLENAVKSVNEEGSEIGEKVRKNHDKWRCVLTDSGSDGIIDKEQNIEIVK...	467
ARABIDOPSIS_3GGT	SVEVKR.EETGVFSKESLSGAVRSVNRDSELGNWARRNHVWKESLLRHGLMSGVLNKVEALEKLVQNI NL	452

B

A

B

Reaction	Substrate	Enzyme	HPLC product
1	Cyanidin 3-glc + UDP-glc	Ib3GGT	Cyanidin 3-glc-2"-glc (100 ± 8.5)
2	Cyanidin 3-glc + UDP-xyl	Ib3GGT	Cyanidin 3-glc-2"-xyl (10.4 ± 1.2)
3	Cyanidin 3-glc + UDP-xyl	Ib3GGT ^{T138I}	Cyanidin 3-glc-2"-xyl (6.1 ± 0.79)
4	Cyanidin 3-glc + UDP-glc	Ib3GGT ^{T138I}	No product
5	Cyanidin 3-glc + UDP-xyl	At3GGT	Cyanidin 3-glc-2"-xyl (100 ± 15.0)
6	Cyanidin 3-glc + UDP-glc	At3GGT	No product
7	Cyanidin 3-glc + UDP-glc	At3GGT ^{T142T}	Cyanidin 3-glc-2"-glc (91.0 ± 12.5)
8	Cyanidin 3-glc + UDP-xyl	At3GGT ^{T142T}	Cyanidin 3-glc-2"-xyl (100 ± 18.3)

A **B** **C** **D** **E**

A

B

Reporter (1)

Effector (2)

Scale is 15 μm