Blocking the alternative sigma factor RpoN reduces virulence of *Pseudomonas aeruginosa* isolated from cystic fibrosis patients and increases antibiotic sensitivity in a laboratory strain

Running Title: Blocking RpoN makes *P. aeruginosa* CF strains avirulent

Lloyd MG¹, Vossler JL², Nomura CT^{3,4}, Moffat JF^{1,#}

¹Department of Microbiology and Immunology, SUNY Upstate Medical University, Syracuse, NY, USA

²Department of Clinical Laboratory Science, SUNY Upstate Medical University, Syracuse, NY, USA

³Department of Chemistry, SUNY College of Environmental Science and Forestry, Syracuse, NY, USA

⁴Center for Applied Microbiology, SUNY College of Environmental Science and Forestry, Syracuse, NY, USA

#Address Correspondence to Jennifer F. Moffat

Mailing Address: 750 East Adams Street, Syracuse, New York 13210

E-mail: moffatj@upstate.edu; Phone: (315) 464-5454

Abstract:

Multidrug-resistant organisms (MDROs) are increasing in the health care setting, and there are few antimicrobial agents available to treat infections caused by these bacteria. *Pseudomonas aeruginosa* is an opportunistic pathogen in burn patients and individuals with cystic fibrosis (CF), and a leading cause of nosocomial infections. *P. aeruginosa* is inherently resistant to many antibiotics and can develop or acquire resistance to others, limiting options for treatment. *P. aeruginosa* has virulence factors that are regulated by sigma factors in response to the tissue microenvironment. The alternative sigma factor, RpoN (σ^{54}), regulates many virulence genes and is linked to antibiotic resistance. Recently, we described a cis-acting peptide, RpoN*, which acts as a “molecular roadblock”, binding RpoN consensus promoters at the -24 site and blocking transcription. RpoN* reduces virulence of *P. aeruginosa* laboratory strains both *in vitro* and *in vivo*, but its effects in clinical isolates was not known. We investigated the effects of RpoN* on phenotypically varied *P. aeruginosa* strains isolated from cystic fibrosis patients. RpoN* expression reduced motility, biofilm formation, and pathogenesis in a *P. aeruginosa* – *C. elegans* infection model. RpoN* expression increased susceptibility to several beta-lactam based antibiotics in the lab strain *P. aeruginosa* PA19660 *Xen5*. Here, we show that using a cis-acting peptide to block RpoN consensus promoters has potential clinical implications in reducing virulence and enhancing the activity of antibiotics.

Introduction:

Multidrug-resistant organisms (MDROs) are an increasing problem in the healthcare setting. Both Gram-negative and Gram-positive MDROs are prevalent globally (1, 2). There are few or no antimicrobial agents available for treatment of infections caused by these bacteria (3). *Pseudomonas aeruginosa*, a Gram-negative, opportunistic pathogen is a leading cause of nosocomial infections and is associated with infections in burn patients (4, 5). *P. aeruginosa* is also responsible for colonizing the respiratory tract and causing chronic infections in individuals with cystic fibrosis (CF) (6). It is the most common pathogen isolated from individuals with CF, and is a major source of morbidity and mortality (7-10).

In CF patients, *P. aeruginosa* undergoes a transformation from a non-mucoid form upon initial colonization of the lungs to a mucoid form as the disease progresses. This results in a chronic debilitating pulmonary infection characterized by the overexpression of alginate. Mucoid strains synthesize large quantities of alginate exopolysaccharide, enhancing biofilm formation and protecting *P. aeruginosa* from antibiotics or the immune response (11), possibly through formation of microcolonies (12, 13). While aggressive prevention regimens have led to a decline in prevalence of *P. aeruginosa* in CF patients, multidrug resistant strains are still prevalent and occurred in 19.4% of CF infections in 2015 (14). *P. aeruginosa* is inherently resistant to a number of antibiotics (15, 16). It can also acquire resistance through exogenous resistance genes via horizontal gene transfer or mutations (17), limiting available treatment options. Antimicrobial development is directed toward alternative treatments and novel targets.

Promising strategies include enhancing the activity of currently available antibiotics and decreasing virulence of the bacteria once an infection occurs (18-25).

P. aeruginosa virulence is caused by many factors, including production of toxins, proteases, phospholipases, the presence of pili and flagella, and biofilm formation (26). This virulence is regulated by a network of transcription factors, such as sigma factors RpoS and RpoN, and quorum sensing regulators (27). The alternative sigma factor, σ^{54} or RpoN, regulates nitrogen assimilation, quorum sensing, motility, and biofilm formation (28-33). RpoN regulation was recently linked to *P. aeruginosa* tolerance to several antibiotics (34-36). RpoN binds to specific promoters with conserved -24, -12 sequences upstream of RpoN-regulated genes throughout the genome and is a key virulence regulator (37). The specific and conserved nature through which RpoN controls its regulon led us to develop the RpoN molecular roadblock, RpoN*. RpoN* is a cis-acting peptide that specifically binds the -24 site of RpoN consensus promoters, blocking transcription by RpoN and other factors (38). When RpoN* is expressed in *P. aeruginosa* laboratory strains, transcription is affected globally and virulence is attenuated (38). RpoN* also affects virulence in an RpoN deletion strain of *P. aeruginosa* PAO1, demonstrating its ability to attenuate gene expression by repressing expression of genes located downstream of RpoN promoters (38). This strategy of blocking multiple promoters throughout the *P. aeruginosa* genome may be an effective method to combat virulence and evade development of resistance.

P. aeruginosa isolated from CF patients are phenotypically and genetically varied (39, 40). Many *P. aeruginosa* clinical isolates have mutations, including deletion or loss of function, in the *rpoN* gene (41, 42). It was not known how the cis-acting RpoN* peptide would affect virulence phenotypes in *P. aeruginosa* clinical isolates, particularly in strains that do not express or have low levels of RpoN. In this study, we describe the effects of RpoN* on *in vitro* and *in vivo* virulence of *P. aeruginosa* isolated from CF patients and its effects on antibiotic resistance. Expression of RpoN* reduced virulence-associated phenotypes in clinical isolates and improved *P. aeruginosa* susceptibility to multiple antibiotics. This study demonstrates that RpoN* has potential clinical applications and potentially represents an effective strategy to combat both antibiotic resistance and infections with *P. aeruginosa* in CF patients.

Results:

Virulence phenotypes were variable in P. aeruginosa isolates from CF patients:

P. aeruginosa isolated from different CF patients or within the same CF patient have varied phenotypes and genotypes (39, 40). *P. aeruginosa* adapts over time, leading to mutations and changes in expression of genes related to motility, quorum sensing, and overall virulence (41, 43). To determine the virulence-related phenotypic profiles of the strains used in this study (Table 1), each *P. aeruginosa* patient isolate was evaluated for motility and biofilm formation, compared to the positive, virulent control strain *P. aeruginosa* PA19660 Xen5. Several patient isolates were highly motile in the swimming assay (flagella), including SCH0057-7, SCH0256-1, SCH0354-1 and UUH0201, while others were nonmotile (Fig 1A). Most strains were motile in the twitching assay (pili) and

produced moderate biofilms, with SCH0254-118 migrating the furthest (Fig 1B) and forming the most extensive biofilm (Fig 1C). SCH0254-116, SCH0397-3, and UUH0202 did not form biofilms.

The pathogenesis of patient isolates was evaluated in a *P. aeruginosa* – *C. elegans* infection model. All patient isolates were compared to *E. coli* OP50, an avirulent negative control. SCH0057-7 was the most pathogenic in the paralytic killing assay, which is mediated by hydrogen cyanide production (44, 45) (Fig 2A). Other strains were moderately pathogenic, including SCH0256-1, SCH0354-1, SCH0397-3, and UUH0202. SCH0057-7, SCH0338-58, and UUH0202 were highly pathogenic in the slow killing assay, which mimics establishment and proliferation of an infection and is mediated by the *lasR*, *gacA*, *lemA*, and *ptsP* genes (46), while UUH0201 was moderately pathogenic (Fig 2B). As expected, the virulence-associated phenotypes of patient isolates varied widely *in vitro* and *in vivo*.

RpoN protein levels varied among patient isolates:

Others reported that the *rpoN* gene was mutated or lost in approximately 20% of *P. aeruginosa* isolates from CF patients (41). Loss or mutation in the *rpoN* gene can result in phenotypes similar to those observed in the patient isolates evaluated here (29, 31, 32). Thus, we evaluated relative protein levels of RpoN in these patient isolates by western blot. RpoN levels were moderately high in the positive control *P. aeruginosa* PAO1-S, while low or minimal protein levels were detected in the isogenic $\Delta rpoN$ mutant negative control (Fig 3). The low level of background in the $\Delta rpoN$ mutant is likely due to

nonspecific antibody binding. RpoN levels varied in the CF patient isolates, with high levels in SCH0057-7, SCH0397-3, and UUH0201; intermediate levels in SCH0254-116, SCH0338-58, and UUH0202; and low levels in SCH0254-23, SCH0254-118, SCH0256-1, SCH0354-1, SCH03269, and UUH0101.

RpoN expressed in CF patient isolates reduced virulence-associated phenotypes in vitro:*

The effect of RpoN* expression on motility and biofilm formation in patient isolates was not known. Unfortunately, some patient isolates could not be transformed, and so only four isolates were evaluated for the effects of RpoN* expressed from a plasmid. SCH0057-7, SCH0256-1, SCH0338-58, and SCH0354-1 were transformed with a plasmid expressing RpoN* or the empty vector and selected with gentamicin. If RpoN* affected transcription of virulence-related genes in different genetic backgrounds as previously reported (38), we expected attenuation of virulence-related phenotypes in *P. aeruginosa* CF patient isolates. RpoN* significantly reduced colony diameter in all four patient isolates in the swimming motility assay (Student's t-test, ** $p \leq 0.01$, *** $p \leq 0.0001$) (Fig 4A). RpoN* significantly reduced colony diameter in SCH0057-7, SCH0256-1, and SCH0338-58 in the twitching motility assay (Student's t-test, ** $p \leq 0.01$, *** $p \leq 0.0001$) (Fig 4B). Colony diameter varied widely in SCH0354-1 when RpoN* was expressed and was always smaller than with empty vector, although the difference was not significant. In the biofilm formation assay, RpoN* significantly reduced biofilm formation by SCH0057-7 and SCH0256-1 (Student's t-test, $p \leq 0.0001$) (Fig 4C). Thus RpoN* reduced virulence-associated phenotypes of *P. aeruginosa* isolated from CF patients.

RpoN expression increased worm survival in P. aeruginosa – C. elegans infection*

model:

Initial evaluation of patient isolates revealed a single *P. aeruginosa* strain, SCH0057-7, that was both transformable and pathogenic in the *P. aeruginosa* – *C. elegans* infection assay. Therefore, effects of RpoN* on pathogenesis of SCH0057-7 were evaluated using the paralytic killing assay, which is based on *P. aeruginosa* hydrogen cyanide production and mimics conditions in the CF lung (44, 45). Wild-type *P. aeruginosa* SCH0057-7 was the positive, virulent control and *E. coli* was the negative, avirulent control. The test conditions were *P. aeruginosa* SCH0057-7 expressing RpoN* or carrying the empty vector plasmid. If RpoN* affected virulence-related phenotypes in *P. aeruginosa* SCH0057-7, then we expected increased survival of *C. elegans*. Wild type SCH0057-7 and with the empty vector killed approximately 80% of *C. elegans* (Fig 5). In contrast, RpoN* expression significantly increased *C. elegans* survival (Mantel-Cox Log-Rank Test, $p \leq 0.0001$). Thus, RpoN* expression reduced pathogenesis of a patient isolate in a *P. aeruginosa* – *C. elegans* infection model.

RpoN increased antibiotic susceptibility in vitro:*

Antibiotic resistance is a problem in CF patients with *P. aeruginosa* infections (47-49). We previously reported that RpoN* alters transcription of several genes involved in multidrug efflux pumps that confer natural resistance (38). Additionally, RpoN is implicated in tolerance to various classes of antibiotics (34-36). We evaluated the effects of RpoN* on antibiotic susceptibility using a MicroScan Neg MIC 43 panel. The

test conditions were *P. aeruginosa* PA19660 *Xen5* that was mock-transformed, or transformed with the empty vector or RpoN* plasmid. We expected that RpoN* would improve antibiotic susceptibility of *P. aeruginosa*. In PA19660 *Xen5* mock-transformed or with the empty vector, antibiotic susceptibility profiles were the same, except for gentamicin, which increased in the empty vector strain due to the GM^R selection marker (data not shown). In PA19660 *Xen5* expressing RpoN*, susceptibility to five beta-lactam antibiotics was improved 2- to 4-fold (Fig 6). These were cefotaxime, cefepime, and ceftazidime (three cephalosporins), piperacillin (a ureidopenicillin), and imipenem (a carbapenem). Susceptibility to some antibiotics was unchanged (data not shown). The results demonstrate that RpoN* expression increased *P. aeruginosa* susceptibility to several antibiotics.

Discussion:

Here, we confirm and expand results of previous studies by showing the ability of RpoN* to abrogate virulence phenotypes in *P. aeruginosa* isolates from CF patients and to improve susceptibility to several antibiotics. Our working model of the mechanism of action of RpoN* is that it binds the -24 promoter consensus sites, blocking transactivation by RpoN and other sigma factors. By altering the transcriptome, RpoN* reduced virulence in well-characterized laboratory strains (38). Thus, the motivation for this study was to understand the clinical relevance of RpoN*. We demonstrated that RpoN* expressed in CF patient isolates reduced motility and biofilm formation *in vitro*, independently of RpoN protein levels. The RpoN* molecular roadblock protected C.

C. elegans from a highly virulent *P. aeruginosa* patient isolate in an *in vivo* infection model.

RpoN* also improved *P. aeruginosa* susceptibility to antibiotics.

P. aeruginosa isolated from CF patients are highly variable (39, 40), with the *rpoN* gene often mutated or lost (41). The patient isolates evaluated in this study had a broad range of motility, biofilm formation, RpoN protein levels, and virulence in *C. elegans*.

There was no correspondence between most *in vitro* phenotypes, *in vivo* pathogenesis, and RpoN levels (Fig. S1). The only correlation observed was between twitching or pili-associated motility and biofilm formation (Supplemental Fig 1F, $p=0.0357$, $R^2=0.37050$).

Other studies suggested that *in vitro* phenotypes of *P. aeruginosa* isolates can be related to disease status in CF patients (50). Patient information and status of *P.*

aeruginosa infections is limited for the isolates described here, so a comparison between phenotypes and patient status is not feasible. Interestingly, two isolates,

UUH0201 and UUH0202, were obtained five months apart from the same patient, with UUH0201 collected first. The UUH0202 strain was less motile and RpoN protein levels dropped compared to UUH0201, but virulence increased. This supports the concept that

in vitro phenotypes reflect *P. aeruginosa* infection status in CF patients (50). Further

work would be needed to fully elucidate such correlations.

The RpoN* molecular roadblock reduced virulence phenotypes in patient isolates with high or low levels of RpoN. For instance, RpoN protein levels were higher in SCH0057-7 than PAO1-S, and RpoN* reduced flagellar and pili motility, biofilm formation and pathogenesis. In contrast, relative RpoN protein levels were low in SCH0256-1 and

SCH0354-1, and yet RpoN* reduced motility. Thus, the roadblock was effective in the presence or absence of the native sigma factor. This confirms our previous findings, which show that RpoN* reduced virulence in a laboratory strain that was deleted for *rpoN* (38). Unfortunately, barriers to transformation precluded evaluating RpoN* in some of the other clinical isolates. However, the strains that were successfully transformed represented much of the diversity across the patient isolates.

The CF patient isolates demonstrated variable pathogenesis in the *C. elegans* paralytic killing model that spans 6 hours. Only one pathogenic isolate, SCH0057-7, was transformable and thus possible to evaluate the effects of RpoN* *in vivo*. This strain and several others were also pathogenic in the 4-day slow killing assay, but this assay was not used to evaluate RpoN* because of difficulty maintaining the plasmid and RpoN* expression. Gentamicin selection and IPTG induction are not durable, we found (38), because the *C. elegans* cuticle is impermeable and the compounds are poorly absorbed in the intestine (51). While it is expected that expressing RpoN* in CF isolates would improve *C. elegans* survival in the slow killing assay, it is not feasible with the current vector. If issues with maintaining the plasmid and expression of the roadblock were resolved, it would be interesting to evaluate RpoN* in this assay using patient isolates.

The molecular roadblock, RpoN*, binds numerous promoters in bacterial genomes, altering the transcriptome. RpoN* expression in *P. aeruginosa* greatly reduced transcription of the *mex* family genes (38), which are involved in multidrug efflux pumps

(20). Increased expression of *mex* genes is linked to increased resistance to antibiotics (17). Therefore, we investigated whether RpoN* alters *P. aeruginosa* susceptibility to antibiotics. We employed a clinical laboratory assay for testing bacterial susceptibility or resistance to antibiotics, and found that RpoN* improved antibiotic susceptibility at least two-fold for five different antibiotics, including imipenem. This agrees with work by others that showed RpoN is involved in *P. aeruginosa* tolerance of carbapenems, quinolones, and tobramycin (34-36). Unfortunately, the commercial assay uses pre-determined antibiotic concentrations in a 96-well plate, limiting the scope of the molecular roadblock's effects. Additionally, the *P. aeruginosa* strain used here is sensitive to quinolones and tobramycin, so the effects of RpoN* expression on resistance to these antibiotics was not evaluated. It will be important to test clinical strains that are resistant to quinolones, carbapenems, and tobramycin to determine the effects of RpoN*. Further studies are needed to uncover the full spectrum of RpoN* effects on antibiotic susceptibility.

Multi-drug resistant organisms (MRDOs) are increasing worldwide, even those with resistance to entire classes of antibiotics. Alarming, nearly all antibiotics brought to market in the past 30 years are variations on existing drugs (52). Research into alternative strategies to treat bacterial infections is a priority, including compounds to enhance the activity of existing antibiotics or neutralize virulence factors. The molecular roadblock falls into the latter type. RpoN* binds consensus promoters throughout the *P. aeruginosa* genome, affecting the transcription of numerous virulence factors. Due to the many binding sites for RpoN*, it is unlikely antibiotic resistance will develop during

treatment. The binding sequence of the RpoN consensus promoter is conserved across gram-negative and gram-positive bacteria (36, 37). We explored the effects of RpoN* on virulence phenotypes of *Pseudomonas putida*, *Burkholderia cepacia*, and *Escherichia coli* (unpublished data), suggesting that RpoN* may reduce virulence in multiple organisms. More studies are needed to identify the spectrum of RpoN* activity and its resistance frequency. Currently, the molecular roadblock is a tool for antimicrobial development and is not a usable drug. However, a small molecule that works in the same cis-acting manner as RpoN* would be an effective, clinically relevant strategy to combat *P. aeruginosa* virulence and antibiotic resistance.

Materials and Methods:

Bacteria and Nematodes:

P. aeruginosa clinical isolates were provided by the Seattle Children's Hospital (SCH strains) and Upstate University Hospital (UUH strains). *P. aeruginosa* PAO1-M was provided by C. Manoil (44), and *P. aeruginosa* PAO1-S and $\Delta rpoN$ were provided by D. Haas (31). *P. aeruginosa* PA19660 *Xen5* was purchased from PerkinElmer. *E. coli* OP50 was provided by D. Pruyne (SUNY Upstate Medical University). All strains are listed in Table 1. For long-term storage, bacteria were grown overnight in LB broth at 37°C with shaking, and frozen in 10% glycerol at -80°C. *Caenorhabditis elegans* N2 was purchased from the *Caenorhabditis* Genetics Center (University of Minnesota, Minneapolis, MN), and maintained on nematode growth media (NGM) seeded with *E. coli* OP50 at 20°C (53). Populations were synchronized via egg lay and grown to the young adult stage at 20°C (54).

297

298 *Plasmids:*

299 RpoN* and empty vector plasmids were previously described (38). Plasmids were
300 maintained in *E. coli* INV110 (Invitrogen) with gentamicin selection (30 mg/L). RpoN*
301 expression was induced with isopropyl β -D-1-thiogalactopyranoside (IPTG, 1 mM).

302

303 *Transformation:*

304 Permissive *P. aeruginosa* patient isolates were transformed by electroporation prior to
305 all experiments, per standard protocol (55). Transformed bacteria were selected on LB
306 agar or BHI agar supplemented with gentamicin (30 mg/L). Individual colonies were
307 picked for each assay.

308

309 *Western Blot Analysis:*

310 Overnight bacteria cultures were treated with Cell Lytic B Lysis Reagent (Sigma) to
311 generate crude cell lysates. The soluble protein fraction was separated on 10% Mini-
312 PROTEAN® TGX Stain-Free™ protein gels (BioRad), activated for 5 minutes with UV
313 light, imaged and transferred via semi-dry apparatus to a PVDF membrane. Membranes
314 were incubated with primary antibody specific for *E. coli* RNA σ^{54} (1:500, BioLegend)
315 overnight, then with secondary antibody HRP goat anti-mouse (1:10,000, Jackson
316 ImmunoResearch). The chemiluminescent signal was generated with the Pierce
317 SuperSignal West Femto substrate kit (Thermo Scientific), and detected with
318 ChemiDoc™ MP Imaging System (Bio-Rad Laboratories). Protein bands and total

protein per lane were measured with Image Lab (Version 5.2.1; Bio-Rad Laboratories).
RpoN bands were then compared to corresponding total detected protein in each lane.

Phenotyping Assays:

Assays to measure swimming, twitching (56), and biofilm formation (57), were conducted according to standard protocols. Transformed *P. aeruginosa* clinical isolates were grown in appropriate media supplemented with gentamicin (30 mg/L), and with or without IPTG (1 mM). Motility assay and microtiter plate biofilm assay were conducted at 37°C for 24h. Images of motility assays were obtained with IVIS-50™ (Perkin Elmer) and colony diameter was measured with Living Image software (Perkin Elmer). Biofilms were stained with 0.1% crystal violet, extracted in 95% ethanol, and absorbance was measured at 550 nm with a μ Quant microplate spectrophotometer (BioTek).

P. aeruginosa – C. elegans infection assays:

For the paralytic killing assay, laboratory strains, clinical isolates, or transformed *P. aeruginosa* were spread on Brain Heart Infusion (BHI) agar (Difco) with, when applicable, gentamicin (30 mg/L) and with or without IPTG (1 mM). *E. coli* was spread on BHI agar. All plates were grown overnight at 37°C. Bacteria colonies were swabbed onto BHI agar, supplemented with gentamicin and/or IPTG (1 mM) when applicable, and grown at 37°C for 24 h (44). Adult *C. elegans* were added to plates and the assay was conducted at room temperature, per standard protocol (44). For the slow killing assay, laboratory strains or clinical isolates of *P. aeruginosa* were grown overnight in LB broth at 37°C with shaking, and cultures were spread on a modified NGM agar (0.35%

bactopeptone, 2% bactoagar) (58). Plates were incubated at 37°C for 24 h, then at room temperature for an additional 24 h. The assay was conducted at 20°C, and worms were scored every 24 h per standard protocol (58).

Antibiotic Sensitivity Testing:

Transformed or mock transformed bacteria were grown overnight in LB broth with gentamicin (30 mg/L) and IPTG (1 mM) or only IPTG (1 mM), respectively. MicroScan Neg MIC 43 panels (Beckman Coulter Inc., Brea, CA) were used. Panels were set up per manufacturer's protocol (MicroScan Gram Negative Procedure Manual, version 09/2016) using the RENOX system (Beckman Coulter Inc., Brea, CA) with a final well concentration of $3-7 \times 10^5$ CFU/mL. The following modifications were made to the manufacturer's protocol: LB broth supplemented with IPTG (1 mM) and with or without gentamicin (30 mg/L) was used in place of saline for whole panel. Plates were incubated at 35°C for 16-20 h, and read using a MicroScan autoSCAN-4 (Beckman Coulter Inc, Brea, CA). Quality control was performed on the panels per manufacturer's protocol.

Statistics:

Data were analyzed using Excel and GraphPad Prism with a significance of $p \leq 0.05$ (Microsoft, Washington; GraphPad Software Inc., California).

Data Availability:

The datasets produced for the current study are included in this manuscript or are available from the corresponding author upon reasonable request.

Acknowledgements

This work was supported by the NIH (CTN: 2R15GM104880) and by the Hill Collaboration [58482 to JFM and CTN]. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. *C. elegans* strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We would like to thank the Clinical Microbiology Lab at the Syracuse VA Medical Center in Syracuse, NY for supplying the antibiotic susceptibility testing panels and for use of their facility for reading the results. We would also like to thank Dr. Ran Anbar and Donna Linder at the Golisano Center at Upstate University Hospital and Marcella Blackledge and Dr. Rafael Hernandez at the Seattle Children's Hospital for supplying the *P. aeruginosa* CF patient isolates (Seattle Children's Hospital CF patient isolates obtained under NIH P30 DK089507).

Competing Financial Interests:

The authors declare no competing financial interests.

Author Contributions:

385 M.G.L. wrote the manuscript. C.T.N. and J.F.M. conceived the study. M.G.L. and J.L.V.
386 conducted the experiments. M.G.L. generated the figures. All authors reviewed the
387 manuscript.
388

Figure 1. Characterization of virulence phenotypes of *P. aeruginosa* strains isolated from cystic fibrosis patients. *P. aeruginosa* CF patient isolates were compared to laboratory strain PA19660 *Xen5*. All assays were conducted at 37°C for 24 h. (A) Colony diameter of swimming, or flagellar, motility assay conducted on soft (0.3%) agar. (B) Twitching, or pili, motility assay conducted on semi-hard (1.3%) agar. Colony diameter was measured across point of inoculation to the edges of bacteria colony. (C) Biofilm formation assay was conducted in 96-well microtiter plates. Biofilms were stained with crystal violet (0.1%), solubilized in ethanol (95%), and absorbance

measured at OD₅₅₀. Bars are the mean $\pm$ SD; n = 5 to 6 replicates in motility assays and n = 10 in biofilm assay. Each assay was performed at least three separate times and representative results are shown.

Figure 2. Pathogenesis of *P. aeruginosa* isolated from cystic fibrosis patients in *P. aeruginosa* – *C. elegans* infection model. Kaplan-Meier survival curves for *P. aeruginosa* – *C. elegans* infection assays. (A) Paralytic killing assay on BHI agar. Assay was conducted at room temperature and worm status scored every 2 h. (B) Slow killing assay on modified NGM agar (0.35% bactopectone, 2% bactoagar). Assay was conducted at 20°C and worm status scored every 24 h. Strains used included CF patient isolates, and *E. coli* for reference. n = 48 to 90 worms per strain.

Figure 3. RpoN protein in *P. aeruginosa* isolates is highly varied. Immunoblot analysis of RpoN expression in *P. aeruginosa* CF patient isolates and laboratory strains PAO1-S and $\Delta rpoN$. Immunoblots performed on 10% Mini-PROTEAN® TGX Stain-Free™ gels (BioRad). RpoN protein levels were calculated by comparing measured total protein in each lane to the measured RpoN band (presented as arbitrary units (AU)). Values were normalized against PAO1-S. This graph is representative of immunoblots from multiple bacterial cultures and western blot analyses.

Figure 4. RpoN* expression decreases motility and biofilm formation. *P.*

aeruginosa CF patient isolates with empty vector (E.V., black bars), or with RpoN*-expression vector (gray bars). Media was supplemented with gentamicin (30 mg/L), and IPTG (1 mM) when applicable, and all assays were conducted at 37°C for 24 h. (A) Colony diameter of swimming, or flagellar, motility assay conducted on soft (0.3%) agar. (B) Colony diameter of twitching, or pili, motility assay conducted on semi-hard (1.3%) agar. (C) Biofilm formation assay conducted in 96-well microtiter plates. Bars are the

mean $\pm$ SD (Student's t-test, *** $p \leq 0.0001$; ** $p \leq 0.01$). $n = 4$ to 5 replicates in motility assays; $n = 12$ in biofilm assays.

Figure 5. RpoN^{*} promotes *C. elegans* survival in paralytic killing assay.

P. aeruginosa CF isolate SCH0057-7 wild type, with empty vector (E.V.), or with RpoN^{*}-expression vector, were compared to avirulent *E. coli* OP50. Paralytic killing assay was conducted on BHI agar supplemented with gentamicin (30 mg/L) and IPTG (1 mM), when applicable. Assay was conducted at room temperature, and worm status scored every 2 h. Kaplan-Meier survival curves represent combined survival of three separate assays (exception: *E. coli*). Mantel-Cox log-rank test was used to analyze E.V. and RpoN^{*} curves (*** $p \leq 0.0001$). $n = 180$ worms per SCH0057-7 condition, $n = 60$ worms for *E. coli*.

Figure 6. RpoN* expression increases susceptibility to antibiotics. Overnight cultures of *P. aeruginosa* PA19660 Xen5 with empty vector or expressing RpoN*, were diluted in fresh LB broth, added to wells of MicroScan Neg MIC 43 panel, and incubated at 35°C for 16-20 h. Media was supplemented with gentamicin (30 mg/L) and IPTG (1 mM). Each point is the average of two biological replicates per experiment and the data shown are representative of three separate experiments.

Table 1. *P. aeruginosa* strains used in this study.

Location	Strain	Source	CF mutations	Reference
<u>Laboratory</u>				
	PAO1-M	---	n/a	C. Manoil (27)
	PAO1-S	---	n/a	D. Haas (15)
	$\Delta rpoN$	---	n/a	D. Haas (15)
	PA19660 Xen5	Septicemia	n/a	Perkin Elmer
<u>Clinical Isolate</u>				
<i>Seattle Children's Hospital (SCH)</i>				
	SCH0057-7	unknown	$\Delta F508$ / $\Delta F508$	This study
	SCH0254-23	unknown	$\Delta F508$ / unknown	This study
	SCH0254-116	unknown	$\Delta F508$ / unknown	This study
	SCH0254-118	unknown	$\Delta F508$ / unknown	This study
	SCH0256-1	sputum	$\Delta F508$ / $\Delta F508$	This study
	SCH0338-38	sputum	unknown / unknown	This study
	SCH0354-1	sputum	$\Delta F508$ / G551D	This study
	SCH0397-3	unknown	$\Delta F508$ / unknown	This study
	SCH03269	sputum	$\Delta F508$ / $\Delta F508$	This study
<i>Upstate University Hospital (UUH)</i>				
	UUH0101	sputum	$\Delta F508$ / unknown	This study
	UUH0201	sputum	$\Delta F508$ / $\Delta F508$	This study
	UUH0202	sputum	$\Delta F508$ / $\Delta F508$	This study

References:

1. Centers for Disease C, Prevention. 2013. Vital signs: carbapenem-resistant Enterobacteriaceae. MMWR Morb Mortal Wkly Rep 62:165-70.
2. Cohen ML. 2000. Changing patterns of infectious disease. Nature 406:762-7.
3. Freire-Moran L, Aronsson B, Manz C, Gyssens IC, So AD, Monnet DL, Cars O, Group E-EW. 2011. Critical shortage of new antibiotics in development against multidrug-resistant bacteria-Time to react is now. Drug Resist Updat 14:118-24.
4. Lyczak JB, Cannon CL, Pier GB. 2000. Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. Microbes Infect 2:1051-60.
5. Bodey GP, Bolivar R, Fainstein V, Jadeja L. 1983. Infections caused by *Pseudomonas aeruginosa*. Rev Infect Dis 5:279-313.
6. Davis PB, Drumm M, Konstan MW. 1996. Cystic fibrosis. Am J Respir Crit Care Med 154:1229-56.
7. Emerson J, Rosenfeld M, McNamara S, Ramsey B, Gibson RL. 2002. *Pseudomonas aeruginosa* and other predictors of mortality and morbidity in young children with cystic fibrosis. Pediatr Pulmonol 34:91-100.
8. Ahlgren HG, Benedetti A, Landry JS, Bernier J, Matouk E, Radzioch D, Lands LC, Rousseau S, Nguyen D. 2015. Clinical outcomes associated with *Staphylococcus aureus* and *Pseudomonas aeruginosa* airway infections in adult cystic fibrosis patients. BMC Pulm Med 15:67.
9. Govan JR, Deretic V. 1996. Microbial pathogenesis in cystic fibrosis: mucoid *Pseudomonas aeruginosa* and *Burkholderia cepacia*. Microbiol Rev 60:539-74.

10. Sanders DB, Fink AK. 2016. Background and Epidemiology. *Pediatr Clin North Am* 63:567-84.
11. Hoiby N. 2002. Understanding bacterial biofilms in patients with cystic fibrosis: current and innovative approaches to potential therapies. *J Cyst Fibros* 1:249-54.
12. Fegan M, Francis P, Hayward AC, Davis GH, Fuerst JA. 1990. Phenotypic conversion of *Pseudomonas aeruginosa* in cystic fibrosis. *J Clin Microbiol* 28:1143-6.
13. Lam J, Chan R, Lam K, Costerton J. 1980. Production of mucoid microcolonies by *Pseudomonas aeruginosa* within infected lungs in cystic fibrosis. *Infection and immunity* 28:546-556.
14. Registry CFFP. 2016. 2015 Annual Data Report. Cystic Fibrosis Foundation, Bethesda, Maryland.
15. Hancock RE, Speert DP. 2000. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment. *Drug Resist Updat* 3:247-255.
16. Porras-Gómez M, Vega-Baudrit J. 2012. Overview of multidrug-resistant *Pseudomonas aeruginosa* and novel therapeutic approaches. *Journal of Biomaterials and Nanobiotechnology* 3:519.
17. Riou M, Avrain L, Carbonnelle S, El Garch F, Pirnay JP, De Vos D, Plesiat P, Tulkens PM, Van Bambeke F. 2016. Increase of efflux-mediated resistance in *Pseudomonas aeruginosa* during antibiotic treatment in patients suffering from nosocomial pneumonia. *Int J Antimicrob Agents* 47:77-83.

18. Dickey SW, Cheung GYC, Otto M. 2017. Different drugs for bad bugs: antivirulence strategies in the age of antibiotic resistance. *Nat Rev Drug Discov* doi:10.1038/nrd.2017.23.
19. Wagner S, Sommer R, Hinsberger S, Lu C, Hartmann RW, Empting M, Titz A. 2016. Novel Strategies for the Treatment of *Pseudomonas aeruginosa* Infections. *J Med Chem* 59:5929-69.
20. Poole K, Srikumar R. 2001. Multidrug efflux in *Pseudomonas aeruginosa*: components, mechanisms and clinical significance. *Curr Top Med Chem* 1:59-71.
21. Stempel N, Strehmel J, Overhage J. 2015. Potential application of antimicrobial peptides in the treatment of bacterial biofilm infections. *Current pharmaceutical design* 21:67-84.
22. Park S-C, Park Y, Hahm K-S. 2011. The role of antimicrobial peptides in preventing multidrug-resistant bacterial infections and biofilm formation. *International journal of molecular sciences* 12:5971-5992.
23. Jakobsen TH, Bjarnsholt T, Jensen PO, Givskov M, Hoiby N. 2013. Targeting quorum sensing in *Pseudomonas aeruginosa* biofilms: current and emerging inhibitors. *Future Microbiol* 8:901-21.
24. Smith DJ, Lamont IL, Anderson GJ, Reid DW. 2013. Targeting iron uptake to control *Pseudomonas aeruginosa* infections in cystic fibrosis. *Eur Respir J* 42:1723-36.
25. Rogers GB, Carroll MP, Bruce KD. 2012. Enhancing the utility of existing antibiotics by targeting bacterial behaviour? *Br J Pharmacol* 165:845-57.

26. Gellatly SL, Hancock RE. 2013. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathog Dis* 67:159-73.
27. Balasubramanian D, Schneper L, Kumari H, Mathee K. 2013. A dynamic and intricate regulatory network determines *Pseudomonas aeruginosa* virulence. *Nucleic Acids Res* 41:1-20.
28. Potvin E, Sanschagrin F, Levesque RC. 2008. Sigma factors in *Pseudomonas aeruginosa*. *FEMS Microbiol Rev* 32:38-55.
29. Totten PA, Lara JC, Lory S. 1990. The *rpoN* gene product of *Pseudomonas aeruginosa* is required for expression of diverse genes, including the flagellin gene. *J Bacteriol* 172:389-96.
30. Damron FH, Owings JP, Okkotsu Y, Varga JJ, Schurr JR, Goldberg JB, Schurr MJ, Yu HD. 2012. Analysis of the *Pseudomonas aeruginosa* regulon controlled by the sensor kinase KinB and sigma factor RpoN. *J Bacteriol* 194:1317-30.
31. Heurlier K, Denervaud V, Pessi G, Reimann C, Haas D. 2003. Negative control of quorum sensing by RpoN (sigma54) in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 185:2227-35.
32. Thompson LS, Webb JS, Rice SA, Kjelleberg S. 2003. The alternative sigma factor RpoN regulates the quorum sensing gene *rhII* in *Pseudomonas aeruginosa*. *FEMS Microbiol Lett* 220:187-95.
33. Caiazza NC, O'Toole GA. 2004. SadB is required for the transition from reversible to irreversible attachment during biofilm formation by *Pseudomonas aeruginosa* PA14. *J Bacteriol* 186:4476-85.

34. Viducic D, Murakami K, Amoh T, Ono T, Miyake Y. 2017. RpoN Promotes *Pseudomonas aeruginosa* Survival in the Presence of Tobramycin. *Front Microbiol* 8:839.
35. Viducic D, Murakami K, Amoh T, Ono T, Miyake Y. 2016. RpoN Modulates Carbapenem Tolerance in *Pseudomonas aeruginosa* through Pseudomonas Quinolone Signal and PqsE. *Antimicrob Agents Chemother* 60:5752-64.
36. Viducic D, Ono T, Murakami K, Katakami M, Susilowati H, Miyake Y. 2007. *rpoN* gene of *Pseudomonas aeruginosa* alters its susceptibility to quinolones and carbapenems. *Antimicrob Agents Chemother* 51:1455-62.
37. Schulz S, Eckweiler D, Bielecka A, Nicolai T, Franke R, Dotsch A, Hornischer K, Bruchmann S, Duvel J, Haussler S. 2015. Elucidation of sigma factor-associated networks in *Pseudomonas aeruginosa* reveals a modular architecture with limited and function-specific crosstalk. *PLoS Pathog* 11:e1004744.
38. Lloyd MG, Lundgren BR, Hall CW, Gagnon LB-P, Mah T-F, Moffat JF, Nomura CT. 2017. Targeting the alternative sigma factor RpoN to combat virulence in *Pseudomonas aeruginosa*. *Scientific Reports* 7.
39. Clark ST, Diaz Caballero J, Cheang M, Coburn B, Wang PW, Donaldson SL, Zhang Y, Liu M, Keshavjee S, Yau YC, Waters VJ, Elizabeth Tullis D, Guttman DS, Hwang DM. 2015. Phenotypic diversity within a *Pseudomonas aeruginosa* population infecting an adult with cystic fibrosis. *Sci Rep* 5:10932.
40. Duong J, Booth SC, McCartney NK, Rabin HR, Parkins MD, Storey DG. 2015. Phenotypic and Genotypic Comparison of Epidemic and Non-Epidemic Strains of

Pseudomonas aeruginosa from Individuals with Cystic Fibrosis. PLoS One
10:e0143466.

41. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. Proc Natl Acad Sci U S A 103:8487-92.
42. Mahenthiralingam E, Campbell ME, Speert DP. 1994. Nonmotility and phagocytic resistance of *Pseudomonas aeruginosa* isolates from chronically colonized patients with cystic fibrosis. Infect Immun 62:596-605.
43. Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* Evolutionary Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol 24:327-37.
44. Darby C, Cosma CL, Thomas JH, Manoil C. 1999. Lethal paralysis of *Caenorhabditis elegans* by *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A 96:15202-7.
45. Gallagher LA, Manoil C. 2001. *Pseudomonas aeruginosa* PAO1 kills *Caenorhabditis elegans* by cyanide poisoning. J Bacteriol 183:6207-14.
46. Tan MW, Rahme LG, Sternberg JA, Tompkins RG, Ausubel FM. 1999. *Pseudomonas aeruginosa* killing of *Caenorhabditis elegans* used to identify *P. aeruginosa* virulence factors. Proc Natl Acad Sci U S A 96:2408-13.
47. Mustafa MH, Chalhoub H, Denis O, Deplano A, Vergison A, Rodriguez-Villalobos H, Tunney MM, Elborn JS, Kahl BC, Traore H, Vanderbist F, Tulkens PM, Van Bambeke F. 2016. Antimicrobial Susceptibility of *Pseudomonas aeruginosa*

- Isolated from Cystic Fibrosis Patients in Northern Europe. Antimicrob Agents Chemother 60:6735-6741.
48. Hill D, Rose B, Pajkos A, Robinson M, Bye P, Bell S, Elkins M, Thompson B, Macleod C, Aaron SD, Harbour C. 2005. Antibiotic susceptibilities of *Pseudomonas aeruginosa* isolates derived from patients with cystic fibrosis under aerobic, anaerobic, and biofilm conditions. J Clin Microbiol 43:5085-90.
49. Tomas M, Doumith M, Warner M, Turton JF, Beceiro A, Bou G, Livermore DM, Woodford N. 2010. Efflux pumps, OprD porin, AmpC beta-lactamase, and multiresistance in *Pseudomonas aeruginosa* isolates from cystic fibrosis patients. Antimicrob Agents Chemother 54:2219-24.
50. Mayer-Hamblett N, Rosenfeld M, Gibson RL, Ramsey BW, Kulasekara HD, Retsch-Bogart GZ, Morgan W, Wolter DJ, Pope CE, Houston LS, Kulasekara BR, Khan U, Burns JL, Miller SI, Hoffman LR. 2014. *Pseudomonas aeruginosa* in vitro phenotypes distinguish cystic fibrosis infection stages and outcomes. Am J Respir Crit Care Med 190:289-97.
51. O'Reilly LP, Luke CJ, Perlmutter DH, Silverman GA, Pak SC. 2014. *C. elegans* in high-throughput drug discovery. Adv Drug Deliv Rev 69-70:247-53.
52. Silver LL. 2011. Challenges of antibacterial discovery. Clin Microbiol Rev 24:71-109.
53. Wood WB. 1988. The Nematode *Caenorhabditis elegans*: Introduction to *C. elegans* biology. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

54. Lionaki E, Tavernarakis N. 2013. Assessing aging and senescent decline in *Caenorhabditis elegans*: cohort survival analysis. *Methods Mol Biol* 965:473-84.
55. Choi KH, Kumar A, Schweizer HP. 2006. A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Methods* 64:391-7.
56. O'Toole GA, Pratt LA, Watnick PI, Newman DK, Weaver VB, Kolter R. 1999. Genetic approaches to study of biofilms. *Methods Enzymol* 310:91-109.
57. O'Toole GA. 2011. Microtiter dish biofilm formation assay. *J Vis Exp* doi:10.3791/2437.
58. Tan MW, Mahajan-Miklos S, Ausubel FM. 1999. Killing of *Caenorhabditis elegans* by *Pseudomonas aeruginosa* used to model mammalian bacterial pathogenesis. *Proc Natl Acad Sci U S A* 96:715-20.