1 Identification of a *Nocardia seriolae* secreted protein targeting 2 host cell mitochondria and inducing apoptosis in 3 fathead minnow (FHM) cells

4
5 Jianlin Chen^{1,2,3¶}, Wenji Wang^{1,2,3¶}, Liqun Xia^{1, 2*}, Zhiwen Wang^{1,2,3}, Yishan Lu^{1, 2, 3,4*},
6 Jiahui Huang^{1,2,3}, Suying Hou^{1,2,3}

7
8 ¹ Shenzhen Research Institute of Guangdong Ocean University, Shenzhen

9 ² Fisheries College of Guangdong Ocean University, Zhanjiang

10 ³ Guangdong provincial engineering research center for aquatic animal health assessment, Shenzhen

11 ⁴ Guangdong Provincial Key Laboratory of Pathogenic Biology and Epidemiology for Aquatic Economic Animals, Zhanjiang

12 ^{#a} Current Address: Shenzhen Research Institute of Guangdong Ocean University, No. 3 of Binhai 2nd Road, Dapeng New District,
13 Shenzhen, Guangdong Province 518108, China. Tel: +86 755 23250356. Fax: +86 755 28380068.

14 ^{*}Corresponding author

15 11465668@qq.com (LX); fishdis@gmail.com (YL)

16 [¶]These authors contributed equally to this work and should be regarded as co-first authors.

17 Abstract

18 *Nocardia seriolae*, is a Gram-positive, partially acid-fast, aerobic, and filamentous bacterium. This
19 bacterium is the main pathogen of fish nocardiosis. A bioinformatic analysis based on the genomic
20 sequence of the *N. seriolae* strain ZJ0503 showed that *ORF3141* encoded a secreted protein with a
21 signal peptide at the N-terminate which may target the mitochondria in the host cell. However, the
22 functions of this protein and its homologs remain unknown. In this study, we experimentally tested
23 the bioinformatic prediction on this protein. Mass spectrometry analysis of the extracellular products
24 from *N. seriolae* showed that ORF3141 was a secreted protein. Subcellular localization of the
25 ORF3141-GFP fusion protein revealed that the green fluorescence protein co-localized with the
26 mitochondria, while ORF3141Δsig-GFP (with the signal peptide deleted) fusion protein was evenly

distributed in the whole cell of fathead minnow (FHM) cells. Thus, the N-terminate signal peptide had a significant impact on mitochondrial targeting. Notably, the expression of ORF3141 protein changed the distribution of mitochondria from perinuclear halo into lumps in the transfected FHM cells. In addition, apoptotic features were found in the transfected FHM cells by overexpression of ORF3141 and ORF3141 Δ sig proteins, respectively. Quantitative assays of mitochondrial membrane potential value, caspase-3 activity and apoptosis-related gene mRNA expression suggested that cell apoptosis was induced in the transfected FHM cells. In conclusion, the ORF3141 was a secreted protein of *N. seriolae* that targeted host cell mitochondria and induced apoptosis in FHM cells. This protein may participate in the cell apoptosis regulation and plays an important role in the pathogenesis of *N. seriolae*.

Keywords: *Nocardia seriolae*; secreted protein; subcellular localization; mitochondrial targeting; cell apoptosis

Author summary

Nocardia seriolae is the causative pathogen responsible for fish nocardiosis. This facultative intercellular bacterium, adapts to survive and colonize by evading intracellular killing after being engulfed with macrophages in the host. Despite considerable economic losses caused by *N. seriolae* in fish infection, the pathogenic mechanism and specific virulence factor of this bacterium remain ambiguous. In this study, the characteristic of ORF3141 protein function was investigated by subcellular localization and its possible contributions on the ability of *N. seriolae* to induce apoptosis in transfected fathead minnow (FHM) cells was investigated. Here, we confirmed that ORF3141 was a secreted protein that targeted host cell mitochondria and induced cell apoptosis in FHM cells. Interestingly, after deleting the signal peptide, ORF3141 Δ sig protein was evenly distributed in the whole host cell and did not co-localize with the mitochondria which could also induce cell apoptosis. Thus, the N-terminate signal peptide played an important role in mitochondrial targeting, and the domain part without the signal peptide had a critical relationship with cell apoptosis. These results demonstrated that ORF3141 may act as a potential virulence factor that induces apoptosis in fish cells. This protein is significant to elucidate the pathogenic

mechanism of *N. seriolae* and this study may provide beneficial insight to prevent and treat fish nocardiosis.

Introduction

Fish nocardiosis is a systemic bacterial disease with three kinds of pathogenic bacteria, namely, *Nocardia seriolae*, *N. samonivida* and *N. asteroides*, which have been isolated from diseased fish [1–3]. The outbreak of fish nocardiosis is frequently reported in global aquaculture industries and has caused substantial commercial losses in Southeast Asia, especially China [4–7]. However, the pathogen of fish nocardiosis that has been reported with considerable frequency is *N. seriolae*. Notably, *N. seriolae* is the most frequently isolated *Nocardia* sp. from diseased fish [7]. This bacterium is an opportunistic pathogen, by infecting immunocompromised fishes through feeds, gills, and wounds and causing chronic systemic granulomatous disease [8]. The clinical signs of infected fish include skin ulcers and numerous white nodular structures on the gills and in the head kidney, trunk kidney, spleen, and liver [9]. *N. seriolae* infections have been documented in more than 39 kinds of fish, both freshwater and especially marine species, such as golden pompano (*Trachinotus ovatus*), snubnose pompano (*T. blochii*), yellow croaker (*Larimichthys crocea*), northern snakehead (*Channa argus*), blotched snakehead (*C. maculata*), largemouth bass (*Micropterus salmoides*), yellow tail (*Seriola quinqueradiata*), amberjack (*S. dumerelli*) and sea bass (*Lateolabrax japonicus*) [5,10,11].

The pathogenic mechanisms and specific virulence factor of *N. seriolae* remain unknown. Beaman found that the virulence of *N. asteroides* was associated with its resistance to oxidative killing by phagocytes, inhibition of phagosome-lysosome fusion, neutralization of phagosomal acidification, and alteration of lysosomal enzymes within phagocytes, as well as the secretion of toxic substances by the nocardia, which are apparently involved in these processes [12]. Previous studies showed that *N. asteroides* might induce apoptosis in host cells [13–15]. The secretome studies of pathogenic actinomycetes have shown that secreted proteins, especially several mitochondrial targeted proteins, are closely related to their pathogenicity and may play an important role in the regulation of cell apoptosis and bacterial pathogenesis [16–19]. The modulation of cell

apoptosis in the host by pathogenic bacteria is a major pathogenicity mechanism. Many bacterial proteins have a highly specific activity with the host mitochondria, which participate in the cell apoptosis by influencing the signaling pathways [20–22]. The mechanism of mitochondrial pathway regulation of apoptosis is very complex. The mitochondrial targeted protein allows the mitochondrial outer membrane permeabilization (MOMP) accompanied with the loss of mitochondrial membrane potential ($\Delta\Psi_m$). Subsequently, various apoptotic mediators (mammalian serine protease (Omi/HtrA2), second mitochondrial activator of caspase (SMAC), direct IAP binding protein with low pI (Diablo), cytochrome C (Cyt C), and apoptosis-inducing factors (AIF)) are released into the cytosol. The releases of Cyt c in cytosol causes the association of apoptosis protease-activating factor-1 (Apaf-1) and ATP/dATP to form apoptotic body, while AIF enters the nucleus acting in a caspase independent manner by causing chromatin condensation and fragmentation. Finally, the downstream procaspase-3 is activated, and cell apoptosis is induced in the host [23–25]. Thus, we speculated whether the mitochondrial targeting secreted proteins from *N. seriolae* are potential virulence factors.

A bioinformatic analysis based on the whole genome sequence data of *N. seriolae* strain ZJ0503 [26] showed that *ORF3141* encoded a secreted protein which might target the mitochondria in the host cell, and the ORF3141 protein had a signal peptide at the N-terminate (residues 1-30, MKLLNPRGFGLVCASAAVAAGLMLAGCANT). Currently, the function of this protein and its homologs are totally unknown. Meanwhile, the homologs of ORF3141 are mainly present in high GC Gram-positive bacteria. We proposed that the secreted protein of ORF3141 was possibly a virulence factor of *N. seriolae* strain ZJ0503. In this study, the function of ORF3141 was explored and whether this protein was the bacterial virulence factor that induced infection in fish was determined. Gene cloning, secreted protein identification, subcellular localization, overexpression, and apoptosis detection assays of ORF3141 and ORF3141 Δ sig proteins were performed. Moreover, the results indicated that *ORF3141* gene encoded a secreted protein which contained a mitochondrial targeting sequence and was involved in apoptosis regulation. These results may lay the foundation for further study on the function of ORF3141 protein and promote the understanding of the molecular pathogenic mechanism of *N. seriolae*.

Results

Cloning and the sequence analysis of ORF3141

The *ORF3141* and *ORF3141Δsig* genes of *N. seriolae* strain ZJ0503 were cloned, and the recombinant plasmids of pEGFP-3141, pEGFP-3141Δsig, pcDNA-3141, and pcDNA-3141Δsig were constructed successfully. Sequence analysis revealed that an open reading frame (ORF) of *ORF3141* gene was 558 bp, and the deduced amino acid sequence had a length of 185 residues (Fig. 1A). The predicted molecular weight of ORF3141 protein was 18.96 kDa, and the theoretical isoelectric point (pI) was 5.85. The instability index of ORF3141 was computed to be 25.43, indicating its stability. The grand average of hydropathicity was -0.112, which signifies that the ORF3141 protein was hydrophilic. ORF3141 was predicted to be a secreted protein with a cleavable signal peptide at N-terminus and co-localized with the mitochondria in the host. ORF3141 was composed of a signal peptide (residues 1-30, MKLLNPRGFGLVCASAAVAAGLMLAGCANT) at N-terminate, two low complexity domains (residues 47-70 and 90-117), and five kinds of functional sites including N-myristoylation, Protein kinase C phosphorylation, Casein kinase II phosphorylation, N-glycosylation, and cAMP- and cGMP-dependent protein kinase phosphorylation sites (Fig. 1B). No protein superfamily members of ORF3141 protein were found. The three-dimensional structure of ORF3141 from *N. seriolae* was predicted by I-TASSER. The structure was closed to that of multidrug ABC transporter Sav1866 from *Staphylococcus aureus* (Fig. 1C), which was classified as a hydrolase [27]. Moreover, the protein contained two ligand binding sites (residues 108 and 111) of coenzyme F430 and an enzyme active site (residues 107) of ATP phosphohydrolase (steroid-exporting).

Protein BLAST showed that the deduced amino acid sequence of ORF3141 protein displayed low identity with homologous sequence from Actinomycetes, with the highest identity of *N. concava* (84%) and the lowest identity of *Mycobacterium marinum* (23%). This protein had a rather high homology in *Nocardia* species (61% - 84%) (S1 Table) and its sequence was also found in other *N. seriolae* stains, such as *N. seriolae* N-2927 [28], *N. seriolae* U-1 [29], *N. seriolae* EM150506, *N. seriolae* CK-14008 and *N. seriolae* UTF-1 [30] (S2 Table). Multiple alignment of the deduced amino acid sequences of the ORF3141 protein showed no putative conserved domains

(Fig. 2A). A phylogenetic tree was constructed to identify the evolutionary relationships between *N. seriolae* with other species of ORF3141 homologs, on the basis of the deduced amino acid sequence of ORF3141. The results showed that ORF3141 proteins in *Nocardia sp.* were clustered in one group (Fig. 2B).

Identification of ORF3141 protein was a secreted protein

Preparation and shotgun mass spectrometry (MS) of the extracellular products of *N. seriolae* strain ZJ0503 showed that the characteristic peptide “TIETELNATR” of ORF3141 was detected with confidence greater than or equal to 99%. This result confirmed that ORF3141 was a secreted protein of *N. seriolae*.

Subcellular localization of ORF3141 and ORF3141 Δ sig proteins in FHM cells

Subcellular localization of ORF3141 and ORF3141 Δ sig proteins in the FHM cells were determined by the expression of ORF3141-GFP and ORF3141 Δ sig-GFP fusion protein, respectively. ORF3141-GFP and ORF3141 Δ sig-GFP fusion protein were detected with strong green fluorescence signal at 48 h post-transfection (hpt). The mitochondria were shown with red fluorescence, and the nucleus were displayed with blue fluorescence. Compared with the location of mitochondria and nucleus in the pEGFP-3141 or pEGFP-3141 Δ sig transfected cells, the ORF3141-GFP fusion protein was exhibited an aggregated distribution and co-localized with the mitochondria in the cytoplasm, while the ORF3141 Δ sig-GFP fusion protein was distributed in the whole FHM cell and did not co-localize with the mitochondria. These contrasting results indicated that the N-terminate signal peptide played an important role in mitochondrial targeting. In addition, the expression of ORF3141 or ORF3141 Δ sig proteins changed the distribution of the mitochondria from perinuclear halo into lumps, and the apoptotic bodies were detected. By contrast, the signal of GFP was distributed in both cytoplasm and nucleus in the control cells which were transfected with pEGFP-N1, and no specific fluorescence co-localized with the mitochondria (Fig. 3).

Apoptosis induced in FHM cells by overexpression of ORF3141 and ORF3141 Δ sig proteins

To investigate whether ORF3141 and ORF3141 Δ sig proteins were involved in the apoptosis of fish cells, plasmids pcDNA-3141 and pcDNA-3141 Δ sig were transfected into the FHM cells, respectively. Several typical apoptotic features, such as cellular shrinkage, blebbing of the nuclear membrane, nuclear pyrosis, and nuclear fragmentation, in ORF3141 and ORF3141 Δ sig proteins overexpressed FHM cells were observed by microscope with fluorescent and white light (Fig. 4). Thus, the ORF3141 and ORF3141 Δ sig proteins might lead to fish cell apoptosis. The number of apoptotic bodies were counted and used to calculate the apoptotic rate (number of apoptotic bodies / total number of cells $\times$ 100%) [31]. The results showed that 19.82 % and 17.05% cells underwent apoptosis in pcDNA-3141 and pcDNA-3141 Δ sig transfected groups respectively, while only 2.75% cells in control group (Fig. 5A). At 48 hpt, the expression of ORF3141 or ORF3141 Δ sig proteins in the plasmids transfected FHM cells were confirmed by the presence of a specific band using RT-PCR (Fig. 5B) and Western blot analysis (Fig. 5C).

Apoptotic detection of ORF3141 and ORF3141 Δ sig overexpressed FHM cells

The $\Delta\Psi_m$ was shown with the evident damage in JC-1 polymer/monomer fluorescence ratio in ORF3141 or ORF3141 Δ sig overexpressed cells. The $\Delta\Psi_m$ values of both pcDNA-3141 and pcDNA-3141 Δ sig transfected groups cells were decreased to minimum values at 72 hpt, which were 0.54- and 0.53-folds lower than that of the control cell group, respectively (Fig. 6A). Meanwhile, measurement of caspase-3 activity showed that caspase-3 was activated in ORF3141 or ORF3141 Δ sig proteins overexpressed cells and reached maximum value at 48 hpt, which were approximately 1.23- and 1.80-folds higher than that of the control cell group, respectively (Fig. 6B). Several apoptosis-related genes of Bcl-2 family mRNA expression were investigated at 0, 24, 48, 72, and 96 hpt by qRT-PCR, including B cell lymphoma-2 (*Bcl-2*), Bcl-2 associated X (*Bax*), Bcl-2 antagonist of cell death (*Bad*), and Bcl-2 interacting domain death agonist (*Bid*) genes. The results showed that pro-apoptotic (*Bad*, *Bid*, and *Bax*) and anti-apoptotic (*Bcl-2*) genes were rapidly activated in both pcDNA-3141 and pcDNA-3141 Δ sig transfected groups cells and their mRNA expression reached peak level at 72 hpt, respectively. Moreover, there were 11.19- and 10.60-folds

of *Bax* to *Bcl-2* genes mRNA expression in pcDNA-3141 and pcDNA-3141 Δ sig transfected groups higher than that of 5.17-fold in the control group at 72 hpt (Fig. 6C). All quantitative assays of $\Delta\Psi_m$ value, caspase-3 activity, and apoptosis-related genes mRNA expression indicated that cell apoptosis could be induced by the overexpression of ORF3141 or ORF3141 Δ sig proteins in the FHM cells.

Discussion

No relevant literature on the ORF3141 protein of *Nocardia* or other species is available, and this protein remains largely unknown. Here, we preliminary investigated the characteristic, structure, and function of ORF3141 from *N. seriolae* strain ZJ0503. Moreover, *ORF3141* and *ORF3141* Δ sig were successfully cloned. The sequence analysis showed an ORF of *ORF3141* gene of 558 bp. This gene encoded a protein with 185 amino acid residues, molecular weight of 18.96 kDa, and isoelectric point of 5.85. A phylogenetic tree based on the deduced amino acid sequence of ORF3141 shared low homology between *N. seriolae* and other homologous sequences from Actinomycetes. However, ORF3141 had a rather high homology in *Nocardia* species. Bioinformatic analysis revealed that ORF3141 had no putative conserved domains and no superfamily. This protein comprised a signal peptide and two low complexity domains, but the function of these domains remains unknown. *ORF3141* was predicted to encode a secreted protein and co-localized with the mitochondria in the host cells. Experimentally, the MS analysis of the extracellular products from *N. seriolae* strain ZJ0503 proved that ORF3141 was a secreted protein. Subcellular localization of ORF3141 in the host cells exhibited an aggregated distribution and coincided with the mitochondria. Thus, ORF3141 co-localized with the mitochondria in the host cells. Distinguishingly, ORF3141 Δ sig protein was evenly distributed in the whole cell and did not co-localize with the mitochondria in the host cells. These contrasting results were due to the presence or absence of the signal peptide. Similar results were found in the subcellular localization study of EGFP-GRP75 fusion proteins; that is, GRP-75 with signal peptide was exclusively co-localized with the mitochondria, whereas GRP-75 without the signal peptide was distributed in the whole host cell [32]. Thus, the N-terminate signal peptide was important for mitochondrial targeting. The mitochondrial targeting protein contains specific

sequences with specific information. Five kinds of mitochondrial targeting signals (MTSSs) are used to determine the protein localization with the mitochondrial outer membranes, inner membranes, or matrix [33]. Many proteins are destined for the mitochondrial matrix typically cleavable targeting signal at the N-terminus [34], which are commonly 20 - 60 amino acids in length and are cleaved upon import into the mitochondrial matrix by the mitochondrial processing peptidase [35,36]. Therefore, ORF3141 was assumed to target the mitochondria matrix.

The function of ORF3141 has not been studied, until now. In our study, we observed that the distribution of mitochondria was altered from perinuclear halo into lumps, and apoptotic bodies appeared during the subcellular localization and overexpression assays, which indicated that cell apoptosis was induced after ORF3141 or ORF3141 Δ sig proteins stimulating in FHM cells. Mitochondria appear to play a central role in the regulation of several cell death pathways, such as apoptosis, autophagic cell death, necrosis, pyroptosis, and pyronecrosis [37,38]. Other characterized bacterial proteins had also been shown to target the mitochondria and induce apoptosis. For example, Omp38 of *Acinetobacter baumannii* was shown to target mitochondria of Hep-2 cells where it induced the release of Cyt c and AIF, to promote apoptosis [39]. When expressed in human cells, PorB of *Neisseria gonorrhoeae* co-localized with the mitochondria and caused dissipation of $\Delta\Psi_m$ but did not induce the release of Cyt c [40]. SipB of *Salmonella enterica* targeted the host cell mitochondria during infection and supported autophagy-mediated cell death in caspase-1 deficient macrophages [41]. As a mitochondrial targeting bacterial secreted protein, ORF3141 might participate in the regulation of cell apoptosis. Distinctively, ORF3141 Δ sig protein was observed to also induce cell apoptosis without targeting the mitochondria in FHM cells. Thus, it was proved that the domain part without the signal peptide of ORF3141 protein had a critical relationship with cell apoptosis whether it was targeted to the mitochondria or existent in the cytoplasm. Moreover, quantitative assays of $\Delta\Psi_m$ value, caspase-3 activity, and apoptosis-related genes mRNA expression were performed to determine how did the ORF3141 and ORF3141 Δ sig protein participate in cell apoptosis in FHM cells. The Bcl-2 family proteins are critical regulators of mitochondrial apoptosis, which include the anti-apoptotic multidomain members (containing all four BH domains), the pro-apoptotic multidomain members (containing three BH domains), and pro-apoptotic BH3-only members (containing the sole BH3 domain). In addition, a large number of

Bcl-2 family protein contain a hydrophobic transmembrane anchoring (TM) domain at the C-terminus which allows them to co-localize to subcellular membranes [42,43]. Among these proteins, Bcl-2 was discovered as an anti-apoptosis protein that blocks cell apoptosis, while Bax was clarified as a pro-apoptosis protein which can form large openings in lipid bilayers to control the MOMP and induce apoptosis [44–46]. Thus, a slight change in the dynamic balance of Bax and Bcl-2 seem to determine the inhibition or promotion of cell apoptosis with apoptotic stimuli [47]. In fact, it is clear that Bcl-2 family proteins can also localize to cytosol and other organelles, including the Golgi apparatus, the endoplasmic reticulum (ER), and the nuclear out membrane (NOM) or the nucleus itself [48]. In non-apoptotic cells, Bax localization is essentially diffuse in the cytosol but it is relocated to mitochondria after cell apoptosis induction [49]. Therefore, it could be speculated that both ORF3141 and ORF3141 Δ sig proteins were initiated by activating the apoptosis-related genes (Bad, Bid, Bax, and Bcl-2) and induced the translocation of Bax from the cytosol to the mitochondria in FHM cells. Besides, the relative ratio of *Bax* and *Bcl-2* genes mRNA expression was out-of-balance, which were responsible for the $\Delta\Psi_m$ values declined and caspase-3 activity increased in this study. Although the subcellular localization of ORF3141 and ORF3141 Δ sig proteins were different, they could induce the cell apoptosis via the mitochondrial pathway.

Further studies are required to verify the mechanisms involved in ORF3141-induced cell apoptosis by yeast two hybrid experiment. Whether the ORF3141 was the major virulence factor of *N. seriolae* remains to be clarified by constructing Δ ORF3141 mutant attenuated and ORF3141 mutant completed *N. seriolae*. The relationship of the interaction between ORF3141 and macrophage also need to be highlighted in future studies. In summary, we cloned an *ORF3141* gene from *N. seriolae* strain ZJ0503 and analyzed its sequence structure. Moreover, MS analysis, subcellular localization, overexpression assay, and apoptosis detection were performed. These results showed that ORF3141 of *N. seriolae* strain ZJ0503 was a secreted protein that targeted the host cell mitochondria and induced apoptosis in transfected FHM cells.

Materials and methods

Bacterial strains, FHM cells, and plasmids

N. seriolae strain ZJ0503 was isolated from diseased golden pompano (*T. ovatus*) in YangJiang City, Guangdong Province, China [26] and was cultured in an optimized medium [glucose 20 g·L⁻¹, yeast extract 15 g·L⁻¹, K₂HPO₄ 0.75 g·L⁻¹, CaCl₂ 0.2 g·L⁻¹ (sterilized separately), and NaCl 5 g·L⁻¹, PH 6.5 ± 0.2] at 28 °C [50]. *Escherichia coli* DH5α was used for gene cloning and it grown in Luria-Bertani (LB) medium with vigorous shaking at 37 °C. Fathead minnow (FHM) epithelial cells [51] were stored at -196 °C in Guangdong Provincial Key Laboratory of Pathogenic Biology and Epidemiology for Aquatic Economic Animals and cultured at 25 °C in Leibovitz's L15 medium containing 10% fetal bovine serum (Invitrogen, USA) in 5% CO₂ incubator. Plasmid pEGFP-N1 was used for subcellular localization, while pcDNA3.1/His A was used for overexpression.

Ethics statement

All animal experiments were handled in accordance with guidelines defined by the Institutional Animal Care Use Committee (IACUC) of Guangdong Ocean University and were approved by Guangdong Provincial Key Laboratory of Pathogenic Biology and Epidemiology for Aquatic Economic Animals, following strict compliance with the regulations of the local government.

Gene cloning and plasmid construction of *ORF3141* and *ORF3141*Δsig

Genomic DNA was extracted from *N. seriolae* strain ZJ0503 using TIANamp Bacteria DNA Kit (Tiangen, Beijing) following the manufacture's instruction. Four pairs of different primers were carefully designed with corresponding restriction enzyme sites using Primer 5.0, on the basis of *ORF3141* and *ORF3141*Δsig genes of the whole genome sequence data of *N. seriolae* strain ZJ0503. The PCR primers of pEGFP-3141F/R and pcDNA-3141F/R (Table 3) were used to amplify the *ORF3141* gene. The PCR primers of pEGFP-3141ΔsigF/R and pcDNA-3141ΔsigF/R (Table 1) were used to amplify the *ORF3141*Δsig gene. The PCR procedures were performed with KOD-

plus-Neo DNA polymerase (Toyobo, Osaka, Japan), using the following PCR program: pre-denaturation at 98 °C for 2 min, 30 cycles at 98 °C for 10 s, 55 °C for 15 s, 69.5 °C for 15 s; and a final extension at 68 °C for 5 min. All PCR products of *ORF3141* and *ORF3141Δsig* genes were electrophoresed on 1% agarose gel and purified using EasyPure PCR Purification Kit (TRANSGEN, Beijing). The purified PCR product of *ORF3141* was digested by corresponding restriction enzymes, ligated into eukaryotic vectors pEGFP-N1 and pcDNA3.1/His A, respectively. And then transformed into competent *E. coli* DH5α cells. The different constructs were confirmed by corresponding restriction enzyme digestion and DNA sequencing by Guangzhou Sangon Biologic Engineering & Technology and Service Co. Ltd. The purified PCR product of *ORF3141Δsig* was processed similarly. Finally, the constructed recombinant plasmids were named as pEGFP-3141, pEGFP-3141Δsig, pcDNA-3141, and pcDNA-3141Δsig.

308

309

Table 1. Primers used for gene cloning

Gene Name	Primer Name	Sequence 5'-3'	Restriction endonucleases
ORF3141	pEGFP-3141F	GGAATTCATGAAGCTGCTGAACCCGCG	<i>Bam</i> H I
	pEGFP-3141R	CGGGATCCCGCTTGCCCGGGCAGGCGTTC	<i>Eco</i> R I
	pcDNA-3141F	CGGGATCCATGAAGCTGCTGAACCCGCG	<i>Eco</i> R I
	pcDNA-3141R	CGGAATTCCTTGCCCGGGCAGGCGTTCCG	<i>Bam</i> H I
ORF3141Δsig	pEGFP-3141ΔsigF	GGAATTCATGGTCGAGGGTACCCCGACGGT C	<i>Eco</i> R I
	pEGFP-3141ΔsigR	CGGGATCCCGCTTGCCCGGGCAGGCGTTC	<i>Bam</i> H I
	pcDNA-3141ΔsigF	CGGGATCCATGGAGGGTACCCCGACGGT	<i>Bam</i> H I
	pcDNA-3141ΔsigR	CGGAATTCCTTGCCCGGGCAGGCGTTCC	<i>Eco</i> R I

310 Bioinformatics analysis, sequence alignments, and phylogenetic analysis

311 Based on the whole genome sequence data of *N. seriola* strain ZJ0503, sequence analysis was
 312 performed with the BLAST program using NCBI (<http://www.ncbi.nlm.nih.gov/BLAST/>). The
 313 amino acid sequence for ORF3141 was deduced, and the physical and chemical properties were

predicted using ExPASy software (<http://www.expasy.org/>). The location of domains was predicted by the InterProScan program (<http://www.ebi.ac.uk/Tools/pfa/iprscan/>). The typical structures of ORF3141 protein were predicted by I-TASSER online software (<http://zhanglab.ccmb.med.umich.edu/I-TASSER/>). The potentially excreted proteins were predicted by LocTree3 (<https://roslab.org/services/loctree/>) and ExPASy-PROSITE. The subcellular localization and signal peptides were predicted with LocTree 3 and SignalP 4.1 Server (www.cbs.dtu.dk/services/SignalP/), respectively. The MTSs were predicted using TargetP (<http://www.cbs.dtu.dk/services/TargetP/>) and PSORT II (<https://psort.hgc.jp/form2.html>). Protein multiple sequence alignments of ORF3141 protein were performed by ClustalX 2.0 program with the default parameters and edited by the GeneDoc software. The phylogenetic tree was generated based on the deduced amino acid sequence of ORF3141 with the neighbor-joining method using MEGA 6 program, in which the Poisson distribution substitution model and bootstrapping procedure with 1000 bootstraps were applied.

Preparation and identification of the *N. seriolae* extracellular products

The extracellular products of *N. seriolae* strain ZJ0503 were obtained by cellophane overlay method [52]. Specifically, *N. seriolae* were grown on optimized medium agar plate at 28 °C for 2 d, and a single colony was prepared for bacterial suspension. Then, 100 µL of the bacterial suspension was spread closely on optimized medium plates covered with sterile cellophane sheet and incubated at 28 °C for 3 - 5 d. The *N. seriolae* cells grown on the cellophane sheet were stripped away from the optimized medium plates, and the extracellular products were washed down with sterilized PBS subsequently. The harvested suspension was centrifuged at 8000×g, 4 °C for 20 min, and the supernatant containing extracellular products was filter sterilized with a 0.2 µm membrane filter. Then, the sterilized supernatant was transferred into a dialysis tubing (3.5k MW) and dialyzed in ultrapure water at 4 °C for 16 - 24 h. During dialysis, the ultrapure water was changed 3 - 4 times. The purified supernatant was transferred into a centrifuge tube after dialysis and frozen under -80 °C. Finally, it was lyophilized using a vacuum freeze dryer to obtain the protein dry powder which was identified using shotgun MS.

FHM cell mediated transient expression and subcellular localization

Plasmids of pEGFP-3141, pEGFP-3141 Δ sig, and pEGFP-N1 were prepared in advance using an endotoxin-free plasmid purification kit (Qiagen Inc., Chatsworth, CA). Subcellular localization of ORF3141 and ORF3141 Δ sig proteins were performed through pEGFP-N1 fusion protein expression. Given that a permanent FHM cell line is presently available, FHM cells were cultured in 24-well plates and grown to 70 % confluency for transfection. The FHM cells were transfected with empty pEGFP-N1 plasmids, recombinant pEGFP-3141 plasmids, and recombinant pEGFP-3141 Δ sig plasmids using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol, respectively. At 48 hpt, the FHM cells were washed with PBS (pH 7.4), stained with 300 nM MitoTracker Red CMXRos dye (Molecular Probes, Carlsbad, CA) at 28 °C for 45 min, fixed with 4% paraformaldehyde for 30 min, and labeled with 1 μ g/mL diamidino-2-phenylindole (DAPI) at room temperature for 10 min. Finally, the cells were rinsed with PBS and mounted with 50% glycerol. The fluorescence exhibited by the transfected FHM cells were observed using a fluorescence microscope (Leica DM IRB).

Overexpression of ORF3141 and ORF3141 Δ sig proteins in FHM cells

The extraction and transfection of recombinant pcDNA-3141 plasmids, recombinant pcDNA-3141 Δ sig plasmids, and control pcDNA3.1/His A plasmid were performed with reference to step 2.5. The transfected FHM cells were stained with DAPI at 48 hpt and microscopically observed to test whether the overexpression of ORF3141 and ORF3141 Δ sig proteins induced apoptosis in fish cells. Meanwhile, to identify the expression of ORF3141 and ORF3141 Δ sig proteins in the transfected FHM cells, the cells were harvested at 48 hpt to extract the total RNA and proteins for RT-PCR and Western blot analysis, respectively. RT-PCR was performed with primers pcDNA-3141F/R and pcDNA-3141 Δ sigF/R (Table 3) following the synthesis of cDNA. Western blot was conducted using mouse anti-His monoclonal antibody (Sigma, St. Louis, MO) as the primary antibody at a dilution of 1:1000 and horseradish peroxidase-conjugated goat anti-mouse IgG (Sigma, St. Louis, MO) as the secondary antibody at a dilution of 1:5000.

Quantitative analysis of $\Delta\Psi_m$ value, caspase-3 activity and apoptosis-related gene mRNA expression

Several quantitative assays of mitochondrial membrane potential ($\Delta\Psi_m$) value, caspase-3 activity, and apoptosis-related gene mRNA expression (pro-apoptotic gene: *Bad*, *Bid*, and *Bax*; anti-apoptotic gene: *Bcl-2*), were performed to confirm the apoptosis caused by overexpression of proteins in the transfected recombinant pcDNA-3141 or pcDNA3141 Δ sig in the FHM cells. The $\Delta\Psi_m$ values were assessed at 24, 48, and 72 hpt using the JC-1 assay kit (Beyotime, Shanghai, China) by the method described previously with minor modification [53]. The positive control was treated with carbonyl cyanide 3-chlorophenylhydrazone (CCCP, 10 mM) to medium at 1:1000 25 °C for 20 min. Then, $\Delta\Psi_m$ value was measured by changes in the 590/530 JC-1 emitted fluorescence with an Enspire 2300 Multilabel Reader (Perkin Elmer, MA, USA). The caspase-3 activity was detected in the FHM cells at 24 and 48 hpt as described previously [54] using a caspase-3 colorimetric assay kit (BioVision, Milpitas, CA).

Three groups of transfected FHM cells were harvested at 0, 12, 24, 48, 60, and 72 hpt to extract the total RNA, and quantitative real-time PCR (qRT-PCR) was performed following the synthesis of cDNA. The cell apoptotic effect of post-transfection on the expression of apoptosis-related genes mRNA, was investigated using real-time SYBR green PCR Master Mix on Applied Biosystems 7500 Real-Time PCR system (ABI, USA). Each assay was performed in triplicate with β -actin gene as the internal control. According to the apoptosis gene sequence, five pairs of specific primer (Table 2) were carefully designed for qRT-PCR investigation. The PCR was performed in a 10 μ L reaction volume containing 0.5 μ L of each primer (10 μ M), 0.3 μ L of cDNA, 3.7 μ L of PCR-grade water, and 5 μ L of SYBR[®] Select Master Mix (ABI, USA) according to the manufacturer's protocol. The PCR conditions were as follows: 95°C for 2 min; 40 cycles of 95 °C for 15 s, and 58 °C for 1 min for four apoptosis-related genes and β -actin, respectively. Melt curve analysis of the amplification products was performed over a range of 60 - 95 °C at the end of each PCR reaction to confirm the single product generation. The relative expression levels of four apoptosis-related genes were calculated using the comparative Ct $2^{(-\Delta\Delta C_t)}$ method [55].

Table 2. Primes used for apoptosis-related genes investigated by qRT-PCR.

Gene Name	Primer Name	Sequence 5'-3'
<i>β-actin</i>	β-actin-F	ACAATCAATACGGCTGCCATGG
	β-actin-R	TTGGCATACAGGTCCTTACTTACGT
<i>Bad</i>	Bad-F	TGATCCTTTCAGGCGGAGATCTCGC
	Bad-R	CAGACTCTTTGTGACTCCAAAGGAA
<i>Bid</i>	Bid-F	CTGCTTCTCCTTTCCTTCTTTGAGC
	Bid-R	GATCAACTCAGCAGCCATATCCCTT
<i>Bax</i>	Bax-F	TGGCACTGTTTCACCTCG
	Bax-R	ATCCTCCTTGCTGTCTGATC
<i>Bcl-2</i>	Bcl-2-F	TGGGACTGTTTGCCTTCG
	Bcl-2-R	TCTGCCGCTGCATCTTTT

395

396 Statistical analysis

397 Data were presented as the means ± standard deviation (SD). Statistical analysis was performed
398 with one-way ANOVA with the SPSS statistics 21.0 software and the figures were edited by
399 GraphPad Prism software. Data represent the means for three independent experiments and
400 statistical significant is highlighted with asterisks in the figures as follows: $p > 0.05$, not significant;
401 $p < 0.05$ (*), significant; $p < 0.01$ (**), extremely significant. The means and p values for pairwise
402 comparisons of all experiments are provided in S3 Table.

403 Acknowledgment

404 We are grateful to all the laboratory members for the discussion and critical readings of the
405 manuscript.

406 References

407 1. Chen SC. Study on the pathogenicity of *Nocardia asteroides* to the Formosa snakehead, *Channa maculata*

- 408 (Lacepède), and largemouth bass, *Micropterus salmoides* (Lacepède). J Fish Dis. 2010;15: 47–53.
409 doi:10.1111/j.1365-2761.1992.tb00635.x
- 410 2. Pei-Chi W, Ming-An T, Yu-Chi L, Yanting C, Shih-Chu C. *Nocardia seriolae*, a causative agent of systematic
411 granuloma in spotted butterflyfish, *Scatophagus argus*, Linn. Afr J Microbiol Res. 2014;8: 3441–3452.
412 doi:10.5897/AJMR2014.6874
- 413 3. Xia L, Zhang H, Lu Y, Cai J, Wang B, Jian J. Development of a Loop-Mediated Isothermal Amplification
414 Assay for Rapid Detection of *Nocardia salmonicida*, the Causative Agent of Nocardiosis in Fish. J Microbiol
415 Biotechnol. 2015;25: 321–327. doi:10.4014/jmb.1406.06052
- 416 4. Vu-Khac H, Duong VQB, Chen S-C, Pham TH, Nguyen TTG, Trinh TTH. Isolation and genetic
417 characterization of *Nocardia seriolae* from snubnose pompano *Trachinotus blochii* in Vietnam. Dis Aquat
418 Organ. 2016;120: 173–177. doi:10.3354/dao03023
- 419 5. Ho PY, Byadgi O, Wang PC, Tsai MA, Liaw LL, Chen SC. Identification, Molecular Cloning of IL-1 β and
420 Its Expression Profile during *Nocardia seriolae* Infection in Largemouth Bass, *Micropterus salmoides*: Int J
421 Mol Sci. 2016;17. doi:10.3390/ijms17101670
- 422 6. Labrie L, Ng J, Tan Z, Komar C, Ho E, Grisez L, et al. Nocardial infections in fish: an emerging problem in
423 both freshwater and marine aquaculture systems in Asia. Dis Asian Aquac VI. 2005;
- 424 7. Elkesh A, Kantham KPL, Shinn AP, Crumlish M, Richards RH. Systemic nocardiosis in a Mediterranean
425 population of cultured meagre, *Argyrosomus regius* Asso (Perciformes: Sciaenidae). J Fish Dis. 2013;36:
426 141–149. doi:10.1111/jfd.12015
- 427 8. Manrique WG, Da SCG, de Castro MP, Petrillo TR, Figueiredo MA, Ma DAB, et al. Expression of Cellular
428 Components in Granulomatous Inflammatory Response in *Piaractus mesopotamicus* Model. Plos One.
429 2015;10: e0121625.
- 430 9. Maekawa S, Yoshida T, Wang P-C, Chen S-C. Current knowledge of nocardiosis in teleost fish. J Fish Dis.
431 2018;41: 413–419. doi:10.1111/jfd.12782
- 432 10. Wang GL, Yuan SP, Jin S. Nocardiosis in large yellow croaker, *Larimichthys crocea* (Richardson). J Fish
433 Dis. 2005;28: 339–345. doi:10.1111/j.1365-2761.2005.00637.x
- 434 11. Wang GL, Yi-Jun XU, Jin S. Identification and phylogenetic analyses of *Nocardia* in snakehead,
435 *Ophiocephalus argus* Cantor. J Fish China. 2008;32: 449–454.
- 436 12. Beaman BL, Beaman L. *Nocardia* species: host-parasite relationships. Clin Microbiol Rev. 1994;7: 213–64.
- 437 13. Barry DP, Beaman BL. *Nocardia asteroides* strain GUH-2 induces proteasome inhibition and apoptotic death
438 of cultured cells. Res Microbiol. 2007;158: 86–96. doi:10.1016/j.resmic.2006.11.001
- 439 14. Camp DM, Loeffler DA, Razoky BA, Tam S, Beaman BL, LeWitt PA. *Nocardia asteroides* culture filtrates
440 cause dopamine depletion and cytotoxicity in PC12 cells. Neurochem Res. 2003;28: 1359–1367.
441 doi:10.1023/A:1024944431725

- 442 15. Loeffler DA, Camp DM, Qu S, Beaman BL, LeWitt PA. Characterization of dopamine-depleting activity of
443 *Nocardia asteroides* strain GUH-2 culture filtrate on PC12 cells. Microb Pathog. 2004;37: 73–85.
444 doi:10.1016/j.micpath.2004.05.001
- 445 16. Lartigue L, Faustin B. Mitochondria: Metabolic regulators of innate immune responses to pathogens and cell
446 stress. Int J Biochem Cell Biol. 2013;45: 2052–2056. doi:10.1016/j.biocel.2013.06.014
- 447 17. Rudel T, Kepp O, Kozjak-Pavlovic V. Interactions between bacterial pathogens and mitochondrial cell death
448 pathways. Nat Rev Microbiol. 2010;8: 693–705. doi:10.1038/nrmicro2421
- 449 18. Saint-Georges-Chaumet Y, Edeas M. Microbiota-mitochondria inter-talk: consequence for microbiota-host
450 interaction. Pathog Dis. 2016;74: ftv096. doi:10.1093/femspd/ftv096
- 451 19. West AP, Shadel GS, Ghosh S. Mitochondria in innate immune responses. Nat Rev Immunol. 2011;11: 389–
452 402. doi:10.1038/nri2975
- 453 20. Rudel T, Kepp O, Kozjak-Pavlovic V. Interactions between bacterial pathogens and mitochondrial cell death
454 pathways. Nat Rev Microbiol. 2010;8: 693–705. doi:10.1038/nrmicro2421
- 455 21. Neely AM, Zhao G, Schwarzer C, Stivers NS, Whitt AG, Meng S, et al. N-(3-Oxo-acyl)-homoserine lactone
456 induces apoptosis primarily through a mitochondrial pathway in fibroblasts. Cell Microbiol. 2018;20: e12787.
457 doi:10.1111/cmi.12787
- 458 22. Karijolic J, Abernathy E, Glaunsinger BA. Infection-Induced Retrotransposon-Derived Noncoding RNAs
459 Enhance Herpesviral Gene Expression via the NF-kappa B Pathway. Plos Pathogens. 2015;11: e1005260.
460 doi:10.1371/journal.ppat.1005260
- 461 23. Chen AW-G, Tseng Y-S, Lin C-C, Hsi Y-T, Lo Y-S, Chuang Y-C, et al. Norcantharidin induce apoptosis in
462 human nasopharyngeal carcinoma through caspase and mitochondrial pathway. Environ Toxicol. 2018;33:
463 343–350. doi:10.1002/tox.22521
- 464 24. Huang T-C, Chiu P-R, Chang W-T, Hsieh B-S, Huang Y-C, Cheng H-L, et al. Epirubicin induces apoptosis
465 in osteoblasts through death-receptor and mitochondrial pathways. Apoptosis. 2018;23: 226–236.
466 doi:10.1007/s10495-018-1450-2
- 467 25. Tang C, Wang J, Wei Q, Du Y-P, Qiu H-P, Yang C, et al. Tropomyosin-1 promotes cancer cell apoptosis via
468 the p53-mediated mitochondrial pathway in renal cell carcinoma. Oncol Lett. 2018;15: 7060–7068.
469 doi:10.3892/ol.2018.8204
- 470 26. Xia L, Cai J, Wang B, Huang Y, Jian J, Lu Y. Draft Genome Sequence of *Nocardia seriolae* ZJ0503, a Fish
471 Pathogen Isolated from *Trachinotus ovatus* in China. Genome Announc. 2015;3:
472 doi:10.1128/genomeA.01223-14
- 473 27. Dawson RJ, Locher KP. Structure of the multidrug ABC transporter Sav1866 from *Staphylococcus aureus* in
474 complex with AMP-PNP. Febs Lett. 2007;581: 935. doi:10.1016/j.febslet.2007.01.073
- 475 28. Imajoh M, Fukumoto Y, Jin Y, Sukeda M, Shimizu M, Ohnishi K, et al. Draft Genome Sequence of *Nocardia*

- 476 *seriolae* Strain N-2927 (NBRC 110360), Isolated as the Causal Agent of Nocardiosis of Yellowtail (*Seriola*
477 *quinqueradiata*) in Kochi Prefecture, Japan. Genome Announc. 2015;3: e00082-15.
- 478 29. Imajoh M, Sukeda M, Shimizu M, Yamane J, Ohnishi K, Oshima S. Draft Genome Sequence of
479 Erythromycin- and Oxytetracycline-Sensitive *Nocardia seriolae* Strain U-1 (NBRC 110359). Genome
480 Announc. 2016;4: e01606-15.
- 481 30. Yasuike M, Nishiki I, Iwasaki Y, Nakamura Y, Fujiwara A, Shimahara Y, et al. Analysis of the complete
482 genome sequence of *Nocardia seriolae* UTF1, the causative agent of fish nocardiosis: The first reference
483 genome sequence of the fish pathogenic *Nocardia* species. Plos One. 2017;12: e0173198.
484 doi:10.1371/journal.pone.0173198
- 485 31. Cai J, Huang Y, Wei S, Ouyang Z, Huang X, Qin Q. Characterization of LPS-induced TNF α factor (LITAF)
486 from orange-spotted grouper, *Epinephelus coioides*. Fish and Shellfish Immunol. 2013;35: 1858–1866.
487 doi:10.1016/j.fsi.2013.09.023
- 488 32. Chen H, Niu X, Gao A, Zhang S. Mitochondrial signal peptide guides EGFP-GRP75 fusion proteins into
489 mitochondria. Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi Chin J Cell Mol Immunol. 2016;32: 1311–1316.
- 490 33. Habib SJ, Neupert W, Rapaport D. Analysis and prediction of mitochondrial targeting signals. In: Pon LA,
491 Schon EA, editors. Mitochondria, 2nd Edition. San Diego: Elsevier Academic Press Inc; 2007. pp. 761–781.
- 492 34. Habib SJ, Neupert W, Rapaport D. Analysis and prediction of mitochondrial targeting signals. Methods Cell
493 Biol. 2007;80: 761–781.
- 494 35. Bohni PC, Daum G, Schatz G. Import of proteins into mitochondria. Partial purification of a matrix-located
495 protease involved in cleavage of mitochondrial precursor polypeptides. J Biol Chem. 1983;258: 4937–43.
- 496 36. Hawlitschek G, Schneider H, Schmidt B, Tropschug M, Hartl FU, Neupert W. Mitochondrial protein import:
497 identification of processing peptidase and of PEP, a processing enhancing protein. Cell. 1988;53: 795–806.
498 doi:10.1016/0092-8674(88)90096-7
- 499 37. Nagai T, Abe A, Sasakawa C. Targeting of enteropathogenic *Escherichia coli* EspF to host mitochondria is
500 essential for bacterial pathogenesis - Critical role of the 16th leucine residue in EspF. J Biol Chem. 2005;280:
501 2998–3011. doi:10.1074/jbc.M411550200
- 502 38. Ma C, Wickham ME, Guttman JA, Deng W, Walker J, Madsen KL, et al. *Citrobacter rodentium* infection
503 causes both mitochondrial dysfunction and intestinal epithelial barrier disruption in vivo: role of
504 mitochondrial associated protein (Map). Cell Microbiol. 2006;8: 1669–1686. doi:10.1111/j.1462-
505 5822.2006.00741.x
- 506 39. Choi CH, Lee EY, Lee YC, Park TI, Kim HJ, Hyun SH, et al. Outer membrane protein 38 of *Acinetobacter*
507 *baumannii* localizes to the mitochondria and induces apoptosis of epithelial cells. Cell Microbiol. 2005;7:
508 1127–1138. doi:10.1111/j.1462-5822.2005.00538.x
- 509 40. Muller A, Rassow J, Grimm J, Machuy N, Meyer TF, Rudel T. VDAC and the bacterial porin PorB of
510 *Neisseria gonorrhoeae* share mitochondrial import pathways. Embo J. 2002;21: 1916–1929.

511 doi:10.1093/emboj/21.8.1916

512 41. Hernandez LD, Pypaert M, Flavell RA, Galan JE. A Salmonella protein causes macrophage cell death by
513 inducing autophagy. *J Cell Biol.* 2003;163: 1123–1131. doi:10.1083/jcb.200309161

514 42. Ola MS, Nawaz M, Ahsan H. Role of Bcl-2 family proteins and caspases in the regulation of apoptosis. *Mol*
515 *Cell Biochem.* 2011;351: 41–58. doi:10.1007/s11010-010-0709-x

516 43. Edlich F. BCL-2 proteins and apoptosis: Recent insights and unknowns. *Biochem Biophys Res Commun.*
517 2018;500: 26–34. doi:10.1016/j.bbrc.2017.06.190

518 44. Pan Y, Ye C, Tian Q, Yan S, Zeng X, Xiao C, et al. miR-145 suppresses the proliferation, invasion and
519 migration of NSCLC cells by regulating the BAX/BCL-2 ratio and the caspase-3 cascade. *Oncol Lett.*
520 2018;15: 4337–4343. doi:10.3892/ol.2018.7863

521 45. Kalkavan H, Green DR. MOMP, cell suicide as a BCL-2 family business. *Cell Death Differ.* 2018;25: 46–
522 55. doi:10.1038/cdd.2017.179

523 46. Shimizu S, Narita M, Tsujimoto Y. Bcl-2 family proteins regulate the release of apoptogenic cytochrome c
524 by the mitochondrial channel VDAC. *Nature.* 1999;399: 483–487. doi:10.1038/20959

525 47. Liu JJ, Huang RW, Lin DJ, Peng J, Wu XY, Lin Q, et al. Expression of survivin and bax/bcl-2 in peroxisome
526 proliferator activated receptor-gamma ligands induces apoptosis on human myeloid leukemia cells in vitro.
527 *Ann Oncol.* 2005;16: 455–459. doi:10.1093/annonc/mdi077

528 48. Popgeorgiev N, Jabbour L, Gillet G. Subcellular Localization and Dynamics of the Bcl-2 Family of Proteins.
529 *Front Cell Dev Biol.* 2018;6: 13. doi:10.3389/fcell.2018.00013

530 49. Wolter KG, Hsu YT, Smith CL, Nechushtan A, Xi XG, Youle RJ. Movement of Bax from the cytosol to
531 mitochondria during apoptosis. *J Cell Biol.* 1997;139: 1281–92.

532 50. Xia L, Wang B, Xia H, Huang Y, Jian J, Yishan LU. Optimal culture conditions and medium of *Nocardia*
533 *serioleae*. *South China Fish Sci.* 2013;9: 51–56.

534 51. Gravell M, Malsberger RG. A permanent cell line from the fathead minnow (*Pimephales promelas*). *Ann N*
535 *Y Acad Sci.* 1965;126: 555–65. doi:10.1111/j.1749-6632.1965.tb14302.x

536 52. Xia L, Liang H, Xu L, Chen J, Bekaert M, Zhang H, et al. Subcellular localization and function study of a
537 secreted phospholipase C from *Nocardia serioleae*. *FEMS Microbiol Lett.* 2017;364.
538 doi:10.1093/femsle/fnx143

539 53. Sun YS, Lv LX, Zhao Z, He X, You L, Liu JK, et al. Cordycepol C induces caspase-independent apoptosis
540 in human hepatocellular carcinoma HepG2 cells. *Biol Pharm Bull.* 2014;37: 608–17.

541 54. Zhao Z, Chen C, Hu C-Q, Ren C-H, Zhao J-J, Zhang L-P, et al. The type III secretion system of *Vibrio*
542 *alginolyticus* induces rapid apoptosis, cell rounding and osmotic lysis of fish cells. *Microbiol-Sgm.* 2010;156:
543 2864–2872. doi:10.1099/mic.0.040626-0

544 55. Yang X. Analysis of the copy number of exogenous genes in transgenic cotton using real-time quantitative
545 PCR and the 2- $\Delta\Delta$ CT method. Afr J Biotechnol. 2012;11. doi:10.5897/AJB11.4117

546

547 **Supporting information**

548 **S1 Table.** Compared with the homology of ORF3141 among other *Nocardia* species.

549 **S2 Table.** The same sequence of ORF3141 between different *N. seriolae* strain.

550 **S3 Table.** Statistical analysis of all experimental data.

551 **Figure captions**

552 **Fig 1. Sequence and structure analysis of ORF3141 from *N. seriolae*.** (A) The full lengths
553 nucleotide and deduced amino acid sequence of *ORF3141* gene. The underline amino acid sequence
554 showed the signal peptide. (B) Schematic representation of the domain topology of ORF3141. The
555 ORF3141 protein was comprised of a signal peptide at N-terminate (residues 1-30), two low
556 complexity domains (residues 47-70, 90-117), and five kinds of functional sites. (C) The three-
557 dimensional structures of ORF3141 protein from *N. seriolae* (Left) and the closest structure to
558 multidrug ABC transporter Sav1866 from *Staphylococcus aureus* (Right). The diagrams were
559 generated using I-TASSER online and PyMOL 1.8 software.

560

561 **Fig 2. Multiple sequence alignment and construction of phylogenetic tree.** (A) Multiple
562 alignment of the deduced amino acid sequences of ORF3141 protein among different species.
563 Shaded regions indicate residues sharing homology, black regions indicate 100 % homology, dark
564 gray regions indicate homology higher than 75 %. (B) Construction of phylogenetic tree among *N.*
565 *seriolae* and other species with ORF3141 protein homology sequences. Protein sequences were
566 aligned with Clustal W, and the nonrooted neighbor-joining tree was generated by MEGA 5.0
567 program. Number at branch points indicate bootstrap support. GenBank accession numbers are
568 shown as followed: *Mycobacterium tuberculosis* H37Ra (ABQ74022.1), *Mycobacterium marinum*
569 M (ACC41759.1), *Mycobacterium rhodesiae* NBB3 (AEV74934.1), *Skermania piniformis*

(WP_066468955.1), *Gordonia soli* NBRC 108243 (GAC69437.1), *Gordonia polyisoprenivorans* NBRC 16320 (GAB24816.1), *Gordonia alkanivorans* NBRC 16433 (GAA10605.1), *Rhodococcus globerulus* (WP_045067034.1), *Rhodococcus opacus* (WP_012688156.1), *Rhodococcus koreensis* (WP_072945555.1), *Nocardia seriolae* ZJ0503 (WP_033086993.1), *Nocardia concave* (WP_040804030.1), *Nocardia pneumonia* (WP_040776180.1), *Nocardia brasiliensis* NBRC 14402 (GAJ80338.1), *Streptomyces ochraceiscleroticus* (WP_031055338.1), *Streptomyces violens* (WP_030248901.1).

Fig 3. Subcellular localization of ORF3141 and ORF3141Δsig proteins in FHM cell. Green fluorescence shows the ORF3141-GFP, ORF3141Δsig-GFP or GFP, red fluorescence shows the mitochondria, blue fluorescence shows the nucleus, and the arrow refers to the nuclear fragmentation and condensation. In the control plasmid pEGFP-N1, the green fluorescence was evenly distributed in the whole cell of FHM cells, mitochondria were annularly distributed in the periphery of nucleus, the edge of nucleus was smooth with uniform dyeing and had no apoptosis characteristics. Subcellular localization of ORF3141-GFP fusion proteins were mainly distributed in the cytoplasm that coincided with the distribution of mitochondria, which indicated that the protein ORF3141 is targeted to mitochondria. Differently, subcellular localization of ORF3141Δsig-GFP fusion proteins were evenly distributed in the whole cell of FHM cells, and were not coincide with the distribution of mitochondria, which indicated that the subcellular localization changed with the excision of signal peptide. The expression of ORF3141 or ORF3141Δsig proteins changed the distribution of mitochondria from perinuclear halo into lumps and apoptotic bodies were detected.

Fig 4. Overexpression of ORF3141 and ORF3141Δsig proteins in FHM cells. The transfected cells were fixed at 48 h post-transfection staining by DAPI. Arrows indicated the apoptotic bodies (fragmented nucleus), arrow heads indicated the apoptotic cells.

Fig 5. Apoptosis nucleus ratio, RT-PCT, and Western blot of pcDNA-3141 and pcDNA-3141Δsig transfected FHM cells. (A) The apoptotic nucleus was counted and the percentages of

apoptotic cells were calculated. Data represent the means for three independent experiments and error bars indicate SD (** $p < 0.01$). (B) Confirmation of the ORF3141 and ORF3141 Δ sig expression in FHM cells by RT-PCR. Lane M, DNA marker; lane 1, pcDNA-3141 Δ sig/FHM; lane 2, pcDNA-3141/FHM. (C) Confirmation of the ORF3141 expression in FHM cells by western blot. Lane M, Protein marker; lane 1, pcDNA-3141/FHM; lane 2, pcDNA 3.1 His/A (Up) and confirmation of the ORF3141 Δ sig expression in FHM cells by western blot. Lane M, protein marker; lane 1, pcDNA 3.1 His/A; lane 2, pcDNA-3141 Δ sig/FHM (Down).

Fig 6. Apoptotic detection of ORF3141 and ORF3141 Δ sig overexpressed FHM cells. (A) Mitochondrial membrane potential assay. FHM cells transfected with pcDNA-3141, pcDNA-3141 Δ sig or pcDNA 3.1 His/A plasmid were collected at indicated time points after transfection and the mitochondrial membrane potential were assessed using the JC-1. Un-transfected cells treated with CCCP was positive control. The data were expressed as the JC-1 polymer/monomer fluorescence ratio and error bars indicate SD (* $p < 0.05$, ** $p < 0.01$). (B) Measurement of caspase-3 activity. FHM cells transfected with pcDNA-3141, pcDNA-3141 Δ sig or pcDNA 3.1 His/A plasmid were collected at indicated time points after transfection and the levels of cleaved caspase-3 were measured. The data were expressed as fold increase compared to the corresponding caspase-3 activity values in un-transfected cells and error bars indicate SD (* $p < 0.05$, ** $p < 0.01$). (C) qRT-PCR analysis of the expression of apoptosis-related genes (*Bad*, *Bid*, *Bax*, and *Bcl-2*) and error bars indicate SD (* $p < 0.05$, ** $p < 0.01$).

A

1 **ATG**AAGCTGCTGAACCCGCGGGGGTTCGGATTGGTTTGCGCATCCGCGGGCCGTGGCGGGCG
 1 V K L L N P R G F G L V C A S A A V A A
 61 GGGTTGATGCTGGCGGGCTGCGCGAACACGGTCGAGGGTACCCCGACGGTCGATCAGGCT
 21 G L M L A G C A N T V E G T P T V D Q A
 121 CAGGTGACCTCGTACCGGGCCGAGGTGTCCTCGTCCGCGGGCGGGCCGCCAGTTCGTCCAAG
 41 Q V T S Y R A E V S S S A A A A S S S K
 181 GCCGCGGCGCAGGTGCGCAAGGCCACGGCCGACAACCTGCGATCCGTTCCGCAAGACCGCC
 61 A A A Q V A K A T A D N C D P F R K T A
 241 bioRxiv preprint doi: <https://doi.org/10.1101/341951>; this version posted June 7, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.
 81 G T A V D R Y N E F V D A H D A S A A D
 301 CAAGTGGCGAAGCGGGACGCGGGCGGGCGGGGCGCTCGAGGACGCGGGCGAAGACGATCGAG
 101 Q V A K R D A A A G A L E D A A K T I E
 361 ACCGAGTTGAACGCGACCCGGGCGGATCTGCCCCGCCGATCTCGCGGGCAAGCTCACCGAC
 121 T E L N A T R A D L P A D L A G K L T D
 421 TATGTGAACGCGGGCGCGCTCGCTGGCCGCGGAGATCCGGAAGATGTCGGGCGGGTCGTCG
 141 Y V N A A R S L A A E I R K M S G G S S
 481 GTGGCGCCGCTGAACGACGCGAGCAAGAAGGTCAACGACGCTCTAACCGCTGTTCGGAAC
 161 V A P L N D A S K K V N D A L T A V R N
 541 GCCTGCCCCGGGCAAG**TGA**
 181 A C P G K *

B

C

A

N. seriolae : -----M¹LLN-----P¹RGF¹CLVC-----ASA¹A¹AAG : 21
N. concava : -----M¹LLN-----P¹RGF¹CLVC-----ASA¹A¹AAG : 21
N. brasili : -----M¹RLD-----P¹RRF¹CLIC-----ATGA¹A¹AVG : 21
M. rhodesia : -----MVTGW-LCP¹R-----R¹LRFAAL-----VVGS¹A¹AAS : 24
M. marinum : -----MLIGWRAVP¹RRHGGA¹FSGS¹RARRGAQ-----ALGC¹A¹AVL : 34
S. piniform : -----M¹TEQR-----Y¹RA¹CLTG-----AMTAAITM : 22
R. globerul : -----MKAGPV¹SGSKSVDRY¹RR¹CLVV-----AIGA¹A¹TVG : 31
R. opcus : -----M¹RTSS-----R¹RW¹CLLS-----AAAAAIVG : 20
G. polyisop : MRRASARRAYRAELRN¹VGGCHSGTEECHQTDADQGRSGVLRVR¹TA¹CGATRVGVTVVLTAVI¹TAA : 66
G. soli : -----MQLGDCAL-----ARDTAPVGGRG-----AVS¹RRGC-----VALA¹A¹VAG : 34

N. seriolae : -----L¹MLAGGANT¹VEG¹IP¹TV¹Q¹AEVTSY¹RAEVSS¹AAA¹SSSKAAAQ-----VAK¹A¹TADN¹CD : 74
N. concava : -----L¹VLAGGSNT¹VEG¹IP¹TV¹Q¹AEVTSY¹KAAAATS¹AAA¹SSAKAAAA-----VAK¹A¹TADN¹CD : 74
N. brasili : -----L¹VVAGCGNRVHG¹IPNP¹TADLAAY¹KTEAAAS¹AAA¹TSSKRAAA-----QTK¹A¹IGDN¹CA : 74
M. rhodesia : -----AM¹TVGGSNVTECKPSV¹DSADAPVY¹RASVSASIQESIASSSARESERQESLTQE¹VHTSCE : 85
M. marinum : -----L¹MG¹IVGGTSVTDG¹IATP¹TKVAPAY¹RSSVSM¹SIS¹SAATSSSVRESQRQQSLTTK¹AIRTSCE : 95
S. piniform : TVAVIA¹IVVAGCGDRV¹DGSARE¹SGEASAY¹SEAAAS¹SS¹AAAAA¹QAAA-----TAK¹A¹KADTCG : 81
R. globerul : -----SV¹V¹IVGGSGAVECSPTP¹NRVQAAAY¹QAEVTAS¹-IA¹SSSKAAAD-----AVETA¹FQTCE : 84
R. opcus : -----AG¹V¹IAGCGGATEG¹IAQQ¹DSQAAEY¹AAEATSS¹SV¹A¹SSSQKAAD-----ERE¹A¹LVEQCT : 74
G. polyisop : -----V¹L¹IFAGCATS¹PCAPT¹V¹SAQVSMY¹SEQSASTS¹AARR-----ELAVT¹LOR : 112
G. soli : -----AV¹IVGGCS-SV¹DGSGTAASGQVAA¹YRAEVTAS¹RAE¹IVQ-----K¹AGTDVCS : 79

bioRxiv preprint doi: <https://doi.org/10.1101/341651>; this version posted June 7, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

N. seriolae : PRRKIAGTAV¹EN¹EV¹DA¹HLASAADQLTKRDA¹ATALEDA¹KGVETQLTATGAA¹PAD¹LAGKLT¹D : 140
N. concava : PFRKSAGNAV¹DR¹NEFV¹DA¹HLASAADQLTKRDA¹ATALEDA¹KGVETQLTATGAA¹PAD¹LAGKFT¹D : 140
N. brasili : QFPTTTGVGVSKYNEF¹IDA¹HLANAGDYAAKREL¹A¹ASTLDDA¹NKVETGISTAKDAMPAD¹LA¹KFT¹D : 140
M. rhodesia : SLSTSSVDA¹ITAVNAYVDA¹FNQSAADAEAKAGP¹IDALNHS¹ADLVG---GSVSDP¹TP¹ELK¹DAMRS : 148
M. marinum : TLATTSKEA¹IDKVNEFVGA¹FNAGRSTGPTEGP¹IEALNNS¹ASSVS---SALNEA¹SP¹DM¹RDAFNA : 157
S. piniform : AFLGRTPDA¹IDA¹NGFVDA¹SNKRADDIAARRSA¹VT¹ELRKS¹DEVDAGVQAAGPP¹DP¹DLAKRFAD : 147
R. globerul : ELIRRSKDDVAV¹FNAYIDA¹NNDES¹SVTSKAQA¹ASSADE¹FG---WLD¹TATAD¹V¹PAS¹LSGLFGD : 146
R. opcus : MFVTRAGET¹IDTYNTFIDA¹NAEAADTGAKASA¹AAALRSA¹ND---GANAATPA¹PP¹DLTGLLT¹D : 136
G. polyisop : QAMSSMVVM¹VRG¹YNTFV¹RLN-AVQSYDRVGD¹LDDRRASLIAGVDQIRAKVTGA¹VTAD¹V¹TGPVNA : 177
G. soli : TSTSAIVVM¹VQGYNTFIRQLN-ATQSYDRIGDSDEKARAGLIAGADQIRGALGDRTP¹T¹VGD¹PART : 144

N. seriolae : VVNAAR¹SLAAE¹IRKMSGGSS¹AP¹INDASKKVNDALTA¹VRNACPGK----- : 185
N. concava : VVTAAAR¹SLAAE¹IRKLSSGAS¹AP¹INDASKKVNDALTA¹VRNACPAK----- : 185
N. brasili : VVNAAR¹SLAETRKM¹TYTAP¹GP¹INDASRRVNDARNA¹VRDACPKR----- : 185
M. rhodesia : VVDAAR¹RLAVA¹IAGNYGPDEFNA¹AITKLNDTKTSALN¹CDAA¹Y----- : 191
M. marinum : VTDAA¹RAVANA¹IGTHAPTGEFNRRVDQLNDTKTKALK¹CMASY----- : 200
S. piniform : VAGAARE¹LATAADQM¹QD-AVQA¹VNQAKERFND¹SLEAV¹RKGCD----- : 188
R. globerul : LADNLR¹SLAGV¹KRDHSPDE¹INS¹ITDTTNSIRDSIRTECGAL----- : 188
R. opcus : VADNYRE¹LAAAVDSGQRGDI¹NT¹ASRGDELSDSIRAACPTS----- : 178
G. polyisop : FLTSSGR¹IGDA¹IG-RRELVG¹NP¹IADTWTRDKQSVLA¹CATYLP¹TPPG¹LAP-----TTTGARPPG : 236
G. soli : FLNTTGE¹LGA¹IG-KRELGG¹NP¹ISDRWTREKNAVLA¹CGKYRPLPPATPSDQGTAPSSGAVPPP : 208

B

pEGFP-3141

pEGFP-3141 Δ sig

pEGFP-N1

GFP

DAPI

Mito Tracker Red

Merge

<https://doi.org/10.1101/341651>bioRxiv preprint doi: <https://doi.org/10.1101/341651>; this version posted May 7, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

CC-BY 4.0 International license

Fluorescent light

White light

pcDNA

pcDNA-3141 Δ sig

pcDNA-3141

A

B

C