Experimental manipulation of nocturnal nest cavity temperature in wild blue tits

Emily G. Simmonds^{1,2}, Ben C. Sheldon¹, Tim Coulson¹, and Ella F. Cole¹*

¹Department of Zoology, University of Oxford, Oxford, UK

²Department of Mathematical Sciences and Centre for Biodiversity Dynamics, Norwegian University of Science and Technology (NTNU), Trondheim, Norway.

* Corresponding author. Any comments or suggestions are much appreciated. emilygsimmonds@gmail.com

Abstract

Advances in the timing of reproduction in temperate species are some of the most well documented biotic responses to increasing global temperatures. However, the magnitude and rate of these advances in timing are not equal across all taxonomic groups. These differences can lead to disruption of interspecific relationships if species respond differently to temperature changes. Understanding the relationship between temperature and phenology is a key step in predicting future population trends for species living in seasonal environments. However, experimentally manipulating temperature in the wild is logistically challenging and has consequently rarely been attempted. In this study we experimentally test whether in-nest temperatures in early spring act as a cue for breeding phenology in a population of wild blue tits (*Cyanistes caeruleus*). We split nests into three treatments; heated, cooled, and control. In-nest temperature in the heated and cooled

25 boxes was manipulated by an average of ± 0.6 °C from control temperatures using heating devices
 26 and ice packs respectively. We assessed the impact of our experimental manipulation on box
 27 occupancy and reproductive timing. We found trends towards earlier phenology in heated nest
 28 boxes in addition to a higher occupancy rate in cooled boxes, however neither of these trends was
 29 found to be statistically significant. Our ability to distinguish statistical signals was hampered by
 30 unexpectedly low occupancy rates across all experimental treatments. Based on the results we
 31 cannot say if nocturnal in-nest temperature is an important cue for nest box choice or the timing of
 32 laying.

33

34 **Keywords:** *blue tits (cyanistes caeruleus), phenology, experimental manipulation, lay date, in-nest*
 35 *temperature, heating and cooling*

Introduction

Advances in phenology, the timing of life history events, have been widely documented amongst temperate species over recent decades, with examples spanning multiple ecosystems and almost all taxonomic groups (Thackeray et al. 2010). These advances have coincided with increases in global temperatures as a result of anthropogenic climate change (Menzel et al. 2006; Parmesan 2006; Singer & Parmesan 2010). Failure to time breeding to match with the peak abundance of resources can result in energetic demands that cannot be met and a consequent reduction in fitness (Reed, Grøtan, et al. 2013; Reed, Jenouvrier, et al. 2013). Therefore understanding how temperature influences breeding phenology is crucial for predicting the impact of climatic change on reproductive success. Despite this importance, the precise role that temperature plays in altering breeding phenology has not yet been determined.

Matching peak energetic demands to a shifting resource peak is particularly challenging for consumer species, which rely on species in lower trophic levels as their resource. The consumer species must respond to temperature in the same way as the resource in order to maintain synchrony, this is challenging for species with temperature independent development, where embryos have a fixed developmental period. In contrast to species with temperature-dependent development, which respond directly to temperature changes (Perrins 1979), temperature-independent species must use predictive cues in order to determine the optimal breeding time (van Noordwijk et al. 1995; Perrins 1979). Not all species respond to the environment in the same way, this can have significant impacts on interspecific interactions (Thackeray et al. 2010; Parmesan 2007; Visser & Holleman 2001; Plard et al. 2014). If interacting species differ in their response to environmental cues or the specific cues they use to time their phenology, they will shift their timing to differing degrees, which can disrupt synchrony between species (Cushing 1969). For instance, if a prey species advances their phenology at a faster rate than a predator species, due to greater responsiveness to an environmental cue this could result in temporal mismatch. Peak abundance of

resources would occur prior to the peak energetic demand, reducing survival and reproductive success of the predator (Reed, Grøtan, et al. 2013). This has been the case for great tits (*Parus major*) in the Netherlands. Great tits in the Hoge Veluwe population have not advanced their phenology as much as resources despite increases in spring temperatures, whereas peak abundance of their prey species, winter moth caterpillars (*Operapthera brumata*) has advanced by over one week in two decades (Visser et al. 1998). This has resulted in fitness reductions for these great tits (Reed, Grøtan, et al. 2013). In order to understand how and where these mismatches may occur, it is important to identify the precise cues that individuals use to time their reproductive behaviour. Only through understanding the mechanisms that control reproductive timing can we predict how timing will change into the future and whether temporal mismatch will be likely.

Attempts to identify the cues that drive reproductive phenology consist of several approaches. The most commonly used approach is to conduct regression based analyses on observational data (van Noordwijk et al. 1995; Charmantier et al. 2008; Menzel et al. 2006; Parmesan & Yohe 2003; Torti & Dunn 2005). This approach typically consists of regressing candidate cues (often temperature) against a phenological event of interest. Either testing different temporal windows or days of temperature (Roberts 2008; Thackeray et al. 2016; van de Pol et al. 2016; Phillimore et al. 2013). The results of studies of this type, have shown that the breeding phenology of northern temperate species is highly correlated with various measures of spring temperature (Visser et al. 1998; van Noordwijk et al. 1995; Charmantier et al. 2008; Cleland et al. 2007; Menzel et al. 2006; Parmesan & Yohe 2003). April temperatures were shown to correlate with clutch initiation date in gulls (Brommer et al. 2008) and temperatures during March and April explained 70 % of the variation in clutch initiation date for great tits and blue tits (*Cyanistes caeruleus*) in the UK (Perrins & McCleery 1989). However, it can be difficult to pinpoint exact temperature cues through these analyses given that temporal windows and the temperature variables tested are all at least partially correlated.

88 Furthermore, regression based studies do not allow the determination of causality, they can only
 89 assess if two variables correlate. In order to establish causal links between environmental cues and
 90 phenological events we must therefore use experimental manipulations are required.
 91
 92 Experimental manipulations of temperature cues pose a logistical challenge, particularly for large
 93 vertebrate species. While sessile species, such as plants, can be manipulated in situ, mobile species
 94 such as most animals, cannot. Animal species tend to cover considerable distances, daily, weekly or
 95 seasonally, experiencing temperature cues across a large spatial and temporal range. Consequently,
 96 manipulating temperature across these scales in the wild is often impossible. As a result, captive
 97 experiments provide one alternative (Schaper et al. 2011; Schaper et al. 2012). These captive
 98 experiments provide a good environment to determine the role of different cues and test causality.
 99 For instance, captive experiments on great tits used temperature controlled aviaries to explore the
 100 influence of temperature on clutch initiation date. This experiment showed that while an increasing
 101 temperature caused clutch initiation date to advance, changes to mean temperature or temperature
 102 variation alone did not impact phenology (Schaper et al. 2012). However, experiments in captivity
 103 do not tell us whether these relationships would hold in the wild. There have been examples of
 104 experimental manipulations failing to capture observed patterns (Lambrechts et al. 1999;
 105 Wolkovich et al. 2012). Captive environments have highly controlled conditions, often
 106 investigating a single focal cue at a time (Schaper et al. 2011; Schaper et al. 2012) and consequently
 107 failing to take account of the influence of other variables. It is unlikely in reality that a single driver
 108 determines all phenological change. Furthermore, captive environments create artificial conditions,
 109 which are potentially stressful for the animals and can lead to adjustment of behaviour away from
 110 what it would be in the wild (Lambrechts et al. 1999).
 111
 112 To identify whether relationships identified during captive experiments or regression analyses also
 113 hold in the wild we need to perform experiments on natural systems. Cavity nesting birds, such as

great tits and blue tits, provide a good study system for such experiments. Reproductive timing plays an important role in reproductive success in these species, as a failure to time the peak energetic requirement of reproduction with the peak availability of food resources can result in reduced reproductive success (Reed, Grøtan, et al. 2013). There have also been many statistical assessments of cues on these populations (Husby et al. 2010; van Noordwijk et al. 1995; Perrins & McCleery 1989; Perrins 1965b) and some captive experiments (Schaper et al. 2012), which can be used to inform field experiments. Additionally nest cavities provide an enclosed environment that can be manipulated. Previous studies that have experimentally heated and cooled nest cavities, have shown that cavity temperature influences nocturnal incubation (Vedder 2012) and the incubation intensity (Ardia et al. 2010) in blue tits and tree swallows (*Tachycineta bicolor*), respectively. In contrast, in spite of observational evidence that suggests nest temperatures influence clutch initiation date (Dhondt & Eyckerman 1979; O'Connor 1978) similar experimental manipulations have been unable to demonstrate this effect. A previous experiment on great tits attempting to manipulate clutch initiation date (Nager & van Noordwijk 1992), through heating and cooling nest boxes in two consecutive years, failed to identify a significant signal of temperature on clutch initiation date. This experiment heated and cooled groups (N = 40 boxes per year) of nest boxes at night by ± 0.75 °C from two to four weeks prior to the median clutch initiation date of each year; however, no difference in phenology was detected between treatments. Here we repeat and extend this study using a population of wild blue tits. We heated and cooled 87 (N = 29 per treatment) nest boxes by an average of ± 0.6 °C from 6 pm to midnight from 17 days before the first egg was laid, and testing whether this manipulation influences nest box choice, commencement of nest building and clutch initiation date. We hypothesise that female blue tits will preferentially choose to nest in warmer nest boxes in order to minimise their nocturnal energy expenditure (Ardia et al. 2009; Vedder 2012; Dhondt & Eyckerman 1979). Furthermore, we predict that the temperature of the nest cavity will also act as a cue for clutch initiation date (Dhondt & Eyckerman 1979; O'Connor 1978),

therefore the warmer nests will also experience the earliest clutch initiation dates and the coolest boxes, the latest clutch initiation date.

Methodology

The Study system

This study was conducted in spring 2015 using a population of wild blue tits in Wytham Woods, UK. The nest box breeding population of blue tits has been studied in detail using standardised procedures (Perrins 1965a; McCleery & Perrins 1985) since 2002. Annually 250-450 pairs of blue tits breed in the woodland (Perrins 1979). 147 blue tit nest boxes were erected across the woodland in 2002, with a further 41 boxes added in 2008. These blue tit boxes have entrance holes with a diameter too small to allow larger species, such as great tits, to enter the boxes. By using these boxes rather than those with larger entrance holes, which are also present in the study area, we were able to exclude great tits from our experimental manipulations. We use 87 of these blue tit nest boxes as the focus of our study (Figure 1). These 87 were chosen because they are located close together in a topographically similar area of the woodland, consequently limiting habitat differences. Proximity of the boxes was essential for this experiment to ensure all experimental treatments could be set running within a two hour window each evening.

Environmental data collection

Altitude data were extracted by Wilkin et al (Wilkin et al. 2006) from an Inverse Distance Weighting interpolation of a 50 m resolution Land Form profile Digital-Terrain-Model data set provided by Ordnance Survey.

Local ambient temperature was collected via a grid of ambient temperature iButtons (DS1923-F5, accurate to ± 0.5 °C, HomeChip Ltd) set to measure absolute temperature every 30 minutes. 200 of these ambient temperature iButtons were distributed in a grid system across Wytham woods with

positions chosen to reflect the density of nest boxes. QGIS (Team 2016) was used to match each experimental nest box to a nearest ambient temperature iButton, therefore allowing the calculation of local ambient temperature for each box. The measure of local ambient temperature used in this study is mean temperature for the experimental period, from 1st April to 6th May or until the first egg is laid, whichever is first.

Long-term temperature data is taken from the Radcliffe Meteorological Station (located 5 km East of Wytham Woods). This station records absolute temperature during the day and calculates daily minimum and maximum temperatures from 1815 to the present day (Radcliffe Meteorological Station n.d.).

Figure 1: Map of position of experimental (filled circles) and non-experimental (open circles) nest boxes in Wytham woods. Experimental treatments are colour coded.

Experimental Manipulation and Phenological Data Collection

Of the 87 experimental boxes, 29 were assigned to each of the three treatment types: heated, cooled, and control. Boxes were assigned to treatments using a stratified sampling technique, taking account of historical box popularity. Historical popularity was determined by calculating the proportion of years in which each box has been occupied since its placement in the woodland. Boxes were then grouped into bands of popularity in increments of 10 %, within each group treatments were assigned randomly using R (Team 2008). This assignment method was used to prevent a clustering of treatments by historical popularity as this metric showed a significant negative correlation with mean clutch initiation date of the nest box (EST = -3.18, SE = 1.46, P = 0.03), with the historically most popular boxes having the earliest clutch initiation dates. This sampling approach also created an even likelihood of box occupancy across treatments, therefore allowing us to explicitly test whether our experimental manipulation altered the popularity of individual boxes. We tested whether other spatial components, such as altitude, influenced the mean clutch initiation date of each box, but this was not statistically significant (EST = 0.02, SE = 0.03, P = 0.52). Consequently, we did not take account of altitude when assigning treatments. This assignment of treatments does mean individuals can move between different treatments if they do not roost in the same nest box every night, however, it also avoids temporal or spatial clustering of treatments and is essential for the exploration of in-nest temperature influence on occupancy.

Manipulations began on the 27th of March 2015, 17 days before the first egg was laid in the experimental boxes. Manipulations occurred every evening, ending when the first egg was laid in a box or on the 6th May 2015 for unoccupied boxes, 30 days after the first egg was laid in the whole woodland. The cut off of 30 days after the first egg is a common measure used to exclude second clutches and repeat clutches which result from early failures (Van Der Jeugd & McCleery 2002).

202 The heating device comprised (see Figure 2 for details of device construction) of a three-volt
 203 filament bulb secured in a mount within a metal topped polystyrene disc (approximately 2.5-3.5 cm
 204 thick), which was placed on top of a plastic ring with a 5 cm gap at the front (also 2.5 cm in height).
 205 The bulb was powered by two rechargeable batteries, size C, located beneath the polystyrene disc
 206 within the plastic ring (see Figure 2). The bulb was covered from above by a 5 cm diameter metal
 207 disc, to hide the light produced. Heating was achieved by the bulb heating the metal disc and
 208 convectional transfer of the heat up into the nest cavity. These heated boxes also contained four,
 209 non-frozen, Thermafreeze ice packs (Thermafreeze EU, black ice packs) in the same configuration
 210 as the cooling device (see below) to ensure that all experimental boxes had the same outward
 211 appearance.

212 **Figure 2: Photograph of experimental set up for a heated box.** Key components of the set up are
 213 numbered 1) Ice packs mounted front and back of the box, 2) battery pack, 3) polystyrene insert, 4) plastic
 214 ring to hold polystyrene above the battery pack, 5) metal disc topping the bulb. In control and cooled boxes,
 215 the battery pack would be replaced by a second polystyrene insert.

216
 217 The cooling device comprised two frozen Thermafreeze ice packs secured at the back of each nest
 218 box and two on the nest box door. Ice packs at the back of the box were secured using the same
 219 metal topped polystyrene disc as used in the heating device, with another polystyrene disc below
 220 (around 2.5 cm thick) to achieve the same thickness as the heating device but without the plastic
 221 ring and batteries. Ice packs on the nest box door were secured using rubber bands. All ice packs
 222 had a black outer side to camouflage them from the birds once the nest box was closed.
 223
 224 The control treatments comprised the same construction as the cooled treatment but with non-frozen
 225 ice packs. Therefore, all boxes had the same appearance, a raised metallic box floor of
 226 approximately 5 cm above the true base, with ice packs mounted front and back.
 227
 228 Both the batteries and frozen ice packs had an effective life of around six hours and therefore were
 229 replaced each evening around 2.5 to 1.5 hours prior to sunset so that the treatments were running
 230 during the night. This period of the day was chosen as female blue tits usually roost overnight
 231 within their nest cavity prior to nest building and therefore should be present within the box to
 232 experience the temperature manipulation (Kluyver 1950; Perrins 1965a; Kidd et al. 2015).
 233 Temperatures were altered on average by ± 0.6 °C from the control boxes from sunset (around 6
 234 pm) to midnight. Batteries and ice packs were replaced each evening; control boxes were also
 235 visited to achieve a uniform disturbance level across boxes. Some damage to equipment did occur
 236 (pecks to polystyrene inserts) and equipment was either repaired or replaced to ensure the
 237 treatments could continue. Experimental treatments could not be changed on two nights during the
 238 experiment (29th March 2015 and 31st March 2015) due to high winds. At the same time that
 239 treatments were changed several aspects of breeding phenology were recorded; signs of nest
 240 building (moss in box), signs of roosting (feathers and faeces), number of eggs, and whether the
 241 eggs were being incubated. From this we could calculate the nest building date (the date when moss

first appeared in the nest box), clutch initiation date (the date on which the first egg was laid in a nest box) and clutch completion date (the date when the last egg was laid in a nest box).

In-nest temperatures were recorded for the entire duration of the experiment using iButton thermometers (DS1921G-F5, accurate to ± 1 °C, HomeChip Ltd) set to record temperature every 30 minutes. The iButtons were secured to the nest box wall using duct tape approximately 5 cm above the heating or cooling devices and half way between the front ice packs and the back ice packs. The effectiveness of the heating and cooling treatments during the experiment was assessed by calculating the mean temperature difference between treatments during the experimental periods (6 pm to 11:59 pm). The nights of the 29th and 31st March were excluded from these calculations because the experimental treatments did not run on those nights. Occupied nests were also excluded from the calculations after the first egg was laid, as the experimental treatments were halted at the appearance of the first egg. This type of experimental design was based on those previously employed on other nest box breeding passerine populations (Ardia et al. 2009; Pérez et al. 2008; YomTov & Wright 1993; Bryan & Bryant 1999; Vedder 2012; Alvarez & Barba 2014; Nager & van Noordwijk 1992). Previous manipulations have taken the form of heating and cooling usually during incubation (after laying has commenced) with heating devices attached to the outside of the nest box (Ardia et al. 2009; Pérez et al. 2008; YomTov & Wright 1993; Bryan & Bryant 1999) or within the nest box, below the nest itself (Vedder 2012; Alvarez & Barba 2014). Nager and Van Noordwijk (Nager & van Noordwijk 1992) heated and cooled nest boxes prior to laying through ice packs and heating devices located at the back of the nest box.

Statistical Analyses

A power analysis was conducted prior to starting the experiment to determine whether differences were likely to be detected between treatments given the number of nest boxes available. Based on the average historical occupancy of the study boxes (67 %) we would expect an occupancy rate of

19 boxes per treatment. With this number the power analysis showed that we could expect a 75 % chance of detecting a mid to strong signal (an effect size of 0.4 or higher). We would expect a mid to high effect size based on previous work that has found a strong correlation between temperature and clutch initiation date (Perrins 1970; Perrins 1973). Furthermore, analyses of the relationship between spring temperature (mean of daily minimum and maximum temperatures for March and April) and clutch initiation date from 1960 to 2011 showed close to a 6 day delay in clutch initiation date for every 1 °C decline in mean temperature (EST = -5.71, SE = 0.08, P = <2e-16), therefore we would expect a strong signal from our experimental manipulation.

We performed statistical analyses to determine the influence of our experimental manipulations on breeding phenology and box occupancy. For the analyses of the influence of experimental treatments on box occupancy, we first tested if there was a difference in the number of boxes occupied between treatments. A chi-squared test of association was performed to determine if the number of occupied boxes differed significantly between treatments. It is also important to consider whether any other environmental factors could be influencing occupancy. Philopatry could have a role here with individuals or pairs returning to the same boxes every year. Across the length of this study 12 % of individuals (8 % of breeding attempts) returned to the same box they had previously bred in, suggesting that for the majority of individuals previous location is not the only driver of nest box choice. Therefore we used one way ANOVAs and a Welch's T-Test to explore the difference in historical popularity, altitude and ambient temperature between occupied and unoccupied boxes. A Welch's T-Test was used for local ambient temperature due to heteroscedasticity of temperature data. The whole dataset was used for these analyses as there was not enough data to split by treatment.

Analyses of the influence of experimental treatments on breeding phenology were performed using one way ANOVAs to determine if there are differences between phenology measures in each

treatment group. The phenology measures considered here were the date of nest building (the date on which moss was first observed in the nest box), the clutch initiation date (date when the first egg is laid), and date of clutch completion (date when the last egg in the clutch is laid). Regression analyses were also performed to assess the influence of environmental variables on clutch initiation date. Environmental variables tested were altitude and local ambient mean temperature in a single linear regression to assess the combined influence of these variables on variance in phenology.

Ethics statement

Work was subject to review by the Department of Zoology ethical committee, University of Oxford. All work adhered to UK standard requirements and was carried out under Natural England licence 20114732. This experimental methodology was discussed with, and approved by the Departmental Animal Welfare Ethical Review Body (AWERB). Field work took place in Wytham Woods (lat. 51°46'N, long. 1°20'W), private land that belongs to the University of Oxford; for permission contact the Conservator, Nigel Fisher. No endangered or protected species were involved in the study. A breakdown of nest building, abandonment and chick mortality by treatment can be found in supporting information Table S1, however these rates did not differ from those in non-experimental boxes.

Results

The influence of experimental treatment on occupancy

Occupancy in our experimental boxes was 16 %, which was a decline of 73 % from the preceding year (2014). Adjacent non-experimental boxes had a blue tit occupancy of 14 % (Table 1), a decrease of 48 % from 2014 occupancy. The number of great tits in these boxes also declined from 2014 to 2015, and to a greater extent, by 54 % (35 occupied in 2014 and 19 in 2015). Occupied boxes had higher historical popularity than unoccupied boxes ($F(1,82) = 6.15$, $P = 0.02$, see Figure 3), and this pattern was consistent across treatments. It should be noted that three of the occupied

experimental boxes (N = 17) were occupied by marsh tits (*Poecile palustris*). As these are a different species to our study species, this could influence some of our results. Therefore we removed these nests from our analyses, leaving a sample size of N = 14.

Table 1: Numbers of occupied and empty boxes by treatment

* Adjacent non-experimental boxes were defined as those boxes within 100 m of an experimental box. These boxes are those with the larger nest box entrance holes and therefore could be occupied by either great tits or blue tits. However we only include blue tits in this analysis. Great tit occupancy of the adjacent non-experimental boxes was 19/95, 20%

Treatment	Occupied boxes	Empty boxes (not occupied by blue tits)	Total
Cooled	7 (24%)	22	29
Control	5 (17%)	24	29
Heated	2 (7%)	27	29
Adjacent non-experimental *	13 (14%)	82	95

Table 2: Numbers of blue tit occupied nest boxes 2014 and 2015

* The whole woodland refers to all boxes in the entirety of Wytham woods (see Fig. 1, N = 1203). This includes all great tit (larger entrance hole, N = 1016) and blue tit only (smaller entrance hole, N = 187) boxes and covers all sections of the woodland. Occupancy numbers are for nests confirmed as blue tits, so including a minimum of 3 eggs. Adjacent non-experimental boxes are defined in using the same criteria as Table 1.

**Excluding marsh tits

	Boxes occupied 2014	Boxes occupied 2015 (year of experiment)	Proportional difference
Whole woodland *	430	258	- 40%
Non-experimental boxes	378	244	- 35%
Experimental boxes	52	14**	- 73%
Adjacent non- experimental boxes	27	13	- 48%

334 *Does in-nest temperature influence nest box choice?*

335 The number of nest boxes occupied during our experimental period varied between treatments
 336 (Table 1). Cooled boxes had the highest proportion (24 %) of boxes occupied, followed by control
 337 boxes (17 %) and heated boxes (7 %) have the lowest levels of occupancy. These differences were
 338 not statistically significant ($\chi^2 = 3.97$, $DF = 2$, $P = 0.14$). We explored additional factors that may
 339 explain variation in occupancy. Neither altitude nor local mean ambient temperature differed
 340 significantly between occupied and unoccupied boxes (Altitude results, $F(1,82) = 0.98$, $P = 0.33$,
 341 local mean ambient temperature results - $T = -1.00$, $P = 0.33$, $DF = 13.74$, see supporting
 342 information, 0).

343
 344 **Figure 3: Boxplots of the historical popularity of occupied and unoccupied boxes by treatment.** Raw
 345 data are plotted as points and the box plots show the median and interquartile and full range for each
 346 treatment. Historical popularity represents the proportion of years a box was occupied since its placement in
 347 the woodland.

348

Does in-nest temperature influence breeding phenology?

For these analyses $N = 13$ (two heated boxes, five control boxes and six cooled boxes), one nest was removed from analyses due to lack of a clutch initiation date as laying occurred after the conclusion of experimental treatments. On average, birds nesting in heated boxes started nest building, laying and completed their clutches earlier than birds nesting in the control or cooled boxes (Figure 4), however none of these differences were statistically significant (nest building $F(2,11) = 1.20$, $P = 0.34$, clutch initiation date $F(2,11) = 0.91$, $P = 0.43$, clutch completion date $F(2,11) = 0.91$, $P = 0.43$) (for full ANOVA tables see supporting information, 0). The influence of environmental variables, altitude (EST = -0.15, SE = 0.08, $P = 0.07$) and local ambient temperature (EST = -2.52, SE = 4.67, $P = 0.60$), were also tested but showed no statistically significant relationship with clutch initiation date, the focal phenological variable here.

Figure 4: Boxplots of the dates of a) nest building, b) clutch initiation and c) clutch completion by treatment. Raw data are plotted as points and the box plots show the median and interquartile range for each treatment. Dates are all plotted in April days (April 1st = 1)

Discussion

Experimental manipulation corresponded with low occupancy of nest boxes

This study experimentally tested whether nocturnal in-nest temperatures act as a cue for nest box choice and clutch initiation date in a population of wild blue tits. The experimental manipulation carried out in this study altered the minimum daily temperature experienced by individuals, either elevating it or reducing it dependent on treatment group. One of the defining findings of this study is a substantial and unforeseen decline in the occupancy of our experimental boxes to 16 %. This is considerably below the expected occupancy levels, which since 2008 have averaged 67 %. This 67 % occupancy would have given us approximately 19 boxes per treatment, which according to the preliminary power analysis conducted, would have been sufficient to detect the mid to high signal we expected (effect size of 0.4 or greater). However, due to circumstances beyond our control, this sample size was not achieved. The potential to increase our sample size was limited by the availability of blue tit only nest boxes. While more blue tit boxes are available in the woodland the remainder are located either in different habitat types or spatially distant from the study boxes. The inclusion of these other boxes would have introduced confounding factors and been logistically difficult to manage. Changes of equipment had to be achieved within a two hour window each evening. However, with the sample size available, we should have been able to detect a signal if one had been present and box occupancy was at the expected levels.

The low occupancy levels achieved in this experiment have resulted from several sources. Firstly, the number of nesting blue tit pairs declined across the whole of Wytham Woods in 2015, reducing by around 40 % (Table 2). Secondly, it was also clear that the parent blue tits found the disturbance and/or altering of the box appearance a deterrent from nesting in these boxes. Evidence that the birds were disturbed by the presence of the equipment was the pecking of polystyrene mounts in many of the boxes. This demonstrates the birds were not deterred from entering the box but also

appeared to want to remove the heating and cooling devices. Evidence that birds entered the experimental boxes was also found in the form of signs of roost (faeces and feathers) found in several of the experimental boxes. However this use of boxes did not translate into nest building. Smaller nest cavities have been linked to higher nestling mortality (Mertens 1977) and could therefore deter parent birds. Additionally, the blue tit occupancy of the experimental nest boxes declined by a larger proportion than the non-experimental boxes (73 % and 48 % respectively, see Table 2). However, this cannot definitively be attributed to our experimental treatment because declines in both blue tit and great tit numbers were seen throughout the whole of Wytham Woods from 2014 to 2015. The lack of increased blue tit occupancy in the surrounding non-experimental boxes, suggests that birds deterred from the experimental boxes did not relocate to nearby unaltered boxes. However, disturbance did not seem to influence marsh tits to the same degree, their numbers remained at a similar level of occupancy to historical records, suggesting they were less deterred by the experimental set up than blue tits. Further refinements of techniques to minimise the amount of equipment required to heat and cool boxes would improve these experimental techniques. Caution should be employed for future blue tit nest box manipulations.

The impact of low sample size on statistical analyses

The sample size achieved in this study was exceptionally low, this heavily shaped our ability to draw statistically supported conclusions from this experimental work. With a sample size of 13 across all three treatments (two heated, five control, and six cooled), very little could be ascertained statistically from the resulting dataset. As analyses were planned prior to running of the experiment, they were duly conducted. However, even if statistically significant results are found, we cannot confidently reject a null hypothesis. The inability to determine statistical results stems from fundamental issues with multiple testing and p-values.

414 The rise of inappropriate and misleading statistical analyses is a growing concern across many
 415 scientific disciplines, in particular psychology, medicine, ecology, and evolution (Mcshane et al.
 416 2017; Horton 2015; Parker et al. 2016; Forstmeier et al. 2016). It is something we must consider
 417 when analysing the data generated in this study and interpreting the results. A focus and reliance on
 418 p-values and null hypothesis testing, passing of a 0.05 threshold indicating a rejection of the null
 419 hypothesis, can be especially problematic for multiple comparisons on small datasets (Gelman &
 420 Loken 2014). When we accept a 5 % change of a Type 1 error (false positive result) over the course
 421 of multiple analyses, either on the same dataset or across studies, then 5 % of statistically
 422 significant results will actually have arisen if the null hypothesis were true (Forstmeier et al. 2016;
 423 Nuzzo 2014). Whether a result is deemed statistically significant or not is particularly changeable
 424 for small datasets where errors are high relative to the signal and the influence of individual data
 425 points is strong. Given this, there have been several calls to abandon p-values in statistics (Mcshane
 426 et al. 2017; Good 1988; Cumming 2014), although this call is not universal (Savalei & Dunn 2015).
 427
 428 Given the small sample size here, and the multiple comparisons conducted, the results should be
 429 analysed with caution. Multiple comparisons had to be conducted as there were multiple response
 430 variables to a focal explanatory variable (treatment), which were all chosen *a priori*. As a result, we
 431 move away from discussion of statistical significance in this study, as it cannot be relied upon in
 432 this instance (Mcshane et al. 2017). Instead we focus on the effect size of any trends found.

433 ***Trends found***

434 Trends shown in this analysis are that historical occupancy levels (popularity) also appear to predict
 435 occupancy during the experimental period. Boxes occupied during our experiment were occupied
 436 for, on average, 15.6 % more of the historical time period than the unoccupied boxes. It seems
 437 logical that the historically most popular boxes are also those occupied most readily during the
 438 experimental period. Neither altitude nor local mean temperature appear to explain popularity,

439 either historically or during this experiment. Further analyses of different box positions would be
 440 required, manipulating specific aspects of the box environment, for example, aspect or oak density,
 441 to determine the drivers of occupancy/popularity of nesting sites. Although cooled boxes had the
 442 highest number of occupied boxes, this was statistically indistinguishable from control or heated.
 443 The difference between treatments was a maximum of four boxes. Past studies have found a slight
 444 preference for cooler nest boxes (Dhondt & Eyckerman 1979), however, this could also be a chance
 445 difference, substantially more analyses would be required to draw definitive conclusions.
 446
 447 No influence was found of in-nest temperature on breeding phenology. There was some overlap
 448 between the cooled and control groups for all measures of phenology, however, mean differences
 449 had effect sizes of several days. On average heated boxes initiated nest building and egg laying
 450 approximately three to four days prior to control boxes and six to four days prior to cooled boxes.
 451 The magnitude of these effect sizes from $< 1^{\circ}\text{C}$ change in in-nest temperature, predominantly driven
 452 by the heated treatment, suggest we cannot rule out an influence on nest box temperature on
 453 breeding phenology even if it could not be statistically supported in this study. No influence was
 454 found of environmental variables on clutch initiation dates. As ambient temperature typically
 455 correlates strongly with clutch initiation dates across our long-term data, we would expect that in
 456 this case the sample size was too low to distinguish a signal or the influence of ambient temperature
 457 was dampened by the experimental manipulation. However, from the current data, we cannot
 458 discriminate between the two possibilities. We were unable to find statistical support for the
 459 relationships observed by Dhondt and Eyckerman (Dhondt & Eyckerman 1979) and O'Connor
 460 (O'Connor 1978), similarly to Nager and Van Noordwijk (Nager & van Noordwijk 1992) we found
 461 little effect of in-nest manipulations on clutch initiation date. Due to the difficulty of wild
 462 experimental analyses and the low sample size achieved here, this result does not necessarily
 463 indicate a statistical null result. We would strongly encourage further replicated analyses of this
 464 type with higher sample sizes and across multiple years, before a robust conclusion can be reached.

Acknowledgements

This study would not have been possible without the hard work of Tony Price and John Hogg, who assisted in the design and manufactured the heating and cooling devices for this experimental work. We would like to extend our gratitude to both Tony and John for all of the time they put in to this project. We are also grateful to all of the Wytham fieldworkers who collected population census data on the Wytham great tits. In particular, we would like to give a special mention to Keith McMahon, Stephen Lang, Koosje Lamers and Nico Alioravainen, who assisted with the changing of batteries and ice packs for this experiment come rain or shine. This work was supported by NERC grant NE/K006274/1 to Ben Sheldon.

References

- Alvarez, E. & Barba, E., 2014. Behavioural responses of great tits to experimental manipulation of nest temperature during incubation. *Ornis Fennica*, 91(4), pp.220–230.
- Ardia, D.R. et al., 2009. Temperature and life history: Experimental heating leads female tree swallows to modulate egg temperature and incubation behaviour. *Journal of Animal Ecology*, 78(1), pp.4–13.
- Ardia, D.R., Pérez, J.H. & Clotfelter, E.D., 2010. Experimental cooling during incubation leads to reduced innate immunity and body condition in nestling tree swallows. *Proceedings. Biological sciences / The Royal Society*, 277, pp.1881–1888.
- Brommer, J.E., Rattiste, K. & Wilson, A.J., 2008. Exploring plasticity in the wild : laying date – temperature reaction norms in the common gull *Larus canus*. *Proceedings of the Royal Society B: Biological Sciences*, (January), pp.687–693.
- Bryan, S.M. & Bryant, D.M., 1999. Heating Nest-Boxes Reveals an Energetic Constraint on Incubation Behaviour in Great Tits, *Parus major*. , 266(1415), pp.157–162.
- Charmantier, A. et al., 2008. Adaptive Phenotypic Plasticity in Response to Climate Change in a Wild Bird Population. *Science*, 320(5877), pp.800–803.

- 490 Cleland, E.E. et al., 2007. Shifting plant phenology in response to global change. *Trends in ecology*
491 *& evolution*, 22(7), pp.357–65.
- 492 Cumming, G., 2014. The new statistics: Why and how. *Psychological Science*, 25(1), pp.7–29.
- 493 Cushing, D.H., 1969. The Regularity of the Spawning Season of some Fishes. *Journal du Conseil*
494 *international pour l'Exploration de la Mer*, 33(1), pp.81–97.
- 495 Dhondt, A.A. & Eyckerman, R., 1979. Temperature and Date of laying by tits parus spp. *Ibis*,
496 121(3), pp.329–331.
- 497 Forstmeier, W., Wagenmakers, E.J. & Parker, T.H., 2016. Detecting and avoiding likely false-
498 positive findings - a practical guide. *Biological Reviews*, pp.000–000.
- 499 Gelman, A. & Loken, E., 2014. Data-dependent analysis—a “garden of forking paths”— explains
500 why many statistically significant comparisons don’t hold up. *American Scientist*, 102(40).
501 Available at: <https://www.americanscientist.org/article/the-statistical-crisis-in-science>.
- 502 Good, I.J., 1988. The Interface Between Statistics and the Philosophy of Science. *Statistical*
503 *Science*, 3(4), pp.386–412.
- 504 Horton, R., 2015. Offline: What is medicine’s 5 sigma? *The Lancet*, 385(9976), p.1380.
- 505 Husby, A. et al., 2010. Contrasting patterns of phenotypic plasticity in reproductive traits in two
506 great tit (*Parus major*) populations. *Evolution*, 64(8), pp.2221–2237.
- 507 Van Der Jeugd, H.P. & McCleery, R.H., 2002. Effects of spatial autocorrelation, natal philopatry
508 and phenotypic plasticity on the heritability of laying date. *Journal of Evolutionary Biology*,
509 15(3), pp.380–387.
- 510 Kidd, L.R. et al., 2015. Who escapes detection? Quantifying the causes and consequences of
511 sampling biases in a long-term field study. *Journal of Animal Ecology*, 84(6), pp.1520–1529.
- 512 Kluijver, H.N., 1950. Daily routines of the Great Tit, *Parus m. major* L. *Ardea*, 38(December),
513 pp.99–135.
- 514 Lambrechts, M.M. et al., 1999. Do experiments with captive non-domesticated animals make sense
515 without population ® eld studies ? A case study with blue tits ’ breeding time. *Proceedings of*

- 516 *the Royal Society B: Biological Sciences*, 266, pp.1311–1315.
- 517 McCleery, R.H. & Perrins, C.M., 1985. Behavioural Ecology. Ecological Consequences of
- 518 Adaptive Behaviour. In Oxford: Blackwell Scientific Publications, pp. 353–373.
- 519 Mcshane, B.B. et al., 2017. Abandon Statistical Significance. *eprint arXiv:1709.07588*, pp.1–12.
- 520 Menzel, A. et al., 2006. European phenological response to climate change matches the warming
- 521 pattern. *Global Change Biology*, 12(10), pp.1969–1976.
- 522 Mertens, J.A.L., 1977. Thermal conditions for successful breeding in Great Tits (*Parus major* L.) - I.
- 523 Relation of growth and development of temperature regulation in nestling great tits.
- 524 *Oecologia*, 28(1), pp.1–29.
- 525 Nager, R.G. & van Noordwijk, A.J., 1992. Energetic Limitation in the Egg-laying Period of Great
- 526 Tits. *Proceedings. Biological sciences / The Royal Society*, 249, pp.259–264.
- 527 van Noordwijk, A.J., McCleery, R.H. & Perrins, C.M., 1995. Selection for the Timing of Great Tit
- 528 Breeding in Relation to Caterpillar Growth and Temperature. *Journal of Animal Ecology*,
- 529 64(4), pp.451–458.
- 530 Nuzzo, R., 2014. Scientific method: Statistical errors. *Nature*, 506(7487), pp.150–152.
- 531 O'Connor, R.J., 1978. Nest-Box Insulation and the Timing of Laying in the Wytham Woods
- 532 Population of Great Tits *Parus Major*. *Ibis*, (120), pp.534–537.
- 533 Parker, T.H. et al., 2016. Transparency in Ecology and Evolution: Real Problems, Real Solutions.
- 534 *Trends in Ecology and Evolution*, 31(9), pp.711–719.
- 535 Parmesan, C., 2006. Ecological and Evolutionary Responses to Recent Climate Change. *Annual*
- 536 *Review of Ecology, Evolution and Systematics*, 37, pp.637–669.
- 537 Parmesan, C., 2007. Influences of species, latitudes and methodologies on estimates of phenological
- 538 response to global warming. *Global Change Biology*, 13(9), pp.1860–1872.
- 539 Parmesan, C. & Yohe, G., 2003. A globally coherent fingerprint of climate change impacts across
- 540 natural systems. *Nature*, 399, pp.579–583.
- 541 Pérez, J.H. et al., 2008. Experimental heating reveals nest temperature affects nestling condition in

- 542 tree swallows (*Tachycineta bicolor*). *Biology letters*, 4(5), pp.468–471.
- 543 Perrins, C.M., 1979. *British Tits*, London: Collins.
- 544 Perrins, C.M., 1965a. Fluctuations and Clutch-Size in the Great Tit, *Parus major* L. *Journal of*
- 545 *animal ecology*, 34(3), pp.601–647.
- 546 Perrins, C.M., 1965b. Population Fluctuations and Clutch-Size in the Great Tit. *Journal of Animal*
- 547 *Ecology*, 34(3), pp.601–647.
- 548 Perrins, C.M., 1973. Some effects of temperature on breeding in the great tit and manx shearwater.
- 549 *Journal of Reproduction and Fertility*, 19, pp.163–173.
- 550 Perrins, C.M., 1970. The timing of birds’ breeding season. *Ibis*, 112, pp.242–255.
- 551 Perrins, C.M. & McCleery, R.H., 1989. Laying Dates and Clutch Size in the Great Tit. *The Wilson*
- 552 *Journal of Ornithology*, 101(2), pp.236–253.
- 553 Phillimore, A.B. et al., 2013. Inferring local processes from macro-scale phenological pattern: A
- 554 comparison of two methods. *Journal of Ecology*, 101(3), pp.774–783.
- 555 Plard, F. et al., 2014. Mismatch between Birth Dates and Vegetation Phenology Slows the
- 556 Demography of Roe Deer. *PLoS Biology*, 12(4), pp.1–8.
- 557 van de Pol, M. et al., 2016. Identifying the best climatic predictors in ecology and evolution.
- 558 *Methods in Ecology and Evolution*, 7(10), pp.1246–1257.
- 559 Radcliffe Meteorological Station, Meteorological data.
- 560 Reed, T., Grøtan, V., et al., 2013. Population growth in a wild bird is buffered against phenological
- 561 mismatch. *Science*, 340, pp.488–491.
- 562 Reed, T., Jenouvrier, S. & Visser, M.E., 2013. Phenological mismatch strongly affects individual
- 563 fitness but not population demography in a woodland passerine. *Journal of Animal Ecology*,
- 564 82(1), pp.131–144.
- 565 Roberts, A.M.I., 2008. Exploring relationships between phenological and weather data using
- 566 smoothing. *International Journal of Biometeorology*, 52(6), pp.463–470.
- 567 Savalei, V. & Dunn, E., 2015. Is the call to abandon p-values the red herring of the replicability

568 crisis? *Frontiers in Psychology*, 6(MAR).

569 Schaper, S. V. et al., 2012. Increasing temperature, not mean temperature, is a cue for avian timing
570 of reproduction. *The American naturalist*, 179(2), pp.E55-69.

571 Schaper, S. V. et al., 2011. Spring phenology does not affect timing of reproduction in the great tit
572 (Parus major). *The Journal of experimental biology*, 214(Pt 21), pp.3664–71.

573 Singer, M.C. & Parmesan, C., 2010. Phenological asynchrony between herbivorous insects and
574 their hosts: signal of climate change or pre-existing adaptive strategy? *Philosophical
575 Transactions of the Royal Society B: Biological Sciences*, 365(1555), pp.3161–3176.

576 Team, Q.D., 2016. QGIS Geographic Information System.

577 Team, R.D.C., 2008. R: A language and environment for statistical computing.

578 Thackeray, S.J. et al., 2016. Phenological sensitivity to climate across taxa and trophic levels.
579 *Nature*, 535(7611), pp.241–245.

580 Thackeray, S.J., Sparks, T.H. & Frederiksen, M. et al., 2010. Trophic level asynchrony in rates of
581 phenological change for marine, freshwater and terrestrial environments. *Global Change
582 Biology*, 16(12), pp.3304–3313.

583 Torti, V.M. & Dunn, P.O., 2005. Variable effects of climate change on six species of North
584 American birds. *Oecologia*, 145, pp.486–495.

585 Vedder, O., 2012. Individual birds advance offspring hatching in response to increased temperature
586 after the start of laying. *Oecologia*, 170(3), pp.619–628.

587 Visser, M.E. et al., 1998. Warmer springs lead to mistimed reproduction in great tits (Parus major
588). *Proceedings. Biological sciences / The Royal Society*, 265(May), pp.1867–1870.

589 Visser, M.E. & Holleman, L.J.M., 2001. Warmer springs disrupt the synchrony of oak and winter
590 moth phenology. *Proceedings. Biological sciences / The Royal Society*, 268(1464), pp.289–94.

591 Wilkin, T.A. et al., 2006. Density effects on life-history traits in a wild population of the great tit
592 Parus major: analyses of long-term data with GIS techniques. *Journal of Animal Ecology*,
593 75(2), pp.604–615.

- 594 Wolkovich, E.M. et al., 2012. Warming experiments underpredict plant phenological responses to
 595 climate change. *Nature*, 485(7399), pp.494–497.
- 596 YomTov, Y. & Wright, J., 1993. Effect of heating nest boxes on egg laying in the blue tit (*Parus*
 597 *caeruleus*). *Auk*, 110(1), pp.95–99.

598

Supporting information

Table of breakdown of nest building, abandonment and chick mortality by

treatment

Table S1: Table of breakdown of nest building, abandonment and chick mortality by treatment

* These figures include great tit and marsh tit nesting attempts. Species identification can only occur after 3 eggs have been laid so prior to this point species distinction is not possible. All other figures are blue tit only.

	Nest building	Abandonment pre-laying	Abandonment post-laying	Chick mortality
Cooled	7/27 (26%)	1/7 (14%)	0	2/61 (3%)
Control	5/27 (19%)	0	0	4/37 (11%)
Heated	2/27 (7%)	0	0	0
Non-experimental	43/50* (86%)	9/43* (21%)	0	17/113 (15%)

Supporting tables: Does in-nest temperature influence nest box choice?

Table S2: Full ANOVA table from analysis of occupancy as a function of historical popularity

Response = historical popularity					
	DF	Sum of squares	Mean squared	F value	P
Occupied	1	0.28	0.28	6.15	0.02
Residuals	82	3.78	0.05		

Table S3: Full ANOVA table from analysis of occupancy as a function of altitude

Response = altitude					
	DF	Sum of squares	Mean squared	F value	P
Occupied	1	125.9	125.92	0.98	0.33
Residuals	82	10533.5	128.46		

610 **Table S4: Full table of Welch's T-Test of occupancy as a function of mean temperature**

Response = mean temperature	
T	-1
DF	13.74
P	0.33

611

612 ***Supporting tables: Does in-nest temperature influence breeding phenology?***

613 **Table S5: Full ANOVA table of nest building date as a function of treatment**

Response = Nest building date					
	DF	Sum of squares	Mean squared	F value	P
Treatment	2	71.63	35.81	1.19	0.34
Residuals	11	331.3	30.12		

614

615 **Table S6: Full ANOVA table of lay date as a function of treatment**

Response = Lay date					
	DF	Sum of squares	Mean squared	F value	P
Treatment	2	22.09	11.05	0.91	0.43
Residuals	11	120.83	12.08		

616

617 **Table S7: Full ANOVA table of clutch completion date as a function of treatment**

Response = Clutch completion date					
	DF	Sum of squares	Mean squared	F value	P
Treatment	2	9.097	4.55	0.91	0.43
Residuals	11	50.13	5.01		

618

619 **Table S8: Table of outputs from phenology linear regression**

	Estimate	SE	P
Intercept	8.81	44.76	0.85
Local mean temperature	2.52	4.67	0.60
Altitude	-0.15	0.08	0.07

620