1 **Genetic Diversity Study of *Fusarium culmorum*: Causal agent of wheat crown rot in**
2 **Iraq**

3 O. N. Matny¹, S. A. Shamsallah², M. Haas³

4 ^{1,2} Department of Plant Protection, College of Agriculture, University of Baghdad, Iraq

5 email: schamsalah2@yahoo.com². ³ University of Minnesota, Department of Plant

6 Pathology, USA, (current affiliation) Domestication Genomics, Leibniz Institute of Plant

7 Genetics and Crop Research (IPK), Corrensstrasse 3, 06466 Gatersleben, Germany.

8 Email: haas@ipk-gatersleben.de³.

9 **Corresponding Author:** odaimatny@coagri.uobaghdad.edu.iq¹, onmatny@umn.edu

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

ABSTRACT

Fusarium crown rot (FCR), caused by *Fusarium culmorum* (Wm.G.Sm) Sacc., is an important disease of wheat both in Iraq and other regions of wheat production worldwide. Changes in environmental conditions and cultural practices such as crop rotation generate stress on pathogen populations leading to the evolution of new strains that can tolerate more stressful environments. This study aims to investigate the genetic diversity among isolates of *F. culmorum* in Iraq. Twenty-nine samples were collected from different regions of wheat cultivation in Iraq to investigate the pathogenicity and genetic diversity of *F. culmorum* using the REP-PCR technique. Among the twenty-nine isolates of *F. culmorum* examined for pathogenicity, 96% were pathogenic to wheat at the seedling stage. The most aggressive isolate, from Baghdad, was IF 0021 at 0.890 on the FCR severity index. Three primer sets were used to assess the genotypic diversity via REP, ERIC and BOX elements. The amplicon sizes ranged from 200-800 bp for BOX-ERIC2, 110-1100 bp for ERIC-ERIC2 and 200-1300 bp for REP. In total, 410 markers were polymorphic, including 106 for BOX, 175 for ERIC and 129 for the REP. Genetic similarity was calculated by comparing markers according to minimum variance (Squared Euclidean). Clustering analysis generated two major groups, group 1 with two subgroup 1a and 1b with 5 and 12 isolates respectively, and group 2 with two subgroups 2a and 2b with 3 and 9 isolates respectively. This is the first study in this field that has been reported in Iraq.

Keyword: Genetic Diversity, *Fusarium*, Pathogenicity, *Triticum aestivum*, REP-PCR.

INTRODUCTION

Fusarium culmorum (Wm.G.Sm) Sacc., a fungal plant pathogen with a wide host range and is the causal agent of several diseases on these plants. On wheat, *F. culmorum* causes two important diseases that can cause serious economic losses on wheat: head blight and crown rot (Burgess et al. 2001; Chakraborty et al. 2006). Reliable estimates for yield loss due to *Fusarium* crown rot (FCR) in Iraq are not available, but where data are available, FCR can be devastating. For example, FCR can reduce yields of winter wheat production in the Pacific Northwest region of the USA by up to 61% (Smiley et al. 2005). FCR also affects grain quality through the production of mycotoxins such as DOV, NIV, ZEN and T2-toxin which can be harmful to human, and livestock health (Pestka and Smolinski 2005; Blandino et al. 2012).

Over the past 5 years, FCR re-emerged as an economically important disease in Iraq, causing significant yield losses to the wheat crop (Matny et al. 2012). A few studies on FCR have been carried out to understand the genetic diversity present in *F. culmorum* populations and to understand why this disease has re-emerged. Drought conditions in Iraq from 2011 to 2016 are likely a contributing factor to the spread of *F. culmorum* in wheat fields since many studies have shown that dry environments are favorable for *F. culmorum* growth and reproduction (Scherf et al. 2013; Balmas et al. 2006). FCR has also been reported in other Middle Eastern countries such as Turkey (Tunali et al. 2006; Emre et al. 2016), Iran (Hajieghrar 2009; Eslahi 2012) and Syria (Khalifeh et al. 2009).

Genetic diversity analyses of microorganisms have demonstrated that pathogen diversity depends on global environmental changes and shifts in agro-ecological systems (Saharan

and Naef 2008; Gurel et al. 2010). *F. culmorum* isolates show high levels of phenotypic and genotypic variability in culture, including colony morphology, pigmentation and sporulation (Puhalla 1981; Kollers et al. 2013; Miedaner et al. 2013). In addition, variation in aggressiveness and mycotoxin production have been found among various isolates that were collected from different geographic locations (Gang et al. 1998; Berna et al. 2012; Winter et al. 2013; Fang et al. 2015).

There are many methods and techniques used for studying the genetic diversity of microorganisms, including Repetitive Polymerase Chain Reaction (REP-PCR), also known as Repetitive DNA-based fingerprinting. The amplification of prokaryotic genomic sequences between the repetitive elements include: Repetitive Extragenic Palindromic (REP) sequences, Entero-bacterial Repetitive Intergenic Consensus (ERIC) sequences and BOX elements. REP-PCR applications are in widespread use among studies of plant pathogenic bacteria, but among eukaryotic microbes, have only been tested in *F. oxysporum* (Edel et al. 1995). The principle aim of this study is to characterize the genetic diversity among *F. culmorum* isolates collected in Iraq through REP-PCR and associate the results with their geographic distribution and pathogenicity toward wheat at the seedling stage.

MATERIALS AND METHODS

Plant material and fungal isolation

Wheat plants exhibiting FCR symptoms were collected from seven provinces representing different agricultural zones in Iraq. All samples were collected in paper

envelopes and necessary data (sample number, place and date of collection and host cultivar name) were recorded. The samples were brought to the laboratory and kept in a well-ventilated area at room temperature ($25\pm 3^{\circ}\text{C}$ and 30% humidity) until the samples could be processed. The crown of wheat plants were cut into 0.5-1.0 cm segments, and treated with a 10% sodium hypochlorite (bleach) solution (diluted from commercially-available concentrated bleach) for 2 min, washed with sterile water and dried with filter paper. All samples were placed into 9 cm petri dishes containing potato dextrose agar (PDA). Fifty milligrams of Agrimycin-343 was added to the medium after autoclaving. Plates were incubated at $25^{\circ}\text{C}\pm 2$ for 5 days and then a single *Fusarium* spp. spore was picked up (under a microscope at 400X) using a needle according to colony and spore characterization methodologies (Booth 1971; John and Brett 2006) and placed into a new petri dish containing PDA for use in pathogenicity assays.

Pathogenicity assay

For the pathogenicity assays, *F. culmorum* isolates were grown on autoclaved millet seed. First, one kilogram of millet seed was soaked in water for 12 h, then the water was drained, and several 250 ml flasks were filled with 50 g each of this millet seed and autoclaved at 121°C and 1.5 kg cm^{-1} pressure for 20 min. One disc (0.5 cm) of a 7-day old *F. culmorum* colony was placed into each flask and incubated at $25\pm 2^{\circ}\text{C}$ for 14 days. Pathogenicity tests were then performed on wheat seedlings. A 1:1 mixture of sterile soil and peat moss was autoclaved at 121°C and 1.5 kg cm^{-1} pressure for 1h. This process was repeated on two separate days. Pots ($5 \times 10\text{ cm}$) used in the greenhouse experiments were filled with the mixture soil and 5 g of *F. culmorum* inoculum was added to each pot in the top 5 cm surface layer of the soil. All pots were watered and placed in the

greenhouse for 2 days 25 ± 2 . Three seeds of *Triticum aestivum* L. cv. Abu-Ghreeb1 (a commonly used cultivar used in Iraq) were sown in each pot, and each treatment repeated three times. Plants were irrigated with sterilized water as needed. After 35 days, crown rot symptoms and stem discoloration characteristic of FCR were apparent on the inoculated plants. FCR Severity was calculated by measuring the length of discoloration relative to seedling height. The FCR severity index was obtained by multiplying this ratio by the number of leaf-sheath layers with necrosis. The FCR index was calculated according to the following formula: (length of stem discoloration/seedling height) $\times$ (number of leaf sheath layers with necrosis) (Mitter et al. 2006).

Fungal growth and DNA extraction

Twenty- nine *Fusarium* spp. isolates were grown on PDA media in 9 cm petri dish for 7 days at $25 \pm 2^\circ\text{C}$. The REDEExtract-N-AMPTM Plant PCR kit (Sigma-Aldrich, St. Louis, MO, USA) was used for DNA extraction according to manufacturer's instructions. Briefly the hyphal tip of the mycelium was harvested with a sterilized needle and placed in 0.2 ml collection tubes, to which 50 μl extraction solution was added, followed by incubation at 95°C for 10 min. The DNA concentrations and quality were measured to ensure quality using a Nanodrop 2000 spectrophotometer (Thermo Fisher Scientific, USA), qualitative analyses of DNA were carried out via agarose gel electrophoresis.

Amplification of fungal DNA

Species identification of *Fusarium* spp. isolates was determined by amplifying and sequencing the Translation Elongation Factor 1 alpha (TEF-1 α) gene. Forward (EF1) 5'-ATGGGTAAGGA(A/G)GACAAGAC-3' and reverse (EF2) 5'-GGA(G/A)GTAC

CAGT(G/C)ATCATG-3' (O'Donnell et al. 2000) primers were used to amplify the TEF-1 α gene. The PCR reaction solution was prepared at total volume of 20 μ l. The PCR conditions were as follows: an initial denaturation at 95°C for 5 min followed by 35 cycles of denaturation at 94°C for 50 s, annealing at 53°C for 50 s, extension at 72°C for 1 min, and a final extension at 72°C for 7 min. Amplification products were visualized on 1.0% agarose gel stained with SYBRTM safe DNA gel stain in 1X TAE (InvitrogenTM, Carlsbad, CA, USA). Amplification of a product approximately 700 bp long was generated by PCR from the DNA template.

DNA sequencing

The PCR products of TEF-1 α were prepared for sequencing by cleaning up with the QiAquick® PCR purification kit (city? Bogota?, Colombia). Sequencing was carried out commercially (ACGT, Inc., Chicago, USA). The sequencing chromatograms were read and aligned using MEGA6 software and the sequences were compared with those in GenBank (<http://www.ncbi.nlm.nih.gov/>) for the TEF-1 α gene using the Basic Local Alignment Search Tool (BLAST). All sequences of the isolates were sent to GenBank to obtain accession numbers.

Genetic diversity study

F. culmorum isolates collected from different regions of Iraq were genotyped using three primer pairs designed to amplify multiple regions of the genome simultaneously: (1) REP1R-Dt (5'- III NCGNCATCNGGC-3') and REP-2G (5'-GCGGCTTATCGG GCCTAC-3') for REP; (2) ERIC 1 (5'-ATG TAAGCTCCTGGGGATTAC-3') and ERIC 2 (5'-AAGTAAGTGACTGG GGTGAGCG-3') for ERIC, and (3) BOX-A1R (5'-

CTACGGCAAG GCGACGCTGACG-3') (Versalovic et al. 1991) for BOX- ERIC2. The PCR conditions were: an initial denaturation at 94°C for 5 min, followed by 40 cycles of 94°C for 1 min, an annealing step of 50°C (BOX and ERIC) or 37°C (REP) for 1 min, and an extension at 72°C for 2 min, followed by a final extension at 72°C for 15 min. Amplification products were visualized on 1.5% agarose gels with SYBR™ safe DNA gel stain in 1X TAE (Invitrogen™). In total, 410 potential markers were generated by this method for genotyping. REP-PCR markers were evaluated together in pair-wise comparisons. Single and shared fragments were analyzed using by Multivariate Statistical Package (MVSP) 3.22 program, the similarity was calculated according to minimum variance (Squared Euclidean) (Kovach 2001).

RESULTS

Pathogenicity tests for *F. culmorum* on wheat seedlings demonstrated that the collected isolates vary in their pathogenicity toward wheat cultivar Abu-Grheeb1, ranging from 0.001 to 0.890 on the FCR severity index; however, some isolates (IF 0003, IF 0013, IF 0024) were non-pathogenic, with a score of 0.00 on the FCR severity index. The isolates that resulted in the highest FCR severity index scores were: IF 0021, IF 0028, IF 0045, IF 0046, IF 0015, and IF 0005. Isolates IF 0021, IF 0028, IF 0045 and IF 0046 originate from Baghdad while IF 0015 and IF 0005 come from Anbar and Diyala provinces, respectively (Table 1).

192 Table 1. *Fusarium culmorum* accessions used in this study including loci information,
193 TEF-1 α gene test and disease severity index.

No.	Accession No.	Isolate No.	Species	Location	TEF-1 α gene	FCR severity index
1	KY205745	IF 0003	<i>F. culmorum</i>	Karbala	+	0.00
2	KY205746	IF 0004	<i>F. culmorum</i>	Karbala	+	0.030
3	KY190104	IF 0005	<i>F. culmorum</i>	Diyala	+	0.286
4	KY190106	IF 0006	<i>F. culmorum</i>	Diyala	+	0.001
5	KY190111	IF 0007	<i>F. culmorum</i>	Diyala	+	0.074
6	KY190107	IF 0008	<i>F. culmorum</i>	Diyala	+	0.002
7	KY190127	IF 0009	<i>F. culmorum</i>	Kirkuk	+	0.098
8	KY190118	IF 0013	<i>F. culmorum</i>	Anbar	+	0.00
9	KY190123	IF 0014	<i>F. culmorum</i>	Anbar	+	0.011
10	KY190116	IF 0015	<i>F. culmorum</i>	Anbar	+	0.309
11	KY190108	IF 0017	<i>F. culmorum</i>	Najaf	+	0.022
12	KY205747	IF 0021	<i>F. culmorum</i>	Baghdad	+	0.890
13	KY190121	IF 0022	<i>F. culmorum</i>	Baghdad	+	0.294
14	KY190126	IF 0024	<i>F. culmorum</i>	Diyala	+	0.00
15	KY190122	IF 0026	<i>F. culmorum</i>	Anbar	+	0.050
16	KY190112	IF 0028	<i>F. culmorum</i>	Baghdad	+	0.543
17	KY205748	IF 0029	<i>F. culmorum</i>	Baghdad	+	0.004
18	KY190109	IF 0030	<i>F. culmorum</i>	Kirkuk	+	0.001
19	KY190113	IF 0031	<i>F. culmorum</i>	Kirkuk	+	0.030

20	KY190114	IF 0032	<i>F. culmorum</i>	Kirkuk	+	0.002
21	KY190117	IF 0033	<i>F. culmorum</i>	Kirkuk	+	0.004
22	KY190124	IF 0040	<i>F. culmorum</i>	Babylon	+	0.076
23	KY190110	IF 0041	<i>F. culmorum</i>	Babylon	+	0.005
24	KY190119	IF 0042	<i>F. culmorum</i>	Babylon	+	0.001
25	KY205749	IF 0044	<i>F. culmorum</i>	Diyala	+	0.019
26	KY190105	IF 0045	<i>F. culmorum</i>	Baghdad	+	0.350
27	KY190125	IF 0046	<i>F. culmorum</i>	Baghdad	+	0.200
28	KY190120	IF 0047	<i>F. culmorum</i>	Baghdad	+	0.008
29	KY190115	IF 0052	<i>F. culmorum</i>	Anbar	+	0.010

194

195 The results of species-specific identification using TEF-1 α demonstrated that all isolates
196 used in this study are in fact *F. culmorum*.

197 The genetic diversity study showed monomorphic and polymorphic bands pattern. The
198 annealing temperature for the BOX, ERIC and REP primers used in this study were
199 different from the output provided by Gurel et al. (2010). In this study, we found the
200 optimum annealing temperature for the primers was 50°C for BOX and ERIC, and 37°C
201 for REP. The PCR bands for the final amplification products were between 200-800 bp
202 for BOX-ERIC2, 110-1100 bp for ERIC-ERIC2, and 200-1300 bp for REP. A total of
203 410 polymorphic markers were identified in this study for the *F. culmorum* isolates,
204 including 106 for BOX, ERIC for 175 and 129 for the REP-PCR. These markers were
205 used to evaluation the minimum variance between the strains (Fig. 2).

206

Minimum variance cluster analysis was used to detect the variance between the *F. culmorum* isolates (fig.1). The dendrogram illustrates separated the isolates into major groups in this present study. Group 1 includes 17 isolates and may be divided into two sub-groups (1A and 1B) consisting of 5 and 11 isolates, respectively. All isolates in this group were collected from northern and central sites in Iraq. Group 2 consists of 12 isolates and may also be divided into two subgroups (2A and 2B) consisting of 3 and 9 isolates, respectively. Group 2A isolates originate from central Iraq while group 2B isolates come from northern, central, and southern Iraq.

Isolates IF0022 and IF0026 share the highest similarity values (100% similar). There is no relationship between the geographic origin of the isolate and its genetic relationship to other isolates (Fig. 1). One of the reasons is because of the use of seed that has not been certified by the Iraqi Ministry of Agriculture and the exchange of seed between farmers across different regions leading to the transfer the pathogen with the seed from one province to another. Also, some farmers obtain their seed from local markets where the seed source is unknow.

Figure 1. Dendrogram of 29 *Fusarium culmorum* isolates generated based on the number of bands and position of appearance for three primers BOX-ERIC2, ERIC-ERIC2 and REP by using Multivariate Statistical Package (MVSP) 3.22 program to show minimum variance (Squared Euclidean).

(A) BOX-ERIC2

(B) ERIC-ERIC2

(C) REP

Figure 2. Fingerprint pattern for 29 *F. culmorum* isolates by using three primer pairs: (A) BOX-ERIC2, (B) ERIC-ERIC2 and (C) REP.

DISCUSSION

Genetic diversity studies of pathogen populations are important for understanding the genetic potential of economically important pathogens to adapt to climate change and the implications for management of diseases caused by these pathogens (McDonald and Linde 2002). *F. culmorum* has also been reported to cause seed-borne diseases of pre- and post-emergence seedling death and is one of the causal species of Fusarium head blight (FHB) (Polley and Turner 1995). In this study, **89.7%** of the *F. culmorum* isolates were pathogenic towards wheat at the seedling stage, while a smaller **10.3%** number of isolates were non-pathogenic.

As the results of our study of the genetic diversity among 29 isolates of *F. culmorum* collected from different regions of Iraq show, there is no relationship between geographic location and genetic similarity of the isolates. We suggest that this means that *F. culmorum* populations have the ability to survive and adapt to different and extreme variations in climate, from the cold area in northern Iraq with temperatures ranging from

-5 to 10°C in winter and 35 to 45°C in summer; to central and southern Iraq where temperatures range from around 5 to 15°C during winter to 45 to 55°C in the summer.

Many studies of DNA analysis have been reported to investigate genetic variability and population structure of *F. culmorum* using a variety of molecular markers, such as random amplified polymorphic DNA (RAPD) (De Nijs et al. 1995; Gargouri et al. 2003; Yörük and Albayrak, 2013) and restriction fragment length polymorphism (RFLP) (Nicholson et al 1993; Llorensa et al. 2006). These studies suggest that there is extensive genetic diversity in *F. culmorum* populations. In addition, Mishra et al. (2003) found a high degree of intra-specific polymorphism among *F. culmorum* isolates using inter-simple sequence repeat (ISSR) analysis. Albayrak et al. (2016) also studied the relationship between *Fusarium* isolates according to their species and geographic regions by using ISSR markers. In another study, Bayraktar and Fatma (2010) found that ISSR markers have a high degree of intra- and interspecific polymorphisms among *Fusarium* spp. Finally, Gargouri et al. (2003) used RAPD markers to study the genetic variability and population structure of *Fusarium culmorum* isolated from wheat stem bases.

In addition to the genetic diversity and variation in pathogenicity towards wheat presented in this study, populations of *F. culmorum* are also characterized by high levels of phenotypic variability in culture, such as sporulation, pigmentation, mycotoxin production, and colony morphology (Puhalla 1981). Gang et al. (1998) found a large variation among *F. culmorum* isolates collected from the various geographic areas for aggressiveness, race designation and mycotoxin production.

CONCLUSION

This is the first report of the genetic diversity of *F. culmorum* populations present in Iraq using the REP-PCR method. Two groups of *F. culmorum* were identified in this study according to minimum variance (Squared Euclidean). This is the first study in this field that has been reported in Iraq. Although the study was limited to 29 isolates due to the difficulty in completing more extensive sampling, this study provides a first glimpse at the genetic diversity and variation in pathogenicity present in *F. culmorum* populations in Iraq.

ACKNOWLEDGMENTS

This work was carried out and supported by the University of Minnesota, Department of Plant Pathology. Thanks to Dr. Scott Bates and Dr. Zewei Song for all the support and help to complete this paper.

REFERENCES

1. Albayrak, G., Emre, Y., Aylin, G., Bahram, S. (2016). Genetic diversity among *Fusarium graminearum* and *F. culmorum* isolates based on ISSR markers, Arch Biol Sci, 68(2), 333-343. doi:10.2298/ABS150630025A
2. Balmas, V., Delogu, G., Sposito, S., Rau, D., Migheli, Q. (2006). Use of a complexation of tebuconazole with b-cyclodextrin for controlling foot and crown rot of durum wheat incited by *Fusarium culmorum*. The Journal of Agricultural and Food Chemistry, 54, 480-484, doi: 10.1021/jf0523014

- 299 3. Bayraktar, H., and Fatma, D. (2010). Molecular Identification and Genetic Diversity
300 of *Fusarium* species Associated with Onion Fields in Turkey, 159 (1), 28–34. DOI:
301 10.1111/j.1439-0434.2010.01715.x
- 302 4. Berna, T., Friday, O., Gul, E., Rhyannyn, W., Julie, N., Sukumar, C. (2012). Fitness of
303 three *Fusarium* pathogens of wheat. FEMS Microbiol Ecol 81 596–609. doi:
304 10.1111/j.1574-6941.2012.01388.x
- 305 5. Blandino, M., Haidukowski, M., Pascale, M., Plizzari, L., Scudellari, D., Reyneri, A.
306 (2012). Integrated strategies for the control of *Fusarium* head blight and
307 deoxynivalenol contamination in winter wheat. Field Crops Research, 133, 139-149,
308 doi: 10.1016/j.fcr.2012.04.004
- 309 6. Booth, C. (1971). The Genus *Fusarium*. Commonwealth Mycological Institute, Kew,
310 Surrey. Farnham Royal, England: Commonwealth Agricultural Bureaux [for the]
311 Commonwealth Mycological Institut, United Kingdom, Pp, 237.
- 312 7. Burgess, L.W., Backhouse, D., Summerell, B.A., Swan, L.J. (2001). Crown rot of
313 wheat In: Summerell BA, Leslie JF, Backhouse D, Bryden WL, Burgess LW (eds),
314 *Fusarium*: Paul E. Nelson Memorial Symposium. APS Press, St. Paul, Minnesota,
315 USA, pp, 271-294.
- 316 8. Chakraborty, S., Liu, C.J., Mitter, V., Scott, J.B., Akinsanmi, O.A., Ali, S., Dill-Macky,
317 R., Nicol, J., Backhouse, D., Simpfendorfer, S. (2006). Pathogen population structure
318 and epidemiology are keys to wheat crown rot and *Fusarium* head blight management.
319 Australasian Plant Pathology, 35, 643-655, doi: 10.1071/AP06068
- 320 9. De Nijs, M., Larson, J. S., Gams, W., Rombouts, F. M., Wernas, K., Thrane, U. (1997).
321 Variation in random amplified polymorphic DNA patterns and secondary metabolic

322 profiles within *Fusarium* species from cereals from various parts of The Netherlands.
 323 Food Microbiology, 14, 449-457.

324 10. Edel, V., Steinberg, C., Avelange, I., Laguerre, G., Alabouvette, C. (1995). Comparison
 325 of three molecular methods for the characterization of *Fusarium oxysporum* strains.
 326 Phytopathology, 85, 579-585, doi: 10.1094/Phyto-85-579

327 11. El-Khalifeh, M., El-Ahmed, A., Al-Saleh, A., Nachit, M. (2009). Use of AFLPs to
 328 differentiate between *Fusarium* species causing root rot disease on durum wheat
 329 (*Triticum turgidum* L. var. *durum*). African Journal of Biotechnology, 8, 4347-4352.

330 12. Emre, Y., Berna, T., Bayram, K., Fatih, Ö., Gülşen, U., Işıl, Z., Ayşegül, S.,
 331 Gonca, M. (2016) . Characterization of high-level deoxynivalenol producer *Fusarium*
 332 *graminearum* and *F. culmorum* isolates caused head blight and crown rot diseases in
 333 Turkey, J Plant Dis Prot, 123: 177-186. doi:10.1007/s41348-016-0027-y

334 13. Eslahi, M., (2012). Fungi associated with root and crown rot of wheat in Khuzestan
 335 province, Iran. Journal of Crop Protection, 1(2), 107-113.

336 14. Fang, Ji., Jirong, W., Hongyan, Z., Jianhong, X., Jianrong, S. (2015) . Relationship of
 337 Deoxynivalenol Content in Grain, Chaff, and Straw with *Fusarium* Head Blight
 338 Severity in Wheat Varieties with Various Levels of Resistance, Toxins, 7, 728-742.
 339 doi:10.3390/toxins7030728

340 15. Gang, G., Miedaner, T., Schuhmacher, U., Schollenberger, M., Geiger, H.H. (1998).
 341 Deoxynivalenol and nivalenol production by *Fusarium culmorum* isolates differing in
 342 aggressiveness toward winter rye. Phytopathology, 88,879-884, doi: 10.1094/PHYTO
 343 .1998.88.9.879

- 344 16. Gargouri, S., Louis, B., Mohamed, H., Mohamed, M. (2003). Genetic Variability and
345 Population Structure of the Wheat Foot Rot Fungus, *Fusarium culmorum*, in Tunisia,
346 European Journal of Plant Pathology, 109, 807- 815. doi:10.1023/A:1026137817723
- 347 17. Gurel, F., Albayrak, G., Diken, O., Cepni, E., Tunali, B. (2010). Use of REP-PCR for
348 Genetic Diversity Analyses in *Fusarium culmorum*. J. Phytopathology, 158,387–389,
349 doi: 10.1111/j.1439-0434.2009 .01630.x
- 350 18. Hajieghrar, B. (2009). Wheat crown and root rotting fungi in Moghan area, Northwest
351 of Iran, African Journal of Biotechnology, 8 (22), 6214-6219. doi:
352 10.5897/AJB09.1087
- 353 19. John, F. L., & Brett, A.S. (2006). The Fusarium Laboratory Manual. Blackwell
354 Publisher, London, pp 388, doi: 10.1002/9780470278376
- 355 20. Kollers, S., Bernd, R., Jie, L., Viktor, K., Erhard, E. , Odile, A., Maike, H., Jorg, P.,
356 Dagmar, K., Martin, G., Marion, S. Roder. (2013). Whole Genome Association
357 Mapping of Fusarium Head Blight Resistance in European Winter Wheat (*Triticum*
358 *aestivum* L.), PLoS One, 8 (2): e57500. doi: 10.1371/journal.pone.0057500
- 359 21. Kovach, WL. (2001). MVSP: Multivariate statistical package v3.12f. Kovach
360 Computing Services, Pentraeth, Anglesey, UK.
- 361 22. Llorensa, A., Hinojoa, M., Mateob, R., González-Jaénc, M., Valle-Algarab,
362 F., Logriecod, A., Jiménez, M. (2006). Characterization of *Fusarium* spp. isolates by
363 PCR-RFLP analysis of the intergenic spacer region of the rRNA gene (rDNA),
364 International Journal of Food Microbiology, 106, (3), 297–306. doi: 10.1016/j.ijf
365 oodmicro.2005.09.005

- 366 23. Matny, O.N., Chakraborty, S., Obanar, F., AL-Ani, R.A. (2012). Molecular
367 identification of *Fusarium* spp. causing crown rot and head blight on winter wheat in
368 Iraq. Journal of Agricultural Technology, 8(5), 1677-1690.
- 369 24. McDonald, B.A., & Linde, C. (2002). The population genetics of plant pathogens and
370 breeding for durable resistance. Euphytica, 124, 163-180, doi: 10.1023/A:10156
371 78432355
- 372 25. Miedaner, T., Würschum, T., Maurer, H., Korzun, V., Ebmeyer, E. (2013). Whole
373 Genome Association mapping for *Fusarium* head blight resistance in European soft
374 winter wheat, Mol Breed, 28, 647–655.
- 375 26. Mishra, P.K., Fox, R.T., & Cutham, A. (2003). Inter-simple sequence repeat and
376 aggressiveness analyses revealed high genetic diversity, recombination and long range
377 dispersal in *Fusarium culmorum*. Annals of Applied Biology, 143, 291-301, doi:
378 10.1111/j.1744-7348.2003.tb00297.x
- 379 27. Mitter, V., Zhang, M. C., Liu, C. J., Ghosh, R., Ghosh, M., & Chakraborty, S. (2006).
380 A high-throughput glasshouse bioassay to detect crown rot resistance in wheat
381 germplasm. Plant Pathology, 55,433-441, doi: 10.1111/j.1365-3059.2006.01384.x
- 382 28. Nicholson, P., Jenkinson, P., Rezanoor, H.N., & Parry, D.W. (1993). Restriction
383 fragment length analysis of variation in *Fusarium* species causing ear blight of cereals.
384 Plant Pathology, 42, 905-914, doi: 10.1111/j.1365-3059.1993.tb02676.x
- 385 29. O'Donnell, K., Kistler, H.C., Tacke, B.K., Casper, H.H. (2000). Gene genealogies
386 reveal global phylogeographic structure and reproductive isolation among lineages of
387 *Fusarium graminearum*, the fungus causing wheat scab. Proceedings of the National
388 Academy of Sciences, USA, 9, 7905-7910.

- 389 30. Pestka, J.J., & Smolinski, A.T. (2005). Deoxynivalenol: toxicology and potential
390 effects on humans. *Journal of Toxicology and Environmental Health. Part B, Critical*
391 *Review*, 8, 39-69, doi: 10.1080/10937400590889458
- 392 31. Polley, R. W., & Turner, J. A. (1995). Surveys of stem base diseases and *Fusarium* ear
393 diseases in winter wheat in England, Wales and Scotland 1989–1990. *Annals of*
394 *Applied Biology*, 126, 49-59, doi: 10.1111/j.1744-7348.1995.tb05002.x
- 395 32. Puhalla, J.E. (1981). Genetic considerations of the genus *Fusarium*. pp. 291–305. In:
396 *Fusarium Diseases, Biology, and Taxonomy* (Nelson P.E., Toussoun T.A, Cook R.J.,
397 ed.),. University Park, Pennsylvania, Pennsylvania State University Press, PA, USA.
- 398 33. Saharan, M., Naef, A. (2008) . Detection of genetic variation among Indian wheat head
399 scab pathogens (*Fusarium* spp. isolates) with microsatellite markers. *Crop Protection*,
400 27, 1148–1154. doi: 10.1016/j.cropro.2008.01.008
- 401 34. Scherm, B., Balmas, V., Spanu, F., Pani, G., Delogu, G., Pasquali, M., Migheli, Q.
402 (2013). *Fusarium culmorum*: Causal agent of foot and root rot and head blight on
403 wheat. *Molecular Plant Pathology*, 14(4), 323-341, doi: 10.1111/mpp.12011
- 404 35. Smiley, R., Gourlie, J., Easley, S., Patterson, L-M., Whittaker, R. (2005). Crop damage
405 estimates for crown rot of wheat and barley in the Pacific Northwest. *Plant Disease*, 89,
406 595-604, doi:10.1094/PD-89-0595
- 407 36. Tunal, B., Nicol, J., Erol, F.Y., Altiparmak, G. (2006). Pathogenicity of Turkish crown
408 and head scab isolates on stem bases on winter wheat under greenhouse conditions.
409 *Plant Pathology Journal*, 5(2), 143-149, doi: 10.3923/ppj.2006.143.149
- 410 37. Versalovic, J., Koeuth, T., Lupski, J.R. (1991). Distribution of repetitive DNA
411 sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic*
412 *Acids Res*, 19, 6823–6831.

- 413 38. Winter, M., Koopmann, B., Döll, K., Karlovsky, P., Kropf, U., Schlüter, K., von
414 Tiedemann, A. (2013). Mechanisms regulating grain contamination with trichothecenes
415 translocated from the stem base of wheat (*Triticum aestivum*) infected with *Fusarium*
416 *culmorum*. Phytopathology 103:682-689. doi: 10.1094/PHYTO-11-12-0296-R
- 417 39. Yörük, E., G. Albayrak. (2013). Genetic characterization of *Fusarium graminearum*
418 and *F. culmorum* isolates from Turkey by using random-amplified polymorphic DNA,
419 Genet. Mol. Res, 12 (2), 1360-1372, doi: 10.4238/2013.April.25.7