1 **Rapid, specific detection and quantification of *Yersinia pestis***
2 **using a species-specific SNP in the ferric uptake regulator**
3 **gene (*furMAMA*)**

4
5 Authors: Pernille Nilsson,^{a#} Shair Gurbanov^b and W. Ryan Easterday^{a#}

6
7 Affiliations:

8 ^a Centre for Ecological and Evolutionary Synthesis, Department of
9 Biosciences, University of Oslo, P.O. Box 1066 Blindern, NO-0316 Oslo,
10 Norway

11 ^b Republican Anti-plague station, Baku, Azerbaijan

12 Email addresses:

13 Pernille Nilsson: pernille.nilsson@ibv.uio.no

14 Shair Gurbanov: shair-gurbanov@mail.ru

15 W. Ryan Easterday: w.r.easterday@ibv.uio.no

16
17 Running head: Specific quantification of *Y. pestis* with TaqMAMA qPCR

18
19 Corresponding authors:

20 #Address correspondence to Pernille Nilsson, pernille.nilsson@ibv.uio.no or

21 W. Ryan Easterday, w.r.easterday@ibv.uio.no

26 **Abstract**

27 *Yersinia pestis*, the causative agent of plague, is responsible for about 700
 28 human cases of bubonic and pneumonic plague each year. Yet the disease is
 29 far more prevalent within rodent reservoirs than in humans. One of the main
 30 means of outbreak prevention is extensive wildlife surveillance, where
 31 accurate and rapid detection is essential to prevent spillover into the human
 32 population from which, it may otherwise spread more rapidly and over larger
 33 distances. Moreover, detection and quantification of the agent aids in
 34 investigative studies to understand aspects of the pathogen such as
 35 transmission mechanics, pathology, contamination risk and more. Partially
 36 based on a previously developed assay by Gabitzsch et al. 2008 we designed
 37 a TaqMan® mismatch amplification mutation assay (TaqMAMA) where a
 38 primer leverages a species-specific SNP in the chromosomal single copy
 39 ferric uptake regulator gene of *Yersinia pestis*. The assay allows for specific,
 40 rapid detection and quantification of *Yersinia pestis* using only a single
 41 species-specific marker in a highly conserved virulence gene. This low-cost
 42 and simple modification of an existing assay eliminates the need for running
 43 multiple molecular markers for pathogen detection or performing time-
 44 consuming culturing and counting of colonies for quantification.

45

46 **Keywords**

47 *Yersinia pestis*, TaqMAMA, quantification, qPCR, pathogen surveillance, SNP
 48 based detection, private allele, bacterial detection.

49

50 **Background**

51 The causative agent of plague, *Yersinia pestis*, reached infamy after causing
 52 three devastating human pandemics during recorded history. The disease
 53 however, is mainly a vector-borne, wildlife disease, circulating within rodent
 54 populations across the world, which occasionally spills over into human
 55 populations. Today there are approximately 700 reported human cases of
 56 plague worldwide each year with more than 100 deaths (World Health
 57 Organization, 2017), most of which occur in Africa. Due to the pandemic
 58 history of the pathogen and the continued potential for human outbreaks,
 59 close surveillance of its presence in wildlife systems has been incorporated
 60 into several local and national surveillance programs, some dating back to the
 61 beginning of the 1900s (Melikishvili, 2006).

62
 63 Plating of infected animal tissues or fleas and subsequent counting of
 64 bacterial colonies is a well-established, sensitive and widely used method in
 65 many surveillance programs and in diagnostics (Bevins et al., 2012). This is a
 66 time-consuming method that exposes the workers to infectious material for
 67 extended time. The culturing of *Y. pestis* has historically had challenges due
 68 to the relatively slow growth of *Yersinia spp.* compared to the near
 69 omnipresent contaminating microbes on traditional media (Dennis et al., 1999;
 70 Gupta et al., 2015; Sarovich et al., 2010). To overcome some of these issues,
 71 new and improved growth media with increased selectivity for *Y. pestis* have
 72 been developed but still struggle with obstacles such as reduced recovery
 73 rate of the pathogen and desired level of selectivity leading to underestimation
 74 of bacterial numbers (Ber et al., 2003; Riehm et al., 2011; Sarovich et al.,

75 2010). As of today, culturing remains a gold standard for detection and in
76 particular quantification.

77

78 The recent development of immunoassay tests for rapid and easy detection in
79 the form of dipsticks have become preferential, particularly for quick
80 diagnostics in challenging field conditions in remote locations (Chanteau et
81 al., 2003; Simon et al., 2013). Yet immunoassays are typically not sensitive
82 enough to pick up cases of low-level infections or early on in infections where
83 only small amounts or no antibody is present (Andrianaivoarimanana et al.,
84 2012; Chanteau et al., 2003; Simon et al., 2013). Other limitations of these
85 assays are potential false negative results due to the lack of detection of
86 strains that have genetic deletions of the assays target (such as the F1
87 capsular antigen) (Anisimov et al., 2004; Eppinger et al., 2010; Perry and
88 Fetherston, 1997). They are also altogether unsuitable for quantification.

89

90 In the era of sequencing and genomics and the power and relative ease of
91 PCR based methods, the gold standard of detection is shifting from culture-
92 based methods towards molecular tools. In this context, PCR based assays
93 are able to detect the pathogen at low quantities making them the most
94 appropriate choice for investigative studies where maximal sensitivity is
95 required. Yet specific detection of bacterial pathogens using PCR or
96 immunoassay tests must often navigate several major obstacles, many being
97 the products of bacterial evolution. Most pathogens evolve out of complexes
98 of related species, where mechanisms like horizontal gene transfer (HGT)
99 allows a constant exchange of genetic elements, often as plasmids or genetic

100 islands (Juhas, 2015). As a result of these evolutionary trends, specific
101 detection of the target of interest, as is the case for many pathogenic bacteria
102 including *Y. pestis* and *Y. pseudotuberculosis*, can be difficult (Achtman et al.,
103 1999; Easterday et al., 2005; U'Ren et al., 2005). For *Y. pestis*, its recent
104 evolution from *Y. pseudotuberculosis*, has been accomplished through a
105 process of functional gene loss and gain through HGT within
106 *Enterobacteriaceae*. This results in *Y. pestis* being, more or less, a clone of *Y.*
107 *pseudotuberculosis* with shared elements from other members of
108 *Enterobacteriaceae* (Achtman et al., 1999; Hinnebusch et al., 2016).

109
110 Historically, discrimination between *Y. pestis* and *Y. pseudotuberculosis* has
111 been challenging for methods targeting chromosomal sequences, due to the
112 high degree of genetic similarity between the two species (Califf et al., 2015;
113 Chain et al., 2004; Gabitzsch et al., 2008; Neubauer et al., 2000). A work-
114 around has been to use multiple assays to detect specific combinations and
115 variants of genes located on the chromosome and/or the plasmids of *Y. pestis*
116 (Stewart et al., 2008; Tomaso et al., 2003). Still, the problems with finding *Y.*
117 *pestis* specific genes on the main chromosome also extends to plasmid
118 targets, such as the pCD1 plasmid, a homologue of pYV1 found in *Y.*
119 *pseudotuberculosis* and *Y. enterocolitica* (Hu et al., 1998; Portnoy et al.,
120 1984).

121
122 The two most conventional targets for detection of *Y. pestis* both under
123 laboratory and field conditions are the Capsule antigen fraction 1 (*caf1*) gene
124 on the pMT1 plasmid and the plasminogen activator gene (*pla*) on the pPCP

125 plasmid. *Pla* is one of the most widely used markers for PCR based detection
 126 of *Y. pestis* in ancient human remains, animals and environmental samples
 127 due to it being present on the pPCP plasmid which is present in multiple
 128 copies in the bacteria and its presumed high specificity for *Y. pestis* (Harbeck
 129 et al., 2013; Stewart et al., 2008; Tomaso et al., 2008). However, recently it
 130 has been established that the gene is present and highly conserved in closely
 131 related bacterial species, and consequently is no longer considered a specific
 132 marker for *Y. pestis* (Armougom et al., 2016; Hänsch et al., 2015). Generally,
 133 a minimum of one other independent molecular marker is required to
 134 positively determine presence of the bacteria; these include genes *caf1* and
 135 Yersinia murine toxin (*ymt*) from the pMT1 plasmid, Low calcium response V
 136 antigen (*lcrV*) from the pCD1 plasmid, pesticin gene on pPCP or
 137 chromosomal targets such as 16S or Inner membrane protein yihN (*yihN*)
 138 gene either in separate or multiplex real-time PCR assays (Iqbal et al., 2000;
 139 Stewart et al., 2008; Tomaso et al., 2008; 2003; Woron et al., 2006).
 140 Furthermore plasmids are subject to variation in copy number during certain
 141 conditions, such as during an infection (Wang et al., 2016). Parkhill et al.
 142 showed that *Y. pestis* on average contain about 200 copies of the pPCP1
 143 plasmid (Parkhill et al., 2001), which is partly the reason for the preference of
 144 *pla* as target for detection as the gene is present in high copy number, yet the
 145 variability in copy number is a severe drawback for quantifying *Y. pestis*
 146 concentrations (Stewart et al., 2008; Tomaso et al., 2003).
 147
 148 By combining the established TaqMan® assay with a mismatch amplification
 149 mutation assay (TaqMAMA) (Cha et al., 1992), Glaab and Skopek (Glaab and

150 Skopek, 1999) developed a powerful and rapid method of discriminating
 151 specific single nucleotide alleles in real-time providing a valuable tool for
 152 several fields of science dealing with discrimination between highly similar
 153 targets (Achtman et al., 1999; Chain et al., 2004; Lärkeryd et al., 2014).
 154 TaqMAMA assays have successfully been designed to specifically amplify,
 155 detect and quantify a particular allele when it is paramount to distinguish
 156 between highly similar DNA sequences. Most of these have been developed
 157 in cancer research (Abbaszadegan et al., 2009; Cha et al., 1992), but also for
 158 work in bioforensics or surveillance when positive detection and discrimination
 159 of a pathogen from close genetic neighbors is vital (Achtman, 2008;
 160 Easterday et al., 2005). Previously, Gabitzsch et al. (Gabitzsch et al., 2008)
 161 developed a TaqMan® assay to detect the ferric uptake regulator (*fur*) gene
 162 for the specific quantification of *Y. pestis*, yet has become outmoded as new
 163 sequences have become available showing the gene is highly conserved in
 164 other *Yersinia* and more widely occurs within *Enterobacteriaceae*.
 165 Fortuitously, a 100% *Y. pestis*-specific SNP exists within this gene inward
 166 from the priming site of the reverse primer (*YpfurR*) from the Gabitzsch assay
 167 (Fig. 1). Here we leverage this species-specific SNP to provide a simple and
 168 cheap augmentation to the original assay creating the only single target assay
 169 for *Y. pestis*, which is quantitative, sensitive and rapid as well as, species-
 170 specific.

FIG 1. Nucleotide alignment of the 87 bp PCR fragment, targeted by the assay, of the *fur* gene of a representative selection of *Yersinia* spp. Only the nucleotide differences between the strains and the generated consensus is shown. Alignment was generated through an initial BLAST search in Geneious, names (NCBI accession numbers) and sequences were subsequently extracted and entered into Jalview where an alignment was automatically performed to generate the consensus sequence and bar plot. The location of the forward and reverse primers and the probe are indicated on the top of the alignment. *pestis*: a selection of *Y. pestis* sequences covering several biovars and geographic locations, *pseudo*.: a selection of *Y. pseudotuberculosis* sequences, *entero*.: a selection of *Y. enterocolitica* sequences, 1: *Y. intermedia*, 2: *Y. aldovae*, 3: *Y. aleksiciae*, 4: *Y. kristensenii*, 5: *Y. frederiksenii*, 6: *Y. ruckeri*.

Results

In silico screen of SNP, a private allele for *Y. pestis*

The ubiquitous presence of the *Y. pestis* allele was confirmed by both megaBLAST and BLASTn searches (NCBI) of the *fur* gene sequence from CO92 and the PCR fragment. In both cases the allele was conserved in all 981 published *fur* gene sequences for *Y. pestis* (a global representation), while the alternate allele was present in all 853 available *fur* gene sequences

190 of *Y. pseudotuberculosis* and all 2950 available *fur* gene sequences of the
191 other members of the *Yersinia* genus (May 2018, Fig. 1). A BLASTn search of
192 the *fur* sequence of CO92 against the genome assembly of the strain used in
193 this study, Az-26 (1102) (not published) confirmed the presence of the *Y.*
194 *pestis* allele in the *fur* gene of this strain, as well as 100% identity with all
195 published *Y. pestis*. The SNP, as to date, is a private allele for *Y. pestis*.

196

197 **Specificity**

198 The specificity of the *fur*MAMA assay was assessed by analyzing serial 10-
199 fold dilutions of genomic DNA in quadruplicates from *Y. pestis* and *Y.*
200 *pseudotuberculosis* alongside the original assay from Gabitzsch et al.
201 (Gabitzsch et al., 2008). We found no difference in the amplification efficiency
202 between *Y. pestis* and *Y. pseudotuberculosis* when using the original reverse
203 primer from the original assay across all DNA concentrations (Fig. 2 (A)). Our
204 augmented assay successfully quantifies concentrations of *Y. pestis* ranging
205 from 5 ng (~100 000 000 genome copies) to 50 fg (~10 genome copies) with a
206 slight shift in amplification efficiency (higher Cq values) compared to when the
207 original reverse primer is used (Fig. 2 (C)). In contrast, we find amplification
208 failure in all dilutions of *Y. pseudotuberculosis* and were only able to achieve
209 false positives using extremely high concentrations of *Y. pseudotuberculosis*
210 template in the *fur*MAMA assay where we saw some weak cross-reactivity in
211 three of four replicates at 5 ng and one of the quadruplicates at 500 pg (Fig. 2
212 (D)). A combined 90 replicates consisting of 64 non-template controls (NTC),
213 and (assumed) negative tissue and soil samples (from non-endemic areas)
214 did not show any amplification.

215

216

217 FIG 2. Real-time plots indicating the amplification efficiency and specificity of the original fur
218 assay vs the *furMAMA* TaqMan® mismatch amplification mutation assay (TaqMAMA). (A)
219 Results of quadruplicate analysis of the 10-fold serial dilutions of *Y. pestis* (blue/teal) and *Y.*
220 *pseudotuberculosis* (red) DNA (left to right: 5 ng – 50 fg) using the original *fur* assay. (B) The
221 standard curve plot of *Y. pestis* DNA with the *furMAMA* assay; $R^2 = 0.99$, slope = -3.2631. (C)
222 Results of quadruplicate analysis of the 10-fold serial dilutions of *Y. pestis* with the original fur
223 assay (blue/teal) and the *furMAMA* assay (orange). Only the first four dilutions are shown.
224 The average Cq values (quadruplicate analysis) in TE buffer were as follows: 5 ng, 17.87;
225 500 pg, 21.80; 50 pg, 25.33 and 5 pg, 28.64 for the original *fur* assay and 5ng, 19.98; 500 pg,
226 23.44; 50 pg, 26.96 and 5 pg, 30.36 for the *furMAMA* assay (curves left to right). (D) Results
227 of quadruplicate analysis of the 10-fold dilutions of *Y. pseudotuberculosis* with the original *fur*
228 assay (red) and the *furMAMA* assay (grey). Only the first four dilutions are shown. The
229 average Cq values (quadruplicate analysis) in TE buffer were as follows: 5 ng, 18.16; 500 pg,
230 21.75; 50 pg, 25.23 and 5 pg, 28.96 for the original *fur* assay while the dilutions show overall
231 amplification failure with the *furMAMA* assay. For the 5 ng *Y. pseudotuberculosis* with
232 *furMAMA* three of the four replicates amplified with the average Cq value of 40.15. For the

233 500 pg only one of the quadruplicates amplified and the Cq value was 43.88. No amplification
234 was seen for either the 50 pg or 5 pg dilutions.

235

236 **Specific quantification and sensitivity of the assay**

237 To evaluate the ability of specific quantification and the sensitivity of the
238 *furMAMA* assay, 10-fold serial dilutions of *Y. pestis* were analyzed. The assay
239 successfully amplified *Y. pestis* over a dynamic range of concentrations with a
240 correlation coefficient (R^2) of 0.99 (Fig. 2 (B)) and was able to consistently
241 quantify as little as 50 fg, which is equivalent to 10 bacterial genomes. The
242 assay showed sporadic amplification at levels of the 5 fg concentration (3 out
243 of 33 reactions).

244

245 **Detection by conventional PCR**

246 The *furMAMA* assay was also run as a standard PCR without probe to test its
247 applicability as a detection only assay for when quantification equipment is not
248 available. The primer pair successfully amplified *Y. pestis* and did not produce
249 a PCR product of the correct size for any of the quadruplicates of *Y.*
250 *pseudotuberculosis* (results not shown).

251

252 **Screening for inhibition by rodent tissue and soil extracts**

253 In order to assess possible effects of foreign DNA on the *furMAMA* assay we
254 ran target-spiked controls over a standard dilution range of concentrations
255 (5ng to 50fg) into DNAs extracted from rodent tissues and soils. *Y. pestis*
256 DNA was spiked into several types of DNA extracts including ear, tail, liver
257 and spleen from several rodent species including, great gerbil (*Rhombomys*

258 *opimus*), lab mouse (*Mus musculus*) and bank vole (*Myodes glareolus*) and
 259 DNA from soil from a previous study (Turner et al., 2016). Regardless of
 260 background DNA (tissue or soil) the assay still detects and quantifies across
 261 the range tested without non-specific cross-reactivity. Environmental samples
 262 such as soil extracts can show a high degree of inhibition depending on the
 263 sample and extraction method used (Whitehouse and Hottel, 2007). Our
 264 assay successfully amplified soil extracts spiked with *Y. pestis* although we
 265 did observe an approximately 10-fold reduction in amplification efficiency that
 266 persisted despite the addition of BSA to the reaction.

267

268 Discussion

269 Although many PCR-based assays exist for *Y. pestis*, none combines species
 270 specificity with accurate quantification. The *fur*MAMA assay is designed to
 271 specifically amplify a fragment of the *fur* gene in *Y. pestis* while preventing
 272 amplification of the gene fragment in other members of the *Yersinia* genus
 273 including its close genetic neighbor *Y. pseudotuberculosis* where the gene
 274 fragment only differs by a single SNP. The SNP in the *fur* gene fragment
 275 targeted by our *fur*MAMA assay is private to, and ubiquitous in all *Y. pestis*.
 276 Specificity is acquired in a single assay through allele-specific amplification of
 277 this SNP in a key virulence gene, *fur*, whose presence is found throughout
 278 *Enterobacteriaceae* without the need for further molecular targets (Fig. 1).
 279 Leveraging a specific SNP within a gene, as done here, is an efficient and
 280 robust method for specific detection of a given bacterial pathogen and can aid
 281 in pathogen detection in metagenomes using next generation sequencing
 282 where close relatives are present in high numbers (Valseth et al., 2017). This

283 circumvents the present issues of specificity when targeting pathogens with
284 high degree of genetic similarity. A recent large-scale comparative study of *Y.*
285 *pestis* and *Y. pseudotuberculosis* genomes showed that fewer genes than
286 previously reported in smaller studies were unique to *Y. pestis* (Califf et al.,
287 2015). However, both specific combinations, variants of genes and SNPs, the
288 last example demonstrated here, will be unique to this slowly mutating
289 pathogen (Cui et al., 2013).

290

291 The original assay published by Gabitzsch et al. (Gabitzsch et al., 2008) was
292 developed to quantify *Y. pestis* in fleas for their work on flea transmission in a
293 laboratory experimental setup and not initially aimed at diagnostics or
294 quantification in field-collected samples. They tested the specificity by running
295 PCR on reference DNA from 28 species, including *E. coli* and other gram-
296 negative bacteria, and reported cross-reactivity for 6 *Yersinia* species,
297 including *Y. enterocolitica*. Surprisingly, they did not report cross-reactivity
298 with *Y. pseudotuberculosis*. BLAST of the derived *fur* gene fragment from the
299 Gabitzsch assay resulted in 99% overall homology with *Y. pseudotuberculosis*
300 sequence and high homology between other *Yersinia* spp. Indeed, we found
301 the original assay amplifies both *Y. pestis* and *Y. pseudotuberculosis* with
302 equal efficiency (Fig. 2 (A)). In contrast, the amplification failure of *Y.*
303 *pseudotuberculosis* with the YpfurR_MAMA primer establishes our assay's
304 ability to distinguish between the two highly genetically similar pathogens, as
305 well as the ability to quantify *Y. pestis* over a broad range of concentrations
306 (Fig. 2(D and C)). The *Y. pestis* SNP in the *fur* gene is universally present in
307 all available *Y. pestis* genomes, which limits the probability of false negative

308 results. Conversely, the absence of the mutation in all other *Yersinia* and
 309 *Enterobacteriaceae* reduce the likelihood of false positive results. Our assay
 310 presented false positives and very low-level cross-reactivity at the highest
 311 concentration of *Y. pseudotuberculosis* DNA produced from whole genome
 312 amplification (WGA). The lack of access to a larger collection of *Y.*
 313 *pseudotuberculosis* prohibited us from further testing of cross-reactivity but
 314 based on the specificity testing *in silico* and by targeting a conserved SNP in
 315 the *Y. pestis* genome we are confident this will not be a major problem.

316

317 **Conclusions**

318 Our assay provides an alternative to the longstanding culturing method and
 319 the immunoassays by providing a highly sensitive and rapid way of both
 320 specifically detecting and quantifying *Y. pestis* in tissue samples. To our
 321 knowledge, this is the first assay to target a species-specific SNP in a
 322 chromosomal gene of *Y. pestis* with the aim of detecting and quantifying the
 323 amount of the bacteria in tissue samples.

324

325 **Materials and Methods**

326 **Bacterial strains and DNA extractions**

327 *Y. pestis* strain Az-26 (1102) was isolated from an organ suspension
 328 of *Meriones vinogradovi* in 1969, 1.5 km southeast of Sirab village in the
 329 Nakhchivan Autonomous Republic of Azerbaijan. The culture has been stored
 330 at the Republican Anti-Plague Station laboratory in Baku, Azerbaijan. The
 331 culture became non-viable, yet it was still possible to isolate DNA from the

332 material in 2012 using Qiagen DNeasy® Blood and Tissue kit following
333 extraction protocol for gram-negative bacteria.
334 *Y. pseudotuberculosis* Type III strain was donated by Jack C. Leo and Dirk
335 Linke (University of Oslo). DNA was isolated from broth culture using Qiagen
336 Blood and Tissue kit following manufacturer's instructions (Qiagen Inc., USA).

337

338 **Rodent DNA**

339 No animals were killed for the purpose of this study. We procured samples of
340 rodent DNA for this study by reaching out to both internal and external
341 colleagues with existing DNA samples stored. The DNA samples stem from
342 unpublished work as detailed below.

343 The great gerbil DNA was provided by colleagues in China where the male
344 individual was captured as part of work carried out in the Junggar Basin of
345 Xinjiang Province, China (unpublished).

346 The C57BL6 mouse DNA stem from work conducted at the Institute of
347 Immunology at Oslo University Hospital, Rikshospitalet, Norway
348 (unpublished). The animal was originally obtained from Janvier labs
349 ([https://www.janvier-labs.com/rodent-research-models-services/research-](https://www.janvier-labs.com/rodent-research-models-services/research-models/per-species/inbred-mice/product/c57bl6jrl.html)
350 [models/per-species/inbred-mice/product/c57bl6jrl.html](https://www.janvier-labs.com/rodent-research-models-services/research-models/per-species/inbred-mice/product/c57bl6jrl.html)).

351 The bank vole DNA belongs to the EcoTick-project on tick-/rodent-borne
352 diseases conducted at the University of Oslo (unpublished).

353

354 **Whole genome amplification (WGA) and standard curves**

355 Whole genome amplification was performed on both bacterial strains using
356 the REPLI-g Mini Kit (Qiagen) following the manufacturer's instructions for

subsequent creation of a dynamic range of DNA starting at high concentrations for standard curves. DNA concentrations after WGA were determined using the Qubit dsDNA BR Assay kit (molecular probes, Life Technologies) and the Qubit 2.0 fluorometer. The standard curves were generated from that starting concentration as serial 10-fold dilutions in TE buffer in the following concentrations: 5 ng, 500 pg, 50 pg, 5 pg, 500 fg, 50 fg and 5 fg.

364

365 **TaqMAMA Primers and Probe design**

Forward primer and probe were identical to oligos designed by Gabitzsch et al. (Gabitzsch et al., 2008) while the reverse primer was modified to a MAMA primer to specifically amplify *Y. pestis*. This required the ultimate 3' base to be complementary to the *Y. pestis* SNP while the penultimate 3' base was designed to mismatch with the shared sequence between *Y. pestis* and *Y. pseudotuberculosis* (underlined in Table 2). Several reverse MAMA primers were designed with different nucleotide mismatches at the 3' penultimate position. These were tested for specificity and amplification efficiency before ultimately choosing the YpfurR_MAMA primer.

375

376 **Table 1. List of Primers and probe sequences assayed in qPCR and PCR**
377 **reactions**

Primer/Probe	Sequence (5'–3')	Product (bp)*
YpfurF	TCTGGAAGTGGTGCAAAATCCTG	
YpfurP	FAM-TGTCACCACGTCAGCGCGGAAGAT-BAMQ1	
YpfurR	AAGCCAATCTCTTCACCAATATCG	90

YpfurR_MAMA	CCAATCTCTTCACCAATATCGATCAGAA	87
-------------	------------------------------	----

* The resulting PCR product size when combining the YpfurF (forward primer) with the respective reverse primer.

380

381 **TaqMAMA PCR protocol**

382 For all experiments, PCR was conducted in 96-well plates (Roche
383 LightCycler® 480 Multiwell Plate 96) using 10-μL reactions that contained final
384 concentrations of 600 nM of both forward and reverse primers, 250 nM probe,
385 1x Express qPCR Supermix (Invitrogen, by Life Technologies)) and 1 μL
386 template DNA. Real-time PCR was performed on a LightCycler® 96
387 Instrument (Roche) as a two-step PCR with the following conditions: 95 °C for
388 2 min and 50 cycles of 95 °C for 15 s and 64 °C for 60 s.

389

390 **Specificity tests**

391 Given limited access to a large strain collection of *Yersinia*, specificity was
392 largely assessed using bioinformatics (BLAST) of available sequence data
393 (see “Bioinformatics” below). Therefore, the specificity and kinetics of the
394 assay was done through analyzing genomic DNA from a representative of
395 both alleles (see Fig. 1); *Y. pestis* strain Az-26 (1102) and *Y.*
396 *pseudotuberculosis* Type III strain running both original reverse primer
397 (YpfurR) and the modified MAMA primer (YpfurR_MAMA) in separate, parallel
398 reactions over a dynamic range of DNA concentrations (5 ng to 5 fg).

399

400 **Sensitivity tests**

401 Sensitivity of the furMAMA assay was determined by analyzing serial 10-fold
402 dilutions of genomic DNA from the *Y. pestis* strain Az-26 (1102) ranging from
403 5 ng to 5 fg with the lowest dilution being equivalent to one bacterial genome.

404

405 **Testing applicability on tissue and environmental samples**

406 The applicability of the assay on rodent tissue samples was tested by running,
407 in parallel, DNA extractions of liver and spleen from a great gerbil
408 (*Rhombomys opimus*), DNA from the tail of a C57BL6 mouse (*Mus musculus*)
409 and DNA extracted from the ear of a Bank vole (*Myodes glareolus*) with and
410 without spike-in of *Y. pestis* DNA standard curve. Real-time PCR was
411 conducted in 96-well plates (Roche LightCycler® 480 Multi-well Plate 96)
412 using 10-µL reactions that contained final concentrations of 600 nM of both
413 forward and reverse primer, 250 nM probe, 1x Express qPCR Supermix
414 (Invitrogen, by Life Technologies)), 1 µL template DNA and for the spiked
415 samples an additional 1 µL DNA from one of the standard curve
416 concentrations.

417 The ability of the assay to detect *Y. pestis* despite the inhibitory effects often
418 seen in environmental samples, was tested by using DNA extractions of soil
419 samples diluted 1:10. The soil samples were run in parallel with non-
420 acetylated Bovine Serum Albumin (New England Biolabs) added, with or
421 without DNA from the *Y. pestis* standard curve in the following manner 1) soil
422 + BSA, 2) soil + standard curve and 3) soil + standard curve + BSA. Real-time
423 PCR was conducted as above with the exception that the reactions containing
424 BSA did so in a final concentration of 1mg/ml.

425

Detection using standard PCR

The primers were also tested with standard PCR to determine their applicability as a pure detection-based assay. PCR was conducted in 8-well PCR strips in 10-μL reactions that contained a final concentration of 600 nM both forward and reverse primer, 1x Express qPCR Supermix and 1 μL template DNA. PCR was performed on a MJ Research PTC-200 Peltier Thermal Cycler Instrument as a two-step PCR with the following conditions: 94 °C for 2 min and 40 cycles of 94 °C for 15 s and 64 °C for 60 s. Successful amplification was confirmed by running a 3% agarose gel stained with GelRed (Biotium) at 75 V for 1 hr and 20 min in TAE buffer.

Bioinformatics

Specificity of the SNP was assessed across available *Yersinia* and other related *Enterobacteriaceae* genomes by performing both megaBLAST and BLASTn searches of the PCR fragment and the whole *fur* gene from *Y. pestis* CO92 (Accession no. NC_003143.1, locus_tag=YPO2634) as query in the NCBI database using default parameters. BLAST hits were inspected to confirm which allele was present in the sequence. Finally, the presence of the *Y. pestis* allele in the strain used in this study (Az-26 (1102)) was confirmed by a BLASTn search using the *fur* gene from *Y. pestis* CO92 as query against a BLAST database generated from the Az-26 (1102) genome assembly. From these data an alignment was made using representative sequences from BLAST searches through Geneious (<https://www.geneious.com>) and aligned using Jalview (Waterhouse et al., 2009).

451 **Abbreviations**

452 Fur gene: Ferric uptake regulator
 453 BSA: Bovine serum albumin
 454 Caf1: Capsule antigen fraction 1
 455 Cq: Quantification cycle
 456 HGT: Horizontal gene transfer
 457 IcrV: Low calcium response V antigen
 458 Pla: Plasminogen activator
 459 SNP: Single nucleotide polymorphism
 460 TAE: Tris-acetate-EDTA
 461 TaqMAMA: TaqMan® mismatch amplification mutation assay
 462 TE: Tris-EDTA
 463 WGA: Whole genome amplification
 464 yihN: Inner membrane protein yihN
 465 Ymt: Yersinia murine toxin

466

467 **Declarations**

468 **Acknowledgements**

469 We would like to thank Jack C. Leo and Dirk Linke at the Department of
 470 Biosciences, University of Oslo for providing the *Y. pseudotuberculosis* Type
 471 III strain. For the use of the great gerbil DNA, the C57BL6 mouse DNA and
 472 the bank vole DNA we would like to thank Ruifu Yang (Beijing Institute of
 473 Microbiology and Epidemiology, China) and Yujiang Zhang (Xinjiang CDC,
 474 China), Shuo-Wang Qiao (Institute of Immunology at Oslo University Hospital,
 475 Rikshospitalet, Norway) and Atle Mysterud (University of Oslo, Norway). We

476 would also like to thank Bradd Haley for bioinformatics support. Lastly, we
477 would like to thank Monica H. Solbakken and Boris V. Schmid for helpful
478 comments on the manuscript.

479 **Authors contributions**

480 WRE and PN conceived and planned the experiments and analyzed the data.
481 PN carried out the experiments and performed bioinformatics. SG procured
482 and isolated the *Y. pestis* strain Az-26 (1102) for this study. The manuscript
483 was written with input from all authors. All authors read and approved the final
484 manuscript.

485 **Availability of data and materials**

486 The *Y. pestis* strain analyzed during this study (Az-26(1102)) is not publicly
487 available due to said strain being non-viable at isolation. The genome
488 sequence is available from the corresponding author upon reasonable
489 request.

490 **Consent for publication**

491 Not applicable.

492 **Competing interests:**

493 The authors declare that they have no competing interests.

494 **Ethics approval and consent to participate**

495 The DNA samples used in this study stems from existing samples stored by
496 internal and external colleagues and hence the appropriate approvals were
497 previously obtained as specified below.

498 The use of great gerbil tissue was approved by the Committee for Animal
499 Welfares of Xinjiang CDC, China.

500 The bank vole was captured as part of EcoTick-project on tick-/rodent-borne
501 diseases led by Atle Mysterud (University of Oslo) and permission for trapping
502 rodents were given by the Norwegian Environmental Agency (ref 2013/11201)
503 and hence conform to the Norwegian laws and regulations.
504 The C57BL6 mouse was used in a project at the Institute of Immunology at
505 Oslo University Hospital, Rikshospitalet, Norway and is subject to protocol
506 number 3775 from the Norwegian Food Safety Authority. This agency
507 evaluates all applications and approvals for ethical handling of animals and
508 animal experiments in Norway.

509 **Funding:**

510 P. N. is supported by a Molecular Life Sciences (MLS) grant (now Uio: Life
511 Science) from the University of Oslo as well as through Colloquia 3 at the
512 Centre for Ecological and Evolutionary Synthesis (CEES) at the University of
513 Oslo. W. R. E. is supported by Colloquia 3 at CEES at the University of Oslo.
514 The funding bodies had no role in the design, analysis or interpretation of the
515 results nor were they involved in writing the manuscript.

516

517 **References**

- 518 Abbaszadegan, M.R., Struewing, J.P., Brown, K.M., Snider, J.V., Goodsaid,
519 F., Gore-Langton, R., Hughes, M.R., 2009. Automated Detection of
520 Prevalent Mutations in BRCA1 and BRCA2 Genes, Using a Fluorogenic
521 PCR Allelic Discrimination Assay. <http://www.liebertpub.com/gte> 1, 171–
522 180. doi:10.1089/gte.1997.1.171
523 Achtman, M., 2008. Evolution, population structure, and phylogeography of
524 genetically monomorphic bacterial pathogens. *Annu. Rev. Microbiol.* 62,
525 53–70. doi:10.1146/annurev.micro.62.081307.162832
526 Achtman, M., Zurth, K., Morelli, G., Torrea, G., Guiyoule, A., Carniel, E., 1999.
527 *Yersinia pestis*, the cause of plague, is a recently emerged clone of
528 *Yersinia pseudotuberculosis*. *Proceedings of the National Academy of*
529 *Sciences* 96, 14043–14048. doi:10.1073/pnas.96.24.14043
530 Andrianainarimanana, V., Telfer, S., Rajerison, M., Ranjalahy, M.A.,
531 Andriamiarimanana, F., Rahaingosoamamitiana, C., Rahalison, L.,

Jambou, R., 2012. Immune responses to plague infection in wild *Rattus rattus*, in Madagascar: a role in foci persistence? *PLoS ONE* 7, e38630. doi:10.1371/journal.pone.0038630

Anisimov, A.P., Lindler, L.E., Pier, G.B., 2004. Intraspecific Diversity of *Yersinia pestis*. *Clinical Microbiology Reviews* 17, 434–464. doi:10.1128/CMR.17.2.434-464.2004

Armougom, F., Bitam, I., Croce, O., Merhej, V., Barassi, L., Nguyen, T.-T., La Scola, B., Raoult, D., 2016. Genomic Insights into a New *Citrobacter koseri* Strain Revealed Gene Exchanges with the Virulence-Associated *Yersinia pestis* pPCP1 Plasmid. *Front Microbiol* 7, 340. doi:10.3389/fmicb.2016.00340

Ber, R., Mamroud, E., Aftalion, M., Tidhar, A., Gur, D., Flashner, Y., Cohen, S., 2003. Development of an improved selective agar medium for isolation of *Yersinia pestis*. *Appl. Environ. Microbiol.* 69, 5787–5792. doi:10.1128/AEM.69.10.5787-5792.2003

Bevins, S.N., Baroch, J.A., Nolte, D.L., Zhang, M., He, H., 2012. *Yersinia pestis*: examining wildlife plague surveillance in China and the USA. *Integrative Zoology* 7, 99–109. doi:10.1111/j.1749-4877.2011.00277.x

Califf, K.J., Keim, P.S., Wagner, D.M., Sahl, J.W., 2015. Redefining the differences in gene content between *Yersinia pestis* and *Yersinia pseudotuberculosis* using large-scale comparative genomics. *Microb Genom* 1, e000028. doi:10.1099/mgen.0.000028

Cha, R.S., Zarbl, H., Keohavong, P., Thilly, W.G., 1992. Mismatch amplification mutation assay (MAMA): application to the c-H-ras gene. *PCR Methods Appl.* 2, 14–20.

Chain, P.S.G., Carniel, E., Larimer, F.W., Lamerdin, J., Stoutland, P.O., Regala, W.M., Georgescu, A.M., Vergez, L.M., Land, M.L., Motin, V.L., Brubaker, R.R., Fowler, J., Hinnebusch, J., Marceau, M., Medigue, C., Simonet, M., Chenal-Francisque, V., Souza, B., Dacheux, D., Elliott, J.M., Derbise, A., Hauser, L.J., Garcia, E., 2004. Insights into the evolution of *Yersinia pestis* through whole-genome comparison with *Yersinia pseudotuberculosis*. *Proceedings of the National Academy of Sciences* 101, 13826–13831. doi:10.1073/pnas.0404012101

Chanteau, S., Rahalison, L., Ralafiarisoa, L., Foulon, J., Ratsitorahina, M., Ratsifasoamanana, L., Carniel, E., Nato, F., 2003. Development and testing of a rapid diagnostic test for bubonic and pneumonic plague. *The Lancet* 361, 211–216. doi:10.1016/S0140-6736(03)12270-2

Cui, Y., Yu, C., Yan, Y., Li, D., Li, Y., Jombart, T., Weinert, L.A., Wang, Z., Guo, Z., Xu, L., Zhang, Y., Zheng, H., Qin, N., Xiao, X., Wu, M., Wang, X., Zhou, D., Qi, Z., Du, Z., Wu, H., Yang, X., Cao, H., Wang, H., Wang, J., Yao, S., Rakin, A., Li, Y., Falush, D., Balloux, F., Achtman, M., Song, Y., Wang, J., Yang, R., 2013. Historical variations in mutation rate in an epidemic pathogen, *Yersinia pestis*. *Proceedings of the National Academy of Sciences* 110, 577–582. doi:10.1073/pnas.1205750110

Dennis, D.T., Gage, K.L., Gratz, N.G., Poland, J.D., Tikhomirov, E., 1999. *Plague manual: epidemiology, distribution, surveillance and control*. World Health Organisation, Geneva, Switzerland.

Easterday, W.R., Van Ert, M.N., Zanecki, S., Keim, P., 2005. Specific detection of *Bacillus anthracis* using a TaqMan mismatch amplification mutation assay. *Biotechniques*.

582 Eppinger, M., Worsham, P.L., Nikolich, M.P., Riley, D.R., Sebastian, Y., Mou,
583 S., Achtman, M., Lindler, L.E., Ravel, J., 2010. Genome sequence of the
584 deep-rooted *Yersinia pestis* strain Angola reveals new insights into the
585 evolution and pangenome of the plague bacterium. *Journal of*
586 *Bacteriology* 192, 1685–1699. doi:10.1128/JB.01518-09
587 Gabitzsch, E.S., Vera-Tudela, R., Eisen, R.J., Bearden, S.W., Gage, K.L.,
588 Zeidner, N.S., 2008. Development of a real-time quantitative PCR assay
589 to enumerate *Yersinia pestis* in fleas. *Am J Trop Med Hyg* 79, 99–101.
590 Glaab, W.E., Skopek, T.R., 1999. A novel assay for allelic discrimination that
591 combines the fluorogenic 5' nuclease polymerase chain reaction
592 (TaqMan®) and mismatch amplification mutation assay. *Mutation*
593 *Research/Fundamental and Molecular Mechanisms of Mutagenesis* 430,
594 1–12. doi:10.1016/S0027-5107(99)00147-5
595 Gupta, V., Gulati, P., Bhagat, N., Dhar, M.S., Viridi, J.S., 2015. Detection of
596 *Yersinia enterocolitica* in food: an overview. *Eur. J. Clin. Microbiol. Infect.*
597 *Dis.* 34, 641–650. doi:10.1007/s10096-014-2276-7
598 Harbeck, M., Seifert, L., Hänsch, S., Wagner, D.M., Birdsell, D., Parise, K.L.,
599 Wiechmann, I., Grupe, G., Thomas, A., Keim, P., Zöller, L., Bramanti, B.,
600 Riehm, J.M., Scholz, H.C., 2013. *Yersinia pestis* DNA from Skeletal
601 Remains from the 6 th Century AD Reveals Insights into Justinianic
602 Plague. *PLoS Pathog.* 9, e1003349. doi:10.1371/journal.ppat.1003349
603 Hänsch, S., Cilli, E., Catalano, G., Gruppioni, G., Bianucci, R., Stenseth, N.C.,
604 Bramanti, B., Pallen, M.J., 2015. The *pla* gene, encoding plasminogen
605 activator, is not specific to *Yersinia pestis*. *BMC Research Notes* 2015 8:1
606 8, 535. doi:10.1186/s13104-015-1525-x
607 Hinnebusch, B.J., Chouikha, I., Sun, Y.-C., 2016. Ecological Opportunity,
608 Evolution, and the Emergence of Flea-Borne Plague. *Infection and*
609 *Immunity* 84, 1932–1940. doi:10.1128/IAI.00188-16
610 Hu, P., Elliott, J., McCready, P., Skowronski, E., Garnes, J., Kobayashi, A.,
611 Brubaker, R.R., Garcia, E., 1998. Structural organization of virulence-
612 associated plasmids of *Yersinia pestis*. *Journal of Bacteriology* 180, 5192–
613 5202.
614 Iqbal, S.S., Chambers, J.P., Goode, M.T., Valdes, J.J., Brubaker, R.R., 2000.
615 Detection of *Yersinia pestis* by pesticin fluorogenic probe-coupled PCR.
616 *Molecular and Cellular Probes* 14, 109–114. doi:10.1006/mcpr.2000.0295
617 Juhas, M., 2015. Horizontal gene transfer in human pathogens. *Crit. Rev.*
618 *Microbiol.* 41, 101–108. doi:10.3109/1040841X.2013.804031
619 Lärkeryd, A., Myrtenäs, K., Karlsson, E., Dwibedi, C.K., Forsman, M.,
620 Larsson, P., Johansson, A., Sjödin, A., 2014. CanSNPer: a hierarchical
621 genotype classifier of clonal pathogens. *Bioinformatics* 30, 1762–1764.
622 doi:10.1093/bioinformatics/btu113
623 Melikishvili, A., 2006. Genesis of the anti-plague system: the Tsarist period.
624 *Crit. Rev. Microbiol.* 32, 19–31. doi:10.1080/10408410500496763
625 Neubauer, H., Meyer, H., Prior, J., Aleksic, S., Hensel, A., Splettstösser, W.,
626 2000. A combination of different polymerase chain reaction (PCR) assays
627 for the presumptive identification of *Yersinia pestis*. *J. Vet. Med. B Infect.*
628 *Dis. Vet. Public Health* 47, 573–580.
629 Parkhill, J., Wren, B.W., Thomson, N.R., Titball, R.W., Holden, M.T.G.,
630 Prentice, M.B., Sebahia, M., James, K.D., Churcher, C., Mungall, K.L.,
631 Baker, S., Basham, D., Bentley, S.D., Brooks, K., Cerdeño-Tárraga, A.M.,

Chillingworth, T., Cronin, A., Davies, R.M., Davis, P., Dougan, G., Feltwell, T., Hamlin, N., Holroyd, S., Jagels, K., Karlyshev, A.V., Leather, S., Moule, S., Oyston, P.C.F., Quail, M., Rutherford, K., Simmonds, M., Skelton, J., Stevens, K., Whitehead, S., Barrell, B.G., 2001. Genome sequence of *Yersinia pestis*, the causative agent of plague. *Nature* 413, 523–527. doi:10.1038/35097083

Perry, R.D., Fetherston, J.D., 1997. *Yersinia pestis* - Etiologic agent of plague. *Clinical Microbiology Reviews* 10, 35–66.

Portnoy, D.A., Wolf-Watz, H., Bolin, I., Beeder, A.B., Falkow, S., 1984. Characterization of common virulence plasmids in *Yersinia* species and their role in the expression of outer membrane proteins. *Infection and Immunity* 43, 108–114.

Riehm, J.M., Rahalison, L., Scholz, H.C., Thoma, B., Pfeffer, M., Razanakoto, L.M., Dahouk, A.I., Neubauer, H., Tomaso, H., 2011. Detection of *Yersinia pestis* using real-time PCR in patients with suspected bubonic plague. *Molecular and Cellular Probes* 25, 8–12. doi:10.1016/j.mcp.2010.09.002

Sarovich, D.S., Colman, R.E., Price, E.P., Chung, W.K., Lee, J., Schupp, J.M., Cobble, K.R., Busch, J.D., Alexander, J., Keim, P., Wagner, D.M., 2010. Selective isolation of *Yersinia pestis* from plague-infected fleas. *J. Microbiol. Methods* 82, 95–97. doi:10.1016/j.mimet.2010.03.019

Simon, S., Demeure, C., Lamourette, P., Filali, S., Plaisance, M., Créminon, C., Volland, H., Carniel, E., 2013. Fast and simple detection of *Yersinia pestis* applicable to field investigation of plague foci. *PLoS ONE* 8, e54947. doi:10.1371/journal.pone.0054947

Stewart, A., Satterfield, B., Cohen, M., O'Neill, K., Robison, R., 2008. A quadruplex real-time PCR assay for the detection of *Yersinia pestis* and its plasmids. *Journal of Medical Microbiology* 57, 324–331. doi:10.1099/jmm.0.47485-0

Tomaso, H., Jacob, D., Eickhoff, M., Scholz, H.C., Dahouk, A.I., Kattar, M.M., Reischl, U., Plicka, H., Olsen, J.S., Nikkari, S., Matero, P., Beuret, C., Ciammaruconi, A., Lista, F., Gala, J.-L., Broll, H., Appel, B., Sellek Cano, R.E., del Carmen Ybarra de Villavicencio, M., Broekhuijsen, M., Indra, A., Petersen, R., Neubauer, H., 2008. Preliminary validation of real-time PCR assays for the identification of *Yersinia pestis*. *Clin. Chem. Lab. Med.* 46, 1239–1244. doi:10.1515/CCLM.2008.251

Tomaso, H., Reisinger, E.C., Dahouk, S., Frangoulidis, D., Rakin, A., Landt, O., Neubauer, H., 2003. Rapid detection of *Yersinia pestis* with multiplex real-time PCR assays using fluorescent hybridisation probes. *Pathogens and Disease* 38, 117–126. doi:10.1016/S0928-8244(03)00184-6

Turner, W.C., Kausrud, K.L., Beyer, W., Easterday, W.R., Barandongo, Z.R., Blaschke, E., Cloete, C.C., Lazak, J., Van Ert, M.N., Ganz, H.H., Turnbull, P.C.B., Stenseth, N.C., Getz, W.M., 2016. Lethal exposure: An integrated approach to pathogen transmission via environmental reservoirs. *Sci Rep* 6, 27311. doi:10.1038/srep27311

U'Ren, J.M., Van Ert, M.N., Schupp, J.M., Easterday, W.R., Simonson, T.S., Okinaka, R.T., Pearson, T., Keim, P., 2005. Use of a real-time PCR TaqMan assay for rapid identification and differentiation of *Burkholderia pseudomallei* and *Burkholderia mallei*. *Journal of Clinical Microbiology* 43, 5771–5774. doi:10.1128/JCM.43.11.5771-5774.2005

682 Valseth, K., Nesbø, C.L., Easterday, W.R., Turner, W.C., Olsen, J.S.,
683 Stenseth, N.C., Haverkamp, T.H.A., 2017. Temporal dynamics in
684 microbial soil communities at anthrax carcass sites. *BMC Microbiology*
685 2008 8:1 17, 206. doi:10.1186/s12866-017-1111-6
686 Wang, H., Avican, K., Fahlgren, A., Erttmann, S.F., Nuss, A.M., Dersch, P.,
687 Fallman, M., Edgren, T., Wolf-Watz, H., 2016. Increased plasmid copy
688 number is essential for *Yersinia* T3SS function and virulence. *Science*
689 353, 492–495. doi:10.1126/science.aaf7501
690 Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., Barton, G.J.,
691 2009. Jalview Version 2--a multiple sequence alignment editor and
692 analysis workbench. *Bioinformatics* 25, 1189–1191.
693 doi:10.1093/bioinformatics/btp033
694 Whitehouse, C.A., Hottel, H.E., 2007. Comparison of five commercial DNA
695 extraction kits for the recovery of *Francisella tularensis* DNA from spiked
696 soil samples. *Molecular and Cellular Probes* 21, 92–96.
697 doi:10.1016/j.mcp.2006.08.003
698 World Health Organization, 2017. Plague [WWW Document].
699 <http://www.who.int/news-room/fact-sheets/detail/plague>. URL
700 <http://www.who.int/news-room/fact-sheets/detail/plague> (accessed
701 5.29.18).
702 Woron, A.M., Nazarian, E.J., Egan, C., McDonough, K.A., Cirino, N.M.,
703 Limberger, R.J., Musser, K.A., 2006. Development and evaluation of a 4-
704 target multiplex real-time polymerase chain reaction assay for the
705 detection and characterization of *Yersinia pestis*. *Diagn. Microbiol. Infect.*
706 Dis. 56, 261–268. doi:10.1016/j.diagmicrobio.2006.06.009
707