Mutations in *Kinesin Family Member 6* Reveal Specific Role in Ependymal Cell Function and Human Neuro-Cranial Development

Mia J. Konjikusic^{1,2}, Patra Yeetong^{3,4,5}, Rungrana Ittiwut^{3,4}, Kanya Suphateetiporn^{3,4}, John B. Wallingford², Christina A. Gurnett⁶, Vorasuk Shotelersuk^{3,4,*}, and Ryan S. Gray^{1,*}

¹ Department of Pediatrics, Dell Pediatric Research Institute, 1400 Barbara Jordan Blvd, The University of Texas at Austin, Dell Medical School, Austin, Texas 78723, USA.

² Department of Molecular Biosciences, Patterson Labs, 2401 Speedway, The University of Texas at Austin, Austin, Texas 78712, USA.

³ Center of Excellence for Medical Genetics, Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

⁴ Excellence Center for Medical Genetics, King Chulalongkorn Memorial Hospital, the Thai Red Cross Society, Bangkok 10330, Thailand.

⁵ Division of Human Genetics, Department of Botany, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

⁶ Department of Neurology, Division Pediatric Neurology, Washington University School of Medicine, 660 S Euclid Ave, St Louis, MO 63110, USA.

*corresponding authors: Ryan S. Gray: ryan.gray@austin.utexas.edu or Vorasuk Shotelersuk: vorasuk.s@chula.ac.th

Keywords: Kinesin, Ependymal cell cilia, intellectual disability

Short running title: KIF6 Specifically Regulates Ependymal Cell Function

ABSTRACT

Cerebrospinal fluid flow is crucial for neurodevelopment and homeostasis of the ventricular system of the brain, with localized flow being established by the polarized beating of the ependymal cell (EC) cilia. Here, we report a homozygous one base-pair deletion, c.1193delT (p.Leu398Glnfs*2), in the *Kinesin Family Member 6 (KIF6)* gene in a child displaying neurocranial defects and intellectual disability. To test the pathogenicity of this novel human *KIF6* mutation we engineered an analogous C-terminal truncating mutation in mouse. These mutant mice display severe, postnatal-onset hydrocephalus. We generated a *Kif6-LacZ* transgenic mouse strain and report expression specifically and uniquely within the ependymal cell (EC) layer of the brain, without labeling other multiciliated mouse tissues. Analysis of *Kif6* mutant mice with scanning electron microscopy (SEM) and immunofluorescence (IF) revealed a reduction in EC cilia, without effect on other multiciliated tissues. Consistent with our findings in mice, defects of the ventricular system and EC cilia were observed in *kif6* mutant zebrafish. Overall, this work describes the first clinically-defined *KIF6* homozygous null mutation in human and defines KIF6 as a conserved mediator of neuro-cranial morphogenesis with a specific role in the maintenance of EC cilia in vertebrates.

AUTHOR SUMMARY

Cerebrospinal fluid flow is crucial for neurodevelopment and homeostasis of the ventricular system of the brain. Localized flows of cerebrospinal fluid throughout the ventricular system of the brain are established from the polarized beating of the ependymal cell (EC) cilia. Here, we identified a homozygous truncating mutation in *KIF6* in a child displaying neuro-cranial defects and intellectual disability. To test the function of *KIF6* *in vivo*, we engineered mutations of *Kif6* in mouse. These *Kif6* mutant mice display severe hydrocephalus, coupled with a loss of EC cilia. Similarly, we observed hydrocephalus and a reduction in EC cilia in *kif6* mutant zebrafish. Overall, this work describes the first clinically-defined *KIF6* mutation in human, while our animal studies demonstrate the pathogenicity of mutations in *KIF6* and establish *KIF6* as a conserved mediator of neuro-cranial development and EC cilia maintenance in vertebrates.

INTRODUCTION

The delicate balance of cerebrospinal fluid (CSF) production and flow is important for the morphogenesis and function of the brain development and homeostasis. CSF circulation in human is largely due to gradients established by the secretion of CSF from the choroid plexuses, and its resorption at the arachnoid granulations [1]. The clinical significance of CSF stasis includes hydrocephalus and intracranial hypertension. Moreover, severely diminished CSF flow combined with increased intracranial pressure can secondarily cause ventriculomegaly, cognitive impairment, as well as degenerative and age-related dementias [2]. For these reasons, the identification of genetic risk factors involved in the pathogenesis of CSF stasis is critical for the development of genetic diagnostics and early interventions for these disorders.

One element for circulation of CSF is the multiciliated ependymal cells (ECs), which are specialized glial cells covering the ventricular walls of the brain and spinal canal [3]. In contrast, to primary cilia which are single, immotile cellular organelles extending from most cell types, ECs contain dozens of apically-arranged motile cilia, which beat in a polarized fashion to generate localized or near-wall CSF flows [4]. Defective EC cilia or loss of their polarized beating causes a disruption of this localized CSF flow leading to increased intracranial pressure, and dilation of ventricles and hydrocephalus in mice [5-8]. Importantly, this EC cilia-driven CSF flow is crucial for regulating brain function and adult neurogenesis [4, 9].

Impaired ciliary motility due to disruptions of the key kinesins, dyneins, and intraflagellar components necessary in most or all cilia, results in a syndromic condition known as primary ciliary dyskinesia (PCD) in humans [10, 11]. While hydrocephalus can occur in some PCD patients, it is a less common manifestation of the disease in humans [11]. In contrast, genes implicated in PCD or mutations which disrupt the structure or motility of all motile cilia are

strongly correlated with hydrocephalus in mouse [8]. Alternatively, some hydrocephalus in mice with dysfunctional cilia may be the result of altered function of the choroid plexus, prior to the onset of cilia-driven CSF flow [7].

KIF6 (Kinesin family member 6, MIM: 613919) encodes a member of the kinesin-9 superfamily of microtubules motor proteins which act predominately as "plus-end" directed molecular motors that generate force and movement across microtubules [12]. Kinesins are critical for numerous cellular functions such as intracellular transport and cell division, as well as for building and maintaining the cilium in a process known as intraflagellar transport [13]. During this process, kinesins have been shown to transport cargo within the ciliary axoneme [14], establish motility and compartmentalization of the axoneme [15], or to facilitate plus-end directed microtubule disassembly and control of axonemal length [16]. As such, multiple kinesins have shown to be associated with monogenic disorders affecting a wide-spectrum of tissues, with several modes of inheritance (www.omim.org). Interestingly, *KIF6* has previously been proposed as locus for susceptibility to coronary heart disease [17], while other studies did not substantiate this association [18]. We previously reported that *kif6* mutant zebrafish are adult viable exhibiting larval-onset scoliosis without obvious heart defects [19]. Because of these conflicting results, and a lack of relevant mouse models, the role of *KIF6* in human disease remains an open question.

Here, we present a patient with consanguineous parents, presenting with neuro-cranial defects and intellectual disability. Homozygosity mapping followed by whole-exome sequencing (WES) identified a novel homozygous frameshift mutation in *KIF6* which is predicted to result in the truncation of the C-terminal cargo-binding domain of the kinesin motor protein. We generated an analogous frameshift mutation in the mouse and found that these mutant mice

displayed progressive, postnatal-onset hydrocephalus with cranial expansion, coupled with a reduction in the quantity of EC cilia. In addition, we observed that *kif6* mutant zebrafish also display dilation of the ventricular system, coupled with reduced EC cilia. Interestingly, we failed to observe additional PCD related defects of other multiciliated tissues in *Kif6* mutant mouse or zebrafish models. Together these results demonstrate that KIF6 function is unique and specific for EC cilia. Finally, we propose that *KIF6* represents a novel gene for neuro-cranial development and intellectual disability in humans.

RESULTS

Clinical features and mutation identification

We identified a Thai boy with intellectual disability and megalencephaly. His parents were first cousins. He was born at 34 weeks gestation with a head circumference of 34 cm (97th centile). APGAR scores were 7 and 9 at 1 and 5 minutes, respectively. Neonatal hypoglycemia (blood sugar of 11 mg/dL) and neonatal jaundice were treated promptly. In the first few months of life, he was found to have delayed neurodevelopment and central hypotonia. He was able to hold his head at 5 months, rolled over at 8 months, walked and had first words at 2 years old. At the age of 9 years and 9 months, an IQ test by Wechsler Intelligence Scale for Children: 4th edition (WISC-IV) revealed that his full-scale IQ was 56, indicating intellectual disability. The patient had possible seizure activity at age 10 described as parasomnias, was found to have intermittent bifrontocentral rhythmic theta activity, and the spells resolved after valproic acid therapy. His height and weight followed the curve of 50th centile, but his head circumference remained at 97th centile (53.5 cm and 55 cm at 6 and 9 years old, respectively). Physical examination was generally unremarkable except macrocephaly and low-set prominent anti-

helical pinnae (Fig 1A). Eye examination, hearing tests, thyroid function tests, chromosomal analysis, and nerve conduction velocity were normal. Both brain CT scans at 4 months and 8 years old and brain MRI at 7 months old showed a slight dolichocephalic cranial shape (cephalic index = 75), without overt structural brain abnormalities (Fig 1B-D). X-ray analysis of the spine showed no obvious scoliosis at 10-years-old (Fig 1E).

To elucidate the genetic etiology, we performed homozygosity mapping, whole genome array comparative genomic hybridization (CGH), and whole exome sequencing (WES). WES identified 83 homozygous variants, which had not been reported as SNPs in dbSNP137 (S1 Table). We then selected only those located within the 63 homozygous regions found by homozygosity mapping (S2 Table). Seven candidate variants (one frameshift and six missense mutations; Table I) were identified. Of the six missense, five were predicted to be either benign by Polyphen-2 or tolerated by SIFT prediction programs. The remaining variant, c.235G>A; p.V79M of the *Carboxypeptidase E (CPE)* gene, was not evolutionarily conserved among diverged species (S1 Fig). We, therefore, decided to further our study on the only candidate truncating mutation, a homozygous one base-pair deletion, c.1193delT (p.Leu398Glnfs*2) in exon 11 of *Kinesin family member 6 (KIF6)* (NM_001289021.2).

KIF6 is located on human chromosome 6p21.2 and comprises 23 exons. The 2.4-kb *KIF6* cDNA encodes a canonical N-terminal kinesin motor domain (amino acid positions 3-353) and three coiled-coil regions (amino acid positions 358-385, 457-493, and 633-683), predicted by SMART server [20]. Segregation of the homozygous sequence variant with the disease phenotype was confirmed by Sanger sequencing (Fig 1F) and by restriction fragment length polymorphism (RFLP) analysis of the pedigree (Fig 1G), while his parents and his unaffected brother were heterozygous for the deletion (Fig 1F, G and data not shown). The deletion was not

147 observed in our 1,600 in-house Thai exomes, the 1000 Genome Database, and the ExAC
148 Database. The pedigree combined with the novelty of the mutation in *KIF6* presented here,
149 strongly suggest this C-terminal truncating mutation in *KIF6* may be etiologic for neuro-cranial
150 developmental defects.

151 **Table 1. Seven candidate variants from WES and homozygosity mapping**

#	Chromosome (position)	Gene	Zygosity	Nucleotide change	Amino acid change	Polyphen-2 (score)	SIFT
1	6 (39513453)	<i>KIF6</i>	Homozygous	c.1193delT	p.L398QfsX399		
2	4 (166300608)	<i>CPE</i>	Homozygous	c.235G>A	p.V79M	Possibly damaging (0.889)	Deleterious
3	6 (41895234)	<i>BYSL</i>	Homozygous	c.391C>T	p.R131C	Probably damaging (0.975)	Tolerated
4	7 (44120345)	<i>POLM</i>	Homozygous	c.359G>A	p.R120Q	Probably damaging (0.999)	Tolerated
5	7 (64167644)	<i>ZNF107</i>	Homozygous	c.962T>G	p.I321S	Probably damaging (0.969)	Tolerated
6	2 (96148317)	<i>TRIM43B</i>	Homozygous	c.146C>T	p.P49L	Benign (0.071)	Tolerated
7	5 (140307142)	<i>PCDHAC1</i>	Homozygous	c.665T>C	p.I222T	Benign (0.001)	Tolerated

152

153 Generation of *Kif6* Mutation in Mouse

154 To test the functional consequence of the C-terminal truncating p.L398fsX2 mutation (Fig 1H),
155 we generated an analogous frameshift mutation in exon14 of the mouse *Kif6*

(ENSMUST00000162854) gene, which is ~150bp downstream of the frameshift mutation found in the patient (Fig 2A). After backcross of founder mice to C57B6/J strain, we identified a nonsense allele with scarless insertion (c.1665ins) of a 3-stop donor cassette -providing integration of an ochre termination codon in all three reading frames into the endogenous *Kif6* locus (S2 Fig). This endonuclease-mediated insertional frameshift mutation (*Kif6*^{em1Rgray}) is predicted to truncate the C-terminal cargo-binding domain of the kinesin motor protein (p.G555+6fs). This novel mutant allele of *Kif6* (hereafter called *Kif6*^{p.G555fs}) is predicted to encode a C-terminal truncated KIF6 protein 168 amino acids longer than is predicted for the human p.L398fsX2 variant (Fig 2A).

Hydrocephalus in *Kif6*^{p.G555fs} Mouse

Intercrossing *Kif6*^{p.G555fs/+} heterozygous animals gave offspring with the expected Mendelian ratios, with typical appearance at birth. However, beginning at postnatal day (P)14-onwards, 100% (n=7) of *Kif6*^{p.G555fs/p.G555fs} homozygous mutant mice displayed classic indications of hydrocephalus including doming of the cranium (Fig 2C), a hunched appearance, and with decreased open field activity. We observed apparent megalencephaly and hemorrhaging in older (P21-P28) *Kif6*^{p.G555fs/p.G555fs} mutant brains (Fig 2D), which likely results from increased intracranial pressure and swelling of the ventricles causing damage to the neural tissue against the cranium. At P14, the body weights were not significantly decreased in *Kif6*^{p.G555fs} mutants (5.8±1.3 (g)rams) compared with littermate controls (7.0±1.2g) (n=5/genotype; *p*=0.17). However, at P28 mutant mice showed decreased weight on average (12.67±1.53 g) compared to littermate controls (15.33±1.15g), although this trend was not statistically significant (n=3/genotype; *p*= 0.07). At P28, extracted whole brain sizes appear to be larger in

Kif6^{p.G555fs/p.G555fs} mutants compared to non-mutant littermate controls (Fig 2D). For these reasons, mutant animals were not maintained for observation past P28. qPT-PCR analysis of several *Kif6* exon-exon boundaries found no evidence for non-sense mediated decay in *Kif6*^{p.G555fs} mutant mice (Fig 2B).

To determine whether a more N-terminal truncated *Kif6* mutation would result in a more severe hydrocephalus phenotype, we isolated a conditional-ready *Kif6* allele, where exon 4 is flanked by LoxP sites (*Kif6*^{tm1c}) (KOMP repository, see Methods and Materials). Recombination of the *Kif6*^{tm1c} allele is predicted to generate a frameshift mutation, which should generate a severely truncated, 89 amino acid, KIF6 protein (p.G83E+6fs) with a non-functional N-terminal motor domain. We generated a whole body conditional knockout by crossing the *Kif6*^{tm1c} mouse to the *CMV-Cre* deleter mouse [21]. We observed postnatal-onset, hydrocephalus in *CMV-Cre*; *Kif6*^{tm1c/tm1c} conditional mutant mice (n=10) analogous to our observations in *Kif6*^{p.G555fs/p.G555fs} mutant mice (data not shown). Interestingly, we find no evidence of non-sense mediated decay in these mutant mice despite the generation of an early premature termination codon (data not shown). Because the onset and progression of hydrocephalus was equivalent comparing the whole-body conditional *CMV-Cre*; *Kif6*^{tm1c/tm1c} and *Kif6*^{p.G555fs/p.G555fs} mutant mice strains we suggest that any KIF6 protein encoded by these mutant mouse strains is likely non-functional. Given its relevance to the human mutation, the majority of experiments were all done using the *p.G555fs* allele.

Mouse brains were analyzed histologically by hematoxylin and eosin (H&E) stained coronal sections. Our analysis of coronal sectioned brain at P14 failed to find significance when comparing the total area in section (499.2+39.9μM (Control) vs. 552.5+50.8μM (*Kif6*^{p.G555fs/p.G555fs}); n=7/genotype; p=0.42). However, lateral and third ventricles (LV and 3V

respectively) were obviously enlarged in *Kif6*^{p.G555fs/p.G555fs} mutants (Fig 2F). Quantitation of LV volumes normalized to total brain volume confirmed ventricular expansion in *Kif6*^{p.G555fs/p.G555fs} mutants (n= 7 for each genotype; $p \leq 0.05$; Fig 2G). No obvious defects of the cortex or development of other brain regions in *Kif6*^{p.G555fs/p.G555fs} mutant mice were apparent at a gross anatomical level (Fig 2E, F). Together these data suggest that *Kif6*^{p.G555fs/p.G555fs} mutant mice display postnatal-onset, progressive hydrocephalus, without obvious overgrowth of neural cortex.

Kif6 is Expressed Specifically in the ECs of the Mouse Brain

To determine the endogenous expression patterns of *Kif6* in the mouse, we isolated a *Kif6-LacZ* reporter mouse (*Kif6-LacZ*^{tm1b}) (KOMP repository, see Methods and Materials). Hemizygous *Kif6-LacZ*^{tm1b/+} mice appeared unremarkable and exhibited no evidence of hydrocephalus. Intercrosses of *Kif6*^{tm1b/+} hemizygous mice failed to generate litters with *Kif6*^{tm1b/tm1b} homozygous mice, suggesting that the homozygosity of the *lacZ* expressing allele is embryonic lethal (data not shown). At P10 and P21, *Kif6*^{tm1b/+} transgenic mice showed *lacZ* expression throughout the ependyma of the ventricular system including the central canal. However, no *lacZ* expression was detected in the choroid plexus or in other regions of the brain (Fig 3A, A' and S3B' Fig), with the exception of a small population of cells flanking the third ventricle (arrows, S3 Fig). Interestingly at P10, other multi-ciliated tissues in these transgenic mice such as the oviduct or trachea were not labeled (Fig 3B-C'). Moreover, no laterality defects or obvious changes to trachea cilia were observed in *Kif6*^{p.G555fs/p.G555fs} mutant mice (S6A, B Fig), suggesting that *Kif6* expression and function are tightly restricted to the multiciliated EC in

mouse. Taken together these data suggested a cellular mechanism centered on defective ECs underlying the development of hydrocephalus in *Kif6*^{p.G555fs/p.G555fs} mutant mice.

Progressive Loss of Cilia in *Kif6*^{p.G555fs/p.G555fs} Mutant Mice

Defects in ECs and their cilia are known to cause hydrocephalus in mouse [8]. To assay EC cilia, we utilized scanning electron microscopy (SEM) to directly visualize the LV. Heterozygous *Kif6*^{p.G555fs/+} mice displayed a high-density of regularly spaced EC multiciliated tufts along the LV surface (Fig 4A-A'), typical for P21 mice [22]. In contrast, homozygous *Kif6*^{p.G555fs/p.G555fs} mutant mice displayed a marked reduction of multiciliated tufts across the LV wall, coupled with a reduction in the density of ciliary axonemes extending from the ECs (Fig 3B-B'). The loss of EC cilia was more severe at P28 (S4 Fig). Together, these data suggested that hydrocephalus may result from either a reduction in EC differentiation and/or defects in EC cilia maintenance during postnatal development.

To address the differentiation status of the ECs, we utilized immunofluorescence (IF) in coronal sectioned brain tissues to image known proteins components of the EC and their cilia. At P21, we observed the expression of the ependymal cell-marker S100B [22] throughout the epithelium lining luminal surface of the ventricles, as well as, the presence of apically localized γ -tubulin-positive basal bodies within these ECs in both WT (Fig 4C-C'') and *Kif6*^{p.G555fs/p.G555fs} mutant mice (Fig 4D-D''). Conversely, we observed a severe reduction in the density of CD133-positive EC axonemes [23] extending into the ventricular lumen in *Kif6*^{p.G555fs/p.G555fs} mutant mice (Fig 4C, C'''), compared with WT (Fig 4D, D'''). Quantitation of several sections from independent mice confirmed a severe reduction of CD133-positive ciliary axonemes due to the loss of KIF6 function (n=5 mice/genotype, p<0.001) (Fig 4E). These results suggest that the

onset of hydrocephalus in *Kif6* mutant mice is primarily due to the loss of EC ciliary axonemes and not the result of defects in the differentiation of these cells.

Ventricular Dilation and Reduced EC Cilia in *kif6* Mutant Zebrafish

Previous studies in *kif6^{sko/sko}* mutant zebrafish found late-onset scoliosis, without obvious hydrocephalus or defects in EC cilia during early embryonic development [19]. Interestingly, reduced CSF flow, ventricular dilation, and loss of EC cilia during larval zebrafish development is associated with scoliosis [24]. In order to determine if *kif6* mutant zebrafish display changes in the ventricular system later in adults, we used iodine contrast-enhanced, micro computed tomography (μ CT) [25] to generate high-resolution (5 μ M) images of the intact zebrafish brain. After reconstruction and alignment of 3D tomographic datasets in the coronal plane, we utilized stereotyped landmarks of the zebrafish brain and spinal cord [26] to compare equivalent axial sections of aged matched (90 days post fertilization (dpf)) *kif6^{sko/+}* heterozygous and *kif6^{sko/sko}* homozygous mutant zebrafish. At each axial level of the brain (Fig 5A), we observed consistent dilation of the ventricular system and central canal in *kif6^{sko/sko}* homozygous mutant zebrafish (yellow arrows; Fig 5C, E, G), compared to a stereotyped anatomy of the wild-type (WT) zebrafish brain (Fig 5B, D, F). Multiple regions of *kif6^{sko/sko}* mutant zebrafish brain were found to be structurally abnormal in *kif6^{sko/sko}* mutants compared to WT zebrafish (S1, 2 Movies). We next quantified the areas of two anatomically distinctive ventricles in our tomographic datasets: (i) the tectal ventricle (TecV) and (ii) a region of the rhombencephalic ventricle (RV) just posterior to the lobus facialis [26]. We found that both the TecV and the posterior RV were significantly more dilated in *kif6^{sko/sko}* mutant zebrafish comparing several optical sections from independent aged-matched zebrafish (n=3 fish/genotype; $p<0.0001$). The central canal was also clearly

dilated in *kif6*^{sko/sko} mutant zebrafish (yellow arrow, Fig 5G). However, we were unable to reliably quantify this area in WT samples at the current resolution. Our previous observations in *kif6*^{sko/sko} mutant zebrafish embryos failed to find phenotypes that are characteristic of cilia defects, such as hydrocephalus, *situs inversus*, or kidney cysts [19]. Moreover, we observed normal development and function of EC cilia in the central canal in embryonic mutant zebrafish [19]. These data, together with our new observations of ventricular dilation in adult *kif6* mutants (Fig 5), suggest that Kif6 is required for the post-embryonic maintenance of the EC cilia as was observed in other zebrafish mutants displaying similar scoliosis as observed in *kif6* mutant zebrafish [19, 24].

In order to assay whether EC cilia were affected in adult (90dpf) *kif6*^{sko/sko} mutant zebrafish, we isolated a stable transgenic allele, *Tg(Foxj1a:GFP)*^{dp1} which effectively labels multiple multiciliated *Foxj1a*-positive cell lineages, including ECs, with cytoplasmic EGFP in zebrafish [24]. We observed no differences in the specification of *Foxj1a:GFP*-positive ECs comparing WT and homozygous *kif6*^{sko/sko} mutant fish (Fig 5I, J). Cytoplasmic GFP can freely diffuse into and label the ciliary axoneme [27]. As such, we were able to observe GFP-positive EC cilia projecting into the ventricular lumen in *Tg(Foxj1a:GFP)*^{dp1}; *kif6*^{sko/+} heterozygous fish (red arrowheads; Fig 5I). In contrast, these GFP-labeled EC cilia were reduced or absent in *Tg[Foxj1a:GFP]*^{dp1}; *kif6*^{sko/sko} mutant fish (Fig 5J, K). Furthermore, SEM analysis of the ventricles in *kif6*^{sko/sko} mutant zebrafish further supported our observations of ventricular dilation and loss of EC cilia in adult *kif6*^{sko/sko} mutant fish (S5 Fig). Akin to our observations in *Kif6* mutant mice trachea, we did not observe defects of other multiciliated tissues such as the nasal cilia (S6C, D Fig) in *kif6*^{sko/sko} mutant zebrafish. Together, these data suggest that Kif6 functions specifically in the maintenance of EC cilia as well as for ventricular homeostasis in zebrafish.

DISCUSSION

This study demonstrates the importance of *KIF6* for neuro-cranial development in vertebrates, and a unique and highly specialized role in ependymal cells where its function is important for maintenance of EC cilia. This is supported by several lines of evidence including the discovery of a novel nonsense-mutation of *KIF6* in a child with neuro-cranial defects and intellectual disability and underscored by functional analysis in both mouse and zebrafish *Kif6* mutant models (Table 2).

Table 2. KIF6 mutations discussed in this paper.

ID/Allele	Species	Nucleotide change	Amino acid change	Phenotype reported	Reference
Patient	Human	c.1193delT	p.L398QfsX399	Delayed neurodevelopment and central hypotonia, neuro-cranial defects, and intellectual disability.	this work
<i>Kif6</i> ^{p.G555fs/p.G555fs}	Mouse	c.1665ins	p.G555+6fs	Severe progressive hydrocephalus, loss of ependymal cell cilia	this work
	Mouse	deletion of exon 4	p.E83+39fs	Severe progressive hydrocephalus, loss of ependymal cell cilia	this work
<i>Kif6</i> ^{Δ3/Δ3}	Mouse	c.177-251_del	p.Lys59Asn, Phe60_Ser84_del	None reported	[28]
<i>kif6</i> ^{sko/sko}	Zebrafish	c.205C>A	p.Tyr53X	Larval onset scoliosis, hydrocephalus, loss of ependymal cilia	[19], this work

We identified a homozygous *KIF6* c.1193delT mutation in a child with macrocephaly and cognitive impairment that segregated with this phenotype in his family, and leads to a loss of the C-terminal second and third coiled-coil regions which are important for dimerization and cargo selectivity of kinesin motors [13]. We engineered an analogous, C-terminal truncating mutation of *KIF6* in mouse, which displays severe hydrocephalus and defects of EC cilia

providing strong evidence for pathogenicity of the mutation in the child. Other than the case described here, no prior mutation directly attributed to human disease has been described for *KIF6*. Taken together, the clinical data reported here suggest that biallelic mutations in *KIF6* may underlie some unexplained intellectual disability and neuro-cranial developmental defects. Future analyses of *KIF6* mutations in these patient groups are warranted.

Further, our analysis of several independent loss-of-function *Kif6* mutant animal models found no evidence of obvious heart abnormalities to explain the prior association of the common variant *KIF6* p.W719R in some [17], but not all [18], studies of coronary heart disease in humans. Because expressed sequence tag clones of *KIF6* have not been reported from cDNAs libraries derived from human heart or vascular tissues (UniGene 1956991 - Hs.588202), any possible functional effects of KIF6 on heart function remains unexplained. However, detailed analysis of coronary function was not explored in our models, therefore it is possible that subtle defects may be present.

An ENU-derived *Kif6* splice acceptor site mutant mouse strain, predicted to delete the 3rd exon of *KIF6* (*Kif6*^{Δ3/Δ3}), also did not show cardiac or lipid abnormalities [28]. Of note this mutant mouse was also not reported to have hydrocephalus. Our analysis shows that the loss of exon 3 in *Kif6*^{Δ3/Δ3} mutant mouse generates an inframe deletion of only 25 amino acids in the N-terminal motor domain of the KIF6 protein, otherwise generating a mostly full-length KIF6 protein (Table 2). In contrast, here we report two novel *Kif6* mutant mice: (i) a C-terminal *Kif6*^{p.G555fs/p.G555fs} deletion mutant, predicted to truncate 248 amino acids of the C-terminal domain, which are important for cargo binding in Kinesin motor proteins [13]; and (ii) a conditional *CMV-Cre;Kif6*^{tm1c/tm1c} mutant which recombines exon 4 leading to an early frame shift mutation predicted to generate a N-terminal truncated 122 amino acid KIF6 protein (Table

2), both of which display indistinguishable progressive, hydrocephalus. The most parsimonious explanation for the difference in phenotypes in these mutant mice is that the *Kif6*^{A3} allele encodes a functional KIF6 protein. Analysis of these mutations in trans or quantitative analysis of these kinesin motor proteins *in vitro* is warranted to more fully address these conflicting observations.

There are marked differences in the phenotypes among the human, mouse, and zebrafish associated with mutations in *KIF6*. For example, *kif6* mutant zebrafish display post-natal onset scoliosis, mirroring adolescent idiopathic scoliosis (IS) in humans [29]. The formation of IS-like defects in zebrafish has been shown to be the result of a loss of CSF flow, associated with a loss of EC cilia and ventricular dilation during a defined window of larval zebrafish development [24]. Interestingly, we did not observe scoliosis in the *Kif6* mutant mice (S7 Fig), despite being of an appropriate age when IS-like scoliosis can manifest in mouse [30]. Moreover, we do not observe scoliosis in the patient at the age of 10 years, though it is possible that he may yet develop scoliosis during adolescence. The mechanism behind these differences may reflect distinctions in the functional input of the ventricular system for spine stability amongst teleosts and amniotes.

Furthermore, while we observe a clear role for KIF6 in maintaining the ventricular system in mouse and zebrafish, the patient does not have obvious hydrocephalus. However, his relative macrocephaly and slightly enlarged ventricles by MRI (Fig 1B-D) may suggest an element of what is commonly referred to as arrested hydrocephalus [31]. The contribution of EC cilia beating to bulk CSF flow might be species dependent. For instance, the majority of CSF flow in humans is thought to occur via the generation of a source-sink gradients; partly from the secretion of the choroid plexus and exchanges of the interstitial fluids, coupled with absorption at the arachnoid villi and lymphatics [32]. In contrast, localized or near-wall CSF flow [4],

generated by polarized beating of EC cilia, are clearly important for the formation of hydrocephalus in rodents [8]; however, there have been limited examples EC cilia defects causing hydrocephalus in humans. Regardless there is growing evidence suggesting that EC cilia dependent CSF flow is crucial for the regulation of brain function and neurogenesis [4], and for adult neural stem cell proliferation [9]. It is possible that a specific loss of EC cilia in humans may only have minor effects on CSF bulk flow and ventricular homeostasis, while causing severe defects of neurogenesis leading to intellectual disability and other neurological disease. It will be important to determine if the loss of KIF6 function in adulthood will contribute to changes in neurodevelopment and behavior, cognitive decline, and ventricular homeostasis.

Finally, *KIF6* now joins five other kinesin genes, *KIF1C*, *KIF2A*, *KIF4A*, *KIF5C* and *KIF7* that were previously reported to be associated with neurological abnormalities in humans [33-36]. Here we suggest that *KIF6* has a uniquely specific function in the EC cilia in vertebrates, resulting in both cognitive impairment and macrocephaly in a child with a homozygous one-base pair deletion. Using a cell biological approach, we identified specific loss of EC cilia in *Kif6* mutant models in both mouse and zebrafish, suggesting a strong conservation of KIF6 function in ventricular system in vertebrates.

Material and Methods

Identification of Mutation in Patient 1

Whole Exome Sequencing (WES)

The patient's genomic DNA of patient was extracted from peripheral blood leukocyte using AchivePure DNA Blood Kit (5 Prime Inc., Gaithersburg, MD). The sample was sent to MacroGen, Inc. (Seoul, Korea) for whole exome sequencing. The 4 ug of DNA sample was enriched by TruSeq Exome Enrichment Kit and was sequenced onto Hiseq 2000. The raw data per exome was mapped to the human reference genome hg19 using CASAVA v1.7. Variants calling were detected with SAMtools.

Homozygosity mapping

The sample was sent to MacroGen, Inc. (Seoul, Korea) for genotyping. The DNA sample was genotyped by HumanOmni 2.5-4v1 DNA BeadChip (Illumina) which contain 2,443,177 SNPs. The experiment was performed by the array protocol. PLINK was used to analyze for the homozygous regions.

Mutation analysis

We performed resequencing of *KIF6* pathogenic region in patient and patient's family. Primers for the amplification of the candidate variant were designed using Primer 3 software (version 0.4.0). Primers KIF6-1193delT-F 5'-CAGCTTGAACATGGCTGAAA-3' and KIF6-1193delT-R 5'-TTCTGTAAAGAGGTGGGAACAA-3' were used to amplify. The 20 ul of PCR reaction contained 50-100 ng of genomic DNA, 200 uM of each dNTP, 150 nM of each primer, 1.5 mM MgCl₂ and 0.5 unit of Taq DNA polymerase (Fermentas Inc., Glen Burnie, MD). The

PCR condition was started with 95 °C for 5 min for pre-denaturation following with the 35 cycles of 94 °C for 30 sec, 55 °C for 30 sec and 72 °C for 30 sec. The product size of these primers is 276 bp. For sequencing, PCR products were treated with ExoSAP-IT (USP Corporation, Cleveland, OH), and sent for direct sequencing at Macrogen Inc. (Seoul, Korea). Bi-directional sequencing was done by using KIF6-1193delT F and R primers. Analyses were performed by Sequencer 4.2 (Gene Codes Corporation, Ann Arbor, MI).

PCR-RFLP

One hundred chromosomes and patient's trio were genotyped by PCR-RFLP. Primer KIF6 *MfeI* F 5'-TGGCTTCACTATAAATTTCACTTTGTCAATG-3' and mutagenic primer KIF6 mutagenic *MfeI* R 5'-TCCTGGTCTTCCAAAAAGGATGCAAT-3' were used to amplify KIF6 T-deletion. The 20 ul of PCR reaction contained 50-100 ng of genomic DNA, 200 uM of each dNTP, 150 nM of each primer, 1.5 mM MgCl₂ and 0.5 unit of Taq DNA polymerase (Fermentas Inc., Glen Burnie, MD). The PCR condition was started with 95 C for 5 min for pre-denaturation following with the 35 cycles of 94 C for 30 sec, 60 C for 30 sec and 72 C for 30 sec. The product size of these primers is around 223 bp. The PCR product was incubated with 10U of *MfeI*-HF (New England Biolabs, Ipswich, MA) at 37 C overnight. Three percent of agarose gel electrophoresis was used to detect the different cut sizes of PCR product. A 196 bp and 26 bp bands were present in one base deletion sample.

Mice

All mouse studies and procedures were approved by the Animal Studies Committee at the University of Texas at Austin (AUP-2015-00185). The *Kif6*^{p.G555fs} mutant mouse were developed

using CRISPR-Cas9-mediated genome editing. Using the CHOP-CHOP online tool [37], we identified a suitable 20-nucleotide site (GGAGATGTCACTGGGACGCC) targeting exon 14 of mouse *Kif6* (ENSMUST00000162854.1) in order to generate a C-terminal truncation allele. The gene specific and universal tracrRNA oligonucleotides (S3 Table) were annealed, filled in with CloneAmp HiFi PCR premix, column purified, and directly used for *in vitro* transcription of single-guide RNAs (sgRNAs) with a T7 Polymerase mix (M0255A NEB). All sgRNA reactions were treated with RNase free-DNase. We utilized a ssDNA oligo (S3 Table) to insert a frameshift mutation in all three reading frames, along with 8-cutter restriction sites for genotyping (3-stop donor) [38] (Fig S2). The *Kif6* ex14 3-stop donor and mKif6-R2-ex14-T7 sgRNA were submitted for pronuclear injection at the University of Texas at Austin Mouse Genetic Engineering Facility (UT-MGEF) using standard protocols (<https://www.biomedsupport.utexas.edu/transgenics>). We confirmed segregation of the *Kif6*^{p.G555fs} allele using several methods including increased mobility on a high percentage electrophoresis gel, donor-specific primer PCR, or PmeI (NEB) digestion of the *Kif6* exon14 amplicon (S2 Fig and S3 Table). PCR products in isolated alleles were cloned to pCRII TOPO (ThermoFisher) to identify scarless integration of the 3-stop donor at the *Kif6* locus using gene specific flanking primers (S3 Table).

Kif6-LacZ^{tm1b} mice were generated by injection of embryonic stem cell clones obtained from the Knockout Mouse Project (KOMP) Repository. Three *Kif6*^{tm1a(KOMP)Mbp} embryonic stem (ES) cell clones (KOMP: EPD0736_3_G01; EPD0736_3_H02; and EPD0736_3_A03) all targeting exon 4 of the *Kif6* gene with a promoter-driven targeting cassette for the generation of a 'Knockout-first allele' [39]. Pronuclear injections of all clones were done using standard procedures established by the UT MGEF. After screening for germline transmission, we isolated

and confirmed a single heterozygous founder male (*Kif6*^{tm1a(KOMP)Mbp}) carrier derived from the G01 clone. We confirmed the locus by long-range PCR, several confirmation PCR strategies targeting specific transgene sequences, and Sanger sequencing of the predicted breakpoints (S3 Table). After several backcrosses to the WT C57BL/6J substrain (JAX), we crossed a hemizygous *Kif6*^{tm1a/+} mutant male to a homozygous *CMV-Cre* female (*B6.C-Tg(CMV-cre)1Cgn/J*) (JAX, 006054) to convert the *Kif6*^{tm1a} allele to a stable LacZ expressing *Kif6*^{tm1b} allele (*Kif6-LacZ*^{tm1b}). Mutant F1 offspring from this cross were backcrossed to WT C57BL6/J mice and the F2 progeny were genotyped to confirm the *Kif6-LacZ*^{tm1b} allele and the presence/absence of the CMV-Cre transgene. A single founder *Kif6-LacZ*^{tm1b} with the desired genotype (*Kif6-LacZ*^{tm1b} hemizygous, Cre transgene absent) was used to expand a colony for spatial expression analysis.

Kif6^{tm1c} conditional ready mice were generated by outcross of the *Kif6*^{tm1a(KOMP)Mbp} allele described above to a ubiquitously expressed Flippase strain (*129S4/SvJaeSor-Gt(ROSA)26Sor*^{tm1(FLP1)Dym/J}) (JAX, 003946). F1 offspring were genotyped and sequenced at several breakpoints to ensure proper flip recombination and a single F1 founder was used to backcross to C57B6/J for propagation of the *Kif6*^{tm1c} strain. Analysis of recombination of the floxed *Kif6*^{tm1c} was performed by crossing homozygous *Kif6*^{tm1c/tm1c} to a compound heterozygous *CMV-Cre; Kif6*^{tm1c/+} mouse. Recombination of the exon 4 of *Kif6* was confirmed by PCR-gel electrophoresis analysis (S3 Table).

LacZ Staining Protocol

Mice were perfused with LacZ fixative and post fixed for 2 hours at RT. Whole brains were then stained in X-gal solution overnight at 37°C followed by post-fixation in 4% PFA overnight at

4°C. The samples were then prepped for cryosectioning in 30% sucrose/OCT and sectioned.

Sections were counter stained in Nuclear Fast Red stain (Sigma).

X-ray Analyses of Mice

Radiographs of the mouse skeleton were generated using a Kubtec DIGIMUS X-ray system

(Kubtec, T0081B) with auto exposure under 25 kV.

Zebrafish Manipulations and Transgenesis

All zebrafish studies and procedures were approved by the Animal Studies Committee at the

University of Texas at Austin (AUP-2015-00187). Adult zebrafish of the AB were maintained

and bred as previously described [40]. Individual fish were used for analysis and compared to

siblings and experimental control fish of similar size and age. Independent experiments were

repeated using separate clutches of animals. Strains generated for this study: Tg(Foxj1a:GFP)^{dp1}.

Previously published strains: *kif6*^{sko} [19]. Transgenic lines were generated using the Tol2-system

as described before [41].

Mouse and Zebrafish Perfusions and Embedding of Brain Tissues

Mice were humanely euthanized by extended CO₂ exposure and transferred to chemical hood

where the mouse was perfused with buffered saline followed by 4% PFA. Whole brains were

placed in 4% PFA 4 hours at RT, then at 4° C overnight. Zebrafish were euthanized by exposure

to lethal, extended dose of Tricane (8%) followed by decapitation. Zebrafish brains were

extracted and fixed in 4% PFA at 4° C overnight. For paraffin embedding, the fixed brains were

embedded and cut using standard paraffin embedding and sectioning protocols. Paraffin sections were stained with standard hematoxylin-eosin solution.

For frozen sections both mouse or zebrafish brains were fixed as above and then equilibrated to 30% or 35% sucrose, respectively at 4° C overnight. Whole brains were then placed in O.C.T. Compound (Tissue-Tek) and flash in cold ethanol bath. All blocks were stored at -80° Celsius until sectioning on a cryostat (Leica). All sections were dried at RT for ~2hrs. and stored at -80°C until use.

Immunofluorescence Protocol for Frozen Brain Sections

Sectioned tissues were warmed at room temperature for ~1 hour, then washed thrice in 1xPBS + 0.1% Tween (PBST). Antigen retrieval was hot citrate buffer (pH6.8). Blocking was done in 10% Normal goat serum (Sigma) in 1xPBST. Primary antibodies (S100B at 1:1,000, ab52642, Abcam; CD133(Prominin-1), 134A, 1/500; Gamma Tubulin, sc-17787, Santa Cruz (C-11), 1/500; Anti-GFP, SC9996, Santa Cruz, 1:1,000) were diluted in 10% NGSS, 1xPBST and allowed to bind overnight at 4°C in a humidified chamber. Secondary fluorophores (Alexa Fluor 488(A-11034); 568(A10042); and 647(A32728), 1:1,000, ThermoFisher) were diluted in 10% NGS; 1xPBST were allowed to bind at RT for ~1hr. We used Prolong gold with DAPI (Cell Signaling Technologies, 8961) to seal coverslips prior to imaging.

Iodine-contrast μ CT

Zebrafish specimens were fixed overnight in 10% buffered formalin, washed thrice in diH₂O and stained ~48 hours in 25% Lugol's solution/75% distilled water. Specimens were scanned by the High-resolution X-ray CT Facility (<http://www.ctlab.geo.utexas.edu/>) on an Xradia at 100kV,

508 10W, 3.5s acquisition time, detector 11.5 mm, source -37 mm, XYZ [816, 10425, -841], camera
 509 bin 2, angles ± 180 , 1261 views, no filter, dithering, no sample drift correction. Reconstructed
 510 with center shift 5.5, beam hardening 0.15, theta -7, byte scaling [-150, 2200], binning 1, recon
 511 filter smooth (kernel size = 0.5).

512

513 Statistical Analysis and image measures

514 GraphPad Prism version 7.0c for Mac (GraphPad Software) was used to analyze and plot data.
 515 Images for measurement were opened in FIJI (Image J) [42], and measures were taken using the
 516 freehand tool to draw outlines on ventricular area or whole brain area. Statistically significant
 517 differences between any two groups were examined using a two-tailed Student's t-test, given
 518 equal variance. P values were considered significant at or below 0.05.

519

ACKNOWLEDGMENTS

We thank members of the Gray and Wallingford labs for helpful discussions and critical reading of the manuscript. We would also like to thank Jin Xiang Ren and William Shawlot for help with mouse engineering, Terry Heckmann and Ryoko Minowa for excellent technical support. We thank Jessie Maisano for her help and expertise with iodine-contrast μ CT for this study. This study was supported (V.S.) by the Thailand Research Fund (DPG6180001) and the Chulalongkorn Academic Advancement into Its 2nd Century Project. The research of M.K. was supported in part by the Provost Graduate Excellence Fellowship, Institute of Cell and Molecular Biology, University of Texas at Austin. The research of R.S.G. was supported start-up funds from the University of Texas at Austin Dell Medical School and by a NIH grant R01AR072009.

Figure Legends

Fig 1. *KIF6* mutation in a child with intellectual disability. (A) A low-set prominent anti-helical left pinna. (B) MRI of the brain at 7 months old shows dolichocephaly with a normal brain structure. (C) and (D) CT of the brain at 8 years old, sagittal and axial views, respectively show dolichocephalic shape of the cranium (cephalic index = 75) without demonstrable intracranial abnormality. (E) X-ray of the spine shows no scoliosis (F) Electropherograms of the patient, a control, and the patient's father in the upper, middle and lower panels, respectively. The patient is homozygous while his father is heterozygous for the c.1193delT. (G) Pedigree and RFLP, using MfeI restriction enzyme: Lane M = 100 bp marker. The arrow head indicates the 500 bp band. Lanes 1-5 are controls. Lanes 6 and 7 are the proband's father and mother, respectively, showing that they are heterozygous. Lane 8 is the proband showing that he is homozygous for the c.1193delT. (H) Representative KIF6 structure. The arrow shows the position of the c.1193delT mutation.

Fig 2. *Kif6*^{p.G555fs} mutant mice display progressive hydrocephaly. (A) Schematic of the non-sense mutation in the patient (*KIF6*^{p.L398fsX2}) and the mouse mutation (*Kif6*^{p.G555fs}), both predicted to truncate the C-terminal domain of KIF6 protein. (B) Fold change qRT-PCR of *Kif6* expression using cDNA libraries derived from lateral ventricles from WT (black bars) and *Kif6*^{p.G555fs/p.G555fs} (gray bars) mutant mice. (C) Lateral X-rays of mouse cranium at P14 and P28 showing the progressive cranial expansion in *Kif6*^{p.G55/p.G555fs} homozygous mutant mice. (D) Ventral view of whole mouse P28 brain to highlight hemorrhaging and slight enlargement of total brain size in *Kif6*^{p.G55/p.G555fs} homozygous mutant mice. (E, F) H&E stained coronal sections of the mouse brain (P14), showing dilation of the lateral (LV) and third (3V) ventricles in *Kif6*^{p.G55/p.G555fs}

homozygous mutant mice (F). (G) Quantitation of ventricular area over total brain area in *Kif6^{p.G55/p.G55fs}* homozygous mutant mice and heterozygous littermate controls (n=7 mice per genotype; two-tailed t-test; p=0.0173). Scale bars: 1cm in (C); 5mm in (D); and 300 μM in (F, F).

Fig 3. *Kif6-LacZ* expression is specific to the ependymal cells. (A-C') Representative LacZ staining in a variety of multiciliated tissues from P10 *Kif6-LacZ^{tm1b/+}* transgenic mice. (A, A') Coronal section at the 4th ventricle showing specific *LacZ* expression in the ependymal cell (EC) layer and stark lack of expression in the choroid plexus (CP) or surrounding neuronal tissues. (B, B') Sectioned oviduct tissue shows no *LacZ* expression. (C, C') Sectioned trachea tissue shows no *LacZ* expression. Scale bars: 300μM in (A-C); and 20μM in (A'-C').

Fig 4. *Kif6* mutant mice have a reduction in ependymal cell cilia. (A-B') SEM of the lateral ventricular wall (*en face* view) in *Kif6^{p.G55/p.G55fs}* homozygous mutant mice and heterozygous littermate controls at P21, showing a reduction in the number and density of EC cilia tufts. (C-D''') Immunofluorescence of the wild-type (WT) and *Kif6^{p.G55/p.G55fs}* homozygous mutant mice at P21. (C-D) Three color merge of (C', D') αS100B (ependymal cells; magenta) channel; (C'', D'') α-γ-tubulin (basal bodies; cyan) channel; and (C''', D''') αCD133 (prominin-1; green). (C', D') αS100B staining showing no changes in ependymal cell specification between *Kif6^{p.G55/p.G55fs}* homozygous mutant and WT mice. (C'', D'') α-γ-tubulin staining showing typical basal body positioning at the apical surface of ECs in both *Kif6^{p.G55/p.G55fs}* homozygous mutant and WT mice. (C''', D''') αCD133 staining reveals a marked of EC cilia projecting into the ventricular lumen in *Kif6^{p.G55/p.G55fs}* homozygous mutant mice compared to WT mice. (E) Quantitation of

fluorescent intensity of the CD133 channel (ciliary axonemes) as the average line scan value along the lumal surface of mouse ventricles. Scale bars: 20μM in (A, B); 2 μM in (A', B'); and 20 μM in (C-D''').

Fig 5. *kif6* mutant zebrafish display dilation of the ventricular system and loss of ependymal cell cilia. (A) Schematic of adult zebrafish brain highlighting the relative sectioning of the zebrafish brain. (B-G) Coronally aligned, optical sectioning of Iodine-contrasted μCT imaging datasets of WT (B, D, F) and *kif6*^{sko/sko} homozygous mutant (C, E, G) zebrafish brain at 90dpf. (B-C) The medial region of the TeO showing the TecV which is dilated in *kif6*^{sko/sko} mutant fish (C) compared to age-matched WT (B). (D-E) Sectioning at the region of the medulla oblongata posterior to the lobus facialis showing dysmorphogenesis and deepening of the RV in *kif6*^{sko/sko} mutants (E) compared with WT (D) zebrafish. (F-G) Spinal cord sectioning showing dilation of the Cc in *kif6*^{sko/sko} mutant (G) compared to WT (F) zebrafish. Scale Bars: 1mm in (B-G); and 20 μM in (I, J). (H) Quantitation of the area of the TecV and the RV posterior to the lobus facialis (pos. RV) in WT and *kif6*^{sko/sko} mutant zebrafish, highlighting the consistent dilation in *kif6*^{sko/sko} mutants (n=11 sections/genotype; two-tailed t-test; ****, p<0.0001). (I-J) Immunofluorescence of *Tg[foxl1a::GFP]* in both heterozygous and homozygous *kif6*^{sko/sko} mutant zebrafish stained with αGFP (green) and DAPI (blue) showing GFP positive ECs in both genotypes. However, *kif6*^{sko/+} heterozygous zebrafish display numerous apical tufts of cilia (red arrows) projecting into the ventricle lumen, which are markedly reduced in *kif6*^{sko/sko} mutant zebrafish. Scale Bars: 1mm in (B-G); and 20 μM in (I, J). *TecV*-tectal ventricle *TeO*-tectum opticum; *CCe*-corpus cerebelli; *RV*- rhombencephalic ventricle ventricle; and *Cc*-central canal.

600

601 **Supporting Information**

602 **S1 Fig Clustal Alignment of CPE (p.V79M) variant.** Hs, *Homo sapiens*; Pt, *Pan troglodytes*;
603 Mc, *Macaca mulatta*; Mu, *Mus musculus*; Rn, *Rattus norvegicus*; Bt, *Bos Taurus*; Cl, *Canis lupus*
604 *familiaris*; Oc, *Oryctolagus cuniculus*; Gg, *Gallus gallus*; Dr, *Danio rerio*; Tn, *Tetraodon*
605 *nigroviridis*; Xt, *Xenopus (Silurana) tropicalis*; Tc, *Tribolium castaneum*; Ce, *Caenorhabditis*
606 *elegans*; Sk, *Saccoglossus kowalevskii*.

607

608 **S2 Fig *Kif6*^{p.G555fs} generation and genotyping.** (A) Schematic of target cut site and insertion
609 cassette into exon 14 of *Kif6* locus. Insertion cassette contains three stop codons, one in each
610 reading frame, and two 8 basepair restriction enzyme cut sites for easy genotyping. (B-C)
611 Agarose gels of PCR products confirming germline transmission of donor cassette in F₁
612 generation from CRISPR injected chimeras. (B) RE digest of PCR product from exon 14
613 flanking target site, shows cutting (asterisks) in heterozygous F₁ mice. Wildtype band (arrow)
614 appears in lane one and all the subsequent lanes. (C) PCR product from donor specific primer
615 and *Kif6* exon 14 reverse primer confirming donor insertion and germline transmission. (D)
616 Table describing CRISPR injected mice, number with detectable indels, total with integration of
617 donor oligo, and total displaying hydrocephaly of chimeric injected CRISPR mice. (E) Germline
618 transmission of donor cassette from chimeric CRISPR F₀ mice to F₁ generation.

619

620 **S3 Fig *LacZ* expression in different ages of *Kif6-LacZ*^{tm1b} transgenic mouse brain.** (A-A'')
621 Coronal sections of P21 mouse brains showing LacZ staining restricted to the EP cell layer in the
622 4th ventricle. Zoom in shows LacZ positive cells have cilia projecting into the lumen (arrows).

(B-B') Coronal sections of P21 mouse brains showing LacZ staining of ventral portion of 3rd ventricle. (B') Some sporadic staining appearing in the nuclei of the hypothalamus (arrows). (C-C') LacZ staining in the fourth ventricle at P10 showing staining specific to ependymal cell layer.

S4 Fig SEM of lateral ventricle in *Kif6*^{p.G555fs/p.G555fs} mutant and control at P28. SEM of *Kif6* wildtype vs. *Kif6*^{p.G555fs/p.G555fs} mutants shows *Kif6* mutants show a complete loss of ependymal cell cilia on the lateral wall by P28. Scale bar 10μM.

S5 Fig SEM of ventricle in *kif6*^{sko/sko} mutant zebrafish display dilation of the ventricular system and loss of ependymal cell cilia. SEM of zebrafish brain shows dilation of brain ventricles indicative of hydrocephaly. Higher magnification images reveal loss of ependymal cell cilia tufts in *kif6* zebrafish mutants when compared with heterozygous counterparts. Scale bars 20μM and 200μM.

S6 Fig Immunofluorescence (IF) of *Kif6* mutant multiciliated tissues in mouse and zebrafish. (A-B) Representative IF of trachea sections in *Kif6*^{p.G555fs/+} and *Kif6*^{p.G555fs/p.G555fs} mice showing no cilia defects present in trachea of *Kif6* mutant mice. Acetylated tubulin (green) marking cilia, DAPI-stained nuclei (magenta) (C-D) Representative IF of zebrafish nasal pit cilia shows typical cilia in *kif6* mutant zebrafish to wildtype counterparts. Acetylated tubulin (magenta) marking cilia, gamma-tubulin marking basal bodies (green). Scale bars are 20μM.

645 **S7 Fig X-rays of *Kif6*^{p.G555fs/p.G555fs} mutant mice.** Representative X-rays of wildtype and
 646 *Kif6*^{p.G555fs/p.G555fs} mutant mice shows no scoliosis at P28. *Kif6*^{p.G555fs/p.G555fs} mice do however
 647 display skull expansion caused by progressive hydrocephalus (Animal #1 and #2).

648

649 **Table SI - Eighty-three homozygous variants from WES**

650

651 **Table SII - Sixty-three homozygous regions from homozygosity mapping**

652

653 **Table SIII - Mouse specific oligos and primers**

654

655 **Movie 1- Representative WT_Danio_Iodine-contrasted microCT transverse**

656

657 **Movie 2- Representative *kif6*^{sko/sko}_Danio_Iodine-contrasted microCT**
 658 **transverse**

659 References

- 660 1. Whedon JM, Glassey D. Cerebrospinal fluid stasis and its clinical significance. *Altern*
661 *Ther Health Med.* 2009;15(3):54-60. PubMed PMID: 19472865; PubMed Central PMCID:
662 PMCPMC2842089.
- 663 2. Rubenstein E. Relationship of senescence of cerebrospinal fluid circulatory system to
664 dementias of the aged. *Lancet.* 1998;351(9098):283-5. doi: 10.1016/S0140-6736(97)09234-9.
665 PubMed PMID: 9457114.
- 666 3. Jacquet BV, Salinas-Mondragon R, Liang H, Therit B, Buie JD, Dykstra M, et al. FoxJ1-
667 dependent gene expression is required for differentiation of radial glia into ependymal cells and a
668 subset of astrocytes in the postnatal brain. *Development.* 2009;136(23):4021-31. doi:
669 10.1242/dev.041129. PubMed PMID: 19906869; PubMed Central PMCID: PMCPMC3118431.
- 670 4. Spassky N, Meunier A. The development and functions of multiciliated epithelia. *Nat*
671 *Rev Mol Cell Biol.* 2017;18(7):423-36. Epub 2017/04/13. doi: 10.1038/nrm.2017.21. PubMed
672 PMID: 28400610.
- 673 5. Gray RS, Roszko I, Solnica-Krezel L. Planar cell polarity: coordinating morphogenetic
674 cell behaviors with embryonic polarity. *Dev Cell.* 2011;21(1):120-33. doi:
675 10.1016/j.devcel.2011.06.011. PubMed PMID: 21763613; PubMed Central PMCID:
676 PMCPMC3166557.
- 677 6. Ohata S, Nakatani J, Herranz-Perez V, Cheng J, Belinson H, Inubushi T, et al. Loss of
678 Dishevelleds disrupts planar polarity in ependymal motile cilia and results in hydrocephalus.
679 *Neuron.* 2014;83(3):558-71. doi: 10.1016/j.neuron.2014.06.022. PubMed PMID: 25043421;
680 PubMed Central PMCID: PMCPMC4126882.
- 681 7. Banizs B, Pike MM, Millican CL, Ferguson WB, Komlosi P, Sheetz J, et al.
682 Dysfunctional cilia lead to altered ependyma and choroid plexus function, and result in the
683 formation of hydrocephalus. *Development.* 2005;132(23):5329-39. doi: 10.1242/dev.02153.
684 PubMed PMID: 16284123.
- 685 8. Lee L. Riding the wave of ependymal cilia: genetic susceptibility to hydrocephalus in
686 primary ciliary dyskinesia. *J Neurosci Res.* 2013;91(9):1117-32. doi: 10.1002/jnr.23238.
687 PubMed PMID: 23686703.
- 688 9. Petrik D, Myoga MH, Grade S, Gerkau NJ, Pusch M, Rose CR, et al. Epithelial Sodium
689 Channel Regulates Adult Neural Stem Cell Proliferation in a Flow-Dependent Manner. *Cell*
690 *Stem Cell.* 2018;22(6):865-78 e8. Epub 2018/05/22. doi: 10.1016/j.stem.2018.04.016. PubMed
691 PMID: 29779889.
- 692 10. Kousi M, Katsanis N. The Genetic Basis of Hydrocephalus. *Annu Rev Neurosci.*
693 2016;39:409-35. doi: 10.1146/annurev-neuro-070815-014023. PubMed PMID: 27145913.
- 694 11. Reiter JF, Leroux MR. Genes and molecular pathways underpinning ciliopathies. *Nat*
695 *Rev Mol Cell Biol.* 2017;18(9):533-47. Epub 2017/07/13. doi: 10.1038/nrm.2017.60. PubMed
696 PMID: 28698599; PubMed Central PMCID: PMCPMC5851292.
- 697 12. Verhey KJ, Kaul N, Soppina V. Kinesin assembly and movement in cells. *Annu Rev*
698 *Biophys.* 2011;40:267-88. doi: 10.1146/annurev-biophys-042910-155310. PubMed PMID:
699 21332353.
- 700 13. Hirokawa N, Noda Y, Tanaka Y, Niwa S. Kinesin superfamily motor proteins and
701 intracellular transport. *Nat Rev Mol Cell Biol.* 2009;10(10):682-96. doi: 10.1038/nrm2774.
702 PubMed PMID: 19773780.

14. Lechtreck KF. IFT-Cargo Interactions and Protein Transport in Cilia. *Trends Biochem Sci.* 2015;40(12):765-78. Epub 2015/10/27. doi: 10.1016/j.tibs.2015.09.003. PubMed PMID: 26498262; PubMed Central PMCID: PMC4661101.
15. Demonchy R, Blisnick T, Deprez C, Toutirais G, Loussert C, Marande W, et al. Kinesin 9 family members perform separate functions in the trypanosome flagellum. *J Cell Biol.* 2009;187(5):615-22. Epub 2009/12/02. doi: 10.1083/jcb.200903139. PubMed PMID: 19948486; PubMed Central PMCID: PMC2806587.
16. Niwa S, Nakajima K, Miki H, Minato Y, Wang D, Hirokawa N. KIF19A is a microtubule-depolymerizing kinesin for ciliary length control. *Dev Cell.* 2012;23(6):1167-75. Epub 2012/11/22. doi: 10.1016/j.devcel.2012.10.016. PubMed PMID: 23168168.
17. Li Y, Iakoubova OA, Shiffman D, Devlin JJ, Forrester JS, Superko HR. KIF6 polymorphism as a predictor of risk of coronary events and of clinical event reduction by statin therapy. *Am J Cardiol.* 2010;106(7):994-8. Epub 2010/09/22. doi: 10.1016/j.amjcard.2010.05.033. PubMed PMID: 20854963.
18. Assimes TL, Holm H, Kathiresan S, Reilly MP, Thorleifsson G, Voight BF, et al. Lack of association between the Trp719Arg polymorphism in kinesin-like protein-6 and coronary artery disease in 19 case-control studies. *J Am Coll Cardiol.* 2010;56(19):1552-63. Epub 2010/10/12. doi: 10.1016/j.jacc.2010.06.022. PubMed PMID: 20933357; PubMed Central PMCID: PMC3084526.
19. Buchan JG, Gray RS, Gansner JM, Alvarado DM, Burgert L, Gitlin JD, et al. Kinesin family member 6 (kif6) is necessary for spine development in zebrafish. *Dev Dyn.* 2014;243(12):1646-57. doi: 10.1002/dvdy.24208. PubMed PMID: 25283277.
20. Schultz J, Milpetz F, Bork P, Ponting CP. SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A.* 1998;95(11):5857-64. Epub 1998/05/30. PubMed PMID: 9600884; PubMed Central PMCID: PMC34487.
21. Schwenk F, Baron U, Rajewsky K. A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res.* 1995;23(24):5080-1. Epub 1995/12/25. PubMed PMID: 8559668; PubMed Central PMCID: PMC307516.
22. Spassky N, Merkle FT, Flames N, Tramontin AD, Garcia-Verdugo JM, Alvarez-Buylla A. Adult ependymal cells are postmitotic and are derived from radial glial cells during embryogenesis. *J Neurosci.* 2005;25(1):10-8. doi: 10.1523/JNEUROSCI.1108-04.2005. PubMed PMID: 15634762.
23. Pfenninger CV, Roschupkina T, Hertwig F, Kottwitz D, Englund E, Bengzon J, et al. CD133 is not present on neurogenic astrocytes in the adult subventricular zone, but on embryonic neural stem cells, ependymal cells, and glioblastoma cells. *Cancer Res.* 2007;67(12):5727-36. Epub 2007/06/19. doi: 10.1158/0008-5472.CAN-07-0183. PubMed PMID: 17575139.
24. Grimes DT, Boswell CW, Morante NF, Henkelman RM, Burdine RD, Ciruna B. Zebrafish models of idiopathic scoliosis link cerebrospinal fluid flow defects to spine curvature. *Science.* 2016;352(6291):1341-4. doi: 10.1126/science.aaf6419. PubMed PMID: 27284198.
25. Metscher BD. MicroCT for developmental biology: a versatile tool for high-contrast 3D imaging at histological resolutions. *Dev Dyn.* 2009;238(3):632-40. doi: 10.1002/dvdy.21857. PubMed PMID: 19235724.
26. Wullmann MF, Rupp B, Reichert H. *Neuroanatomy of the zebrafish brain : a topological atlas.* Basel ; Boston: Birkhäuser Verlag; 1996. vi, 144 p. p.

27. Kee HL, Dishinger JF, Blasius TL, Liu CJ, Margolis B, Verhey KJ. A size-exclusion permeability barrier and nucleoporins characterize a ciliary pore complex that regulates transport into cilia. *Nat Cell Biol.* 2012;14(4):431-7. doi: 10.1038/ncb2450. PubMed PMID: 22388888; PubMed Central PMCID: PMC3319646.
28. Hameed A, Bennett E, Ciani B, Hoebers LP, Milner R, Lawrie A, et al. No evidence for cardiac dysfunction in Kif6 mutant mice. *PLoS One.* 2013;8(1):e54636. doi: 10.1371/journal.pone.0054636. PubMed PMID: 23355886; PubMed Central PMCID: PMC3552957.
29. Liu Z, Gray RS. Animal models of idiopathic scoliosis. In: Kusumi K, Dunwoodie SL, editors. *The Genetics and Development of Scoliosis*. New York, NY: Springer Nature: Cham, Switzerland. ; 2018.
30. Karner CM, Long F, Solnica-Krezel L, Monk KR, Gray RS. Gpr126/Adgrg6 deletion in cartilage models idiopathic scoliosis and pectus excavatum in mice. *Hum Mol Genet.* 2015;24(15):4365-73. doi: 10.1093/hmg/ddv170. PubMed PMID: 25954032; PubMed Central PMCID: PMC4492399.
31. Schick RW, Matson DD. What is arrested hydrocephalus? *J Pediatr.* 1961;58:791-9. Epub 1961/06/01. PubMed PMID: 13747587.
32. Brinker T, Stopa E, Morrison J, Klinge P. A new look at cerebrospinal fluid circulation. *Fluids Barriers CNS.* 2014;11:10. Epub 2014/05/13. doi: 10.1186/2045-8118-11-10. PubMed PMID: 24817998; PubMed Central PMCID: PMC4016637.
33. de Ligt J, Willemsen MH, van Bon BW, Kleefstra T, Yntema HG, Kroes T, et al. Diagnostic exome sequencing in persons with severe intellectual disability. *N Engl J Med.* 2012;367(20):1921-9. Epub 2012/10/05. doi: 10.1056/NEJMoa1206524. PubMed PMID: 23033978.
34. Najmabadi H, Hu H, Garshasbi M, Zemojtel T, Abedini SS, Chen W, et al. Deep sequencing reveals 50 novel genes for recessive cognitive disorders. *Nature.* 2011;478(7367):57-63. Epub 2011/09/23. doi: 10.1038/nature10423. PubMed PMID: 21937992.
35. Poirier K, Lebrun N, Broix L, Tian G, Saillour Y, Boscheron C, et al. Mutations in TUBG1, DYNC1H1, KIF5C and KIF2A cause malformations of cortical development and microcephaly. *Nat Genet.* 2013;45(6):639-47. Epub 2013/04/23. doi: 10.1038/ng.2613. PubMed PMID: 23603762; PubMed Central PMCID: PMC3826256.
36. Willemsen MH, Ba W, Wissink-Lindhout WM, de Brouwer AP, Haas SA, Bienek M, et al. Involvement of the kinesin family members KIF4A and KIF5C in intellectual disability and synaptic function. *J Med Genet.* 2014;51(7):487-94. Epub 2014/05/09. doi: 10.1136/jmedgenet-2013-102182. PubMed PMID: 24812067.
37. Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool for the next generation of CRISPR genome engineering. *Nucleic Acids Res.* 2016;44(W1):W272-6. doi: 10.1093/nar/gkw398. PubMed PMID: 27185894; PubMed Central PMCID: PMC4987937.
38. Gagnon JA, Valen E, Thyme SB, Huang P, Ahkmetova L, Pauli A, et al. Efficient mutagenesis by Cas9 protein-mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. *PLoS One.* 2014;9(5):e98186. doi: 10.1371/journal.pone.0098186. PubMed PMID: 24873830; PubMed Central PMCID: PMC4038517.
39. Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, et al. A conditional knockout resource for the genome-wide study of mouse gene function. *Nature.*

794 2011;474(7351):337-42. doi: 10.1038/nature10163. PubMed PMID: 21677750; PubMed Central
795 PMCID: PMCPMC3572410.
796 40. Gray RS, Wilm TP, Smith J, Bagnat M, Dale RM, Topczewski J, et al. Loss of col8a1a
797 function during zebrafish embryogenesis results in congenital vertebral malformations. Dev Biol.
798 2014;386(1):72-85. doi: 10.1016/j.ydbio.2013.11.028. PubMed PMID: 24333517; PubMed
799 Central PMCID: PMC3938106.
800 41. Kawakami K. Tol2: a versatile gene transfer vector in vertebrates. Genome Biol. 2007;8
801 Suppl 1:S7. doi: 10.1186/gb-2007-8-s1-s7. PubMed PMID: 18047699; PubMed Central PMCID:
802 PMCPMC2106836.
803 42. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
804 open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. doi:
805 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMCPMC3855844.
806

Figure 1
Konjikusic et al.,

Figure 2
Konjikusic et al.,

Ventricle

Oviduct

Trachea

Figure 3
Konjikusic et al.,

Figure 4
Konjikusic et al.,

Figure 5
Konjikusic et al.,

Hs	YPELREALVSVWLQCTAISRIYTVGRS-FEGRELLVIELSDNPGVHEPGE
Pt	YPELREALVSVWLQCTAISRIYTVGRS-FEGRELLVIELSDNPGVHEPGE
Mc	YPELREALVSVWLQCTAISRIYTVGRS-FEGRELLVIELSDNPGVHEPGE
Mm	YPELREALVSVWLQCTAISRIYTVGRS-FEGRELLVIELSDNPGVHEPGE
Rn	YPELREALVSVWLQCTAISRIYTVGRS-FEGRELLVIELSDNPGVHEPGE
Bt	YPELREALVSVWLQCAAVSRIYTVGRS-FEGRELLVLELSDNPGVHEPGE
Cl	-----
Oc	GKSAVHSTATEWFLPGCS----MERP-VLG---CCPLMPKSLALFQEGE
Gg	YAE LREALVAVWLQCPAISRIYTVGRS-SEGRELLVIEVSDRPGHEPGE
Dr	YEEMRKSLVSVWLQCP SITRIYTVGES-FEGRELLVLEMSDNPGIHEPGE
Tn	YEELRKALVSVWLQCP TITRIYTIGES-FEGRELLVLEMSDNPGTHEPGE
Xt	YPEMRDALVAVWLQCP SISR IYTVGRS-FEGRELLVIEISDNPGHEPGE
Tc	NDELVQVLQDVNSRCPNITRVYTLTETSVLGLPLYLIEFSTKPGHHEIMK
Ce	QAQLEAKLGEINEKCPEITTLYEIGQS-VEGRPLVVIQFSTTPGEHIPTK
Sk	HEELKKVLDDTAAKCPDITRIYSPGQS-VEKRELWTIEISDKPGQHELGE

CPE (p.V79M)

D

CRISPR + donor injection F0 n (%)	
Total F0 mice weaned	64 (100)
Total genotyped with mutations	54 (84.4)
Total w/ integration of donor oligo	14 (22)
Total w/ hydrocephalus	5 (8)

E

F1 outcrosses with postive integration of donor oligo			
Mouse	ratio litter1 (%): ratio litter 2 (%)	Total pups in litter 1, litter 2	Total transmission %
Female 1 (MK53)	0 (0) : 0 (0)	4, 9	0
Female 2 (MK63)	5 (62.5) : 3 (50)	8, 6	57.1
Female 3 (MK61)	1 (11) : 5 (50)	9, 10	31.6
Male 1 (MK68)	2 (50) : 0 (0)	4, 4	25
Male 2 (MK86)	4 (57) : 4 (44)	7, 4	72.7

Control

Kif6^{p.G555fs/p.G555fs}

Supp. Fig. SF4
Konjikusic et al.,

Supp. Fig. SF5
Konjikusic et al.,

----- WT -----

----- Animal #1 -----

--- Animal #2 ---

Supplementary Table SI. Eighty three homozygous variants from WES

Chromosome	Position_start	Position_end	Gene	Change
1	26608891	26608896	<i>UBXN11</i>	nonframeshift_deletion
1	54605319	54605319	<i>CDCP2</i>	frameshift_insertion
1	152681694	152681694	<i>LCE4A</i>	nonframeshift_insertion
1	156565050	156565050	<i>GPATCH4</i>	frameshift_insertion
1	160650923	160650923	<i>CD48</i>	frameshift_deletion
2	96148317	96148317	<i>TRIM43B</i>	nonsynonymous_SNV
2	130902532	130902532	<i>CCDC74B</i>	nonsynonymous_SNV
2	240981597	240981597	<i>PRR21</i>	nonsynonymous_SNV
2	240981627	240981627	<i>PRR21</i>	nonsynonymous_SNV
2	240981655	240981655	<i>PRR21</i>	nonsynonymous_SNV
2	241621800	241621800	<i>AQP12B</i>	frameshift_deletion
3	40503552	40503552	<i>RPL14</i>	nonframeshift_insertion
3	53324829	53324834	<i>DCP1A</i>	nonframeshift_deletion
3	56650056	56650056	<i>CCDC66</i>	nonframeshift_insertion
3	75786586	75786586	<i>ZNF717</i>	nonsynonymous_SNV
3	75786662	75786662	<i>ZNF717</i>	nonsynonymous_SNV
3	75786672	75786672	<i>ZNF717</i>	nonsynonymous_SNV
3	75786684	75786684	<i>ZNF717</i>	nonsynonymous_SNV
3	75786764	75786764	<i>ZNF717</i>	frameshift_deletion
3	75790810	75790810	<i>ZNF717</i>	frameshift_insertion

3	195505930	195505930	<i>MUC4</i>	nonsynonymous_SNV
3	195506099	195506099	<i>MUC4</i>	nonsynonymous_SNV
3	195506147	195506147	<i>MUC4</i>	nonsynonymous_SNV
3	195506156	195506156	<i>MUC4</i>	nonsynonymous_SNV
3	195506267	195506267	<i>MUC4</i>	nonsynonymous_SNV
3	195506507	195506507	<i>MUC4</i>	nonsynonymous_SNV
3	195506530	195506530	<i>MUC4</i>	nonsynonymous_SNV
3	195506704	195506704	<i>MUC4</i>	nonsynonymous_SNV
3	195507226	195507226	<i>MUC4</i>	nonsynonymous_SNV
3	195510310	195510310	<i>MUC4</i>	nonsynonymous_SNV
3	195510613	195510613	<i>MUC4</i>	nonsynonymous_SNV
3	195512042	195512042	<i>MUC4</i>	nonsynonymous_SNV
3	195514558	195514558	<i>MUC4</i>	nonsynonymous_SNV
3	195514733	195514733	<i>MUC4</i>	nonsynonymous_SNV
3	195514768	195514768	<i>MUC4</i>	nonsynonymous_SNV
4	166300608	166300608	<i>CPE</i>	nonsynonymous_SNV
5	139931628	139931628	<i>SRA1</i>	frameshift_insertion
5	140307142	140307142	<i>PCDHAC1</i>	nonsynonymous_SNV
5	140725635	140725635	<i>PCDHGA3</i>	nonsynonymous_SNV
5	149512332	149512332	<i>PDGFRB</i>	nonsynonymous_SNV
6	16327953	16327955	<i>ATXN1</i>	nonframeshift_deletion
6	39513453	39513453	<i>KIF6</i>	frameshift_deletion
6	41895234	41895234	<i>BYSL</i>	nonsynonymous_SNV

6	43970526	43970531	<i>C6orf223</i>	nonframeshift_deletion
7	1586661	1586661	<i>TMEM184A</i>	nonframeshift_insertion
7	15725824	15725826	<i>MEOX2</i>	nonframeshift_deletion
7	44040763	44040763	<i>SPDYE1</i>	nonsynonymous_SNV
7	44120345	44120345	<i>POLM</i>	nonsynonymous_SNV
7	64167644	64167644	<i>ZNF107</i>	nonsynonymous_SNV
7	64169019	64169019	<i>ZNF107</i>	frameshift_insertion
7	100639147	100639147	<i>MUC12</i>	nonsynonymous_SNV
7	100639153	100639153	<i>MUC12</i>	nonsynonymous_SNV
7	100646440	100646440	<i>MUC12</i>	nonsynonymous_SNV
7	100647338	100647338	<i>MUC12</i>	nonsynonymous_SNV
7	100647339	100647339	<i>MUC12</i>	nonsynonymous_SNV
7	100647376	100647376	<i>MUC12</i>	nonsynonymous_SNV
7	143270465	143270465	<i>CTAGE15P</i>	nonsynonymous_SNV
8	7629232	7629232	<i>FAM90A10</i>	nonsynonymous_SNV
8	144511981	144511983	<i>MAFA</i>	nonframeshift_deletion
9	79318378	79318392	<i>PRUNE2</i>	nonframeshift_deletion
9	90534190	90534202	<i>FAM75C1</i>	frameshift_substitution
9	90746922	90746922	<i>FAM75C2</i>	nonsynonymous_SNV
9	97080947	97080949	<i>FAM22F</i>	nonframeshift_deletion
9	107367666	107367667	<i>OR13C2</i>	frameshift_deletion
10	27342292	27342294	<i>ANKRD26</i>	nonframeshift_deletion
10	50535007	50535007	<i>C10orf71</i>	frameshift_insertion

10	119302954	119302956	<i>EMX2</i>	nonframeshift_deletion
10	135438687	135438687	<i>FRG2B</i>	nonsynonymous_SNV
10	135438929	135438929	<i>FRG2B</i>	nonsynonymous_SNV
10	135438943	135438943	<i>FRG2B</i>	nonsynonymous_SNV
10	135438961	135438961	<i>FRG2B</i>	nonsynonymous_SNV
11	65636053	65636053	<i>EFEMP2</i>	nonsynonymous_SNV
13	72440664	72440664	<i>DACHI</i>	nonsynonymous_SNV
15	34825154	34825154	<i>GOLGA8B</i>	nonsynonymous_SNV
15	82637448	82637448	<i>GOLGA6L10</i>	nonsynonymous_SNV
19	1556482	1556482	<i>MEX3D</i>	nonsynonymous_SNV
19	13318707	13318712	<i>CACNA1A</i>	nonframeshift_deletion
19	20807178	20807178	<i>ZNF626</i>	frameshift_insertion
19	22156850	22156850	<i>ZNF208</i>	nonsynonymous_SNV
19	46627337	46627337	<i>IGFL3</i>	nonsynonymous_SNV
20	126314	126315	<i>DEFB126</i>	frameshift_deletion
20	238439	238444	<i>DEFB132</i>	nonframeshift_deletion
20	32664865	32664865	<i>RALY</i>	nonframeshift_insertion

Supplementary Table SII. Sixty three homozygous regions from homozygosity mapping

Chromosome	SNP1	SNP2	Position 1	Position 2	Size (Kb)
1	kgp9226796	kgp15281431	42,272,542	43,304,041	1031.499
1	rs12043872	kgp2176506	50,644,891	52,222,874	1577.983
1	kgp22807508	kgp15446070	175,901,363	177,085,495	1184.132
2	rs13383059	kgp1338077	60,542,657	61,840,186	1297.529
2	kgp22835335	rs1345516	63,283,437	64,703,286	1419.849
2	kgp10464820	kgp11645441	72,356,303	73,389,904	1033.601
2	kgp22748117	rs11683207	95,350,864	98,333,290	2982.426
2	kgp11241972	kgp14721953	110,452,502	111,457,965	1005.463
2	kgp14255701	kgp4421106	155,497,134	156,765,691	1268.557
3	rs2242150	kgp920744	48,505,964	50,176,739	1670.775
3	rs1528197	kgp5972268	62,404,892	63,669,341	1264.449
3	kgp7194326	rs17024881	84,835,064	86,118,949	1283.885
3	rs6804377	kgp10736192	89,271,087	90,501,225	1230.138
3	kgp17732198	kgp3463695	186,780,268	195,465,310	8685.042
3	rs3747673	kgp22812098	195,611,844	197,891,568	2279.724

4	rs922333	kgp20990514	64,121,298	65,135,489	1014.191
4	kgp12219459	kgp21246176	155,003,651	182,666,898	27663.247
5	rs11948368	kgp6573465	4,225,004	10,668,729	6443.725
5	kgp3769852	rs13184580	42,912,540	43,918,326	1005.786
5	kgp3714973	kgp12345716	132,628,429	141,015,519	8387.09
5	rs41098	kgp22527688	141,020,100	149,585,074	8564.974
5	kgp22340708	kgp10344299	149,587,580	172,127,143	22539.563
5	kgp22126080	kgp3444559	172,127,834	173,730,182	1602.348
6	rs12663002	rs3094575	28,441,634	29,515,802	1074.168
6	kgp9471913	kgp17000708	34,187,366	43,010,582	8823.216
6	rs9342711	kgp8490913	69,197,250	70,393,636	1196.386
6	rs6926330	kgp17199538	91,038,726	108,320,258	17281.532
6	kgp6823635	kgp10978549	126,186,643	127,216,752	1030.109
7	rs17714729	kgp13497896	40,925,373	44,100,951	3175.578
7	kgp13393169	kgp1758811	44,282,204	58,042,660	13760.456
7	kgp13613108	kgp7238344	61,055,273	63,041,496	1986.223
7	kgp22797960	kgp1628657	63,234,502	64,474,466	1239.964

7	kgp8836768	kgp6578278	64,476,704	66,158,076	1681.372
7	kgp11959682	kgp731589	66,835,845	73,110,455	6274.61
7	kgp13574770	kgp13241424	73,254,871	75,614,264	2359.393
7	rs782487	rs41542	76,161,400	93,722,036	17560.636
8	rs35292150	rs10097659	7,154,036	8,241,316	1087.28
8	kgp20383958	kgp20415047	47,840,086	50,247,347	2407.261
8	kgp4568187	kgp4989542	50,287,180	51,332,763	1045.583
8	kgp11163141	kgp8120576	83,652,903	84,739,570	1086.667
8	kgp1752758	rs6468704	99,405,919	100,994,389	1588.47
8	rs1318739	kgp3429874	104,122,872	105,175,235	1052.363
8	kgp20054401	kgp10669054	114,857,226	115,996,248	1139.022
9	kgp3876298	kgp18506353	66,771,643	71,032,042	4260.399
10	kgp22751182	kgp22827934	73,974,125	75,401,246	1427.121
10	kgp28675	kgp1752039	128,613,842	133,791,976	5178.134
11	rs7129994	kgp12754305	84,380,034	85,620,335	1240.301
12	kgp1178574	rs7311759	111,816,925	113,261,665	1444.74
13	kgp7808335	rs8002509	36,452,369	47,822,976	11370.607

13	rs9568798	kgp16817285	53,614,554	54,687,807	1073.253
13	kgp9865216	kgp16686988	61,860,814	62,932,723	1071.909
13	kgp7743232	rs9523513	65,234,035	92,781,385	27547.35
13	kgp1128060	rs16953570	96,133,164	97,438,801	1305.637
15	rs4508402	rs4779824	30,361,587	31,404,294	1042.707
15	kgp19953534	kgp19917981	64,125,067	65,168,281	1043.214
16	kgp6578578	kgp22822345	31,888,867	33,579,417	1690.55
16	kgp10923077	kgp16403964	46,920,923	48,045,778	1124.855
17	kgp22779423	kgp14115230	57,513,947	59,233,842	1719.895
18	kgp9512908	kgp6363611	51,184,594	52,469,636	1285.042
19	rs1144539	kgp7929656	37,331,614	38,475,123	1143.509
20	kgp19251773	rs6061136	29,423,716	30,644,465	1220.749
20	kgp19300395	rs2425193	32,830,486	34,848,116	2017.63
21	kgp6761551	kgp7702343	41,544,215	44,477,135	2932.92

<i>Kif6^{tm1a(KOMP)Wtsi}</i> Cassette primers	
CSD-lacF	GCTACCATTACCAGTTGGTCTGGTGTC
CSD-neoF	GGGATCTCATGCTGGAGTTCTTCG
CSD-loxF	GAGATGGCGCAACGCAATTAATG
<i>Kif6^{tm1a(KOMP)Wtsi}</i> Gene Specific Primers	
CSD-Kif6-R	GGTTAGGAGGAAGAGAAGGGCATCC
CSD-Kif6-ttR	ACAGATGCTGGAGATCACACTCTCG
CSD-Kif6-F	ACTCTCTTCAAAGCCCACATCATGC
Mouse CRISPR oligos/genotyping primers	
mKif6-R2-ex14-T7	TAATACGACTCACTATAGGAGATGTCACTGGGACGCCGTTTTAGAGCTAGAAATAGC
Universal T7 tracer	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTAACTTGC TATTTCTAGCTCTAAAC
Ms_ <i>Kif6</i> crispr_r2_do nor oligo	GAGAGGAGATGTCACTGGGAGTCATGGCGTTTAAACCTTAATTAAGCTGTTGTAGCGCCAG GAGGCTTTTGAGAT
Mus_ <i>Kif6</i> _ex14F	TCCCAAAATGATGTGACTGAAG
Mus_ <i>Kif6</i> _ex14R	AGTCTCTGGACTGGCTTACCTG
<i>Kif6p.G555fs</i> 3stopDonor_FWD	CATGGCGTTTAAACCTTAATTAAGCTG
<i>Kif6p.G555fs</i> 3stopDonor_REV	CAGCTTAATTAAGGTTTAAACGCCATG
Mouse Kif6 qPCR primer sets	
Ms_ <i>Kif6</i> qPCR_exon8_ FWD1	TCGGAAAAACACCGTACACA
Ms_ <i>Kif6</i> qPCR_exon10_ _REV1	CTTTTGCAAGCGAACAATCA
Ms_ <i>Kif6</i> qPCR_exon7_ FWD2	TTCAACCCGGTCACACTGTA
Ms_ <i>Kif6</i> qPCR_exon8_ REV2	TACGGTGTTTTCCGAAAGG
Ms_ <i>Kif6</i> qPCR_exon6_ FWD3	TGGAGGACCCTGATCAGAAC
Ms_ <i>Kif6</i> qPCR_exon7_ REV3	TACAGTGTGACCGGGTTGAA