SpoT and GppA hydrolases prevent the gratuitous activation of RelA by pppGpp in *Escherichia coli*.

Rajeshree Sanyal^{1,2} and Rajendran Harinarayanan^{1,*}

¹Laboratory of Bacterial Genetics, Centre for DNA Fingerprinting and Diagnostics, Hyderabad, India.

²Graduate studies, Manipal Academy of Higher Education, Manipal, India.

Short title: RelA activation by pppGpp.

Keywords: GppA / (p)ppGpp / RelA / SpoT / stringent response

* Corresponding author

E-mail: hari@cdfd.org.in (RH)

20 **Summary**

21 Stringent response, a conserved regulation seen in bacteria, is effected through the modified
 22 nucleotides (p)ppGpp. The metabolic cycle of these molecules is driven by the synthase
 23 activity of RelA and SpoT and the hydrolase activity of SpoT and GppA which together sets
 24 the basal (p)ppGpp pool. Growth arrest due to (p)ppGpp accumulation from basal RelA
 25 activity apparently explained the essentiality of SpoT hydrolase function. We found, pppGpp
 26 degradation was enhanced when the SpoT hydrolase activity was lowered or eliminated and
 27 when this was alleviated by inactivation of the GppA hydrolase, gratuitous synthesis of
 28 (p)ppGpp by RelA was activated, leading to growth arrest. The RelA-ribosome interaction
 29 was not mandatory for these phenotypes. Our results show, for the first time, elevated
 30 pppGpp promoted the amplification of RelA-mediated stringent response in the absence of
 31 established RelA activating signals in the cell and the SpoT and GppA hydrolases prevented
 32 this. The accumulation of pppGpp inhibited the SpoT hydrolase activity. We propose this
 33 autocatalytic activation of RelA by pppGpp is likely to be an allosteric regulation and can
 34 result in a bistable switch.

35 **Introduction**

36 Intracellular signaling molecules play a key role in the physiology of organisms by regulating
 37 key cellular processes and coordinating them with extracellular or intracellular signals
 38 (Pesavento and Hengge, 2009). In eubacteria, (p)ppGpp is a signaling molecule that
 39 accumulates during starvation, switching the balance of metabolism from growth and cell
 40 division to survival and stress response (Chatterji and Kumar Ojha, 2001; Braeken *et al.*,
 41 2006; Potrykus and Cashel, 2008; Hauryliuk *et al.*, 2015). (p)ppGpp, also sometimes referred
 42 to as an ‘alarmone’, consist of a pair of molecules – guanosine tetraphosphate (ppGpp) and
 43 guanosine pentaphosphate (pppGpp), which are synthesized by the transfer of a

44 pyrophosphate moiety from ATP to GDP or GTP respectively (Potrykus and Cashel, 2008;
45 Atkinson *et al.*, 2011). The gram-negative bacteria *Escherichia coli* is a well-studied model
46 organism with regard to the role of (p)ppGpp in bacterial physiology. β - and γ -proteobacteria,
47 including *E. coli*, have two proteins which are involved in stress response and (p)ppGpp
48 metabolism – RelA and SpoT (Mittenhuber, 2001; Atkinson *et al.*, 2011). Both proteins are
49 members of the Rel/Spo homolog (RSH) family and share similar domain architecture. The
50 N-terminal half of the protein has the catalytic domain with the (p)ppGpp synthase and
51 hydrolase functions in the case of SpoT and only synthase function in case of RelA. The C-
52 terminal half of the protein has the regulatory domains important for sensing stress and
53 starvation signals. RelA is a ribosome-bound protein that is activated by the “hungry” codons
54 that appear following amino acid starvation and the consequent increase in the cellular
55 concentration of uncharged t-RNA (Wendrich *et al.*, 2002). Recent cryo-electron microscopy
56 studies have provided important insights into the structural basis for RelA activation by the
57 entry of uncharged tRNA into the A-site of an elongating ribosome (Arenz *et al.*, 2016;
58 Brown *et al.*, 2016; Loveland *et al.*, 2016). SpoT has a weak synthetic activity and a strong
59 (p)ppGpp hydrolase activity and is an essential gene (An *et al.*, 1979; Xiao *et al.*, 1991). It
60 not only regulates the basal (p)ppGpp levels in the cell (Sarubbi *et al.*, 1988), but also
61 responds to various stress and starvation signals such as carbon (Xiao *et al.*, 1991), fatty acid
62 (Seyfzadeh *et al.*, 1993) and iron (Vinella *et al.*, 2005) limitation. The SpoT hydrolase
63 activity was inhibited in the presence of uncharged tRNA and more severely in the presence
64 of ribosomes (Richter, 1980), conditions that mimic amino acid starvation. Various factors
65 interact with SpoT and regulate the balance between its synthase and hydrolase functions.
66 CgtA, a G-protein interacts with SpoT to up-regulate its hydrolase activity under nutrient-rich
67 condition (Jiang *et al.*, 2007). The acyl carrier protein was reported to interact with SpoT and
68 up-regulates its synthase activity during fatty acid starvation (Battesti and Bouveret, 2006). In

addition to SpoT, pppGpp is hydrolyzed by GppA, a pentaphosphate phosphohydrolase that converts it into ppGpp (Somerville and Ahmed, 1979; Harat and Sy, 1983; Keasling *et al.*, 1993). The physiological relevance of this reaction is not clear. ppGpp being the predominant stringent nucleotide in the cell is considered to be the principal effector of stringent response (Mechold *et al.*, 2013).

The primary and well-studied target for (p)ppGpp is RNA polymerase, to which it binds independently or aided by the protein factor DksA (Ross *et al.*, 2013; Zuo *et al.*, 2013; Ross *et al.*, 2016) and alters the global transcriptome so as to promote survival under starvation or stress (Durfee *et al.*, 2008; Traxler *et al.*, 2012). Various models exist to explain the effects of (p)ppGpp at the level of transcription initiation or sigma factor competition (Barker *et al.*, 2001; Paul *et al.*, 2004; Magnusson *et al.*, 2005). Together with DksA, (p)ppGpp inhibits transcription of rRNA operons and ribosomal protein genes and activates amino acid biosynthetic genes (Paul *et al.*, 2004; Paul *et al.*, 2005). Studies have shown alternative targets for (p)ppGpp in *E. coli* and in other organisms (Kanjee *et al.*, 2012). This includes enzymes from the nucleotide biosynthesis pathway, GTPases, particularly ObgE, EFG, EFTu, RF3 and IF2 (Miller *et al.*, 1973; Hamel and Cashel, 1974; Rojas *et al.*, 1984; Milon *et al.*, 2006). DNA replication and cell division are regulated by (p)ppGpp (Schreiber *et al.*, 1995; Joseleau-Petit *et al.*, 1999; Wang *et al.*, 2007; Ferullo and Lovett, 2008).

Due to the essential nature of the SpoT function, studies have generally been performed using point mutants that altered the steady-state levels of ppGpp. Using a SpoT depletion system, changes in the (p)ppGpp metabolic pattern and the associated growth response can be monitored. In this study, we report, during SpoT depletion or in the hydrolase mutants of *spoT*, there was reduction in the pppGpp pool and this was required to support growth. Our results show, GppA and an uncharacterized activity together lowered the pppGpp pool and prevented the gratuitous activation of RelA-dependent (p)ppGpp synthesis that conferred

growth arrest. Two hypomorphic *relA* alleles that allowed GppA-dependent survival in the absence of SpoT function were isolated and characterized. Using one of the alleles, we show that the accumulation of only ppGpp was insufficient to confer sustained growth arrest. Our results show pppGpp can serve as a positive signal for the amplification of RelA-mediated stringent response in the absence of known documented signals and therefore its level is tightly regulated by the SpoT and GppA hydrolases.

Results

Depletion of SpoT results in the enhanced degradation of pppGpp

The *spoT* gene function that is essential in the wild-type *E. coli* was dispensable in the *relA* mutant (Xiao *et al.*, 1991), and therefore, the essential *spoT* function was inferred as the degradation of (p)ppGpp synthesized through the basal RelA activity. To test this, (p)ppGpp accumulation was followed as the cellular SpoT activity was reduced using a system designed to gradually deplete it. To do this, the chromosomal *spoT* gene was knocked out and SpoT was expressed in a regulated fashion from a plasmid (see methods). For ease of genetic manipulations, either of the two null mutant alleles reported in the literature, namely, *spoT212*, a markerless *spoT* deletion or *spoT207::Cm* have been used in strain constructions. In the $\Delta spoT/pRCspoT$ strain wherein the chromosomal *spoT* gene was deleted, growth was expected to be dependent on the IPTG-driven expression of the *spoT* gene present in the single copy unstable plasmid pRC*spoT* (Nazir and Harinarayanan, 2016). When the $\Delta spoT/pRCspoT$ strain, cultured in the presence of IPTG was washed and sub-cultured, IPTG-dependent growth was observed (Fig. 1A) indicating that the system can be used to study the consequences of SpoT depletion.

To monitor (p)ppGpp levels, cells cultured in the presence or absence of IPTG were allowed to undergo at least two divisions in the low phosphate medium containing $^{32}\text{P-H}_3\text{PO}_4$ and

subjected to thin layer chromatography. This method allowed (p)ppGpp levels to be measured when the growth of the strain was retarded due to the depletion of SpoT (Fig. 1B and 1C). Following SpoT depletion, ppGpp accumulated, but pppGpp was not detectable. We tested different dilutions in order to deplete SpoT to varying extents, and in each case, the result was similar (data not shown). Unlike the $\Delta spoT/pRCspoT$ strain, the growth of the $\Delta relA \Delta spoT/pRCspoT$ strain was less significantly inhibited following SpoT depletion (Fig. S1) and (p)ppGpp accumulation was not detected here (data not shown). Nevertheless, the growth rate of the strain was reduced in the absence of IPTG, which could be due to the composition of the growth medium used (all amino acids at 40 μ g/ml). It has been reported, the growth of the ppGpp⁰ strain in a defined medium was sensitive to amino acid composition (Potrykus *et al.*, 2011).

Due to the presence of amino acids in the growth medium, the signal for RelA activation would be low, this, and the conversion of pppGpp to ppGpp by GppA can together account for the lack of detectable pppGpp accumulation. To test this, isoleucine starvation was provoked by the addition of valine (Leavitt and Umbarger, 1961) before or after SpoT depletion and the (p)ppGpp synthesis was studied. In the culture induced for *spoT* expression, a typical stringent response, that is, accumulation of ppGpp and pppGpp and depletion of GTP was observed (Fig. 1D, lanes 1 – 4). This indicated, the SpoT expression from plasmid did not significantly perturb the stringent response. Following SpoT depletion, a significant accumulation of ppGpp was seen prior to starvation and this further increased with the onset of amino acid starvation; interestingly, pppGpp was still not detectable (Fig. 1D, lanes 5 – 8). This indicated, pppGpp accumulation was especially reduced, as compared to ppGpp during SpoT depletion. The absence of pppGpp could be due to its reduced synthesis or increased degradation or both. The results described below show the GppA hydrolase function was partly required to observe this phenotype. In early studies (Laffler and

Gallant, 1974; Fiil *et al.*, 1977), it was reported that the hydrolase deficient *spoT1* allele accumulated mainly ppGpp during amino acid starvation. We confirmed that following amino acid starvation, the *spoT1* mutant, like the SpoT depleted strain, showed RelA-dependent accumulation of ppGpp, but not pppGpp (Fig. S2). These results suggest, counter-intuitively, lowering the SpoT hydrolase activity lowered the pppGpp accumulation.

Hypomorphic relA alleles provide evidence for concomitant synthesis and degradation of ppGpp during stringent response following amino acid starvation

To identify mutations that suppressed the growth defect of the $\Delta spoT$ strain, transposon-mediated mutagenesis was carried out in the $\Delta spoT::Cm/pRCspoT$ strain and mutants that survived the loss of pRC*spoT* were identified as white colonies from the LB IPTG X-Gal plates. In order to further identify the loss of function *relA* mutants amongst these, the white colonies were screened using the SMG plate test for RelA functionality (Uzan and Danchin, 1976). The *relA* null mutant or mutants that exhibit very low activity (such as *relA1*) do not grow on minimal medium with glucose and the amino acids serine, methionine, and glycine (SMG) (Fig. S3, rows 1 and 2). About 50% of the white colonies screened showed SMG sensitivity and were excluded from further analysis. The insertions in two separate SMG-resistant mutants were mapped and used for further studies. Somewhat surprisingly, one of the transposon insertions was within the *relA* gene (after the 496th codon) and the other at the end of the *rlmD* gene that is immediately upstream of *relA* (Fig. 2A). These insertions, referred to as *relA::Tn10dTet* and *rlmD::Tn10dKan*, when individually introduced into the $\Delta spoT::Cm/pRCspoT$ strain by phage P1 transduction, allowed the segregation of white colonies on LB agar plates with IPTG and X-Gal (Fig. 2B, panels ii and iii), suggesting the presence of each mutation was sufficient to support growth in the absence of SpoT function. The *relA::Tn10dTet* $\Delta spoT::Cm$ and *rlmD::Tn10dKan* $\Delta spoT::Cm$ strains grew on minimal glucose medium with or without SMG, while the *spoT*⁺ derivatives did not grow in the

presence of SMG (Fig. S3, rows 3, 4, 9 and 10). These results showed that a certain threshold of RelA activity and possibly (p)ppGpp accumulation can be tolerated in the absence of SpoT function. The *relA1* allele also survives the *spoT* deletion and the *relA1 ΔspoT* strain grows in minimal medium (Xiao *et al.*, 1991) suggesting the presence of (p)ppGpp in this strain. *relA1* is a naturally selected hypomorphic allele present in the laboratory strains, wherein two polypeptides come together to reconstitute a weak (p)ppGpp synthase activity, which is evident upon over-expression (Metzger *et al.*, 1989). Unlike *relA1*, the *relA1 ΔspoT* strain could grow slowly on SMG plate, with single colonies observed after 72 hrs (Fig. S4A). These results indicated that the basal (p)ppGpp pool increased in the absence of SpoT function in all the hypomorphic *relA* alleles tested.

The transposon insertion after the 496th codon of *relA* generated a UAA stop codon at the site of insertion and therefore was expected to produce a truncated polypeptide containing the 496 amino-terminal amino acids (RelA496Δ). A plasmid-encoded RelA polypeptide with the 455 amino-terminal amino acids, expressed from the IPTG-inducible P_{tac} promoter showed constitutive (p)ppGpp synthesis and was incapable of sensing amino acid starvation; the polypeptide had a half-life of 7.5 minutes as compared to the 2-3 hours for the full-length RelA protein (Schreiber *et al.*, 1991; Svitils *et al.*, 1993). To compare the phenotypes of the RelA truncations, precise deletions were engineered in the chromosomal *relA* gene. These alleles referred as *relA496Δ::Kan* and *relA455Δ::Kan*, when introduced in the *ΔspoT/pRCspoT* strain could support growth in the absence of SpoT function, which was evident from the appearance of white colonies (Fig. 2B, panel iv, and v) in the plasmid segregation assay. The white colonies with the *relA496Δ::Kan ΔspoT* and *relA455Δ::Kan ΔspoT* genotypes exhibited SMG-resistance unlike their *spoT*⁺ counterparts (Fig. S3, rows 5 – 8), and suggested that the absence of SpoT function elevated the cellular (p)ppGpp pool in these strains. Indeed, a slightly elevated basal ppGpp pool was observed in the SMG-resistant

relA::Tn10dTet ΔspoT::Cm strain when compared to the SMG-sensitive *relA::Tn10dTet spoT⁺* strain (Fig. 2C (i), lanes 1 and 5). Amino acid starvation did not elicit stringent response in either strain (Fig. 2C (i), lanes 2 – 4 and 6 – 8), and this is consistent with the reports that the RelA-CTD is required for sensing the ribosome-dependent amino acid starvation signals (Arenz *et al.*, 2016; Brown *et al.*, 2016; Loveland *et al.*, 2016). These results suggest, RelA' (truncated RelA), when expressed at physiological levels could not elevate the (p)ppGpp pool in the presence of SpoT hydrolase activity.

The transposon insertion *rlmD::Tn10dKan*, 13-bp upstream from the *rlmD* stop codon (Fig. 2A) can potentially alter the RelA expression and also the RlmD activity. To test if the suppression of *ΔspoT* growth defect resulted from the loss of RlmD function, the *ΔrlmD::Kan* allele from the Keio collection (Baba *et al.*, 2006) was introduced in the wild-type and *ΔspoT/pRCspoT* strains by Phage P1 transduction. Suppression was not evident in the *ΔrlmD::Kan ΔspoT/pRCspoT* strain as it did not segregate white colonies on LB IPTG X-Gal plate (Fig. 2B, panel vi), however, flip-out of the Kanamycin marker supported the segregation of white colonies (Fig. 2B, panel vii) and the growth of these colonies on minimal medium containing SMG (Fig. S3, rows 11 – 13). These results showed that the loss of RlmD function was not necessary for suppression, while the altered *relA* expression after the flip-out of the Kan cassette was sufficient for the suppression of the *ΔspoT* growth defect. We propose the *rlmD::Tn10dKan* and *ΔrlmD::FRT* mutations reduced the *relA* expression, the former possibly terminated transcripts originating from the promoters within and upstream of *rlmD* (Metzger *et al.*, 1988; Nakagawa *et al.*, 2006; Brown *et al.*, 2014), and the latter eliminated the promoters reported within the *rlmD* ORF. Making use of *lacZ* reporter fusions at the chromosomal locus of these genes, the relative contribution of the promoters located within the *rlmD* ORF to those upstream was studied (Fig. S5). The promoters within the *rlmD* ORF increased the expression 75-fold, indicating that they predominantly

contributed to the RelA expression. The reason, the $\Delta rlmD::Kan$ allele failed to suppress the $\Delta spoT$ growth defect could be because promoter(s) within the Kan^R cassette partially or fully compensated for the loss of expression from the native *relA* promoters.

SMG resistance was observed in the *rlmD::Tn10dKan* $\Delta spoT::Cm$ strain but not in the *rlmD::Tn10dKan* strain (Fig. S3, rows 9 and 10), however, no significant difference in basal ppGpp is evident between the two strains (Fig. 2C (ii), lanes 1 and 5). Similarly, the $\Delta rlmD::FRT$ strain showed SMG-resistance (Fig. S3, row 12), but increase in the basal ppGpp pool was not evident (Fig. 2C (iii) lane 1). Our results suggest, marginal increase in the ppGpp pool less than that which can be reliably detected by 1D-TLC was sufficient to confer SMG-resistance. The $\Delta rlmD::FRT$ $\Delta spoT$ strain showed SMG-resistance (Fig. S3, row 13) and a slightly elevated basal ppGpp pool (Fig. 2C (iii), lane 5). Amino acid starvation did not cause (p)ppGpp accumulation in the *rlmD::Tn10dKan*, *rlmD::Tn10dKan* $\Delta spoT::Cm$ and $\Delta rlmD::FRT$ strains (Fig. 2C (ii) lanes 2 – 4 & 6 – 8; Fig. 2C (iii) lanes 2 – 4). However, stringent response was elicited in the $\Delta rlmD::FRT$ $\Delta spoT$ strain and accumulation of ppGpp, but not pppGpp was observed (Fig. 2C (iii), 6 – 8); the rate of ppGpp accumulation was slower as compared to the wild-type or *spoT1* strain (compare Fig. 1D, lane 2; Fig. S2 lane 2 and Fig. 2C (iii) lane 6). This result provides evidence for the SpoT-mediated degradation of ppGpp being concomitant with the RelA-mediated synthesis of (p)ppGpp during stringent response. This was not entirely expected, because, the SpoT hydrolase activity was reported to be inhibited during the amino acid-induced stringent response (Richter, 1980). The concomitant synthesis and degradation of (p)ppGpp imply the concentration of these molecules at any given time is set by the sum total of their synthesis and degradation rates. This would have implications for the amplification of the stringent response, because, it was reported that (p)ppGpp increased the rate of their own ribosome-dependent synthesis by RelA *in vitro* (Shyp *et al.*, 2012; Kudrin *et al.*, 2018). The results imply, the SpoT hydrolase

activity can directly contribute to the regeneration of GDP from stringent nucleotides during the stringent response and indirectly to GTP as it was proposed that GDP would be converted to GTP by Ndk (Kari *et al.*, 1977; Heinemeyer and Richter, 1978).

GppA activity prevented gratuitous (p)ppGpp synthesis and growth arrest in strains with reduced spoT hydrolase activity

The accumulation of ppGpp but not pppGpp, after SpoT depletion or reduction in the hydrolase activity (*spoT1* mutant), suggested altered (p)ppGpp metabolism. One possibility was, RelA preferentially synthesized ppGpp under these conditions. Results in the previous section highlighted the role of SpoT hydrolase activity in the regeneration of GTP during the stringent response, therefore it was possible, pppGpp synthesis was limited by GTP availability. Furthermore, there was evidence *in vitro* that the *E. coli* RelA prefers GDP as a substrate over GTP (Sajish *et al.*, 2009). We, therefore, asked if increasing the cellular GTP pool in the *spoT1* mutant would support pppGpp accumulation during stringent response. We used the *gsk3* allele that codes for the feedback-resistant guanosine kinase and reported to increase the intracellular GTP pool in the presence of guanosine (Petersen, 1999). Normally there is a growth inhibition associated with GTP accumulation, but this can be overcome by histidine and tryptophan supplementation (Petersen, 1999). Stringent response in the *spoT1 gsk3* strain, in the absence of guanosine, was similar to that seen in the *spoT1* mutant. In the presence of guanosine, the only significant difference observed was the absence of GTP depletion (Fig. S6). Assuming this reflected an increase in the GTP pool, the result suggested an increase in the GTP level was insufficient to support pppGpp synthesis in the *spoT1* mutant.

Another possible reason for the absence of pppGpp accumulation could be, the pppGpp synthesized was rapidly turned over into ppGpp by GppA, a cytoplasmic pppGppase that is

not ribosome-associated (Somerville and Ahmed, 1979). To test this, the $\Delta gppA::Kan$ allele, sourced from the Keio collection (Baba *et al.*, 2006), was introduced by phage P1 transduction to construct the $\Delta spoT::Cm \Delta gppA::Kan/pRCspoT$ and $spoTl \Delta gppA::Kan/pRCspoT$ strains. During SpoT depletion, growth of the $\Delta spoT::Cm \Delta gppA::Kan/pRCspoT$ strain was more severely inhibited as compared to the $\Delta spoT::Cm/pRCspoT$ strain (Fig. 3A (i)). The plating efficiency dropped at least three orders of magnitude in the former strain as compared to the latter strain (Fig. 3A (ii)). When the stringent nucleotides pool was determined in the $\Delta spoT::Cm \Delta gppA::Kan/pRCspoT$ strain after the growth retardation that followed SpoT depletion (Fig. S7), accumulation of ppGpp, pppGpp, and depletion of GTP was observed (Fig. 3B). This showed the absence of pppGpp accumulation during SpoT depletion in the $\Delta spoT::Cm/pRCspoT$ strain was partly due to the GppA activity. In addition to pppGpp accumulation there was also a significant decrease in the GTP pool when compared to the SpoT depletion in $gppA^+$ strain (compare Fig. 3B, lane 2 with Fig. 1C, lane 3).

We asked if the lack of pppGpp accumulation during the stringent response in the hydrolase deficient $spoTl$ strain (Fig. S2) was also due to the GppA activity. Interestingly, synthetic growth defect was evident when the $\Delta gppA::Kan$ allele was introduced into the $spoTl$ genetic background. The $spoTl \Delta gppA::Kan/pRCspoT$ strain failed to segregate white colonies ($spoTl \Delta gppA::Kan$ genotype) in the minimal glucose medium with or without casamino acids and the white colonies were slow growing in LB medium (Fig. 4A, panels ii, vi and x). Deletion of the *relA* gene suppressed the synthetic growth defect (white colonies in panels iv, viii and xii). The slow-growing $spoTl \Delta gppA::Kan$ strain could be maintained on LB plate but failed to grow on minimal glucose medium with or without casamino acids (data not shown). Apparently, an elevated basal pppGpp pool and lowered *spoT* hydrolase activity together conferred synthetic growth inhibition in the presence of RelA activity, while

individually they affected the growth marginally at best (Fig. 4A, the white colonies in panels i, v and ix have the *spoTl* genotype, blue colonies in panels ii, vi and x have the $\Delta gppA::Kan$ genotype). Another hydrolase deficient *spoT* allele, *spoT202* (Sarubbi *et al.*, 1989), also showed synthetic growth defect with the $\Delta gppA::Kan$ allele and the growth defect was more severe than seen for the *spoTl* allele as no white colonies could be recovered on LB medium in addition to minimal medium (data not shown). This suggested, the synthetic growth defect with $\Delta gppA$ arose specifically from the SpoT hydrolase deficiency. As described below, the synthetic growth inhibition was associated with accumulation of (p)ppGpp and the reduced degradation of (p)ppGpp.

To find out if the *spoTl* $\Delta gppA::Kan$ growth defect was accompanied by a change in the stringent nucleotides pool, SpoT depletion was carried out by IPTG withdrawal in the *spoTl* $\Delta gppA::Kan$ /pRC*spoT* strain. As a control, depletion was carried out in the *spoTl*/pRC*spoT* strain (Fig. 4B). Before and after SpoT depletion, the *spoTl*/pRC*spoT* strain exhibited stringent response like the wild-type and *spoTl* strains respectively. That is, isoleucine starvation induced ppGpp and pppGpp synthesis following growth in the presence of IPTG (Fig. 4B, lanes 1–4) and mainly ppGpp, following growth in the absence of IPTG (Fig. 4B, Lanes 5–8) similar to the *spoTl* mutant (Fig. S2, lanes 1 – 4). This indicated, IPTG withdrawal sufficiently depleted the plasmid-encoded protein that the phenotype of the chromosomal allele was evident. As expected, IPTG withdrawal also induced growth inhibition in the *spoTl* $\Delta gppA::Kan$ /pRC*spoT* strain, and associated with the growth inhibition the (p)ppGpp pools were elevated (Fig. 4B, lane 13) and which increased further following amino acid starvation (Fig. 4B, lanes 14–16). The results indicated that elevated pppGpp pool activates RelA-dependent (p)ppGpp synthesis in the *spoTl* strain without amino acid starvation. During stringent response in the presence of IPTG, the *spoTl* $\Delta gppA$ /pRC*spoT* strain accumulated similar levels of ppGpp and pppGpp (Fig. 4B, lanes 10 –

12) as reported for a *gppA* mutant (Somerville and Ahmed, 1979). However, after IPTG withdrawal, that is, in the phenotypically *spoT1 ΔgppA* cells, pppGpp accumulated less than ppGpp (Fig. 4B, lanes 14-16), the plausible reasons for this are discussed later.

Since *relA* deletion suppressed the *spoT1 gppA* synthetic growth defect (Fig. 4A, panels iv, viii and xii), we tested the effect of the hypomorphic *relA* alleles. Plasmid-free derivatives of *relA496Δ::Kan spoT1 ΔgppA::Kan/pRCspoT* and *ΔrlmD::FRT spoT1 ΔgppA::Kan/ pRCspoT* strains (white colonies) were recovered on LB and as well as minimal glucose medium with or without casamino acids (data not shown). This suggested wild-type level expression of full-length RelA was needed for gratuitous (p)ppGpp synthesis and growth arrest in the *spoT1 ΔgppA::Kan* background. Since the *relA496Δ::Kan* and *ΔrlmD::FRT* alleles supported growth in the complete absence of *spoT* function (Fig. 2B, panels iv and vii) we asked if the growth of these strains were dependent on the GppA function. Plasmid free derivatives (white colonies) could not be recovered from the *relA496Δ::Kan ΔspoT ΔgppA::Kan/pRCspoT* and *ΔrlmD::FRT ΔspoT ΔgppA::Kan /pRCspoT* strains on LB (Fig. 4C) and in the minimal glucose medium as well (data not shown). SpoT depletion was carried out in these strains to ask if the growth arrest was accompanied by (p)ppGpp accumulation. Gratuitous synthesis of ppGpp and to a lesser extent, pppGpp, was observed in both strains (Fig. 4D). Notably, (p)ppGpp synthesis by the *relA496Δ::Kan* allele indicated that ribosome binding was not necessary for the pppGpp dependent activation. These results indicate, the residual hydrolase activity in the *spoT1* allele (as compared to the *ΔspoT* allele) prevented the activation of hypomorphic *relA* alleles by pppGpp to gratuitously synthesize (p)ppGpp. We tested, if the *relA1* allele, possibly the weakest of the hypomorphic allele studied here also exhibited synthetic growth defect in the *ΔspoT ΔgppA::Kan* background. Plasmid-free derivatives could be recovered from the *relA1 ΔspoT ΔgppA::Kan / pRCspoT* strain and no growth phenotype was evident in this strain as compared to the parental strain, *relA1 ΔspoT*, in LB or minimal

medium (Fig. S4B). This suggested, the *relA1* allele, unlike the *relA496Δ* or *ΔrlmD::FRT* alleles was not activated by pppGpp to carry out gratuitous (p)ppGpp synthesis. It is possible, in RelA1, the structural determinants required for allosteric activation by pppGpp may not be present since the (p)ppGpp synthase activity was reconstituted by two polypeptides in trans.

SpoT mediated degradation of (p)ppGpp is inhibited by pppGpp

To understand the *ΔgppA spoT1* synthetic growth defect, the contribution of GppA and SpoT, individually and together, to the turnover of the stringent nucleotides was studied. Isoleucine starvation was induced with valine to allow the accumulation of (p)ppGpp and the kinetics of degradation was monitored after the reversal of starvation by isoleucine addition. As reported, the GTP pool decreased and that of the stringent nucleotides increased following amino acid starvation in the wild-type strain (Fig. 5A). In the *ΔgppA* strain, following starvation, the GTP and pppGpp pools were, respectively, 2.5-fold lower and 3.8-fold higher than in the wild-type; the ppGpp pool was not significantly altered (Fig. 5A & B, table S2). Consistent with the biochemical activity of GppA, an increase in the pppGpp pool was expected, but the decrease in GTP pool was unexpected. We suspected this could be due to the reduced regeneration of GTP from the stringent nucleotides following the inhibition of SpoT hydrolase activity by the increase in pppGpp. As seen from Fig. 2C (iii), SpoT activity contributes to the degradation of ppGpp to GDP, during amino acid starvation; GTP can then be made from GDP. Indeed, in the *ΔgppA* strain, following the inhibition of RelA activity by the reversal of amino acid starvation, we observed the rate of degradation of the stringent nucleotides and the accumulation of GTP was slightly reduced as compared to the wild-type strain (Fig. 5A & B, table S2). It is conceivable, pppGpp may be poorly hydrolyzed relative to ppGpp, by SpoT, and therefore, an increase in the pppGpp pool inhibited GTP generation by competitively inhibiting the ppGpp degradation to GDP by SpoT (Fig. 7).

The effect of the increased pppGpp pool on (p)ppGpp degradation by the *spoT1* allele was studied. We confirmed the ppGpp degradation rate following amino acid starvation was reduced in the *spoT1* strain as reported (Fig. 5C). Quantification showed 54% degradation of ppGpp fifteen minutes after reversal of amino acid starvation (Fig. 5E, Table S3) while it was virtually undetectable after 1 minute in the *spoT*⁺ strain (Fig. 5B, Table S2). When the pppGpp pool was elevated in the *spoT1* $\Delta gppA::FRT/pRCspoT$ strain by SpoT depletion, the (p)ppGpp degradation was more severely impaired (Fig. 5D). There was only 22% degradation of ppGpp and 19% degradation of pppGpp after 15 minutes (Fig. 5E and Table S4). It is possible, the increase in (p)ppGpp pool from the turnover defect contributes to the allosteric activation of RelA and the growth arrest of *spoT1* $\Delta gppA$ strain.

Synergistic inhibition of growth by ppGpp and pppGpp

As described above, the RelA-dependent growth arrest in the *spoT1*, *relA496* $\Delta::FRT$ $\Delta spoT$ and $\Delta rlmD::FRT$ $\Delta spoT$ strains were contingent on the loss of GppA function. Based on the principal biochemical activity of GppA, which is the conversion of pppGpp to ppGpp, and the elevated pppGpp pool observed during the growth inhibition, it was deduced as the primary determinant driving RelA activation, although it was less abundant than ppGpp during the growth arrest (Fig. 3B, lane 2; Fig. 4B, lane 13; Fig. 4D, lanes 2 & 4; Fig. 5D, lane 1). We asked, if elevated ppGpp, in the absence of pppGpp, inhibited growth, and if so, whether it was different from that seen in the additional presence of pppGpp. We took advantage of the phenotype of the $\Delta rlmD::FRT$ $\Delta spoT$ strain, which accumulated ppGpp but no detectable pppGpp during amino acid starvation (Fig. 2C (iii), lanes 6 to 8) and the ppGpp degradation rate was greatly diminished after the reversal of amino acid starvation (Fig. 6A), due to the absence of SpoT function. The ppGpp pool size was virtually unaltered for up to 60 minutes and reduced 28% and 37% respectively 120 and 180 minutes after the reversal of starvation (Fig. 6B and table S5). These results show SpoT as the primary ppGpp hydrolase.

391 Growth arrest and recovery associated with amino acid starvation and its reversal were
392 monitored in the $\Delta rlmD::FRT \Delta spoT$ mutant and compared with that seen in the wild-type,
393 $spoT1$, and $\Delta rlmD::FRT$ strains. As expected, growth ceased and resumed immediately
394 following amino acid starvation and its reversal in the wild-type strain (Fig. S8A). In the
395 $spoT1$ strain, the starvation-induced growth arrest was similar to the wild-type, but growth
396 resumed after a lag of ~30 minutes following the reversal of starvation (Fig. S8B), which
397 correlated with the reduced rate of ppGpp degradation observed in the $spoT1$ mutant (Fig.
398 5C). Isoleucine starvation in the $\Delta rlmD::FRT \Delta spoT$ strain caused immediate growth arrest,
399 but after the reversal of starvation, growth inhibition persisted for 120 minutes and then the
400 growth resumed. After 120 minutes, the ppGpp pool was ~54% of GTP (Fig. 6B, 6C & Table
401 S5) and this can be taken as the lowest pool size at which ppGpp conferred growth inhibition.
402 One hour after growth resumed, that is, 180 after isoleucine addition, the ppGpp pool was
403 ~45% of GTP (Fig. 6C and Table S5). Isoleucine starvation and its reversal in the
404 $\Delta rlmD::FRT$ strain, which did not accumulate ppGpp (Fig. 2C(iii), lanes 2 to 4), produced
405 growth arrest and reversal similar to the wild-type strain (Fig. S8C), and indicated that the
406 accumulation of ppGpp was responsible for the transient growth arrest in the $\Delta rlmD::FRT$
407 $\Delta spoT$ strain. When growth inhibition from ppGpp and (p)ppGpp accumulation was
408 compared, it was more severe when both pppGpp and ppGpp are present. For instance, in the
409 $spoT1 \Delta gppA$ /pRCspoT strain, where the growth arrest was severe and prolonged following
410 SpoT depletion, ppGpp and pppGpp accumulated to 37% and 13% of GTP respectively after
411 SpoT depletion (Fig. 5D, lane 1, table S4). These results showed that the growth inhibition
412 conferred by ppGpp in isolation was transient, and this was strongly accentuated in the
413 presence of pppGpp. While we did not measure ppGpp levels beyond 180 minutes, the
414 continued growth of the $\Delta rlmD::FRT \Delta spoT$ strain suggested that there could progressive

decrease in the ppGpp pool and that eventually it may be degraded, although very inefficiently, by hydrolases other than SpoT.

Discussion

Autocatalytic activation of RelA by pppGpp

This study was initiated to test the prediction that the stringent nucleotides would accumulate in the cell following the depletion of SpoT activity. While an increase in the ppGpp pool was observed, accumulation of pppGpp was not seen and furthermore the pppGpp levels was not completely restored by inactivation of the pppGpp hydrolase, GppA. The results indicated that the degradation of pppGpp was enhanced during SpoT depletion and this was necessary to prevent RelA activation. RelA activation by pppGpp was clearly manifest in the *relA496Δ ΔspoT* and *ΔrlmD::FRT ΔspoT* strains, where, the slightly elevated ppGpp pool (Fig. 2C(i) lane 5 and 2C(iii) lane 5), increased several-fold after inactivation of GppA and pppGpp synthesis was evident (Fig. 4D lane 2 and 4). Since the (p)ppGpp pool increased in the absence of SpoT protein, and therefore totally independent of the hydrolase or synthase functions of SpoT, the increase in (p)ppGpp pool can be attributed entirely to the activation of RelA by the elevated basal pppGpp pool (from the loss of GppA activity). While the increase in the pppGpp pool may be important to initiate the allosteric activation of RelA, once initiated, the autocatalytic reaction may be sustained by both stringent nucleotides. ppGpp was reported to allosterically activate its own ribosome-dependent synthesis *in vitro* (Shyp *et al.*, 2012). However, given that a C-terminal His₆-tagged protein was used in that study and RelA's C-terminus resides well inside the ribosomal complex, it was likely that the functionality of the protein was compromised. In a recent report using untagged RelA protein, it was shown that pppGpp as an allosteric regulator and GDP as substrate synergize for the maximum enzymatic activity of RelA *in vitro* (Kudrin *et al.*, 2018). Allosteric

activation of Rel protein by pppGpp was not restricted to full-length RelA, it has also been noted for RelQ, an RNA binding small alarmone synthase from *Enterococcus faecalis* (Beljantseva *et al.*, 2017) and as well as the small alarmone synthase, SAS1, from *Bacillus subtilis* (Steinchen *et al.*, 2015). While RelA is catalytically active as a monomer, SAS1 functions as a homotetramer and tetramerization was found to be important for the allosteric activation by pppGpp. A pppGpp binding site at the C-terminal region of the Rel enzyme from *Mycobacterium smegmatis* has been reported (Syal *et al.*, 2015). The physiological relevance of these findings remain unclear.

Our results provide experimental evidence, the first to our knowledge, that at least two levels of positive control on RelA activation exist *in vivo* (Fig. 7). In addition to the well-studied ribosome-dependent activation of the RelA catalytic activity by the hungry codons, there is a positive regulation of the RelA activity by pppGpp and which is likely to be an allosteric activation. While the activation from amino acid starvation requires RelA interaction with the ribosome the latter was also evident in the absence of ribosome binding (Fig. 4D). Further studies are needed to understand the physiological significance of the pppGpp mediated allosteric activation of RelA. By its very nature, an autocatalytic reaction would result in the generation of a bistable switch (Dubnau and Losick, 2006). We propose fluctuations in the SpoT and GppA hydrolase activities could contribute to the non-uniform (p)ppGpp synthesis in the cells within a population. It has been noted that the formation of persister cells within a population is associated with these cells having very high levels of (p)ppGpp (Harms *et al.*, 2016).

SpoT and GppA hydrolases – the negative regulators of the stringent response

Genetic evidence was used to postulate that the essential function of SpoT was the degradation of (p)ppGpp synthesized by RelA (Xiao *et al.*, 1991) in response to signal(s) the

nature of which is not well understood. To our knowledge, this was not experimentally tested. Our results are in accord with this postulation but have revealed additional layers of regulation. While (p)ppGpp accumulation was anticipated from the depletion of SpoT, only ppGpp accumulation was observed, a phenotype reminiscent of that seen following amino acid starvation in the *spoT1* mutant. Although the molecular basis for this response is not clear, our results have eliminated the possibility that the pppGpp depletion arose solely from the activation of GppA, because, if that were to be the case, the pppGpp pool size would be completely restored in the $\Delta gppA$ background. However, the pppGpp pool observed in the $\Delta gppA$ *spoT1* and $\Delta gppA$ $\Delta spoT$ strains were less than that observed in the $\Delta gppA$ mutant (Fig. 3B lane 2, Fig. 4B lanes 10 – 16). Either the decreased synthesis of pppGpp or an increased degradation of pppGpp by one or more pppGppase, other than GppA, that was reported (Somerville and Ahmed, 1979) and references therein) may contribute to the depletion of pppGpp pool in the *spoT* hydrolase mutant. We favor the latter possibility, because, increasing the GTP pool did not increase the pppGpp synthesis (Fig. S6). Although the molecular mechanism is unclear, decrease in the pppGpp pool was an important adaptation needed to support the growth of strains deficient in *spoT* hydrolase activity.

Negative regulation of the stringent response by SpoT, consequent to amino acid starvation was evident in the $\Delta rlmD::FRT$ background (a hypomorphic allele with reduced expression of full-length RelA protein). Apparently, when there are fewer RelA molecules, signaling through the hungry codons was insufficient to amplify the (p)ppGpp pool faster than their turn over by the SpoT hydrolase activity (Fig. 2C (iii)). Further studies are needed to establish and understand the physiological relevance of such a regulation in the *relA*⁺ background. These results do not support the observations made *in vitro* that the uncharged tRNA's inhibit the SpoT hydrolase activity (Richter, 1980), perhaps because the interaction between SpoT and uncharged tRNA possible *in vitro* may not occur in the cellular milieu. We

propose, this kind of regulation could ensure that the amplification of the stringent response happens after a quorum of hungry codons is available to be sensed by RelA and tight negative regulation is maintained below the quorum through the SpoT and GppA hydrolase activities that lower the basal (p)ppGpp pool and prevent the activation of RelA by the nucleotides. Since, (p)ppGpp accumulation signals stress, such a regulation would ensure stress related adaptations are not provoked in response to minor fluctuations in the amino acid pool but only when there was sufficient starvation to allow a quorum of hungry codons to accumulate. When the rate of RelA-mediated (p)ppGpp synthesis exceeded the rate of SpoT/GppA mediated hydrolysis, the (p)ppGpp concentration can increase rapidly aided by the allosteric activation of RelA by pppGpp.

Effect of (p)ppGpp versus ppGpp on growth

(p)ppGpp synthesis leads to growth inhibition. It has been reported that, in the absence of stress, progressively increasing the basal ppGpp level using hydrolase deficient *spoT* mutants progressively increased the growth inhibition (Fiil *et al.*, 1977; Sarubbi *et al.*, 1988; Xiao *et al.*, 1991). As noted previously (Laffler and Gallant, 1974; Fiil *et al.*, 1977) and as well as in this study, reduction in the *spoT* hydrolase activity decreased the pppGpp pool and therefore, the slow growth of the *spoT* hydrolase mutants are primarily due to elevated ppGpp pool and despite the reduced pppGpp pool. Since the elimination of the GppA activity increased the pppGpp pool in the SpoT hydrolase deficient strains (Fig 3B, 4B, and 4D), we compared the relative growth inhibition conferred by ppGpp with that of (p)ppGpp. While (p)ppGpp accumulation conferred severe and prolonged growth inhibition (Figs. 4A, 5D lane 1, Table S4) at comparable concentration, growth inhibition by ppGpp was transient (Figs. 6A lanes 9 - 11, 6B, 6C and Table S5). We propose, this could be because the two nucleotides regulate cellular functions with different efficiencies and which then synergistically contribute to growth inhibition. For example, it was reported that the inhibition of replication elongation

was more pronounced when the basal pppGpp pool was increased (Denapoli *et al.*, 2013) while RNAP activity was more strongly inhibited by ppGpp (Mechold *et al.*, 2013). Our results also show that an important function of GppA is to lower the pppGpp pool and alleviate the gratuitous activation of RelA especially under conditions that lower the SpoT hydrolase activity. In other words, the SpoT and GppA hydrolases together maintain a low pppGpp pool so as to prevent the accumulation of (p)ppGpp through the autocatalytic amplification of basal RelA activity. The molecular mechanism of allosteric activation of RelA by pppGpp is being investigated.

Experimental Procedures.

Growth conditions

LB medium (1% tryptone, 0.5% yeast extract, 1% NaCl), MOPS buffered minimal medium (Neidhardt *et al.*, 1974) or minimal A medium (Miller, 1992) were used, the latter was supplemented with 0.5% glucose and 20 amino acids, each at a final concentration of 40 $\mu\text{g ml}^{-1}$. In plates, glucose and cas-amino acids were supplemented at 0.2% final concentration. Antibiotics and the concentrations at which they were used are, ampicillin (Amp), 50 $\mu\text{g/ml}$, kanamycin (Kan), 25 $\mu\text{g ml}^{-1}$; tetracycline (Tet), 10 $\mu\text{g ml}^{-1}$ and chloramphenicol (Cm), 15 $\mu\text{g ml}^{-1}$. Isopropyl β -D-thiogalactopyranoside (IPTG) and 5-Bromo-4-chloro 3-indolyl- β -D-thiogalactoside (X-gal) were used at a final concentration of 1 mM and 50 $\mu\text{g ml}^{-1}$, respectively.

Construction of strains and plasmids

The strains used in this study are derivatives of the *E. coli* K12 strain MG1655, referred as wild-type. Strains and primers are listed in Table S1. Mutations were introduced by P1vir mediated transduction using standard protocols. Gene deletions have been sourced from the Keio collection (Baba *et al.*, 2006) and when required, the kanamycin resistance cassette was

flipped out using FLP recombinase expressed from a pCP20 plasmid (Cherepanov and Wackernagel, 1995). The plasmid pRC*spoT* is from the lab collection (Nazir and Harinarayanan, 2016) and was constructed from pRC7 (Bernhardt and De Boer, 2004). Deletions in the *relA* gene on the chromosome, to generate *relA455Δ::Kan* and *relA496Δ::Kan*, wherein all codons after 455 and 496 were deleted and replaced with a TAG stop codon and the kanamycin cassette from pKD13 (Datsenko and Wanner, 2000) was performed by recombineering (Thomason *et al.*, 2014). Forward primers JGOrelA496aaPS4, JGOrelA455aaPS4, and the reverse primer JGOrelAPS1 were used. These constructs generated were verified by sequencing. The transposon insertions *relA::Tn10dTet* and *rlmD::Tn10dKan* were mapped by inverse PCR (Higashitani *et al.*, 1994). The β-galactosidase reporter fusions *relA'-lac* and *rlmD'-lac* within the *relA* and *rlmD* genes respectively, were constructed using the knockout alleles available in the Keio collection and plasmid pKG137 using a published protocol (Ellermeier *et al.*, 2002). The new junctions generated were verified by sequencing.

Plasmid loss and viability measurements

Using a published assay (Bernhardt and De Boer, 2004) the ability of strains to grow following loss of the single-copy, unstable, *amp^r*-plasmid pRC*spoT* was assessed in the *Δlac* genetic background. The plasmid carries the *lacZ* gene. Strains containing the plasmid pRC*spoT* were grown overnight in LB broth containing ampicillin and IPTG, the latter used at a final concentration of 1 mM unless mentioned otherwise. Overnight cultures were washed with minimal A medium and serially diluted to 10⁻⁶ dilution. Dilutions were spread on IPTG X-gal containing plates to obtain ~300 colonies and incubated for 24 to 72 hours at 37°C depending on the growth medium. Blue and white colonies indicating cells that retained and lost the plasmid respectively were counted and expressed as the percentage of white colonies. White colonies were tested for growth on the same medium to confirm viability.

The efficiency of plating determination

Strains were grown overnight in the permissive medium. Overnight cultures were washed with minimal A and serially diluted to 10^{-6} dilution. From each dilution, 10 μ l was spotted, allowed to dry and plates incubated at 37°C. To test for RelA function or elevated (p)ppGpp level, strains were spotted on minimal A medium containing glucose with or without the amino acids serine (S), methionine (M) and glycine (G) (100 μ g/ml each). The growth of the *relA* mutant was inhibited in the presence of SMG (Uzan and Danchin, 1976).

Depletion of SpoT

In strains carrying the single-copy plasmid pRC*spoT*, the chromosomal *spoT* gene was replaced with the $\Delta spoT::Cm$ or the $\Delta spoT$ allele. SpoT expression in pRC*spoT* was induced using IPTG as it was expressed from the *lac* promoter and the plasmid also carried the *lacI* gene. Cultures grown overnight in the presence of ampicillin and IPTG (1mM) in LB or MOPS buffered medium containing glucose and 20 amino acids, were washed and sub-cultured at different dilutions in appropriate liquid medium or plates containing ampicillin and different concentrations of IPTG or without IPTG.

(p)ppGpp estimation by thin layer chromatography

(p)ppGpp estimations were carried out by growing cultures in MOPS buffered medium containing 0.5% glucose and when necessary the 20 amino acids were added, each to a final concentration of 40 μ g/ml. Overnight cultures and the initial growth following dilution were carried out in the presence of 1.32 mM K_2HPO_4 . At an A_{600} of ~ 0.4 to 0.5, the cultures were diluted 10-fold into the low phosphate medium (0.4 mM K_2HPO_4) and allowed to undergo at least two doublings in the presence of 100-200 μ Ci/ml of ^{32}P - H_3PO_4 before sample collection began at $\sim 0.2 A_{600}$. An unlabelled culture was used to monitor A_{600} at periodic time intervals. Valine was added at 100 μ g/ml to induce isoleucine starvation and which was reversed by

adding isoleucine at 100 µg/ml. Samples were collected in tubes containing an equal volume of 2N HCOOH kept chilled on ice. The samples were subjected to three cycles of freeze-thaw and centrifuged at 10000 rpm for 5 minutes at 4°C and 10 µl of the supernatant was applied on PEI cellulose sheets and resolved in 1.5M KH₂PO₄, pH3.4. The nucleotide spots were visualized by phosphorimager (Typhoon FLA 9500) and quantified using multi-gauge software (Fujifilm). The values are expressed as the ratio of a nucleotide (GTP or ppGpp or pppGpp) to total (GTP + ppGpp + pppGpp).

Growth kinetics of amino acid starvation and recovery

Overnight cultures were grown in minimal A glucose medium and sub-cultured in the same medium in three separate flasks. The cultures were incubated at 37 °C, 200 rpm and growth was monitored periodically. At OD₆₀₀ between 0.15 – 0.25, valine was added to two of the cultures and 1 hour later isoleucine was added to one of the cultures having valine. Growth kinetics in the three flasks was compared.

Acknowledgment

We acknowledge strains used from the Keio collection (NBRP, Japan). We thank members of the Laboratory of Bacterial Genetics for their suggestions. We thank Mike Cashel for providing strains used in this study and suggestions. This work was primarily funded by the Centre of Excellence in Microbial Biology – Phase 2 research grant of the Department of Biotechnology, Government of India. R.S is a recipient of DBT-JRF fellowship.

Conflict of interest

The authors declare they have no conflict of interest.

References

- An, G., Justesen, J., Watson, R.J., and Friesen, J.D. (1979) Cloning the spoT Gene of Escherichia coli : Identification of the spoT Gene Product. *J Bacteriol* **137**: 1100–1110
<http://jb.asm.org/content/137/3/1100.full.pdf>.
- Arenz, S., Abdelshahid, M., Sohmen, D., Payoe, R., Starosta, A.L., Berninghausen, O., *et al.* (2016) The stringent factor RelA adopts an open conformation on the ribosome to stimulate ppGpp synthesis. *Nucleic Acids Res* **44**: 6471–6481.
- Atkinson, G.C., Tenson, T., and Hauryliuk, V. (2011) The RelA/SpoT homolog (RSH) superfamily: distribution and functional evolution of ppGpp synthetases and hydrolases across the tree of life. *PLoS One* **6**: e23479
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3153485&tool=pmcentrez&rendertype=abstract>.
- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., *et al.* (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**: 2006.0008
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1681482&tool=pmcentrez&rendertype=abstract>.
- Barker, M.M., Gaal, T., Josaitis, C.A., and Gourse, R.L. (2001) Mechanism of regulation of transcription initiation by ppGpp. I. Effects of ppGpp on transcription initiation in vivo and in vitro. *J Mol Biol* **305**: 673–688
<http://www.ncbi.nlm.nih.gov/pubmed/11162084>
<http://www.ncbi.nlm.nih.gov/pubmed/11162085>.
- Battesti, A., and Bouveret, E. (2006) Acyl carrier protein/SpoT interaction, the switch linking

- 631 SpoT-dependent stress response to fatty acid metabolism. *Mol Microbiol* **62**: 1048–1063.
- 632 Beljantseva, J., Kudrin, P., Andresen, L., Shingler, V., Atkinson, G.C., Tenson, T., and
- 633 Hauryliuk, V. (2017) Negative allosteric regulation of *Enterococcus faecalis* small alarmone
- 634 synthetase RelQ by single-stranded RNA. *Proc Natl Acad Sci* **114**: 3726–3731
- 635 <http://www.pnas.org/lookup/doi/10.1073/pnas.1617868114>.
- 636 Bernhardt, T.G., and Boer, P.A.J. De (2004) Screening for synthetic lethal mutants in
- 637 *Escherichia coli* and identification of EnvC (YibP) as a periplasmic septal ring factor with
- 638 murein hydrolase activity. *Mol Microbiol* **52**: 1255–1269.
- 639 Braeken, K., Moris, M., Daniels, R., Vanderleyden, J., and Michiels, J. (2006) New horizons
- 640 for (p)ppGpp in bacterial and plant physiology. *Trends Microbiol* **14**: 45–54.
- 641 Brown, A., Fernández, I.S., Gordiyenko, Y., and Ramakrishnan, V. (2016) Ribosome-
- 642 dependent activation of stringent control. *Nature* **534**: 277–80
- 643 <http://dx.doi.org/10.1038/nature17675>.
- 644 Brown, D.R., Barton, G., Pan, Z., Buck, M., and Wigneshweraraj, S. (2014) Nitrogen stress
- 645 response and stringent response are coupled in *Escherichia coli*. *Nat Commun* **5**: 1–8
- 646 <http://dx.doi.org/10.1038/ncomms5115>.
- 647 Chatterji, D., and Kumar Ojha, A. (2001) Revisiting the stringent response, ppGpp and
- 648 starvation signaling. *Curr Opin Microbiol* **4**: 160–165.
- 649 Cherepanov, P.P., and Wackernagel, W. (1995) Gene disruption in *Escherichia coli*: TcR and
- 650 KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance
- 651 determinant. *Gene* **158**: 9–14.
- 652 Datsenko, K., and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in

653 *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci USA* **97**: 6640–6645

654 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=18686&tool=pmcentrez&rendert](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=18686&tool=pmcentrez&rendertype=abstract)

655 [ype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=18686&tool=pmcentrez&rendertype=abstract).

656 Denapoli, J., Tehranchi, A.K., and Wang, J.D. (2013) Dose-dependent reduction of

657 replication elongation rate by (p) ppGpp in *Escherichia coli* and *Bacillus subtilis*. *Mol*

658 *Microbiol* **88**: 93–104.

659 Dubnau, D., and Losick, R. (2006) Bistability in bacteria. *Mol Microbiol* **61**: 564–572.

660 Durfee, T., Hansen, A.M., Zhi, H., Blattner, F.R., and Ding, J.J. (2008) Transcription

661 profiling of the stringent response in *Escherichia coli*. *J Bacteriol* **190**: 1084–1096.

662 Ellermeier, C.D., Janakiraman, A., and Slauch, J.M. (2002) Construction of targeted single

663 copy lac fusions using λ Red and FLP-mediated site-specific recombination in bacteria. *Gene*

664 **290**: 153–161.

665 Ferullo, D.J., and Lovett, S.T. (2008) The stringent response and cell cycle arrest in

666 *Escherichia coli*. *PLoS Genet* **4**: 21–23.

667 Fiil, N.P., Willumsen, B.M., Friesen, J.D., and Meyenburg, K. von (1977) Interaction of

668 alleles of the *relA*, *relC* and *spoT* genes in *Escherichia coli*: Analysis of the interconversion

669 of GTP, ppGpp and pppGpp. *MGG Mol Gen Genet* **150**: 87–101.

670 Hamel, E., and Cashel, M. (1974) Guanine nucleotides in protein synthesis. Utilization of

671 pppGpp and dGTP by initiation factor 2 and elongation factor Tu. *Arch Biochem Biophys*

672 **162**: 293–300.

673 Harat, A., and Sy, J. (1983) Guanosine 5'-Triphosphate, 3'-Diphosphate 5'-

674 Phosphohydrolase: Purification and substrate specificity. *J Biol Chem* **258**: 1678–1683.

675 Harms, A., Maisonneuve, E., and Gerdes, K. (2016) Mechanisms of bacterial persistence
676 during stress and antibiotic exposure. *Science* (80-) **354**: aaf42681-9
677 <http://www.sciencemag.org/cgi/doi/10.1126/science.aaf4268>.

678 Hauryliuk, V., Atkinson, G.C., Murakami, K.S., Tenson, T., and Gerdes, K. (2015) Recent
679 functional insights into the role of (p)ppGpp in bacterial physiology. *Nat Rev Microbiol* **13**:
680 298–309 <http://dx.doi.org/10.1038/nrmicro3448>. Accessed April 14, 2015.

681 Heinemeyer, E., and Richter, D. (1978) Characterization of the Guanosine 5'-Triphosphate
682 3'-Diphosphate and Guanosine 5'-Diphosphate 3'-Diphosphate Degradation Reaction
683 Catalyzed by a Specific Pyrophosphorylase from Escherichia coli. *Biochemistry* **17**: 5368–
684 5372.

685 Higashitani, A., Higashitani, N., Yasuda, S., and Horiuchi, K. (1994) A general and fast
686 method for mapping mutations on the Escherichia coli chromosome. *Nucleic Acids Res* **22**:
687 2426–2427.

688 Jiang, M., Sullivan, S.M., Wout, P.K., and Maddock, J.R. (2007) G-protein control of the
689 ribosome-associated stress response protein SpoT. *J Bacteriol* **189**: 6140–6147.

690 Joseleau-Petit, D., Vinella, D., and D'Ari, R. (1999) Metabolic alarms and cell division in
691 Escherichia coli. *J Bacteriol* **181**: 9–14.

692 Kanjee, U., Ogata, K., and Houry, W.A. (2012) Direct binding targets of the stringent
693 response alarmone (p)ppGpp. *Mol Microbiol* **85**: 1029–1043.

694 Kari, C., Török, I., and Travers, A. (1977) ppGpp cycle in Escherichia coli. *MGG Mol Gen*
695 *Genet* **150**: 249–255.

696 Keasling, J.D., Bertsch, L., and Kornberg, A. (1993) Guanosine pentaphosphate

697 phosphohydrolase of *Escherichia coli* is a long-chain exopolyphosphatase. *Proc Natl Acad*
698 *Sci U S A* **90**: 7029–33.
699 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC47069/pdf/pnas01472-0144.pdf>.

700 Kudrin, P., Dzhygyr, I., Ishiguro, K., Beljantseva, J., Maksimova, E., Oliveira, S.R.A., *et al.*
701 (2018) The ribosomal A-site finger is crucial for binding and activation of the stringent factor
702 RelA. *Nucleic Acids Res* **46**: 1973–1983 [https://academic.oup.com/nar/advance-](https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gky023/4829690)
703 [article/doi/10.1093/nar/gky023/4829690](https://academic.oup.com/nar/advance-article/doi/10.1093/nar/gky023/4829690).

704 Laffler, T., and Gallant, J. (1974) SPOT , a New Genetic Locus Involved in the Stringent
705 Response in *E. coli*. *Cell* **1**: 27–30.

706 Leavitt, R., and Umbarger, H.E. (1961) Isoleucine and Valine Metabolism in *Escherichia*
707 *Coli*. *J Bacteriol* **83**: 624–630.

708 Loveland, A.B., Bah, E., Madireddy, R., Zhang, Y., Brilot, A.F., Grigorieff, N., and
709 Korostelev, A.A. (2016) Ribosome.RelA structures reveal the mechanism of stringent
710 response activation. *Elife* **5**: e17029.

711 Magnusson, L.U., Farewell, A., and Nyström, T. (2005) ppGpp: A global regulator in
712 *Escherichia coli*. *Trends Microbiol* **13**: 236–242.

713 Mechold, U., Potrykus, K., Murphy, H., Murakami, K.S., and Cashel, M. (2013) Differential
714 regulation by ppGpp versus pppGpp in *Escherichia coli*. *Nucleic Acids Res* **41**: 6175–6189.

715 Metzger, S., Dror, I.B., Aizenman, E., Schreiber, G., Toone, M., Friesen, J.D., *et al.* (1988)
716 The nucleotide sequence and characterization of the relA gene of *Escherichia coli*. *J Biol*
717 *Chem* **263**: 15699–704 <http://www.ncbi.nlm.nih.gov/pubmed/2844820>.

718 Metzger, S., Schreiber, G., Aizenman, E., Cashel, M., and Glaser, G. (1989) Characterization

719 of the relA1 mutation and a comparison of relA1 with new relA null alleles in Escherichia
720 coli. *J Biol Chem* **264**: 21146–21152.

721 Miller, D.L., Cashel, M., and Weissbach, H. (1973) The interaction of guanosine 5'-
722 diphosphate, 2' (3')-diphosphate with the bacterial elongation factor Tu. *Arch Biochem*
723 *Biophys* **154**: 675–682.

724 Miller, J.H. (1992) *A short course in bacterial genetics: A laboratory manual and handbook*
725 *for and related bacteria*. Cold Spring Harbor Laboratory Press, .

726 Milon, P., Tischenko, E., Tomsic, J., Caserta, E., Folkers, G., Teana, A. La, *et al.* (2006) The
727 nucleotide-binding site of bacterial translation initiation factor 2 (IF2) as a metabolic sensor.
728 *Proc Natl Acad Sci U S A* **103**: 13962–7 <http://www.pnas.org/content/103/38/13962.long>.

729 Mittenhuber, G. (2001) Comparative genomics and evolution of genes encoding bacterial
730 (p)ppGpp synthetases/hydrolases (the Rel, RelA and SpoT proteins). *J Mol Microbiol*
731 *Biotechnol* **3**: 585–600
732 [http://www.ncbi.nlm.nih.gov/sites/entrez?db=PubMed&cmd=retrieve&dopt=AbstractPlus&li](http://www.ncbi.nlm.nih.gov/sites/entrez?db=PubMed&cmd=retrieve&dopt=AbstractPlus&list_uids=11545276%5Cnpapers2://publication/uuid/2F1746BF-5E50-456B-9BC5-A709DE6050A7)
733 [st_uids=11545276%5Cnpapers2://publication/uuid/2F1746BF-5E50-456B-9BC5-](http://www.ncbi.nlm.nih.gov/sites/entrez?db=PubMed&cmd=retrieve&dopt=AbstractPlus&list_uids=11545276%5Cnpapers2://publication/uuid/2F1746BF-5E50-456B-9BC5-A709DE6050A7)
734 [A709DE6050A7](http://www.ncbi.nlm.nih.gov/sites/entrez?db=PubMed&cmd=retrieve&dopt=AbstractPlus&list_uids=11545276%5Cnpapers2://publication/uuid/2F1746BF-5E50-456B-9BC5-A709DE6050A7).

735 Nakagawa, A., Oshima, T., and Mori, H. (2006) Identification and characterization of a
736 second, inducible promoter of relA in Escherichia coli. *Genes Genet Syst* **81**: 299–310.

737 Nazir, A., and Harinarayanan, R. (2016) Inactivation of cell division protein FtsZ by SulA
738 makes Lon indispensable for the viability of a ppGpp0 strain of Escherichia coli. *J Bacteriol*
739 **198**: 688–700.

740 Neidhardt, F.C., Bloch, P.L., and Smith, D.F. (1974) Culture medium for enterobacteria. *J*
741 *Bacteriol* **119**: 736–747.

765 Ross, W., Sanchez-Vazquez, P., Chen, A.Y., Lee, J.H., Burgos, H.L., and Gourse, R.L.
766 (2016) ppGpp Binding to a Site at the RNAP-DksA Interface Accounts for Its Dramatic
767 Effects on Transcription Initiation during the Stringent Response. *Mol Cell* **62**: 811–823
768 <http://dx.doi.org/10.1016/j.molcel.2016.04.029>.

769 Ross, W., Vrentas, C.E., Sanchez-Vazquez, P., Gaal, T., and Gourse, R.L. (2013) The magic
770 spot: A ppGpp binding site on E. coli RNA polymerase responsible for regulation of
771 transcription initiation. *Mol Cell* **50**: 420–429 <http://dx.doi.org/10.1016/j.molcel.2013.03.021>.

772 Sajish, M., Kalayil, S., Verma, S.K., Nandicoori, V.K., and Prakash, B. (2009) The
773 significance of EXDD and RXKD motif conservation in Rel proteins. *J Biol Chem* **284**:
774 9115–9123.

775 Sarubbi, E., Rudd, E., Xiao, H., Ikeharat, K., Kalman, M., and Cashell, M. (1989)
776 Characterization of the spoT gene of Escherichia coli. *J Biol Chem* **264**: 15074–15082.

777 Sarubbi, E., Rudd, K.E., and Cashel, M. (1988) Basal ppGpp level adjustment shown by new
778 spoT mutants affect steady state growth rates and rrnA ribosomal promoter regulation in
779 Escherichia coli. *MGG Mol Gen Genet* **213**: 214–222.

780 Schreiber, G., Metzger, S., Aizenman, E., Roza, S., Cashel, M., and Glaser, G. (1991)
781 Overexpression of the relA gene in Escherichia coli. *J Biol Chem* **266**: 3760–3767.

782 Schreiber, G., Ron, E.Z., and Glaser, G. (1995) ppGpp-Mediated Regulation of DNA
783 Replication and Cell Division in. *Curr Microbiol* **30**: 27–32.

784 Seyfzadeh, M., Keener, J., and Nomura, M. (1993) spoT-dependent accumulation of
785 guanosine tetraphosphate in response to fatty acid starvation in *Escherichia coli*. *Proc Natl*
786 *Acad Sci U S A* **90**: 11004–11008.

787 Shyp, V., Tankov, S., Ermakov, A., Kudrin, P., English, B.P., Ehrenberg, M., *et al.* (2012)
788 Positive allosteric feedback regulation of the stringent response enzyme RelA by its product.
789 *EMBO Rep* **13**: 835–839 <http://embor.embopress.org/cgi/doi/10.1038/embor.2012.106>.

790 Somerville, C.R., and Ahmed, A. (1979) Mutants of *Escherichia coli* Defective in the
791 Degradation of Guanosine 5'-Triphosphate, 3'-Diphosphate (pppGpp). *MGG* **169**: 315–
792 323.

793 Steinchen, W., Schuhmacher, J.S., Altegoer, F., Fage, C.D., Srinivasan, V., Linne, U., *et al.*
794 (2015) Catalytic mechanism and allosteric regulation of an oligomeric (p)ppGpp synthetase
795 by an alarmone. *Proc Natl Acad Sci U S A* **112**: 13348–53
796 <http://www.ncbi.nlm.nih.gov/pubmed/26460002> [http://www.pubmedcentral.nih.gov/arti](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4629338)
797 [clerender.fcgi?artid=PMC4629338](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4629338).

798 Svtils, A.L., Cashel, M., and Zyskind, J.W. (1993) Guanosine tetraphosphate inhibits
799 protein synthesis in vivo. *J Biol Chem* **268**: 2307–2311.

800 Syal, K., Joshi, H., Chatterji, D., and Jain, V. (2015) Novel pppGpp binding site at the C-
801 terminal region of the Rel enzyme from *Mycobacterium smegmatis*. *FEBS J* **282**: 3773–3785.

802 Thomason, L.C., Sawitzke, J.A., Li, X., Costantino, N., and Court, D.L. (2014)
803 Recombineering: Genetic engineering in bacteria using homologous recombination. *Curr*
804 *Protoc Mol Biol* 10.1002/0471142727.mb0116s106.

805 Traxler, M.F., Zacharia, V.M., Marquardt, S., Summers, S.M., Nguyen, T., Stark, S.E., and
806 Conway, T. (2012) Discretely calibrated regulatory loops controlled by ppGpp partition gene
807 induction across the 'feast to famine' gradient in *Escherichia coli*. *Mol Microbiol* **79**: 830–
808 845.

809 Uzan, M., and Danchin, A. (1976) A rapid test for the RelA mutation in *E. coli*. *Biochem*

810 *Biophys Res Commun* **69**: 751–758.

811 Vinella, D., Albrecht, C., Cashel, M., and D'Ari, R. (2005) Iron limitation induces SpoT-
812 dependent accumulation of ppGpp in *Escherichia coli*. *Mol Microbiol* **56**: 958–970.

813 Wang, J.D., Sanders, G.M., and Grossman, A.D. (2007) Nutritional Control of Elongation of
814 DNA Replication by (p)ppGpp. *Cell* **128**: 865–875.

815 Wendrich, T.M., Blaha, G., Wilson, D.N., Marahiel, M.A., and Nierhaus, K.H. (2002)
816 Dissection of the mechanism for the stringent factor RelA. *Mol Cell* **10**: 779–788.

817 Xiao, H., Kalman, M., Ikehara, K., Zemel, S., Glaser, G., and Cashel, M. (1991) Residual
818 guanosine 3',5'-bispyrophosphate synthetic activity of relA null mutants can be eliminated by
819 spoT null mutations. *J Biol Chem* **266**: 5980–5990.

820 Zuo, Y., Wang, Y., and Steitz, T.A. (2013) The Mechanism of *E. coli* RNA Polymerase
821 Regulation by ppGpp is suggested by the structure of their complex. *Mol Cell* **50**: 430–436
822 <http://dx.doi.org/10.1016/j.molcel.2013.03.020>.

823 **Figure Legends**

824 **Figure 1 - Depletion of SpoT is associated with growth retardation and accumulation of**
825 **ppGpp, but not pppGpp.**

826 **A.** $\Delta spoT/pRCspoT$ (RS260) strain cultured in MOPS buffered medium with glucose, 20
827 amino acids, ampicillin and 1mM IPTG was washed and sub-cultured in the same medium
828 with (●, ■) or without (▽, ◇) IPTG at the dilutions indicated and the growth was
829 monitored. **B.** $\Delta spoT::Cm/pRCspoT$ (RS14) was sub-cultured in the same medium used for
830 Fig. 1A at the indicated dilutions. Growth after the subculture into the medium containing the
831 $^{32}\text{P-H}_3\text{PO}_4$ is shown, the solid arrows indicate the time points at which the samples were

collected for (p)ppGpp estimation. **C.** Samples collected as shown in Fig. 1B were subjected to PEI-TLC. Valine-induced stringent response in the wild type is provided as a reference (lane 1). Ratios of a nucleotide to the total (GTP+ppGpp+pppGpp) was calculated for each lane. **D.** Strain $\Delta spoT/pRCspoT$ (RS260) was cultured after 10^3 -fold dilution in MOPS glucose medium containing ampicillin and with (+) or without (–) IPTG, the stringent response was induced with valine (arrow), samples were collected immediately before valine addition or subsequently at the time indicated.

Figure 2 - Hypomorphic *relA* alleles suppress the $\Delta spoT$ growth defect.

A. Cartoon of the transposon insertions *rlmD*::Tn10dKan and *relA*496::Tn10dTet. Bent arrows represent known or annotated promoters, arrow within the transposon indicate the direction of transcription of the antibiotic marker. **B.** Segregation of the unstable plasmid *pRCspoT* was assayed as described in the methods. Relevant genotype of the strain, the percentage of white colonies and the total number of colonies (blue + white) used to calculate the ratio are indicated. **C.** Strains whose relevant genotypes are indicated were cultured in MOPS glucose medium, isoleucine starvation was induced with valine (arrow) and samples were collected immediately before valine addition and subsequently at the times indicated above the lanes and subjected to PEI-TLC. The line between lanes in the TLC indicates that internal lanes from a single TLC have been removed.

Figure 3 - SpoT depletion in the *gppA* mutant was associated with severe growth inhibition, depletion of GTP and accumulation of ppGpp and pppGpp

A. (i). $\Delta spoT/pRCspoT$ (RS260) and $\Delta spoT \Delta gppA::Kan/pRCspoT$ (RS478) strains cultured in the MOPS buffered medium containing glucose, 20 amino acids, ampicillin and 1mM IPTG were washed and sub-cultured in the same medium with (○,△) or without (●,▼) IPTG at the dilutions indicated and the growth monitored **A. (ii)** $\Delta spoT::Cm/pRCspoT$ (RS14)

and $\Delta spoT::Cm \Delta gppA::Kan/pRCspoT$ (RS206) strains cultured in LB containing ampicillin and 1mM IPTG were washed, serially diluted in LB medium and the dilutions spotted on LB agar plates containing Amp and 100 μ M, 10 μ M or no IPTG and incubated for 16 hr at 37°C.

B. $\Delta spoT::Cm \Delta gppA::Kan/pRCspoT$ (RS206) cultured in the medium described in 3A (i) was diluted 200-fold in the same medium or without IPTG and sampled after at least two divisions in the presence of P^{32} (arrows in Fig. S7) and subjected to TLC. The samples were part of a single TLC with the lanes in between deleted.

Figure 4 – Lowering the SpoT hydrolase activity in the absence of GppA function conferred growth arrest from gratuitous RelA activation.

A. The loss or retention of the unstable plasmid $pRCspoT$ was assayed as described in the methods. Relevant genotype of the strain, the percentage of white colonies and the total number of colonies (blue+white) used to calculate the ratio are indicated. The panels within a column represent single genotype. LB and minimal glucose cas-amino acids plates were incubated at 37°C for 24 hours and minimal glucose plates for 48 hours. **B.** Stringent nucleotides pool was measured by TLC following growth in the presence of P^{32} . Overnight cultures of $spoTI/pRCspoT$ (RS40) (Lanes 1 – 8) and $spoTI \Delta gppA/ pRCspoT$ (RS194) (Lanes 9 – 16) were grown in MOPS medium containing glucose and ampicillin in the presence or absence of IPTG as indicated after 1 in 1000 dilution. Amino acid starvation was induced with valine (arrow) and samples collected just before or subsequently at the indicated time. **C.** Plasmid segregation was studied in strains whose relevant genotypes are indicated and as described in methods. **D.** Stringent nucleotides pool was measured by TLC for the strains whose relevant genotypes are indicated. Cultures were inoculated at 1:1000 dilution in MOPS medium containing glucose, 20 amino acids and ampicillin with or without IPTG as indicated below the lanes and at least two doubling were allowed in the presence of P^{32} .

Figure 5 - (p)ppGpp degradation is reduced by elevated pppGpp.

A. TLC was performed for wild-type and $\Delta gppA::FRT$ (RS307, white colony) strains grown in MOPS glucose. Isoleucine starvation was induced with valine (arrow, solid line), and reversed by isoleucine addition (arrow, dotted line) and samples were collected at the indicated time and just before valine addition (lanes 1&7). **B.** The ratio of a nucleotide to total (GTP + ppGpp + pppGpp) was plotted for the samples collected before or after valine (Val) or after valine and isoleucine (Ile) addition. Time after valine and subsequent isoleucine addition are indicated below the bars. Data provided are the mean of two independent experiments along with the standard deviation of the mean. **C.** TLC was performed for the *spoT1* strain (RS24); growth condition and sample collection are as described in 5A. **D.** *spoT1* $\Delta gppA/pRCspoT$ (RS194) strain was grown in the MOPS glucose medium containing ampicillin and 1mM IPTG and sub-cultured 1 in1000 in the same medium but without IPTG. Following growth inhibition, TLC was performed as described in Fig. 5A. **E.** The ratio of a nucleotide to the total was plotted from the TLC's shown in Fig. 5C and Fig. 5D. Sample collection times are indicated below the bars.

Figure 6 - Accumulation of ppGpp is associated with transient growth arrest.

A. TLC was performed for the strain $\Delta rlmD \Delta spoT$ (RS361, white colony); growth and sampling are as described for fig. 5A. **B.** The nucleotide pools were quantified from the TLC shown in Fig. 6A as described in Fig. 5B. **C.** Growth inhibition after valine addition (arrow, solid line) and recovery after isoleucine addition (arrow, dotted line) were determined for the $\Delta rlmD \Delta spoT$ strain (RS361, white colony).

Figure 7 - A model based on the results of this study.

RelA activation from amino acid starvation is well established while the signals responsible for basal RelA activity are not clear. The stringent nucleotides generated are turned over by the SpoT and GppA hydrolases. When the SpoT hydrolase activity is lowered, two or more activities prevent the accumulation of pppGpp, one is GppA, the nature of the other one is unclear. The GppA and SpoT hydrolase activities together prevent the activation of RelA-mediated stringent response and growth arrest. The preeminent contribution of pppGpp to the activation of stringent response is because of the inhibition of the SpoT mediated degradation of (p)ppGpp and the allosteric activation of RelA (thick arrow); ppGpp may also allosterically activate RelA. See text for details. Ndk – Nucleoside diphosphate kinase. An arrow indicates activation or products of a biochemical activity, a line ending with a '⊥' indicates inhibition. A thick arrow refers to the strength /importance of the pathway.

Supporting information.

S1 Table. List of strains, plasmids, and primers.

S2 Table. The ratio of the nucleotide to total in the WT and $\Delta gppA$ mutant following amino acid starvation and reversal of starvation.

S3 Table. The ratio of the nucleotide to total in the *spoT1* mutant following amino acid starvation and reversal of starvation.

S4 Table. The ratio of the nucleotide to total in the *spoT1* $\Delta gppA$ /pRC*spoT* strain after SpoT depletion and followed by amino acid starvation and reversal of starvation.

S5 Table. The ratio of nucleotides to total in the $\Delta rlmD::FRT$ $\Delta spoT$ strain following amino acid starvation and the reversal of starvation.

925 **SI references.**

926 **Fig. S1.** SpoT depletion in the $\Delta relA \Delta spoT$ /pRC*spoT* strain. The $\Delta relA \Delta spoT$ (*spoT212*)/
927 pRC*spoT* (AN120) strain cultured overnight in MOPS medium containing glucose, 20 amino
928 acids, ampicillin and 1mM IPTG was washed and sub-cultured at the dilution indicated in the
929 same medium (●) or without IPTG (○).

930 **Fig. S2.** RelA-dependent accumulation of ppGpp but not pppGpp in the *spoT1* strain. *spoT1*
931 (RS24) and $\Delta relA::Kan spoT1$ (RS485, white colony) strains were cultured in MOPS glucose
932 medium and isoleucine starvation was induced with valine.

933 **Fig. S3.** SMG resistance of the hypomorphic RelA alleles is modulated by SpoT function.
934 Saturated cultures were washed, serially diluted and spotted on plates with or without SMG
935 (SMG refers to serine, methionine, and glycine; see methods) and photographed after 20 hrs
936 at 37°C. Strains and their relevant genotypes are indicated. * growth in the presence of SMG
937 was retarded but not abolished.

938 **Fig. S4.** The effect of *spoT* and *gppA* deletions on the growth of the *relA1* strain. Growth was
939 monitored by streaking the indicated strains on Minimal glucose SMG plates and incubating
940 them for the time indicated (A) or in other plates as indicated (B). LB and Minimal Glucose
941 CAA plates were incubated for 24 hours and Minimal glucose plates were incubated for 48
942 hours. The strains were MG1655, *relA1* (RS39), *relA1 ΔspoT* (RS31 white colony), and *relA1*
943 $\Delta spoT \Delta gppA::Kan$ (RS35 white colony).

944 **Fig. S5.** Expression levels of *relA'-lac* and *rlmD'-lac* fusions in LB. β -galactosidase assay
945 was carried out from mid-log phase cultures of the strains RS692 (*relA'-lac*) and RS693
946 (*rlmD'-lac*) grown in LB. The β -galactosidase activity in Miller units was plotted on a log

scale. Values obtained from three independent experiments were used to calculate the mean and standard deviation.

Fig. S6. The absence of pppGpp accumulation in the *spoT1* strain is not corrected by an increase in the GTP level. The *spoT1 gsk3* (RS697 white colony) strain was cultured in MOPS glucose medium containing histidine, tryptophan (each at 100 µg/ml) without guanosine (lanes 1 to 4) and with guanosine (lanes 5 to 8). Isoleucine starvation was induced by the addition of valine (indicated by arrow). Samples were collected immediately before valine addition or subsequently at the time indicated.

Fig. S7. SpoT depletion in *spoT207::Cm ΔgppA::Kan/pRCspoT* (RS206). The *spoT207::Cm ΔgppA::Kan/pRCspoT* (RS206) grown in MOPS medium containing glucose, 20 amino acids, ampicillin and 1mM IPTG was washed and sub-cultured in the same medium (●) or without (○) IPTG at the dilution indicated. Growth after the sub-culture into the medium containing ³²P-H₃PO₄ is shown. The arrows refer to the points at which samples were collected for the TLC shown in Fig. 3B.

Fig. S8. Growth inhibition by valine and the recovery after isoleucine addition were assayed in **A.** MG1655, **B.** *spoT1* (RS24), and **C.** *ΔrlmD* (RS316). Valine (solid arrow) and isoleucine (dashed arrow) were added as indicated.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

46

Fig. 6

Fig. 7