Diverse *Escherichia coli* lineages, from domestic animals and humans in a household, carry colistin resistance gene *mcr-1* in Ecuador

María Fernanda Loayza, ^{a#}, Liseth Salinas ^a, Fernando Villavicencio ^b, Tamayo Rafael ^b, Stephanie Salas ^b, José Villacís ^b, Carolina Satan ^b, Liliana Ushiña ^b, Ruth Rivera ^b, Olga Muñoz ^b, Jeannete Zurita ^c, Tijet Nathalie ^d, Roberto Melano ^{d,e}, Jorge Reyes ^f, Gabriel Trueba ^a

Instituto de Microbiología, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito, Ecuador ^a; Instituto Nacional de Investigación en Salud Pública “Dr. Leopoldo Izquieta Pérez”, Quito, Ecuador ^b; Unidad de Investigaciones en Biomedicina Zurita & Zurita Laboratorios, Quito, Ecuador ^c; Public Health Ontario Laboratory, Toronto, Ontario, Canada ^d; Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada ^e; Facultad de Ciencias Químicas, Universidad Central del Ecuador, Quito, Ecuador ^f

Running Head: Diverse commensal *E. coli* lineages carrying *mcr-1* gen

#Address correspondence to María Fernanda Loayza, mfloayzav@usfq.edu.ec

22

23 **Abstract**

24 The aim of this study was to investigate the presence of *Escherichia coli* carrying *mcr-1* gene
 25 in domestic animals close to a child who suffered a peritoneal infection by a *mcr-1* positive
 26 *E. coli*. Rectal or cloacal swabs and fecal samples from domestic animals were plated on
 27 selective media to isolate colistin-resistant *E. coli* and isolates were submitted to detection of
 28 *mcr-1* gene, pulsed field gel electrophoresis (PFGE), multi-locus sequence typing (MLST),
 29 replicon typing and S1-PFGE. Four *mcr-1* positive *E. coli* isolates (from chicken, turkey and
 30 dog) were recovered. No shared PFGE pattern or MLST sequence type were observed among
 31 isolates. A 60Kb IncI1γ *mcr-1*-carrying plasmid was detected in all isolates. Our results
 32 suggest that *mcr-1* gene was horizontally disseminated amongst different lineages of *E. coli*
 33 from domestic animals in the child's household.

34 **Importance**

35 Horizontally transferable colistin resistance (*mcr-1* gene) is thought to have originated in
 36 domestic animals and transferred to humans through meat and dairy products. In the present
 37 report we show evidence that the *mcr-1* gene could be transferred to different *E. coli* strains
 38 colonizing different hosts (humans and pets) in the same household.

39 **Introduction**

40 Domestic animals are important source of antibiotic resistant bacteria (and genes) which can be
 41 transmitted to humans through the food chain and direct contact (1). Plasmids and other mobile
 42 genetic elements (MGEs) are involved in the transmission of antimicrobial resistance among
 43 different bacterial genera colonizing different animal species (2, 3).

The first report of a colistin resistance (CR) gene carried by plasmids (*mcr-1*) came from China in 2015 (4). This gene codes for a phosphoethanolamine transferase (MCR-1), which modifies the lipid A moiety in the outer membrane of Gram negative bacteria and confers resistance to polymyxins (4, 5). Among Enterobacteriaceae different *mcr* gen groups (1-5) could be transferred by mobile elements (6–8) and have been detected in humans, food animals, environmental samples in several countries, in different bacterial species and in several plasmid types (9). Enterobacteriaceae has been found as the main *mcr* reservoir (6–8). The gene is easily transferred between commensal *E. coli* colonizing domestic animals and humans (7, 10) and opportunistic pathogens such as *E. coli* ST3941 (11) or *Klebsiella pneumoniae* ST512 KPC-3 (7, 12, 13).

In Ecuador, Ortega *et al.* have described the isolation of a colistin resistant *E. coli* carrying *mcr-1* from the peritoneal liquid in a child with complicated peritonitis (14). We investigated the *mcr-1* gene in *E. coli* isolates from domestic animals in this child's household.

Results

Four colistin resistant *E. coli* isolates were recovered from a dog (2 strains), a turkey (1 strain) and a chicken (1 strain). Susceptibility test for each strain is shown in **Table 1**. All CR isolates (colistin (MIC >4 µg/mL) were also resistant to ceftriaxone (MIC ≥64 µg/mL) and ciprofloxacin (MIC ≥4 µg/mL). Additionally, dog isolates were resistant to ampicillin/sulbactam (MIC ≥32 µg/mL) and one of their isolates were resistant to gentamicin (MIC ≥16 µg/mL).

Sequenced PCR products showed to be identical among them and close related to *mcr-1* sequences that were previously reported (NCBI Accession number: KX11520.1, KX011521, KU935449.1, KU935446.1, KU935447, NG050417, KY013597).

Both PFGE and MLST indicated that CR *E. coli* isolates from domestic animals and the child (14) were different (Figure 1, Table 1). Sequence types (STs) carrying the *mcr-1* gene in the patient was ST609 and in the animal samples ST3941, 2170 and 1630. The isolates recovered from chicken and turkey also harbored *bla*_{CTX-M-65} (the one from turkey was *bla*_{TEM-1} as well), and the 2 *E. coli* from the dog were positive for *bla*_{CTX-M-3} and *bla*_{TEM-1} genes. All CR isolates from domestic animals in the household and the isolate from the child had an IncI1 γ plasmid (Table 1). We were unable to transfer the *mcr-1* gene using the conditions described in methods section. S1-PFGE showed a $\approx$ 60Kb plasmid positive for *mcr-1* gene in all isolates (from animal and human) (Figure 1).

Discussion

E. coli genetic diversity has been studied in different environments showing no clear association patterns among different sources (15). In this study, colistin resistant *E. coli* strains isolated from different sources belonged to different lineages and no one was comparable with child isolate despite of they have an identical sequence.

E. coli ST 609 (child strain) has been previously detected in human (commensal and opportunistic pathogens), and water isolates in Spain, Argentina, United States, Vietnam, Netherlands, Australia, and Norway. ST 3941(chicken strain) has been identified in commensal bacteria from livestock and poultry in China and Vietnam farms, and *E. coli* ST 2170 (turkey strain) has been reported as an opportunistic pathogen causing septicemia in Japan and in healthy poultry farms in Vietnam. ST 1630 (Dog strains) was isolated from chickens in Denmark and United States (reported data in <http://mlst.ucc.ie/mlst/dbs/Ecoli>; last accessed December 8, 2017). Moreover, ST 1630 was also isolated from poultry in Japan (16). These data show the diverse and spread sources of *E. coli* strains detected and support the idea that successful mobile genetic elements could be responsible of

antimicrobial resistance transference among dominant strains of commensal bacteria in different niches.

We were unable to transfer the *mcr-1* gene to an acceptor, sodium azide resistant *E. coli*. Previous reports have described a low transfer frequency of *mcr-1*-carrying plasmids that could depends of plasmid type (16). However, we cannot discard that the conditions used in our study were not the optimal for their horizontal transfer.

Using a specific *mcr-1* probe, one of the isolates (from turkey) showed two positive bands by Southern blot analysis: the common plasmid (~61 kb) and another one at ~140 kb. Li R *et al* (2016), have reported double plasmid carrying *mcr-1* gene in commensal *E. coli* strains isolated from healthy pigs in China farms (17, 18). The genetic analysis of those plasmids showed a composite transposon Tn6330 (2 copies of IS*ApII* flanking the *mcr-1* gene) that can form a circular intermediate which mediates the insertion of the *mcr-1* gene cassette into the IncHI2 plasmid.

Further analysis is required to describe the *mcr-1* genetic environment and complete structure of plasmids from *E. coli* strains isolated in our study.

Both S1-PGFE and replicon typing suggest that the *mcr-1* gene was present in the same plasmid in all CR isolates (from domestic animals and human). It is probable that ≈ 60Kb plasmids were responsible for the transference of colistin resistance; previous reports have shown transmission of antibiotic resistance from bacteria in domestic animals to human bacteria is carried out by plasmids and not by antibiotic resistant clones (3).

Our study corroborates the notion that Enterobacteriaceae colonizing intestines in domestic animals (including companion animals) could be reservoir of *mcr-1* gene (2, 19–22). There was no evidence of colistin treatment in animals sampled, however, the presence of this type of antibiotic

in animal feed and supplements in Ecuador prevents us from rule out antibiotic selection of CR clones.

Materials and Methods

2.1 *E. coli* strain isolation and MIC determination

A cross – sectional study was conducted to detect commensal *E. coli* carrying *mcr-1* gene. Thirty-two fecal samples from soil and ten rectal or cloacal swabs were taken from rabbits (n= 2), guinea pigs (n= 2), dogs (n= 2) and chickens (n= 4). Fecal samples (n=32) from soil were placed in sterile reservoirs and swabs were placed in Tryptic Soy Broth (TSB, BD™) (19). Samples were transported to the Antimicrobial Resistance Laboratory in the Instituto Nacional de Investigación en Salud Pública "Dr. Leopoldo Izquieta Perez", Quito. Samples were plated on Mac Conkey Agar plates (MKL, BD™) supplemented with 2 µg/mL of colistin methansulfonate (RICHET®) (20). Identification and antimicrobial susceptibility profiles of the CR isolates were performed using the VITEK®2 compact (bioMérieux) with AST 272 card. Colistin minimal inhibitory concentration (MIC) was performed using Sensititre™ (23).

2.2 Conjugation assay

Conjugation was performed using CR *E. coli* isolates as donor strains and *E. coli* J53 strain (sodium azide-resistant) as recipient (17). Trans-conjugant selection was performed in Trypticase™ soy agar (Difco BD) with colistin (0.5µg/mL) and sodium azide (100 µg/mL) (24).

2.3 Molecular typing

PCR was performed to detect *mcr* gene (4); amplicons were sequence and aligned using *mcr*-1 (NG_055582.1), *mcr*-2 (NG_051171.1), *mcr*-3 (NG_056184.1), *mcr*-4(MG822665.1), *mcr*-5 (MG384740.1) accession numbers with Geneius software. Pulsed field gel electrophoresis (PFGE) (25) and multilocus sequence typing (MLST) was performed on seven housekeeping genes to define clonal relatedness (26). Replicon typing was performed using a commercial kit (PBRT KIT, DIATHE, Fano, Italy) (20, 27, 28). β -lactamase genes (*bla*_{CTX-M-1}, *bla*_{TEM}, *bla*_{SHV}) were detected using primers previously described (29). Briefly, 12,5mL of GoTaq ® Green Master Mix (Promega, Madison, USA) were mixed with 1μL of upstream primer, 10μM, 1μL of downstream primer, 10μM, 1μL of DNA template and 9,5 μL of Nuclease-free water to complete a 25μL of reaction volume. The reaction mix were amplified using 2-minute of initial denaturation at 94°C followed by 40 cycles of DNA denaturation at 94°C (40 sec), annealing at 60°C (40sec) and extension at 72°C (1min). The final elongation step at 72°C for 5 min. Amplicons were detected by electrophoresis in a 2% agarose gel. For complete amplification and sequencing of the detected resistance genes, primers and conditions previously described were used (30).

S1-PFGE and Southern blot.

Plasmid content of each *E. coli* isolate and their estimated sizes were determined by S1 endonuclease-digested genomic DNA and PFGE (S1-PFGE). Briefly, genomic DNA agarose plugs of each isolate were partially digested with the endonuclease S1(31). DNA bands were separated by PFGE under previously described conditions (32). Plasmids were transferred and immobilized on a nylon membrane and identified by Southern blot analysis, using specific *mcr*-1 digoxigenin-labeled probes (Roche Diagnostics).

Conclusions

Our study suggests a polyclonal dissemination of *mcr-I* gene in *E. coli* domestic animals and humans in an Ecuadorian household. A 60Kb IncI1 γ plasmid carried the *mcr-I* gene in this study, which may have been transferred among different strains colonizing different hosts. This study supports the current notion that mobile gene elements are more important than bacterial clones in the transmission of antibiotic resistance genes from the microbiota in domestic animals to human microbiota (33).

Our results highlight the importance of controlling the use of antibiotics in domestic animals and possibly the need for more studies of AR in isolates and mobile genetic elements in them.

Funding

This study was funded by The Instituto Nacional de Investigación en Salud Pública "Dr. Leopoldo Izquieta Perez", Quito – Ecuador and Instituto de Microbiología, Universidad San Francisco de Quito.

Ethical approval: No required

Competing interest: The authors declare that they don't have any conflict of interests.

References

1. Pompa C, Rantala M, Greko C, Baptiste K, Catry B, Van Duijkeren E, Mateus ALP, Moreno MA, Pyörälä S, Ruzauskas M, Sanders P, Teale C, Threlfall J, Kunsagi Z, Torren_Edo J, Jukes H, Törneke K. 2016. Public health risk of antimicrobial resistance transfer from companion animals. J Antimicrob Chemother 1–12.
2. Mukerji S, O'Dea M, Barton M, Kirkwood R, Lee T, Abraham S. 2017. Development and transmission of antimicrobial resistance among Gram-negative

bacteria in animals and their public health impact. *Essays Biochem* 61:23–35.

3. De Been M, Lanza V, De Toro M, Scharringa J, Dohmen W, Du Y, Hu J, Lei Y, Li N, Tooming-Klunderud A, Heederik DJJ, Fluit AC, Bonten MJM, Willems RJL, Cruz F De, Schaik W Van. 2014. Dissemination of Cephalosporin Resistance Genes between *Escherichia coli* Strains from Farm Animals and Humans by Specific Plasmid Lineages. *Plos Genet* 10.
4. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: A microbiological and molecular biological study. *Lancet Infect Dis* 16:161–168.
5. Poirel L, Jayol A, Nordmann P. 2017. Polymyxins: Antibacterial activity, susceptibility testing and resistance mechanisms encoded by plasmid or chromosomes. *Clin Microbiol Rev* 30:557–596.
6. Schwarz S, Johnson AP. 2016. Transferable resistance to colistin: a new but old threat. *J Antimicrob Chemother* dkw274.
7. Gao R, Hu Y, Li Z, Sun J, Wang Q, Lin J, Ye H, Liu F, Srinivas S, Li D, Zhu B, Liu Y-H, Tian G-B, Feng Y. 2016. Dissemination and mechanism for the MCR-1 colistin resistance. *PLOS Pathog* 12:e1005957.
8. Lescat M, Poirel L, Nordmann P. 2018. Rapid multiplex PCR for detection of mcr-1 to 5 genes. *Diagn Microbiol Infect Dis*.
9. Skov RL, Monnet DL. 2016. Plasmid-mediated colistin resistance (mcr-1 gene); three months later, the story unfolds. *Euro Surveill* 21:1–6.
10. Poirel L, Nordmann P. 2016. Emerging plasmid-encoded colistin resistance: the

- 199 animal world as the culprit? J Antimicrob Chemother dkw074.
- 200 11. Corbella M, Mariani B, Ferrari C, Comandatore F, Scaltriti E, Marone P, Cambieri
201 P. Three cases of mcr-1-positive colistin-resistant *Escherichia coli* bloodstream
202 infections in Italy.
- 203 12. Di Pilato V, Arena F, Carlo T, Cannatelli A, Angelis LH De, Fortunato S, Giani T,
204 Menichetti F, Rossolini GM. 2016. mcr-1,2, a new mcr variant carried a transferable
205 plasmid from a colistin-resistant KPC Carbapenemase-producing *Klebsiella*
206 *pneumoniae* strain of sequence type 512. Antimicrob Agents Chemother 60:25–34.
- 207 13. Castanheira M, Griffin MA, Deshpande LM, Mendes RE, Jones RN, Flamm RK.
208 2016. Detection of mcr-1 among *Escherichia coli* clinical isolates collected
209 worldwide as part of the SENTRY Antimicrobial Surveillance Program during 2014-
210 2015. Antimicrob Agents Chemother 316:806–807.
- 211 14. Ortega D, Barba P, Zurita J. 2016. Colistin-resistant *Escherichia coli* clinical isolate
212 harbouring the mcr-1 gene in Ecuador. Epidemiol Infect 1–4.
- 213 15. Tenaillon O, Skurnik D, Picard B, Denamur E. 2010. The population genetics of
214 commensal *Escherichia coli*. Nat Rev Microbiol 8.
- 215 16. Hiki M, Usui M, Akiyama T, Kawanishi M, Tsuyuki M, Imamura S, Sekiguchi H,
216 Kojima A, Asai T. 2014. Phylogenetic grouping, epidemiological typing, analysis of
217 virulence genes, and antimicrobial susceptibility of *Escherichia coli* isolated from
218 healthy broilers in Japan. Ir Vet J 67:14.
- 219 17. Li R, Xie M, Lv J, Wai-Chi Chan E, Chen S. 2016. Complete genetic analysis of
220 plasmids carrying mcr-1 and other resistance genes in an *Escherichia coli* isolate of
221 animal origin. J Antimicrob Chemother dkw509.
- 222 18. Li R, Xie M, Zhang J, Yang Z, Liu L, Liu X, Zheng Z, Chan EW-CC, Chen S. 2017.

223 Genetic characterization of mcr-1-bearing plasmids to depict molecular mechanisms
 224 underlying dissemination of the colistin resistance determinant. J Antimicrob
 225 Chemother 72:393–401.

226 19. Agga GE, Schmidt JW, Arthur TM. 2016. Effects of In - Feed Chlortetracycline
 227 prophylaxis of beef cattle on animal health and Antimicrobial -Resistant *Escherichia*
 228 *coli*. Appl Environ Microbiol.

229 20. do Monte DFM, Mem A, Fernandes MR, Cerdeira L, Esposito F, Galvão JA, Franco
 230 BDGM, Lincopan N, Landgraf M. 2017. Chicken Meat as Reservoir of Colistin-
 231 Resistant *Escherichia coli* Carrying mcr-1 Genes in South America. Antimicrob
 232 Agents Chemother AAC.02718-16.

233 21. Brown DFJ, Wootton M, Howe RA. 2016. Antimicrobial susceptibility testing
 234 breakpoints and methods from BSAC to EUCAST 3–5.

235 22. Lei L, Wang Y, Schwarz S, Waish T, Ou yanran, Wu Y, Li M, Shen Z. 2017. mcr-1
 236 in Enterobacteriaceae from companion animals, beijing, China, 2012-2016. Emerg
 237 Infect Dis 23:710–1.

238 23. Committee TE, Testing AS, Changes N, Pseudomonas E. 2015. European
 239 Committee on Antimicrobial Susceptibility Testing Breakpoint tables for
 240 interpretation of MICs and zone diameters European Committee on Antimicrobial
 241 Susceptibility Testing Breakpoint tables for interpretation of MICs and zone
 242 diameters.
 243 [http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_50_Breakpoint_Table_01.pdf)
 244 [v_50_Breakpoint_Table_01.pdf](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_50_Breakpoint_Table_01.pdf) 0–77.

245 24. Poirel L, Kieffer N, Brink A, Coetze J, Jayol A, Nordmann P. 2016. Genetic features
 246 of MCR-1-producing colistin-resistant *Escherichia coli* isolates in South Africa.

247 Antimicrob Agents Chemother 60:4394–4397.

248 25. Pulsenet International. 2013. Standard Operating Procedure for Pulsenet Pfg of

249 *Escherichia coli* O157:H7, *Escherichia coli* Non-O157 (Stec), Salmonella Serotypes,

250 Shigella Sonneiand Shigella Flexneri 157:1–14.

251 26. Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, Karch H, Reeves PR,

252 Maiden MCJ, Ochman H, Achtman M. 2006. Sex and virulence in *Escherichia coli*:

253 An evolutionary perspective. Mol Microbiol 60:1136–1151.

254 27. Carattoli A. 2013. Plasmids and the spread of resistance. Int J Med Microbiol

255 303:298–304.

256 28. Tijet N, Faccone D, Rapoport M, Seah C, Pasterán F, Ceriana P, Albornoz E, Corso

257 A, Petroni A, Melano RG. 2017. Molecular characteristics of mcr-1-carrying

258 plasmids and new mcr-1 variant recovered from polyclonal clinical *Escherichia coli*

259 from Argentina and Canada. PLoS One 12:e0180347.

260 29. Mohammed H, Fakhr AE, Mohammed H, Al E, Abdel W, Hassanein G. 2016.

261 Spread of TEM, VIM, SHV, and CTX-M b- lactamases in Imipenem-Resistant

262 Gram-Negative Bacilli Isolated from Egyptian Hospitals. Int J Microbiol 2016:15.

263 30. Tijet N, Andres P, Chung C, Lucero C, Group W-A, Low DE, Galas M, Corso A,

264 Petroni A, Melano RG. 2011. rmtD2, a new allele of a 16S rRNA methylase gene,

265 has been present in Enterobacteriaceae isolates from Argentina for more than a

266 decade. Antimicrob Agents Chemother 55:904–909.

267 31. Barton BM, Harding GP, Zuccarelli AJ. 1995. A General Method for Detecting and

268 Sizing Large Plasmids. Anal Biochem 226:235–240.

269 32. Borgia S, Lastovetska O, Richardson D, Eshaghi A, Xiong J, Chung C, Baqi M,

270 McGeer A, Ricci G, Sawicki R, Pantelidis R, Low DE, Patel SN, Melano RG. 2012.

271 Outbreak of Carbapenem-Resistant Enterobacteriaceae Containing blaNDM-1,
 272 Ontario, Canada. Clin Infect Dis 55:e109–e117.
 273 33. Szmolka A, Nagy B. 2013. Multidrug resistant commensal *Escherichia coli* in
 274 animals and its impact for public health. Front Microbiol 4:1–13.
 275
 276

277 **Table 1.** Antibiotic susceptibility profiles and molecular analyses (PCR for *bla* genes detection, MLST and Replicon typing) of *mcr-1*-positive *E. coli*
 278 isolates.

Strain origin	Antimicrobial resistance	MLST	PCR	Plasmid Inc group										
		ST	<i>bla</i> genes	HI1	I2	N	FIA	FIB	I1y	FIIS	R	X1	Y	FII
Chicken	COL, CRO, CIP	3941	<i>bla</i> _{CTX-M-65}	-	+	-	+	-	+	-	-	-	-	-
Turkey	COL, CRO, CIP	1630	<i>bla</i> _{TEM-1} ; <i>bla</i> _{CTX-M-65}	+	-	-	-	-	+	-	-	-	-	-
Dog1	COL, SAM, CRO, CIP	2170	<i>bla</i> _{TEM-1} ; <i>bla</i> _{CTX-M-3}	-	+	-	-	+	+	+	-	-	+	+
Dog2	COL, SAM, CRO, CIP, GN	2170	<i>bla</i> _{TEM-1} ; <i>bla</i> _{CTX-M-3}	-	-	-	-	-	+	-	-	-	-	-
Child*	COL, CAZ, CRO, FEP, CIP	609 [13]	<i>bla</i> _{CTX-M-55} [13]	-	+	+	-	-	+	-	+	+	-	-

279 COL, colistin; SAM, ampicillin/sulbactam; CAZ, ceftazidime; CRO, ceftriaxone; FEP, cefepime; GN, gentamicin; CIP, ciprofloxacin.

280 ST, sequence type obtained by MLST analysis; *bla* genes, β -lactamase genes; Poultry TC, Transconjugant strain obtained with conjugation using

281 chicken CR *E. coli* isolate as donator and *E. coli* J53 as receptor strain. Replicon typing analysis was performed with PBRT KIT (DIATHEVA):

282 only positive Inc groups in the overall analysis are included. * isolate reported by Ortega *et al*

283

Figure 1. Identification of *mcr-1*-harboring plasmids. (A) S1- PFGE plasmid profiles of the patient and animals *mcr-1* positive *E. coli*. (B) Southern blot using *mcr-1* probe. Arrows indicate the plasmids carrying *mcr-1* gene. Ladder, reference standard *Salmonella enterica* serotype Braenderup strain H9812 restricted with *Xba*I (sizes are given in kilobases)