Polyploidy and elevation contribute to opposing latitudinal gradients in diversification and species richness in lady ferns (Athyriaceae)

Ran Wei¹, Richard H. Ree², Michael A. Sundue³ and Xian-Chun Zhang¹

¹State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, the Chinese Academy of Sciences, Beijing 100093, China; ²Life Sciences Section, Integrative Research Center, The Field Museum, Chicago, IL 60605, USA; ³The Pringle Herbarium, University of Vermont, 27 Colchester Drive, Burlington, VT 05405, USA

Author for correspondence:

Ran Wei

Tel: +86 10 62836479

Email: weiran@ibcas.ac.cn

Short running head: Diversification of *Athyrium* and *Diplazium*

Number of words in the main text: 4563;

Number of words in Introduction: 885;

Number of words in Materials & Methods: 1513;

Number of words in Results: 645;

Number of words in Discussion: 1423;

Number of words in Acknowledgements: 97;

Number of figures: 4 (3 are in color);

Number of tables: 2;

Number of supporting information: Five figures and six tables.

34

35 **Summary**

- 36 • In ferns, the temperate-tropical sister clades *Athyrium* and *Diplazium* present an
37 opportunity to study a latitudinal contrast in diversification dynamics.
- 38 • We generated a taxonomically expanded molecular chronogram and used
39 macroevolutionary models to analyze how diversification rates have changed through
40 time, across lineages, and in concert with changes in elevation and ploidy. We tested a
41 novel model of cladogenetic state-change in which polyploidy can arise as an
42 infraspecific polymorphism, with diversification parameters distinct from those of
43 pure diploids and polyploids.
- 44 • Both *Athyrium* and *Diplazium* accelerated their diversification near the Oligocene-
45 Miocene transition. In *Diplazium*, the rate shift is older, with subsequent net
46 diversification somewhat slower and suggestive of diversity-dependence. In *Athyrium*,
47 diversification is faster and associated with higher elevations. In both clades,
48 polyploids have the highest rate of net accumulation but lowest (negative) net
49 diversification, while the converse is true for polymorphic species; diploids have low
50 rates of both net accumulation and diversification.
- 51 • Diversification in *Athyrium* may have responded to ecological opportunities in
52 expanding temperate habitats during the Neogene, especially in mountains, while the
53 pattern in *Diplazium* suggests saturation in the tropics. Neopolyploids are generated
54 rapidly, primarily through accelerated cladogenesis in polymorphic species, but are
55 evolutionary dead ends.

56 **Key words:** Athyriaceae, elevation, latitudinal gradient, macro-evolution, SSE
57 models, polyploidy

58

59

60

61

62

63

64

65

66

67

68

69 **Introduction**

70 How does the tempo (rate) and mode (process) of lineage diversification vary across
 71 latitude? Consideration of this question has largely focused on explaining the
 72 latitudinal gradient in species richness, especially with respect to how temperate-
 73 tropical contrasts in time-integrated area, temperature and metabolic rate, and
 74 ecological opportunity might influence the rate of speciation (reviewed in Schluter,
 75 2015). A general prediction is that, all else being equal, tropical clades should be
 76 older, and/or have higher net diversification, compared to temperate clades (Wiens &
 77 Donoghue, 2004; Mittelbach *et al.*, 2007). In this context, temperate-tropical sister
 78 clades that otherwise share similar ecologies offer opportunities to study the correlates
 79 and potential drivers of diversification across latitudes over a common timescale.

80 One such opportunity presents itself in the lady-fern family Athyriaceae, in which
 81 the primarily temperate genus *Athyrium*, including allied genera, is sister to the
 82 primarily tropical *Diplazium* (Wei *et al.*, 2013, 2018). These clades respectively
 83 include about 220 and 350 species that occur in habitats that range from wet and
 84 shaded streambanks at lower altitudes to dry and open grasslands higher up (Tryon &
 85 Tryon, 1982; Wang *et al.*, 2013). Both clades are more or less globally distributed,
 86 with the majority of species in tropical and temperate Asia (Wang *et al.*, 2013) and
 87 the tropical Andes (Tryon & Tryon, 1982), and each has been the subject of recent
 88 systematic and biogeographic studies (Wei *et al.*, 2013, 2015, 2018). Here, we
 89 generate a more comprehensive dataset for comparative analyses of diversification: in
 90 particular, how the timing, extent, and phylogenetic distribution of rate heterogeneity
 91 differ in these clades, and what this reveal about their responses to potential drivers of
 92 diversification.

93 In considering such drivers, a common theme that emerges from prior studies of
 94 fern diversification is ecological opportunity. Seminal work at broad taxonomic and
 95 deep temporal scales emphasized how fern lineages, particularly epiphytes,
 96 proliferated ‘in the shadow of angiosperms’, i.e. in novel habitats created as the latter
 97 ascended in the late Cretaceous/early Cenozoic (Schneider *et al.*, 2004; Schuettpelz &
 98 Pryer, 2009; but see Testo & Sundue, 2016). More recently, birth–death models fitted
 99 to phylogenies and the fossil record showed evidence for diversity-dependent

100 origination, interpreted as reflecting opportunistic niche-filling, and extinction driven
101 by environmental change (Lehtonen *et al.*, 2017). For example, in Polypodiaceae,
102 faster diversification is correlated with changes in elevation, a proxy for habitat type,
103 suggesting that niche shifts drive lineage proliferation (Sundue *et al.*, 2015). In the
104 case of *Athyrium-Diplazium*, one might predict that diversification in *Athyrium* was
105 accelerated by ecological opportunities arising during the expansion of temperate
106 habitats during the Neogene, compared to *Diplazium*, which might be in the later
107 stages of diversity-dependent clade growth, reflecting its persistence in more
108 continuously stable tropical environments. Both clades might show responses to
109 elevation, reflecting their respective colonization of temperate and tropical mountains,
110 such as temperate Hengduan Mountains and the tropical Kinabalu and Andes.

111 Another potential driver of diversification that is not directly related to latitude, but
112 demands attention due to its prevalence in ferns, is polyploidy (e.g. Schneider *et al.*,
113 2017). There is evidence that more than 50% of the species in the *Athyrium*-
114 *Diplazium* clade are polyploid (e.g. Praptosuwiryo, 2008; Bir & Verma, 2010). The
115 effect of polyploidy on plant diversification has received much attention from both
116 theoretical and empirical perspectives (e.g. Mayrose *et al.*, 2011; Soltis *et al.*, 2016;
117 reviewed in Vamosi *et al.*, 2018). In particular, polyploid formation as a mode of
118 speciation motivated the development of cladogenetic state-dependent diversification
119 models, in which the rate of speciation coincident with state change is parameterized
120 (Mayrose *et al.*, 2011; Zhan *et al.*, 2016; see also Goldberg & Igić, 2012). These have
121 since been used to show that neopolyploids tend to be evolutionary 'dead ends'
122 (Mayrose *et al.*, 2011; Arrigo & Barker, 2012) due to effects such as gene imbalance
123 of the sex chromosomes during the meiosis (Orr, 1990), increased abortion in
124 heteroploid hybrids (Ramsey & Schemske, 2002), and inefficiency of selection for
125 multi-copy genes in polyploids (Wright, 1969). We wish to know if this holds true in
126 the diversification dynamics of *Athyrium* and *Diplazium*.

127 Previous studies of neopolyploidy using cladogenetic state-change models only
128 considered two states for species, diploid and polyploid (Zhan *et al.*, 2016). However,
129 infraspecific variation in ploidy is common in plants (Wood *et al.*, 2009); in the
130 *Athyrium-Diplazium* clade, 5–10% of species have polyploid individuals recorded
131 within one or more diploid populations (Walker, 1966; Tryon & Tryon, 1982;
132 Takamiya *et al.*, 1999, 2000; Takamiya & Ohto, 2001; Praptosuwiryo, 2008; Bir &

Verma, 2010). Such cases of polymorphism may represent incipient speciation via polyploidization. To test this idea, we constructed cladogenetic state-change models that include three states, diploid, polyploid, and a polymorphic diploid/polyploid state, with free parameters assigned to the rate at which polymorphic species give rise to new neopolyploids via cladogenesis events.

Our study can be summarized as a temperate-tropical contrast in diversification dynamics, focusing on how sister clades have responded to common potential drivers of diversification. Specifically, our aims are to reconstruct a time-scaled global phylogeny of the *Athyrium-Diplazium* clade, estimate variation in diversification rates through time, and infer the effects of elevation and polyploidy. Our intent is to shed light on latitudinal differences in the tempo and mode of diversification, and contribute to a more general understanding of the latitudinal diversity gradient.

Materials and Methods

Taxon sampling and molecular data

Our sampling strategy followed the most recent systematic studies of Athyriaceae (Wei *et al.*, 2013, 2015, 2018; PPG I, 2016). In this study, the circumscription of *Athyrium* includes four closely related genera (*Anisocampium*, *Athyrium*, *Cornopteris* and *Pseudathyrium*) and ten sections (sects. *Athyrium*, *Biserrulata*, *Dissitifolia*, *Mackinnoniana*, *Otophora*, *Polystichoides*, *Rupestria*, *Spinulosa*, *Wallichiana* and *Yokoscentia*) (Supporting Information Table S1). Since no sectional classification in the *Diplazium* clade (especially in subgenus *Callipteris*), we tentatively subdivided *Diplazium* into 11 clades (CHI, DIP, LEP, MET, MON, NEO, PSE, SEA, SIB, SPE and UNK) according to their morphological similarity, phylogenetic relationship and biogeographical affinity (Table S1).

In total, 85 out of *c.* 220 species of *Athyrium* (39%) and 129 out of *c.* 350 species of *Diplazium* (37%) on the global scale were sampled, including representatives from all taxonomic subdivisions or infrageneric groups covering the entire geographical distribution range and ecological habitats of these two genera. Voucher information is listed in Table S2. We included 26 representatives from Aspleniaceae, Athyriaceae, Blechnaceae, Cystopteridaceae, Desmophlebiaceae, Diplaziopsidaceae, Dryopteridaceae, Hemidictyaceae, Onocleaceae, Rhachidosoraceae, Thelypteridaceae and Woodsiaceae as outgroups and to provide appropriate nodes for fossil calibrations.

We assembled a dataset of eight chloroplast regions (*atpA*, *atpB*, *matK*, *rbcL*, *rpl32-trnP*, *rps4*, *rps4-trnS*, *trnL-F*) with a total aligned length of 8297 bp. We used 1129 previously published sequences and generated 336 new sequences (Table S2). DNA extraction, amplification, sequencing, and alignment protocols followed Wei *et al.* (2013, 2018).

171

172 **Divergence time estimation**

We inferred the posterior distribution of chronograms using BEAST 1.8.4 (Drummond & Rambaut, 2007) with the alignment partitioned by protein coding regions and intergenic spacers. Preliminary analyses rejected a strict molecular clock, so we used the uncorrelated lognormal relaxed clock model with a birth–death prior on tree shape. Eight independent runs of 30 million generations, sampling every 1000 generations, were carried out on the Cyberinfrastructure for Phylogenetic Research (CIPRES) Science Gateway (<http://www.phylo.org>; Miller *et al.*, 2010). The resulting log files were combined (with the first 50% samples discarded as burn-in) using LOGCOMBINER 1.8.4 (Drummond & Rambaut, 2007) and checked in TRACER 1.6 to make sure the effective sampling sizes for most of the relevant estimated parameters were well above 200. The tree files were resampled from each run using the same burn-in strategy in LOGCOMBINER. The maximum clade credibility (MCC) topology with a posterior probability limit of 0.5 and mean branch lengths was summarized using TREEANNOTATOR 1.8.4 (Drummond & Rambaut, 2007).

Four calibration points were used. For the root node the prior was a Normal distribution with a mean age of 107.29 Myr and a SD of 10 (with 95% highest posterior density [HPD]: 90.84–123.7 Myr) to cover the range of published split times from 91 Myr to 125 Myr including the 95% HPD intervals (Schneider *et al.*, 2004; Schuettpelz & Pryer, 2009; Rothfels *et al.*, 2015; but see Testo & Sundue, 2016). We used the Paleocene fossil of *Onoclea* (Rothwell & Stockey, 1991) to constrain the minimum age of the Onocleaceae crown group with a lognormal prior distribution with an offset of 54.5 Myr and mean of 1.0 (with 95% HPD: 55.02–68.58 Myr). We used earliest fossil record of *Woodwardia*, known from Paleocene deposits of North America, to constrain the crown node of Blechnaceae (Collinson, 2001), using the same prior settings. We assigned the *Diplazium*-like fossil reconstructions (*Makopteris princetonensis*, Stockey *et al.*, 1999; *Dickwhitea allenbyensis*, Karafit

199 *et al.*, 2006) found in the Middle Eocene deposits of British Columbia in North
200 America to the stem lineage of *Diplazium* (Wei *et al.*, 2015), yielding a lognormal
201 prior with an offset of 36.5 Myr and mean of 2.0 (with 95% HPD: 37.93–74.78 Myr)
202 for the crown node of the *Athyrium-Diplazium* clade.

203

204 **State-independent diversification analysis**

205 We used Bayesian Analysis of Macroevolutionary Mixtures (BAMM) 2.2.2 (Rabosky,
206 2014) to estimate rates of speciation (λ) and extinction (μ) through time and across
207 clades on the MCC tree, pruned to the ingroup *Athyrium-Diplazium* clade. To account
208 for incomplete sampling, we estimated the sampling fractions of infrageneric groups
209 (Table S1) from our own taxonomic knowledge and the literature (e.g. Ching, 1964;
210 Kato, 1977; Tryon & Tryon, 1982; Hsieh, 1986; Tryon & Stolze, 1991; Zhang, 1992;
211 Stolze *et al.*, 1994; Wang, 1997; Mickel & Smith, 2004; Y.C. Liu *et al.*, 2011; Wang
212 *et al.*, 2013; Wei *et al.*, 2013, 2018). Four MCMC analyses were run for 3 million
213 generations each, sampled every 1000 generations. Each run was checked to ensure
214 that the effective sample size (ESS) exceeded 200, and the first 10% of samples were
215 discarded as burn-in. Results were summarized using BAMMTOOLS (Rabosky *et al.*,
216 2014). To evaluate the best model generated by BAMM (compared with a null model
217 with no diversification rate shifts), we relied on Bayes Factors calculated with the
218 COMPUTEBAYESFACTOR function of BAMMTOOLS.

219

220 **State-dependent diversification analyses**

221 We tested two variables for an association with diversification: mean elevation and
222 ploidy. Data were obtained from a variety of sources, including online databases
223 (Global Plants, <https://plants.jstor.org>; Tropicos, <http://www.tropicos.org>; IPCN,
224 <http://www.tropicos.org/Project/IPCN>; CCDB,
225 <http://ccdb.tau.ac.il/Pteridophytes/Athyriaceae>), herbarium specimens, and cytological
226 reports (e.g. Mehra & Bir, 1960; Walker, 1966, 1973; Tryon & Tryon, 1981;
227 Takamiya *et al.*, 1999, 2000; Takamiya & Ohta, 2001; Praptosuwiryo, 2008; Bir &
228 Verma, 2010), and are available in Table S3.

229 To test the hypothesis of net diversification accelerated by divergence in elevation
230 (Sundue *et al.*, 2015), we used BAMM to estimate the rates of elevation change (β) at
231 the tips of the MCC tree, and performed phylogenetic generalized least squares

(PGLS) regression of β on tip values of $r = \lambda - \mu$ using GEIGER (Harmon *et al.*, 2008). In addition, we tested for a direct effect of elevation using the quantitative-state speciation-extinction (QuaSSE) model (FitzJohn, 2010). We considered seven candidate relationships between elevation (log-transformed) and λ : 1) λ is constant and independent of elevation; 2) linear; 3) sigmoid; 4) unimodal, represented by a vertically offset Gaussian function; and another three models (linear, sigmoid, and unimodal) with a directional tendency (Table S4). We ran each model on the entire ingroup clade as well as separately on *Athyrium* and *Diplazium* using DIVERSITREE 0.9-10 (FitzJohn, 2012). We compared models using the Akaike information criterion (AIC).

To explore the effect of ploidy on diversification we used the cladogenetic state change speciation-extinction (ClaSSE) model (Goldberg & Igic, 2012). Species were scored as 1 (polymorphic, i.e. diploid with records of intermingled polyploids), 2 (diploid), 3 (polyploid), or NA (unknown). We considered reports of $2n = 80$ for *Athyrium* and $2n = 82$ for *Diplazium* as diploid. All other species data were multiples of these numbers (e.g. $2n = 160$, $2n = 123$, $2n = 164$, $2n = 205$, $2n = 246$, $2n = 328$) and were thus scored as polyploids (Table S3). We constructed a variety of models with up to seven cladogenetic rates, denoted as a triplet of values for the ancestral and descendant states: λ_{111} , λ_{112} , λ_{113} , λ_{123} , λ_{222} , λ_{223} , λ_{333} , three extinction rates (μ_1 , μ_2 , μ_3), and four transition rates (q_{12} , q_{13} , q_{21} , q_{23}) (Fig. 1). We disallowed anagenetic change from polyploidy (3) to diploidy (1, 2) because no diploidization events were reported according to previous cytological studies and our CHROMEVOLE analysis (see results). After some preliminary analyses, we settled on six candidate models: 1) a full model with all 13 free parameters; 2) a null model with three parameters, in which diversification is state-independent and anagenetic change is symmetric; 3) a ‘cladogenetic’ model with 11 parameters and a single rate of anagenetic change; 4) an ‘anagenetic’ model with five parameters and a single rate for all speciation and extinction events; 5) an ‘extinction’ model with five parameters and 1 rate each for speciation and anagenetic change; and 6) a ‘cladogenetic-anagenetic’ model with 11 parameters, and a single rate for extinction (Fig. 1; Table S5). Models were run on the whole ingroup clade and separately on each genus, and compared using AIC. Parameters of the best-fit models were estimated using MCMC for 100,000 generations in DIVERSITREE.

265 To test the reliability of our ClaSSE analysis given that nearly 40% of the sampled
266 species have unknown ploidy, we simulated 100 trees of the same size as our ingroup
267 clade using the maximum likelihood parameter values of the best-fit model. For each
268 tree we ‘impoverished’ the tip states by randomly setting 40% to the unknown state,
269 NA, and performed model selection using the candidate pool as described above. We
270 also compared the parameters estimated from the impoverished data to their
271 simulation values. All procedures were carried out using DIVERSITREE 0.9-10.

272

273 **Reconstruction of polyploidy evolution**

274 To infer the evolutionary history of polyploidy along the phylogeny, we carried out
275 an ancestral state reconstruction using CHROMEVOLE 2.0 (Glick & Mayrose, 2014)
276 implemented in RASP 4.0 (Yu *et al.*, 2015). To shorten the calculation time, we used
277 the ‘Auto_run’ option, which automatically optimized all the parameters for
278 chromosome evolution models, including rates of single ascending dysploidy (γ),
279 descending dysploidy (δ), whole-genome duplication (ρ) and demi-polyploidy (μ_d).
280 After model optimization, 10,000 simulations were performed using both maximum
281 likelihood and Bayesian approaches to summary the final reconstruction result.

282

283 **Results**

284 **Divergence time estimation**

285 Our molecular dating analysis yielded age estimates that are broadly congruent with
286 those of previous studies with the variations less than 10 Myr (Schneider *et al.*, 2004;
287 Schuettelpelz & Pryer, 2009; Rothfels *et al.*, 2015; Wei *et al.*, 2015; but see Testo &
288 Sundue, 2016 using a different strategy of divergence time estimation based on a time
289 calibrated phylogeny) (Fig. S1). Those that differed tended to be older (e.g. the crown
290 node of the *Diplazium* clade, 47.9 Myr [95% HPD: 41.1–59.4 Myr] in the present
291 study; 41.7 Myr [95% HPD: 33.6–49 Myr] in Wei *et al.*, 2015). This could be due, at
292 least in part, to our much denser sampling of *Athyrium* and *Diplazium* (214 species
293 versus 92 in Wei *et al.*, 2015).

294

295 **Diversification analyses: BAMM, QuaSSE and ClaSSE**

296 The BAMM analysis indicated significant increases in net diversification rate in both
297 *Athyrium* and *Diplazium* (Figs 2a and S2; Table S6). In *Athyrium*, two shifts were

inferred at 16.86 Myr and 5.02 Myr, with the latter leading to a rate 17 times higher than the root rate (0.044–0.76 events Myr^{-1} per lineage; Fig. 2a). In *Diplazium* one shift was inferred at 23.43 Myr (Fig. 2a). Rates-through-time (RTT) plots revealed contrasting diversification dynamics between *Athyrium* and *Diplazium* following their initial shifts (Fig. 2b). In *Athyrium*, the net diversification rate increased continuously, while in *Diplazium* the rate plateaued to a level about half that of *Athyrium* at the present.

Diversification is associated with elevation itself but not its evolutionary rate. The PGLS regression of the rate of change in elevation on net diversification was nonsignificant ($r^2 = 0.286$, $P = 0.379$). By contrast, the best-fit QuaSSE model ($w_i = 0.88$; Table S4) for the *Athyrium-Diplazium* clade included a unimodal relationship between λ and elevation, with λ highest between 1440 m to 3500 m and constant μ (Fig. 2a). There was a negative directional tendency in *Athyrium* (-0.091) and the entire ingroup clade (-0.086), but a relatively low negative directional tendency in *Diplazium* (-0.0013) (Table 1). Separate analyses of *Athyrium* and *Diplazium* showed differences in their unimodal curves for λ . In *Diplazium* the peak is at a lower rate and elevation, and the curve closely matches the frequency distribution of species by elevation, while in *Athyrium* the peak is higher in rate and elevation, and the curve is shifted right relative to the species' frequency distribution (Fig. 3). The best-fit model for *Athyrium* includes a negatively valued directional tendency parameter, but the one for *Diplazium* does not (Table 1).

Our ClaSSE analyses yielded strong support for ploidy-dependent diversification dynamics. The 'cladogenetic' model, with 10 free parameters for speciation and extinction but no differences in q was selected by AIC (Fig. 4a,b,c,d; Table 2). With this model, rates of speciation in polyploid ($\lambda_3 = 0.7054$ events Myr^{-1} per lineage) and polymorphic species ($\lambda_1 = 1.388$ events Myr^{-1} per lineage) exceed rates for diploids ($\lambda_2 = 0.1993$ events Myr^{-1} per lineage), with the total rate being highest for polymorphic species (Fig. 4a,d; Table 2). Extinction rates are low for diploid and polymorphic species but high for polyploids (Fig. 4c); as a result, the net diversification rate of the polymorphic state is the highest, the rate for diploids is lower but positive, and the rate for polyploids is lowest and slightly negative (Fig. 4e). However, the net accumulation rate of polyploids is highest, followed by diploids and polymorphic species (Fig. 4f). In our simulation analysis of the effect of missing data

on model selection, 93% of replicates recovered the correct cladogenetic model (Fig. S3; Table S7), indicating that in this case, model selection is not obviously biased by 40% missing data.

Our CHROMEOL analysis inferred no dysploidization or diploidization events, and at least 15 polyploidization events indicating that all polyploid species in *Athyrium* and *Diplazium* are neopolyploids (Fig. S4).

Discussion

Latitudinal contrasts in diversification between temperate *Athyrium* and tropical *Diplazium*

Our dated phylogeny and BAMM analysis showed that species diversity in the primarily temperate genus *Athyrium* has accumulated relatively recently, with an increasing net rate through the Neogene, compared to its primarily tropical sister clade *Diplazium*, which showed dynamics more suggestive of diversity-dependence. Similar temperate-tropical rate contrasts that run counter to the latitudinal diversity gradient have been found in fishes, birds, mammals, and seed plants (Weir & Schluter, 2007; Schluter, 2015; Spriggs *et al.*, 2015), and contradict the hypothesis that higher diversity in the tropics is the result of faster per-lineage diversification (e.g. Mittelbach *et al.*, 2007). The *Athyrium-Diplazium* contrast may reflect how diversification has responded at a coarse scale to waxing and waning ecological opportunities in temperate and tropical habitats, respectively. In this view, the relatively recent and rapid radiation in *Athyrium* was driven by increasing niche availability in expanding temperate habitats, while for *Diplazium*, the contraction of tropical forests had the opposite effect, as global cooling began in the Late Oligocene (Zachos *et al.*, 2001; Morley, 2003). The plateau of net diversification rate in *Diplazium* suggests that the clade is closer to ecological saturation than *Athyrium*, in which the upward trend in rate suggests a clade in the early stages of rapid growth, further from equilibrium (Fig. 2b). These dynamics are consistent with the idea that species diversity is predicted by area, and other proxies for carrying capacity, integrated over time (Fine & Ree, 2006; Jetz & Fine, 2012). In addition, the relatively high turnover rate μ/λ in *Athyrium* (Fig. S5) may be due to more severe climatic oscillations in the temperate zone than in the comparatively stable tropics over the past 20 Myr or so (Tiffney & Manchester, 2001; Morley, 2003), consistent with the

latitudinal gradient in turnover found in previous studies (Jablonski *et al.*, 2006; Weir & Schluter, 2007; but see Rabosky *et al.*, 2015). This high rate of species turnover, in turn, suggests a rapid species replacement through non-adaptive evolution in *Athyrium* (Rundell & Price, 2009).

Additional support for the idea that diversification dynamics in *Diplazium* are closer to an equilibrium state than *Athyrium* is provided by our QuaSSE analysis of elevation. In *Diplazium* the inferred response of speciation rate to elevation closely matches the frequency distribution of species along elevation (also is reflected by a low negative directional tendency; Table 1), suggesting a stable state, while in *Athyrium* the rate curve is right-shifted: speciation is fastest at elevations higher than the peak of species richness (Fig. 3). The relative lack of high-elevation species of *Athyrium* could be explained partly by the inferred negative directional tendency in the model – i.e. species originated at high elevations but evolved anagenetically toward lower elevations – but it seems more plausible that *Athyrium* has simply colonized mountains too recently for the number of high-elevation species to reach equilibrium. We would expect such a lag time to be lengthened by the relatively high extinction rate inferred for *Athyrium* versus *Diplazium* (Table 1).

Our analyses of elevation stand in contrast to those in the epiphytic fern family Polypodiaceae, in which faster diversification was positively correlated with the rate of change in elevation, not elevation itself, suggesting a process of adaptive divergence and niche-filling along elevation gradients (Sundue *et al.*, 2015). We were unable to replicate this result from our data, possibly because of insufficiently dense taxon sampling. Nevertheless, our results suggest that diversification is accelerated by occurring in mountains, especially for *Athyrium*. However, it is worth noting that even within *Diplazium*, the highest net diversification was inferred in the lone Andean clade (Fig. 2a). Left open are questions about how and why, such as to what degree is speciation accelerated by non-adaptive processes (such as allopatric genetic drift) versus those involving adaptation? More focused studies of possible mechanisms are needed. One might look for interactions between traits with environmental tolerances and mountain buildings in tropical and temperate zones (Xing & Ree, 2017; Hughes & Atchison, 2015). This has been shown in studies of other plant groups have found that the evolution of traits such as growth form, leaf size, pollination syndromes, and metabolic pathways may be linked to faster diversification in mountains where

397 seasonally cold and arid habitats are dominant (Arakaki *et al.*, 2011; Drummond *et al.*,
398 2012; Roquet *et al.*, 2013; Schwery *et al.*, 2015; Lagomarsino *et al.*, 2016).

399

400 **Polyploidy-driven diversification in the *Athyrium-Diplazium* clade**

401 In ferns, several traits, including growth habit, leaf size, leaf morphology and
402 gametophyte morphology, have been studied for their effect on diversification
403 (Sundue *et al.*, 2015; Ramírez-Barahona *et al.*, 2016; Testo & Sundue, 2018), but
404 polyploidy, despite its ubiquity and precedent in being implicated in diversification
405 (Wood *et al.*, 2009), has thus far escaped intensive quantitative analysis. While the
406 macroevolutionary consequences of polyploidy remain a subject of debate (e.g.
407 Mayrose *et al.*, 2011, 2015; Zhan *et al.*, 2014; Soltis *et al.*, 2014), our inference of
408 rapid turnover with negative net diversification in polyploids (Fig. 3d) is consistent
409 with the ‘extinction-risk’ hypothesis, that neopolyploids suffer deleterious genetic
410 effects (Wright, 1969; Orr, 1990; Ramsey & Schemske, 2002), and thus are often
411 evolutionary dead ends (Mayrose *et al.*, 2011; Arrigo & Baker, 2012). This effect is
412 counterbalanced by the high net diversification rate of polymorphic species (Fig. 4d),
413 which rapidly produce polyploids through cladogenesis events and thereby contribute
414 to polyploids having the highest net accumulation rate (Fig. 4e). Rapid and recurrent
415 neopolyploidy in *Athyrium* and *Diplazium* (Fig. S4) is distinct from *Asplenium*, which
416 is characterized by a dominance of paleopolyploids (Schneider *et al.*, 2017). Taken
417 together, these results imply that although neopolyploids in *Athyrium* and *Diplazium*
418 tend to be short-lived, they play a key role in diversification dynamics (Vamosi *et al.*,
419 2018).

420 Intraspecific variation in ploidy is common in plants, and many case studies have
421 examined the origins of polyploidy at the population level, especially in genera of
422 Asteraceae such as *Tragopogon* (Symonds *et al.*, 2010), *Artemisia* (Pellicer *et al.*,
423 2010), *Helianthus* (Bock *et al.*, 2014), as well as in fern genera *Asplenium* (e.g. Dyer
424 *et al.*, 2012) and *Dryopteris* (e.g. Ekrt & Koutecký, 2016), and *Pteris* (Chao *et al.*,
425 2012). This study is the first to explicitly model how such polymorphic species may
426 contribute to diversification dynamics at a macroevolutionary scale, and so it seems
427 noteworthy to find that despite occurring at low frequency, their net diversification
428 rates are in fact higher than pure diploids or polyploids. This is largely driven by λ_{113} ,
429 the rate at which a polymorphic ancestor splits into polymorphic and polyploid

430 descendants (Fig 4a). Mechanistically, it seems reasonable to predict that this
 431 parameter reflects how newly formed polyploids in a population are likely to be
 432 reproductively isolated from their diploid progenitors, and thus form a new lineage.
 433 Intraspecific polyploidy can predispose species to be better adapted to harsher
 434 conditions in novel environments (Levin, 1983; Ramsey & Schemske, 2002; te Beest
 435 *et al.*, 2012). These results could be framed as a ‘ploidy-polymorphism compensation’
 436 hypothesis, wherein polymorphic species play a central role in the maintenance of
 437 neopolyploid diversity. The *Athyrium-Diplazium* clade is not likely to be unique in
 438 supporting this hypothesis; we predict that other lineages such as *Asplenium*
 439 (*Aspleniaceae*) and *Dryopteris* (*Dryopteridaceae*) and *Pteris* (*Pteridaceae*), in which
 440 autopolyploidy, hybridization and apomixis are prevalent (Chao *et al.*, 2012; H.M.
 441 Liu *et al.*, 2012), may have similar compensatory dynamics mitigating the high
 442 extinction risk of polyploids.

443 Why might polyploids have high speciation rates (Fig. 4a)? It is possible that some
 444 factors (a novel physical trait or a property of the environment) associated with
 445 polyploidization may play a role (Zhan *et al.*, 2016; Vamosi *et al.*, 2018). A few case
 446 studies have linked polyploidization to trait evolution and diversification (Schrantz *et*
 447 *al.*, 2011; Tank *et al.*, 2015). In ferns, polyploids may facilitate long-distance
 448 dispersal via tolerance of inbreeding, i.e. gametophytic selfing, increasing the success
 449 rate of single-spore colonization (e.g. Tryon, 1985; Testo *et al.*, 2015; Sessa *et al.*,
 450 2016). This in turn might favor colonization of different ecological niches, and thus
 451 increase the probability for peripatric speciation. In *Deparia*, the species-rich genus in
 452 *Athyriaceae* that is sister to *Athyrium-Diplazium*, polyploid species are characterized
 453 by increased dispersal abilities and greater range expansion than sexual diploids (Kuo
 454 *et al.*, 2016). This seems to be true as well for *Athyrium* and *Diplazium*, in which
 455 polyploid species are often found more broadly distributed than their diploid relatives
 456 (Tryon & Tryon, 1982; Takamiya *et al.*, 1999, 2000; Takamiya & Ohta, 2001; Bir &
 457 Verma, 2010). It thus seems necessary for future studies to target other traits
 458 potentially linked polyploidy that could enhance dispersal ability.

459

460 **Acknowledgements**

461 We thank Qin Li, Matthew Nelsen, Shrabya Timsina, and Charles Bell for helpful
 462 discussions and Jen-Pan Huang for technical assistance and advice about ClaSSE. We

thank Harald Schneider, Hong-Mei Liu, Yue-Hong Yan, Hui Shang, Atsushi Ebihara, Michael Kessler, Marcus Lehnert, Peter Hovenkamp, Ronald Viane, Sabine Hennequin and Claudine Mynssen for providing valuable materials. This study was supported by grants from the National Science Foundation of China Grant (No. 31600175), the Beijing Natural Science Foundation Grant (No. 5162020) and the Visiting Scholar of Bureau of Personnel Chinese Academy of Sciences to the Field Museum awarded to R.W.

470

471 **Author contributions**

472 X.-C.Z. and R.W. conceived the idea of this study; R.W. performed the experiments;
473 M.A.S. provided important Neotropical materials; R.W. and R.H.R. analyzed data;
474 and R.W., R.H.R., and M.A.S. wrote the paper, with significant contribution from X.-
475 C.Z.

476

477 **References**

- 478 Arakaki M, Christin PA, Nyffeler R, Lendel A, Eggli U, Ogburn RM, Spriggs E,
479 Moore MJ, Edwards EJ. 2011. Contemporaneous and recent radiations of the
480 world's major succulent plant lineages. *Proceedings of the National Academy of*
481 *Sciences, USA* **108**: 8379–8384.
- 482 Arrigo N, Barker MS. 2012. Rarely successful polyploids and their legacy in plant
483 genomes. *Current Opinion in Plant Biology* **16**: 140–146.
- 484 Bir SS, Verma SC. 2010. *Chromosome atlas of the Indian pteridophytes*. Dehra Dun,
485 India: Gajendra Singh Gahlot.
- 486 Bock DG, Kane NC, Ebert DP, Rieseberg LH. 2014. Genome skimming reveals the
487 origin of the Jerusalem Artichoke tuber crop species: neither from Jerusalem nor an
488 artichoke. *New Phytologist* **201**: 1021–1030.
- 489 Ching R-C. 1964. On some confused genera of the family Athyriaceae. *Acta*
490 *Phytotaxonomica Sinica* **9**: 41–84.
- 491 Chao Y-S, Liu H-Y, Chiang Y-C, Chiou W-L. 2012. Polyploidy and speciation in
492 *Pteris* (Pteridaceae). *Journal of Botany* **2012**: 817920.
- 493 Collinson ME. 2001. Cainozoic ferns and their distribution. *Brittonia* **53**: 172–235.
- 494 Drummond AJ, Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by
495 sampling trees. *BMC Evolutionary Biology* **7**: 214.

496 Drummond CS, Eastwood RJ, Miotto STS, Hughes CE. 2012. Multiple continental
497 radiations and correlates of diversification in *Lupinus* (Leguminosae): testing for
498 key innovation with incomplete taxon sampling. *Systematic Biology* **61**: 443–460.
499 Dyer RJ, Savolainen V, Schneider H. 2012. Apomixis and reticulate evolution in the
500 *Asplenium monanthes* fern complex. *Annals of Botany* **110**: 1515–1529.
501 Ekrt L, Koutecký P. 2016. Between sexual and apomictic: unexpectedly variable
502 sporogenesis and production of viable polyhaploids in the pentaploid fern of the
503 *Dryopteris affinis* agg. (Dryopteridaceae). *Annals of Botany* **117**: 97–106.
504 Fine PVA, Ree RH. 2006. Evidence for a time-integrated species-area effect on the
505 latitudinal gradient in tree diversity. *The American Naturalist* **168**: 796–804.
506 FitzJohn RG. 2012. Diversitree: comparative phylogenetic analyses of diversification
507 in R. *Methods in Ecology and Evolution* **3**: 1084–1092.
508 Glick L, Mayrose I. 2014. ChromEvol: assessing the pattern of chromosome number
509 evolution and the inference of polyploidy along a phylogeny. *Molecular Biology*
510 *and Evolution* **31**: 1914–1922.
511 Goldberg EE, Igić B. 2012. Tempo and mode in plant breeding system evolution.
512 *Evolution* **66**: 3701–3709.
513 Harmon LJ, Weir JT, Brock CD, Glor RE, Challenger W. 2008. GEIGER:
514 investigating evolutionary radiations. *Bioinformatics* **24**: 129–131.
515 Hsieh Y-T. 1986. The classification of *Athyrium* Roth. *Bulletin of Botanical Research*
516 **6**, 129–135.
517 Hughes CE, Atchison GW. 2015. The ubiquity of alpine plant radiations: from the
518 Andes to the Hengduan Mountains. *New Phytologist* **207**: 275–282.
519 Jablonski D, Roy K, Valentine JW. 2006. Out of the tropics: evolutionary dynamics
520 of the latitudinal diversity gradient. *Science* **314**: 102–106.
521 Jetz W, Fine PVA. 2012. Global gradients in vertebrate diversity predicted by
522 historical area-productivity dynamics and contemporary environment. *PLoS*
523 *Biology* **10**: e1001292.
524 Karafit SJ, Rothwell GW, Stockey RA, Nishida H. 2006. Evidence for sympodial
525 vascular architecture in a filiclean fern rhizome: *Dickwhitea allenbyensis* gen. et
526 sp. nov. (Athyriaceae). *International Journal of Plant Sciences* **167**: 721–727.
527 Kato M. 1977. Classification of *Athyrium* and allied genera of Japan. *Botanical*
528 *Magazine (Tokyo)* **90**: 23–40.

529 Kuo L-Y, Ebihara A, Shinohara W, Rouhan G, Wood KR, Wang C-N, Chiou W-L.
530 2016. Historical biogeography of the fern genus *Deparia* (Athuriaceae) and its
531 relation with polyploidy. *Molecular Phylogenetics and Evolution* **104**: 123–134.

532 Lagomarsino LP, Condamine FL, Antonelli A, Mulch A, Davis CC. 2016. The abiotic
533 and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae).
534 *New Phytologist* **210**: 1430–1442.

535 Lehtonen S, Silvestro D, Karger DN, Scotese C, Tuomisto H, Kessler M, Peña C,
536 Wahlberg N, Antonelli A. 2017. Environmentally driven extinction and
537 opportunistic origination explain fern diversification patterns. *Scientific Reports* **7**:
538 4831.

539 Levin DA. 1983. Polyploidy and novelty in flowering plants. *The American*
540 *Naturalist* **122**: 1–25.

541 Liu H-M, Dyer RJ, Guo Z-Y, Meng Z, Li J-H, Schneider H. 2012. The evolutionary
542 dynamics of apomixis in ferns: a case study from polystichoid ferns. *Journal of*
543 *Botany* **2012**: 510478.

544 Liu Y-C, Chiou W-L, Kato M. 2011. Molecular phylogeny and taxonomy of the fern
545 genus *Anisocampium* (Athuriaceae). *Taxon* **60**: 824–830.

546 Mayrose I, Zhan S-H, Rothfels CJ, Magnuson-Ford KS, Barker MS, Rieseberg LH,
547 Otto SP. 2011. Recently formed polyploid plants diversify at lower rates. *Science*
548 **333**: 1257.

549 Mayrose I, Zhan S-H, Rothfels CJ, Arrigo N, Barker MS, Rieseberg LH, Otto SP.
550 2015. Methods for studying polyploid diversification and the dead end hypothesis:
551 a reply to Soltis *et al.* (2014). *New Phytologist* **206**: 27–35.

552 Mehra PN, Bir SS. 1960. Cytological observations on the Himalayan species of
553 *Athyrium* and comments on the evolutionary status of the genus. *American Fern*
554 *Journal* **50**: 276–295.

555 Mickel J, Smith AR. 2004. *Diplazium* Sw. In: Buck WR, ed. *Pteridophytes of Mexico*,
556 *part 1*. New York, NY, USA: New York Botanical Garden Press, 251–268.

557 Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for
558 inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing*
559 *Environments Workshop (GCE), 14 November 2010*. New Orleans, LA, USA, 1–8.

560 Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB,
561 Harrison SP, Hurlbert AH, Knowlton N, Lessios HA *et al.* 2007. Evolution and the

latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters* **10**: 315–331.

Morley RJ. 2003. Interplate dispersal paths for megathermal angiosperms. *Perspectives in Plant Ecology, Evolution and Systematics* **6**: 5–20.

Orr HA. 1990. ‘Why polyploidy is rarer in animals than in plants’ revisited. *The American Naturalist* **6**: 759–770.

Pellicer J, Garcia S, Canela MA, Garnatje T, Korobkov AA, Twibell JD, Vallès J. 2010. Genome size dynamics in *Artemisia* L. (Asteraceae): following the track of polyploidy. *Plant Biology* **12**: 820–830.

Perrie LR, Shepherd LD, De Lange PJ, Brownsey PJ. 2010. Parallel polyploid speciation: distinct sympatric gene-pools of recurrently derived allo-octoploid *Asplenium* ferns. *Molecular Ecology* **19**: 2916–2932.

PPG I. 2016. A community-derived classification for extant lycophytes and ferns. *Journal of Systematics and Evolution* **54**: 563–603.

Praptosuwiryo TN. 2008. *Biosystematic study of the fern genus Diplazium in west Malesia*. PhD thesis, Bogor Agricultural University, Bogor, Indonesia.

Rabosky DL, Adams DC. 2012. Rates of morphological evolution are correlated with species richness in salamanders. *Evolution* **66**: 1807–1818.

Rabosky DL. 2014. Automatic detection of key innovations, rate shifts, and diversity-dependence on phylogenetic trees. *PLoS ONE* **9**: e89543.

Rabosky DL, Goldberg EE. 2015. Model inadequacy and mistaken inferences of trait-dependent speciation. *Systematic Biology* **64**: 340–355.

Rabosky DL, Title PO, Huang H. 2015. Minimal effects of latitude on present-day speciation rates in New World birds. *Proceedings of the Royal Society B: Biological Sciences* **282**: 20142889.

Ramsey J, Schemske WD. 2002. Neopolyploidy in flowering plants. *Annual Review of Ecology and Systematics* **33**: 589–639.

Ramírez-Barahona S, Barrera-Redondo J, Eguiarte LE. 2016. Rates of ecological divergence and body size evolution are correlated with species diversification in scaly tree ferns. *Proceedings of the Royal Society B: Biological Sciences* **283**: 20161098.

Roquet C, Boucher FC, Thuiller W, Lavergne S. 2013. Replicated radiations of the alpine genus *Androsace* (Primulaceae) driven by range expansion and convergent

key innovations. *Journal of Biogeography* **40**: 1874–1886.

Rothfels CJ, Li F-W, Sigel EM, Huiet L, Larsson A, Burge DO, Ruhsam M, Deyholos M, Soltis DE, Stewart CN *et al.* 2015. The evolutionary history of ferns inferred from 25 low-copy nuclear genes. *American Journal of Botany* **102**: 1–19.

Rothwell GW, Stockey RA. 1991. *Onoclea sensibilis* in the Paleocene of North America, a dramatic example of structural and ecological stasis. *Review of Palaeobotany and Palynology* **70**: 113–124.

Schluter D. 2015. Speciation, ecological opportunity, and latitude. *The American Naturalist* **187**: 1–18.

Schneider H, Schuettzel E, Pryer KM, Cranfill R, Magallón S, Lupia R. 2004. Ferns diversified in the shadow of angiosperms. *Nature* **428**: 553–557.

Schneider H, Liu H-M, Chang Y-F, Ohlsen D, Perrie LR, Shepherd L, Kessler M, Karger DN, Hennequin S, Marquardt J *et al.* 2017. Neo- and Paleopolyploidy contribute to the species diversity of *Asplenium* – the most species-rich genus of ferns. *Journal of Systematics and Evolution* **55**: 353–364.

Schranz ME, Mohammadin S, Edger PP. 2012. Ancient whole genome duplications, novelty and diversification: the WGD Radiation Lag-Time Model. *Current Opinion in Plant Biology* **15**: 147–153.

Schuettzel E, Pryer KM. 2009. Evidence for a Cenozoic radiation of ferns in an angiosperm-dominated canopy. *Proceedings of the National Academy of Sciences, USA* **106**: 11200–11205.

Schwery O, Onstein RE, Bouchenak-Khelladi Y, Xing Y-W, Carter RJ, Linder HP. 2015. As old as the mountains: the radiations of the Ericaceae. *New Phytologist* **207**: 355–367.

Sessa EB, Testo WL, Watkins JE Jr. 2016. On the widespread capacity for, and functional significance of, extreme inbreeding in ferns. *New Phytologist* **211**: 1108–1119.

Soltis DE, Segovia-Salcedo MC, Jordon-Thaden I, Majure L, Miles NM, Mavrodiev EV, Mei W-B, Cortez MB, Soltis PS, *et al.* 2014. Are polyploids really evolutionary dead-ends (again)? A critical reappraisal of Mayrose *et al.* (2011). *New Phytologist* **202**: 1105–1117.

Soltis DE, Visger CJ, Marchant DB, Soltis PS. 2016. Polyploidy: pitfalls and paths to a paradigm. *American Journal of Botany* **103**: 1223–1235.

- 628 Spriggs EL, Clement WL, Sweeney PW, Madriñán S, Edwards EJ, Donoghue MJ.
629 2015. Temperate radiations and dying embers of a tropical past: the diversification
630 of *Viburnum*. *New Phytologist* **207**: 340–354.
- 631 Stockey RA, Nishida H, Rothwell GW. 1999. Permineralized ferns from the Middle
632 Eocene Princeton chert. I. *Makotopteris princetonensis* gen. et sp. nov.
633 (Athyriaceae). *International Journal of Plant Sciences* **160**: 1047–1055.
- 634 Stolze RG, Pacheco L, Øllgaard B. 1994. Physematieae. In: Harling G, Andersson L,
635 eds, *Flora of Ecuador, vol. 49*. Gothenburg, Sweden: University of Gothenburg, 1–
636 108.
- 637 Sundue MA, Testo W, Ranker TA. 2015. Morphological innovation, ecological
638 opportunity, and the radiation of a major vascular epiphyte lineage. *Evolution* **69**:
639 2482–2495.
- 640 Symonds VV, Soltis PS, Soltis DE. 2010. Dynamics of polyploid formation in
641 *Tragopogon* (Asteraceae): recurrent formation, gene flow, and population structure.
642 *Evolution* **64**: 1984–2003.
- 643 Takamiya M, Takaoka C, Ohta N. 1999. Cytological and reproductive studies on
644 Japanese *Diplazium* (Woodsiaceae; Pteridophyta): apomictic reproduction in
645 *Diplazium* with evergreen bi- to tripinnate leaves. *Journal of Plant Research* **112**:
646 419–436.
- 647 Takamiya M, Ohta N, Fujimaru-Takaoka C, Uki K. 2000. Cytological and
648 reproductive studies of Japanese *Diplazium* (Woodsiaceae; Pteridophyta). II.
649 Polyploidy and hybridity in the species group with summer-green bi- to tripinnate
650 leaves. *Journal of Plant Research* **113**: 203–215.
- 651 Takamiya M, Ohta N. 2001. Cytological and reproductive studies of Japanese
652 *Diplazium* (Woodsiaceae; Pteridophyta). III. The cytological complexity of species
653 groups with simply pinnate to bipinnatifid leaves. *Journal of Plant Research* **114**:
654 443–457.
- 655 Tank DC, Eastman JM, Pennell MW, Soltis PS, Soltis DE, Hinchliff CE, Brown JW,
656 Sessa EB, Harmon LJ. 2015. Nested radiations and the pulse of angiosperm
657 diversification: increased diversification rates often follow whole genome
658 duplications. *New Phytologist* **207**: 454–467.
- 659 te Beest M, Roux JJJ, Richardso DM, Brysting AK, Suda J, Kubešová M, Pyšek P.
660 2012. The more the better? The role of polyploidy in facilitating plant invasions.

- 661 *Annals of Botany* **109**: 19–45.
- 662 Testo WL, Watkins JE Jr, Barrington DS. 2015. Dynamics of asymmetrical
663 hybridization in North American wood ferns: reconciling patterns of inheritance
664 with gametophyte reproductive biology. *New Phytologist* **206**: 785–795.
- 665 Testo W, Sundue MA. 2016. A 4000-species dataset provides new insight into the
666 evolution of ferns. *Molecular Phylogenetics and Evolution* **105**: 200–211.
- 667 Testo W, Sundue MA. 2018. Are rates of species diversification and body size
668 evolution coupled in the ferns? *American Journal of Botany* **105**: 1–11.
- 669 Tiffney BH, Manchester SR. 2001. The use of geological and paleontological
670 evidence in evaluating plant phylogeographic hypotheses in the northern
671 hemisphere tertiary. *International Journal of Plant Sciences* **162**: S3–S17.
- 672 Tryon RM. 1985. Fern speciation and biogeography. *Proceedings of the Royal Society*
673 *of Edinburgh. Section B. Biological Sciences* **86**: 353–360.
- 674 Tryon RM, Stolze RG. 1991. *Pteridophytes of Peru, part 4*. Chicago, IL, USA: Field
675 Museum of Natural History.
- 676 Tryon RM, Tryon AF. 1982. *Ferns and allied plants with special reference to the*
677 *tropical America*. New York, NY, USA: Springer-Verlag.
- 678 Tryon RM. 1985. Fern speciation and biogeography. *Proceedings of the Royal Society*
679 *of Edinburgh. Section B. Biological Sciences* **86**: 353–360.
- 680 Vamosi JC, Magallón S, Mayrose I, Otto SP, Sauquet H. 2018. Macroevo-lutionary
681 patterns of flowering plant speciation and extinction. *Annual Review of Plant*
682 *Biology* **69**: 685–706.
- 683 Walker TG. 1966. A cytotaxonomic survey of the pteridophytes of Jamaica.
684 *Transactions of the Royal Society of Edinburgh* **66**: 169–237.
- 685 Walker TG. 1973. Additional cytotaxonomic notes on the pteridophytes of Jamaica.
686 *Transactions of the Royal Society of Edinburgh* **69**: 109–135.
- 687 Wang Z-R. 1997. A revision of the Chinese *Athyrium* Roth (I) – subgeneric
688 classification of the genus. *Bulletin of Botanical Research* **17**: 274–300.
- 689 Wang Z-R, He Z-R, Kato M. 2013. Athyriaceae. In: Wu Z, Raven PH, Hong D, eds.
690 *Flora of China, vol. 2–3*. Beijing: Science Press; St. Louis: Missouri Botanical
691 Garden Press, 418–534.
- 692 Wei R, Schneider H, Zhang X-C. 2013. Toward a new circumscription of the
693 twinsorus-fern genus *Diplazium* (Athyriaceae): a molecular phylogeny with

694 morphological implications and infrageneric taxonomy. *Taxon* **62**: 441–457.

695 Wei R, Xiang Q-P, Schneider H, Sundue MA, Kessler M, Kamau PW, Hidayat A,
696 Zhang X-C. 2015. Eurasian origin, boreotropical migration and transoceanic
697 dispersal in the pantropical fern genus *Diplazium* (Athuriaceae). *Journal of*
698 *Biogeography* **42**: 1809–1819.

699 Wei R, Ebihara A, Zhu Y-M, Zhao C-F, Hennequin S, Zhang X-C. 2018. A total-
700 evidence phylogeny of the lady fern genus *Athyrium* Roth (Athuriaceae) with a new
701 infrageneric classification. *Molecular Phylogenetics and Evolution* **119**: 25–36.

702 Weir JT, Schluter D. 2007. The latitudinal gradient in recent speciation and extinction
703 rates of birds and mammals. *Science* **135**: 1574–1576.

704 Wiens JJ, Donoghue MJ. 2004. Historical biogeography, ecology and species richness.
705 *Trends in Ecology and Evolution* **19**: 639–644.

706 Wood TE, Takebayashi N, Barker MS, Mayrose I, Greenspoon PB, Rieseberg LH.
707 2009. The frequency of polyploid speciation in vascular plants. *Proceedings of the*
708 *National Academy of Sciences, USA* **106**: 13875–13879.

709 Wright S. 1969. *Evolution and Genetics of Populations. II. The Theory of Gene*
710 *Frequencies*. Chicago: University of Chicago Press.

711 Xing Y-W, Ree RH. 2017. Uplift-driven diversification in the Hengduan Mountains, a
712 temperate biodiversity hotspot. *Proceedings of the National Academy of Sciences,*
713 *USA* **114**: 3444–3451.

714 Yu Y, Harris AJ, Blair C, He X-J. 2015. RASP (Reconstruct Ancestral State in
715 Phylogenies): A tool for historical biogeography. *Molecular Phylogenetics and*
716 *Evolution* **87**: 46–49.

717 Zachos J, Pagani M, Sloan L, Thomas E, Billups, K. 2001. Trends, rhythms, and
718 aberrations in global climate 65 Ma to present. *Science* **292**: 686–693.

719 Zhan S-H, Glick L, Tsigenopoulos CS, Otto SP, Mayrose I. 2014. Comparative
720 analysis reveals that polyploidy does not decelerate diversification in fish. *Journal*
721 *of Evolutionary Biology* **27**: 391–403.

722 Zhan S-H, Drori M, Goldberg EE, Otto SP, Mayrose I. 2016. Phylogenetic evidence
723 for cladogenetic polyploidization in land plants. *American Journal of Botany* **103**:
724 1252–1258.

725 Zhang X-C. 1992. Taxonomy of *Athyrium* Roth subgen. *Athyrium* in Yunnan. *Acta*
726 *Phytotaxonomica Sinica* **30**: 245–255.

727

728 **Supporting Information**

729 Additional Supporting Information may be found online in the Supporting
730 Information tab for this article:

731

732 **Fig. S1** Chronogram of the *Athyrium-Diplazium* clade with four calibration points.

733 **Fig. S2** Positions of regime shifts in the maximum a posteriori configuration inferred
734 from the Bayesian analysis of macro-evolutionary mixtures (BAMM).

735 **Fig. S3** Rate simulations and their estimates of confidence intervals for each of the
736 states under 100 replicates of simulated trees and random assignment of 'NA' to tip
737 states in ClaSSE.

738 **Fig. S4** CHROMEOL inferences for the *Athyrium-Diplazium* clade.

739 **Fig. S5** Rate-through-time plot of rates of speciation, extinction and species turnover
740 (termed ' μ/λ ') obtained from BAMM.

741 **Table S1** Species sampling fraction used in macroevolution analyses.

742 **Table S2** Taxa, voucher information and GenBank accession numbers of specimens
743 used in this study.

744 **Table S3** Traits information and coding in QuaSSE, CHROMEOL and ClaSSE.

745 **Table S4** Model selection and results of QuaSSE on the entire ingroup clade as well
746 as separately on *Athyrium* and *Diplazium*.

747 **Table S5** Model selection and results of ClaSSE based on the entire ingroup clade as
748 well as separately on *Athyrium* and *Diplazium*.

749 **Table S6** Bayes factor of different diversification rate models fitted on the *Athyrium-*
750 *Diplazium* dataset.

751 **Table S7** ClaSSE model simulation based on 100 simulated trees with random
752 assignment of missing information to tips.

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

Table 1 Parameter comparison of the best-fit model of quantitative state speciation and extinction (QuaSSE) analysis based on the entire ingroup clade as well as separately on *Athyrium* and *Diplazium*. λ_0 , rate at lowest values of the log elevation for sigmoid and unimodal models; λ_1 , maximum speciation rate for the unimodal models; x_{mid} , inflection point of the sigmoid or the place of the maximum for modal models; s^2 , width (variance) of the Gaussian function; μ , rate of extinction; σ^2 , Brownian diffusion rate of trait evolution.

	<i>Athyrium</i> clade	<i>Diplazium</i> clade	The entire ingroup clade
Rates and Parameters	Unimodal with directional tendency	Unimodal with no tendency	Unimodal with directional tendency
λ_0	1.00e-04	0.0322	0.0303
λ_1	0.8484	0.4197	0.5643
μ	0.5375	0.1617	0.3315
x_{mid}	7.83	6.965	7.712
s^2	0.1954	0.1066	0.4128
σ^2	0.0482	0.0524	0.0481
Directional	-0.091	-0.0013	-0.086

Table 2 Parameter comparisons of the best-fit model in cladogenetic state change speciation and extinction (ClaSSE) analysis based on the entire ingroup clade as well as separately on *Athyrium* and *Diplazium*. Both containing diploidy and polyploidy is coded as 1; diploidy is coded as 2; polyploidy is coded as 3. λ , speciation rate; μ , rate of extinction; q , transition rate between two states. All rates are presented with median values and 95% highest posterior density (HPD) intervals in brackets based on 100 000 MCMC optimization.

	<i>Athyrium</i> clade	<i>Diplazium</i> clade	The entire ingroup clade
Rates	Best-fit model: Cladogenetic speciation and extinction ($\lambda_{111} \neq \lambda_{222} \neq \lambda_{333} \neq \lambda_{112} \neq \lambda_{113} \neq \lambda_{123} \neq \lambda_{223}$, $\mu_1 \neq \mu_2 \neq \mu_3$, $q_{12} = q_{21} = q_{13} = q_{23}$)		
λ_{111}	0.5221 [0.2089–1.3218]	0.2128 [0.0951–0.4404]	0.334 [0.1902–0.6158]
λ_{222}	0.1758 [0.0822–0.3604]	0.0898 [0.0467–0.1573]	0.1326 [0.0741–0.2332]
λ_{333}	0.9462 [0.1513–2.727]	0.7056 [0.2735–1.279]	0.7054 [0.2575–1.299]
λ_{112}	0.1758 [0.01–0.6164]	0.0864 [0.0045–0.3042]	0.0836 [0.0061–0.229]
λ_{113}	1.6779 [0.261–4.305]	0.9392 [0.3241–2.023]	0.8974 [0.4328–1.7195]
λ_{123}	0.173 [0.008–0.6014]	0.063 [0.0032–0.2097]	0.073 [0.0038–0.2145]
λ_{223}	0.3929 [0.032–1.485]	0.0707 [0.0028–0.4607]	0.0667 [0.0029–0.3541]
μ_1	0.3658 [0.023–1.263]	0.0672 [0.0027–0.3211]	0.1658 [0.0107–0.49]
μ_2	0.2075 [0.022–0.4686]	0.0258 [0.001–0.1228]	0.0809 [0.0054–0.2366]
μ_3	1.427 [0.5903–3.247]	0.7513 [0.3637–1.336]	0.829 [0.4372–1.432]
q	0.0421 [0.004–0.1926]	0.0276 [0.0076–0.0704]	0.0255 [0.0081–0.0614]

796 **Figure Legends:**

797 **Fig. 1** The cladogenetic-state change speciation and extinction (ClaSSE) models with
 798 cladogenetic and anagenetic changes. (a) The full model including cladogenetic
 799 diversification and anagenetic state change is depicted. In the diagram, lineages
 800 presented by both diploid and polyploid populations have state 1, and those with only
 801 diploids have state 2, and those with only polyploids have state 3. Each speciation
 802 event gives rise to two daughters (as shown by arrows), either in the same state (at
 803 rates λ_{111} , λ_{222} , and λ_{333}) or in different states (λ_{112} , λ_{113} , λ_{123} , and λ_{223}). Thus, changes
 804 in ploidy can occur through the anagenetic (q_{12} , q_{13} , q_{21} , and q_{23}) or cladogenetic
 805 pathway (λ_{112} , λ_{113} , λ_{123} , and λ_{223}). Extinction rates in each state are represented by μ_1 ,
 806 μ_2 and μ_3 . The rates in gray indicate equal rates, respectively. (b) The null model with
 807 equal speciation, extinction and transition rates, respectively. (c) The ‘cladogenetic’
 808 model. (d) The ‘anagenetic’ model. (e) The ‘Extinction’ model. (f) The ‘cladogenetic-
 809 anagenetic’ model.

810
 811 **Fig. 2** Results of the Bayesian analysis of macro-evolutionary mixtures (BAMM) and
 812 quantitative-state speciation and extinction (QuaSSE) analyses. (a) Chronorate plot
 813 with branches colored by speciation rate (events Myr^{-1} per lineage) as indicated by
 814 the scale bar, representing a summary of the full post-burn-in Markov chain Monte
 815 Carlo (MCMC) sample of the BAMM analysis. Red circles indicate the positions of
 816 regime shifts in the maximum *a posteriori* (MAP) configuration. Numbers beneath
 817 the shifts indicate the marginal probability of a shift occurring along that branch. Log
 818 elevation is shown by the horizontal bar for each species. The vertical dashed lines
 819 indicate the approximate ranges of altitudes in which elevated speciation rates were
 820 inferred by QuaSSE analysis, and extant species whose elevation falls in this range
 821 have their data colored red. (b) Rate-through-time plots for speciation rate (events
 822 Myr^{-1} per lineage) with 95% confidence interval indicated by yellow shaded areas.
 823 Red, the rate across the phylogeny (‘background rate’); blue, the rate of *Athyrium*
 824 clade; green, the rate of *Diplazium* clade.

825
 826 **Fig. 3** Relationship between the elevation and speciation rates as inferred from
 827 QuaSSE analysis, with comparison to the elevational species richness histograms of
 828 the *Athyrium* clade, the *Diplazium* clade and the whole phylogeny, respectively. Red,

829 the rate across the phylogeny; blue, the rate of *Athyrium* clade; green, the rate of
830 *Diplazium* clade.

831

832 **Fig. 4** Posterior distributions of macroevolutionary rates under the best-fitting model
833 of the ClaSSE analysis based on the MCC tree, allowing cladogenetic speciation and
834 extinction of diploids and polyploids. (a) The cladogenetic speciation rates of diploid-
835 polyploid mediated speciation events. (b) The extinction rates of three ploidy states. (c)
836 The transition rate of all anagenetic change (equal). (d) Summary of speciation rates
837 of each state. (e) The net diversification rates of ploidy in cladogenetic speciation. (f)
838 The net accumulation rates of ploidy in cladogenetic speciation.

839

(a)

Full model

No. of parameters: 14

(b)

Null model

No. of parameters: 3

(c)

'Cladogenetic' model

No. of parameters: 11

(d)

'Anagenetic' model

No. of parameters: 6

(e)

'Extinction' model

No. of parameters: 5

(f)

'Cladogenetic-anagenetic' model

No. of parameters: 12

(a)

Mackinnonia
clade

Core *Athyrium*

Athyrium

Andean *Diplazium*
clade

Subgenus *Callipteris*

Diplazium

(a)

(b)

(a)

(b)

(c)

(d)

(e)

(f)