Inositol 1,4,5-trisphosphate receptor type2-independent Ca^{2+} release from the endoplasmic reticulum in astrocytes

Abbreviated title: $\text{IP}_3\text{R2}$ -independent Ca^{2+} release in astrocytes

Yohei Okubo^{1*}, Kazunori Kanemaru^{1,2}, Junji Suzuki³, Kenta Kobayashi^{4,5}, Kenzo Hirose⁶ & Masamitsu Iino^{2*}

¹Department of Pharmacology, Graduate School of Medicine, The University of Tokyo, Tokyo 133-0033, Japan

²Department of Cellular and Molecular Pharmacology, Nihon University School of Medicine, Tokyo 173-8610, Japan.

³Department of Physiology, University of California San Francisco, San Francisco, CA 94158, USA

⁴Section of Viral Vector Development, National Institute for Physiological Sciences, Okazaki 444-8585, Japan.

⁵The Graduate University for Advanced Studies (SOKENDAI), Hayama 240-0193, Japan.

⁶Department of Neurobiology, Graduate School of Medicine, The University of Tokyo, Tokyo 133-0033, Japan

*Corresponding authors: Yohei Okubo & Masamitsu Iino

YO: Department of Pharmacology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

E-mail: yoheio@m.u-tokyo.ac.jp, Tel: +81-3-5841-3414, Fax: +81-3-5841-3390

MI: Department of Cellular and Molecular Pharmacology, Nihon University School of Medicine, 30-1, Oyaguchi Kami-cho, Itabashi-ku, Tokyo 173-8610, Japan.

E-mail: iino@m.u-tokyo.ac.jp, Tel: +81-3-3972-8111

Acknowledgements

We thank Y. Kawashima for technical assistance and Dr. B. Khakh (University of California at San Francisco) for providing the pZac2.1-gfaABC1D-Lck-GCaMP3 plasmid. This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI [Grant Numbers JP16K08543 (Y.O.), JP15H05648 (K.K.), and JP25117002 and JP25221304 (M.I.)] and grants from the Tokyo Society of Medical Sciences (Y.O.).

Conflict of Interest Statement

The authors declare no competing financial interests.

Abstract

Accumulating evidence indicates that astrocytes are actively involved in the physiological and pathophysiological functions of the brain. Intracellular Ca^{2+} signaling, especially Ca^{2+} release from the endoplasmic reticulum (ER), is considered to be crucial for the regulation of astrocytic functions. Mice with genetic deletion of inositol 1,4,5-trisphosphate receptor type 2 ($\text{IP}_3\text{R2}$) are reportedly devoid of astrocytic Ca^{2+} signaling, and thus widely used to explore the roles of Ca^{2+} signaling in astrocytic functions. While functional deficits in $\text{IP}_3\text{R2}$ -knockout (KO) mice have been found in some reports, no functional deficit was observed in others. Thus, there remains a controversy regarding the functional significance of astrocytic Ca^{2+} signaling. To address this controversy, we re-evaluated the assumption that Ca^{2+} release from the ER is abolished in $\text{IP}_3\text{R2}$ -KO astrocytes using a highly sensitive imaging technique. We expressed the ER luminal Ca^{2+} indicator G-CEPIA1 er in cortical and hippocampal astrocytes to directly visualize spontaneous and stimulus-induced Ca^{2+} release from the ER. We found attenuated but significant Ca^{2+} release in response to application of norepinephrine to $\text{IP}_3\text{R2}$ -KO astrocytes. This $\text{IP}_3\text{R2}$ -independent Ca^{2+} release induced only minimal cytosolic Ca^{2+} transients but induced robust Ca^{2+} increases in mitochondria that are frequently in close contact with the ER. These results indicate that ER Ca^{2+} release is retained and is sufficient to increase the Ca^{2+} concentration in close proximity to the ER in $\text{IP}_3\text{R2}$ -KO astrocytes.

Keywords

Astrocyte; Endoplasmic reticulum; Mitochondria; Calcium; Inositol 1,4,5-trisphosphate

Introduction

Accumulating evidence indicates that astrocytes play not only trophic and supportive roles but also active roles in the physiological and pathophysiological functions of the brain (Volterra & Meldolesi, 2005; Barres, 2008). Therefore, clarifying the mechanisms for regulation of astrocytic functions has become a crucial issue in neuroscience. Intracellular Ca^{2+} signaling has attracted attention because astrocytes show robust spontaneous and stimulus-induced Ca^{2+} transients (Verkhratsky, Orkand, & Kettenmann, 1998; Bazargani & Attwell, 2016).

Ca^{2+} release from the endoplasmic reticulum (ER) via inositol 1,4,5-trisphosphate receptor (IP_3R) has been recognized as a major component of spontaneous and G_q -coupled receptor-induced Ca^{2+} signaling in astrocytes. Therefore, to explore the significance of astrocytic Ca^{2+} signaling, mice with genetic deletion of IP_3R type 2 ($\text{IP}_3\text{R}2$) (Li et al., 2005), which is enriched in astrocytes (Sharp et al., 1999; Holtzclaw et al., 2002; Zhang et al., 2014), have been widely used. In $\text{IP}_3\text{R}2$ -knockout (KO) mice, Ca^{2+} transients in astrocytes were considered to be absent based on an imaging study of cytosolic Ca^{2+} transients by fluorescent Ca^{2+} indicator dyes (Petravicz, Fiacco, & McCarthy, 2008). Physiological and pathophysiological phenotypes have been observed in $\text{IP}_3\text{R}2$ -KO mice, including plasticity and learning (Takata et al., 2011; Chen et al., 2012; Navarrete et al., 2012; Perez-Alvarez et al., 2014; Padmashri, Suresh, Boska, & Dunaevsky, 2015; Kim et al., 2016; Monai et al., 2016; Yang et al., 2016), homeostasis of K^+ (Wang et al., 2012a; b), pathogenesis of stroke, traumatic brain injury and neurodegenerative disease (Dong, He, & Chai, 2013; Kanemaru et al., 2013; Li et al., 2015; Rakers & Petzold, 2016; Saito et al., 2018), and depression-like behaviors (Cao et al., 2013). However, several studies have reported no changes in basal synaptic activity and behavior (Petravicz, Fiacco, & McCarthy, 2008; Agulhon et al., 2013; Petravicz, Boyt, & McCarthy, 2014), hippocampal synaptic plasticity (Agulhon, Fiacco, &

McCarthy, 2010), and neurovascular coupling (Nizar et al., 2013; Takata et al., 2013; Bonder & McCarthy, 2014) in IP₃R2-KO mice. These contradictory results have caused confusion about the functional significance of Ca²⁺ signaling in astrocytes (Hamilton & Attwell, 2010; Fiacco & McCarthy, 2018; Savtchouk & Volterra, 2018).

One of the possible explanations that can resolve this controversy is that IP₃R2-independent Ca²⁺ signaling exists and contributes to the functions of IP₃R2-KO astrocytes. In fact, recent studies using genetically encoded Ca²⁺ indicators (GECIs) have clearly shown the presence of Ca²⁺ signals in IP₃R2-KO astrocytes (Kanemaru et al., 2014; Srinivasan et al., 2015; Rungta et al., 2016; Agarwal et al., 2017). These IP₃R2-independent Ca²⁺ transients were thought to be mediated by Ca²⁺ influx via the plasma membrane (Shigetomi et al., 2012; Srinivasan et al., 2015; Rungta et al., 2016) or Ca²⁺ release from mitochondria (Agarwal et al., 2017). These studies suggest that increases in the concentration of Ca²⁺ originating from sources other than the ER may regulate the functions of astrocytes. However, it remains elusive whether ER Ca²⁺ release is completely abolished in IP₃R2-KO astrocytes.

We thus examined whether IP₃R2-independent Ca²⁺ release from the ER is present in IP₃R2-KO astrocytes. Because cytosolic Ca²⁺ transients can be generated by Ca²⁺ derived from sources other than the ER, we directly detected ER Ca²⁺ release as a decrease in the Ca²⁺ concentration within the ER ([Ca²⁺]_{ER}). We recently developed a series of calcium-measuring organelle-entrapped protein indicators (CEPIAs) that allow visualization of Ca²⁺ dynamics within the ER and mitochondria with a high signal-to-noise ratio (Suzuki et al., 2014; Okubo et al., 2015; Suzuki, Kanemaru, & Iino, 2016). In this study, we expressed the ER Ca²⁺ indicator G-CEPIA1_{er} in cortical and hippocampal astrocytes of IP₃R2-KO mice. We found attenuated but significant Ca²⁺ release from the ER. We also used the mitochondrial Ca²⁺ indicator CEPIA2_{mt} and successfully observed robust Ca²⁺ transfer into mitochondria from

the ER in IP₃R2-KO astrocytes. These findings allowed us to reinterpret the results from IP₃R2-KO mice and shed new light on the significance of astrocytic Ca²⁺ signaling.

Materials and Methods

Animals

All animal experiments were carried out in accordance with the regulations and guidelines of the Institutional Animal Care and Use Committee at The University of Tokyo and were approved by the Institutional Review Committee of the Graduate School of Medicine, The University of Tokyo. C57BL/6 mice were used as wild-type (WT) mice. IP₃R2-KO mice (Li et al., 2005) were obtained from J. Chen (University of California at San Diego). Mice were kept under a 12 h light/dark cycle with *ad libitum* access to food and drinking water.

Preparation of viral vectors

To generate adeno-associated viruses (AAVs) for astrocyte-specific expression of GECIs, the cytomegalovirus promoter of pAAV-MCS (AAV Helper Free Expression System, Cell Biolabs, Inc., San Diego, CA) was replaced with the gfaABC₁D astrocyte-specific promoter derived from pZac2.1-gfaABC₁D-Lck-GCaMP3 (Shigetomi et al., 2013). G-CEPIA1*er*, CEPIA2*mt*, GCaMP6f, and ER-YFP were inserted downstream of the gfaABC₁D promoter to generate pAAV-gfaABC₁D-G-CEPIA1*er*, pAAV-gfaABC₁D-CEPIA2*mt*, pAAV-gfaABC₁D-GCaMP6f, and pAAV-gfaABC₁D-ER-YFP. AAV5 vectors were packaged using the AAV Helper Free Expression System. The packaging plasmids (pAAV-RC5 and pHelper) and transfer plasmid (pAAV-gfaABC₁D-G-CEPIA1*er*, pAAV-gfaABC₁D-CEPIA2*mt*, pAAV-gfaABC₁D-GCaMP6f, or pAAV-gfaABC₁D-ER-YFP) were transfected into HEK293T cells using the calcium phosphate method. The medium was replaced at 18 h after transfection with fresh medium, and the cells were incubated for 48 h. Harvested cells were lysed by repeated freezing and thawing, and a crude cell extract containing AAV5 vector particles was obtained. AAV5 vector particles were purified by ultracentrifugation with cesium chloride. The purified particles were dialyzed against PBS

and then concentrated by ultrafiltration using an Amicon 10K MWCO filter (Merck Millipore, Darmstadt, Germany). The copy number of the viral genome (vg) was determined by real-time quantitative PCR.

AAV injection

Male C57BL/6 mice or IP₃R2-KO mice (postnatal day 56–120) were anesthetized with isoflurane (induction at 5%, maintenance at 1–2%, vol/vol, MK-A100, Muromachi, Kyoto, Japan). The mice were placed in a stereotaxic frame (SR-5M-HT, Narishige, Tokyo, Japan). The skull was thinned (about 1 mm in diameter) above the right parietal cortex using a burr powered by a high speed drill (ULTIMATE XL-D, NSK, Kanuma, Japan). AAV5-gfaABC₁D-G-CEPIA1_{er} (0.98 or 1.3×10^{13} vg ml⁻¹), AAV5-gfaABC₁D-CEPIA2_{mt} (0.98×10^{13} vg ml⁻¹), AAV5-gfaABC₁D-GCaMP6f (1.1×10^{13} vg ml⁻¹), or pAAV-gfaABC₁D-ER-YFP (3.0×10^{13} vg ml⁻¹) was unilaterally injected into the cortex (1.5–2 mm posterior to the bregma, 1–1.5 mm lateral to the midline, and 300 μm from the surface) or hippocampus (2 mm posterior to the bregma, 1.5 mm lateral to the midline, and 1.6 mm from the surface) through glass pipettes. A viral solution (1 μL) was delivered at a rate of 200 nL min⁻¹ using a micropump (Legato 130, KD scientific, Holliston, MA). Glass pipettes were left in place for at least 10 min. Mice were sacrificed at 14–28 days after AAV injection for imaging and immunohistochemistry.

Preparation of brain slices and imaging

Coronal cortical or hippocampal slices (300 μm in thickness) were prepared as described previously (Edwards, Konnerth, Sakmann, & Takahashi, 1989). Slices were prepared in ice-cold artificial cerebrospinal fluid (ACSF) bubbled with 95% O₂ and 5% CO₂ using a vibrating slicer (PRO7, Dosaka, Kyoto, Japan). Slices were incubated in a holding chamber

containing ACSF bubbled with 95% O₂ and 5% CO₂ at 35°C for 1 h and then returned to 23°C. ACSF contained (in mM) 125 NaCl, 2.5 KCl, 2 CaCl₂, 1 MgSO₄, 1.25 NaH₂PO₄, 26 NaHCO₃, and 20 glucose. Imaging was carried out with a two-photon microscope (TSC MP5, Leica, Wetzlar, Germany) equipped with a water immersion objective (×25, NA 0.95, HCS IR APO, Leica) and Ti:sapphire laser (MaiTai DeepSee; Spectra Physics, Santa Clara, CA). Slices were transferred to a recording chamber under a microscope and continuously perfused with ACSF bubbled with 95% O₂ and 5% CO₂. Tetrodotoxin (1 μM) was added to ACSF to inhibit neuronal activities throughout imaging. Norepinephrine (NE, 10 μM) and cyclopiazonic acid (CPA, 50 μM) were dissolved in ACSF and administered through the perfusion system of the recording chamber. The excitation wavelength was 900–920 nm. Emitted fluorescence was filtered by a 500–550 nm barrier filter and detected with photomultiplier tubes. Data were acquired in time lapse XY-scan mode (0.2–0.5 Hz). Experiments were carried out at room temperature (22–24°C).

Data analysis

Data were analyzed using ImageJ software. Astrocytes were analyzed in layer 2/3 of the cortex or in the CA1 stratum radiatum of the hippocampus. Fluorescence intensities were corrected for background fluorescence by measuring a non-fluorescent area and normalized by the average of the first 10 or 20 frames to calculate the fractional changes in fluorescence intensity ($\Delta F/F_0$). Spontaneous G-CEPIA1*er* responses were manually detected and analyzed. Fluorescence changes with a rate of decrease faster than $-0.015 \Delta F/F_0 \text{ s}^{-1}$ and amplitude larger than $-0.05 \Delta F/F_0$ were defined as responses. Synchronous decreases in G-CEPIA1*er* fluorescence throughout a single astrocyte were defined as global responses. Localized fluorescence decreases with about 10–15 μm in diameter at the process region were defined as process-localized responses. The amplitude of spontaneous G-CEPIA1*er* responses was

measured after smoothing time courses of $\Delta F/F_0$ by a moving average with five frame windows. The amplitude of NE-induced G-CEPIA1*er*, CEPIA2*mt*, GCaMP6f, or ER-YFP responses was defined as the average of $\Delta F/F_0$ within the 3-min time window starting from the start of NE application. The amplitude of the CPA-induced G-CEPIA1*er* fluorescence decrease was defined as the $\Delta F/F_0$ value in a steady state. For statistical analyses, Student's *t*-test was performed. The asterisks in the figures indicate the following: **P* < 0.05, ***P* < 0.01, and ****P* < 0.001.

Results

G-CEPIA1er detects Ca^{2+} release from the ER in cortical astrocytes

We used a serotype 5 AAV carrying the minimal astrocyte-specific gfaABC₁D promoter to express G-CEPIA1er in astrocytes of the adult mouse cortex (Shigetomi et al., 2013).

G-CEPIA1er-expressing astrocytes in acute cortical slice preparations were imaged using a two-photon microscope (Fig. 1A). At subcellular levels, expression of G-CEPIA1er was observed throughout the soma and processes (Fig. 1A). This G-CEPIA1er distribution was consistent with the distribution pattern of the ER in astrocytes (Patrushev, Gavrilov, Turlapov, & Semyanov, 2013).

G-CEPIA1er-expressing layer 2/3 astrocytes were imaged to analyze Ca^{2+} release from the ER. We first applied CPA, a sarco/endoplasmic reticulum Ca^{2+} -ATPase (SERCA) inhibitor, to confirm G-CEPIA1er responses to $[\text{Ca}^{2+}]_{\text{ER}}$ decreases. Inhibition of SERCA-dependent Ca^{2+} uptake results in Ca^{2+} depletion of the ER. Indeed, CPA treatment resulted in a marked decrease in the G-CEPIA1er fluorescence intensity of both WT and IP₃R2-KO astrocytes, confirming that G-CEPIA1er reflects $[\text{Ca}^{2+}]_{\text{ER}}$ (Fig. 1B). There was no significant difference in the amplitudes of the CPA-induced G-CEPIA1er fluorescence decrease between WT and IP₃R2-KO astrocytes, indicating that the basal $[\text{Ca}^{2+}]_{\text{ER}}$ in IP₃R2-KO astrocytes was similar to that in WT astrocytes (Fig. 1B). However, the rate of decrease in $[\text{Ca}^{2+}]_{\text{ER}}$ was often greater in WT astrocytes than in IP₃R2-KO astrocytes (Fig. 1B). In the majority of WT astrocytes (17 cells among 23 cells), $[\text{Ca}^{2+}]_{\text{ER}}$ decreased in a stepwise manner as shown in Fig. 1B. Each step appeared to reflect spontaneous Ca^{2+} release events that are described in the following (Fig. 3). In the remaining WT astrocytes (6 cells among 23 cells) and all of IP₃R2-KO astrocytes (21 cells) showed a gradual decrease in $[\text{Ca}^{2+}]_{\text{ER}}$ due to Ca^{2+} leakage from the CPA-treated ER (Carreras-Sureda, Pihán, & Hetz, 2018).

Ca²⁺ release from the ER is retained in IP₃R2-KO astrocytes

We next investigated G_q-coupled receptor-induced Ca²⁺ release from the ER in astrocytes. Norepinephrine (NE), which is released from the axons of locus coeruleus neurons, induces cell-wide Ca²⁺ release from the ER through activation of α_1 -adrenergic receptors and following IP₃ signaling in WT astrocytes *in vivo* (Ding et al., 2013; Paukert et al., 2014; Srinivasan et al., 2015). Bath application of NE (10 μ M) induced a large decrease in [Ca²⁺]_{ER} throughout the soma and processes of WT astrocytes, indicating robust and cell-wide Ca²⁺ release from the ER (Fig. 2A and C). Following washout of NE, there was slow recovery of [Ca²⁺]_{ER} due to Ca²⁺ reuptake by the ER. The time course was consistent with ER Ca²⁺ dynamics induced by IP₃R-mediated Ca²⁺ release in other cell types shown in our previous studies (Suzuki et al., 2014; Okubo et al., 2015). In some WT astrocytes, [Ca²⁺]_{ER} undershot the dynamic range of G-CEPIA1*er* (Fig. 2A and C). In such cases, response amplitude would be underestimated. Most GECIs including G-CEPIA1*er* are sensitive to pH. To exclude the possibility that the observed changes in the G-CEPIA1*er* fluorescence intensity was influenced by a change in pH within the ER, we expressed ER-localized YFP that is sensitive to pH but insensitive to Ca²⁺ (Suzuki et al., 2014). ER-YFP showed no response to NE, confirming that NE-induced G-CEPIA1*er* responses reflected decreases in [Ca²⁺]_{ER} (Fig. 2C).

We then studied the ER Ca²⁺ dynamics in IP₃R2-KO astrocytes. Although previous studies have shown that NE-induced Ca²⁺ release is absent in IP₃R2-KO astrocytes, NE application induced a smaller but significant decrease in [Ca²⁺]_{ER} throughout the soma and processes of IP₃R2-KO astrocytes, while there was no change in ER-YFP fluorescence (Fig. 2B and C). Thus, contrary to the previous assumption, this observation clearly indicates that ER Ca²⁺ release is retained in IP₃R2-KO astrocytes.

During our observation of [Ca²⁺]_{ER} in WT astrocytes, we found spontaneous decreases

in $[Ca^{2+}]_{ER}$ (Fig. 3). There were two types of spontaneous events: a “global” response that showed a synchronous decrease in $[Ca^{2+}]_{ER}$ throughout the astrocyte (Fig. 3A), and a “process-localized” response that was localized at processes with diameters of about 10–15 μm (Fig. 3B). However, we did not find either “global” or “process-localized” responses in IP_3R2 -KO astrocytes (Fig. 3C). Thus, in IP_3R2 -KO astrocytes, spontaneous Ca^{2+} release events, if any, appeared to be attenuated to a level undetectable by G-CEPIA1*er*. These spontaneous Ca^{2+} release dynamics may at least partly correspond to cell-wide and localized Ca^{2+} signals imaged by cytosolic Ca^{2+} indicators in previous studies (Kanemaru et al., 2014; Srinivasan et al., 2015), although precise correlations require further analyses.

IP_3R2 -independent Ca^{2+} release in hippocampal astrocytes

Regional differences in the properties of Ca^{2+} signaling in astrocytes have been reported (Chai et al., 2017). To investigate whether IP_3R2 -independent Ca^{2+} release could be observed in astrocytes of brain regions other than the cortex, we visualized Ca^{2+} release from the ER in hippocampal astrocytes. G-CEPIA1*er* was expressed in hippocampal astrocytes using the AAV. G-CEPIA1*er*-expressing astrocytes in the CA1 stratum radiatum of acute hippocampal slice preparations were imaged by two-photon microscopy. Application of NE induced large G-CEPIA1*er* responses in WT astrocytes (Fig. 4A and B), whereas attenuated G-CEPIA1*er* responses were observed in IP_3R2 -KO astrocytes (Fig. 4A and B). Therefore, IP_3R2 -independent Ca^{2+} release was commonly observed in cortical and hippocampal astrocytes of IP_3R2 -KO mice. “Global” and “process localized” spontaneous G-CEPIA1*er* responses were observed in WT astrocytes, but not in IP_3R2 -KO astrocytes, which was consistent with the results in cortical astrocytes (Fig. 4).

Minimal Ca^{2+} transients in the cytosol mediated by IP_3R2 -independent Ca^{2+} release

Having found significant IP₃R2-independent Ca²⁺ release in IP₃R2-KO astrocytes, we decided to reinvestigate changes in the cytosolic Ca²⁺ concentration ([Ca²⁺]_{Cyt}) in response to NE application (Fig. 4). Cytosolic Ca²⁺ indicator GCaMP6f (Chen et al., 2013) was expressed in cortical astrocytes using an AAV, showing a diffuse distribution throughout the cytosol (Fig. 4A). In WT astrocytes, application of NE induced large [Ca²⁺]_{Cyt} responses that were clearly detected by $\Delta F/F_0$ traces of GCaMP6f averaged within the entire cell (Fig. 4A). Responses were abolished after treatment with CPA, indicating a critical contribution of ER Ca²⁺ release to NE-induced [Ca²⁺]_{Cyt} transients in WT astrocytes (Fig. 4B and C). On the other hand, NE induced very little [Ca²⁺]_{Cyt} responses in IP₃R2-KO astrocytes (Fig. 4A). Most IP₃R2-KO astrocytes showed no or minimal responses, and only very few IP₃R2-KO astrocytes showed clear but relatively small responses (Fig. 4B). The [Ca²⁺]_{Cyt} responses in IP₃R2-KO astrocytes were also abolished after treatment with CPA (Fig. 4B and C). These results indicate that IP₃R2-independent Ca²⁺ release, which was detected by G-CEPIA1er, did not always increase [Ca²⁺]_{Cyt} to a level detectable by GCaMP6f.

Robust Ca²⁺ transients in mitochondria mediated by IP₃R2-independent Ca²⁺ release

The results so far indicate that IP₃R2-independent Ca²⁺ release is clearly present in astrocytes, but that it induces only very little cytosolic Ca²⁺ signals. Does it mean IP₃R2-independent Ca²⁺ release has no physiological role? It has been shown that the ER and mitochondria form close contacts where Ca²⁺ is efficiently transferred from the ER to mitochondria without a global increase in [Ca²⁺]_{Cyt} (Rizzuto et al., 1998; de Brito & Scorrano, 2010; Hirabayashi et al., 2017). This mitochondrial Ca²⁺ signaling regulates various physiological and pathophysiological functions of mitochondria (Rizzuto, De Stefani, Raffaello, & Mammucari, 2012). We, therefore, examined whether IP₃R2-independent Ca²⁺ release would contribute to the ER-mitochondrial coupling. To study changes in the Ca²⁺ concentration within

mitochondria ($[Ca^{2+}]_{Mito}$), we used CEPIA2*mt*, a Ca^{2+} indicator for the mitochondrial matrix (Suzuki et al., 2014) (Fig. 5). AAV-mediated expression of CEPIA2*mt* in cortical astrocytes resulted in a punctate distribution of fluorescence, which was consistent with the morphology of mitochondria (Fig. 5A). Application of NE to release Ca^{2+} from the ER induced an increase in $[Ca^{2+}]_{Mito}$ of WT astrocytes (Fig. 5B). Although a subpopulation of WT astrocytes showed no or very little responses (Fig. 5C), such intercellular heterogeneity of mitochondrial Ca^{2+} signaling has been similarly observed in HeLa cells (Suzuki et al., 2014). The NA-induced $[Ca^{2+}]_{Mito}$ response was abolished after treatment with CPA to deplete the ER (Fig. 5C and D). These results indicate that Ca^{2+} enters mitochondria following Ca^{2+} release from the ER in WT astrocytes.

Of note, NE also induced $[Ca^{2+}]_{Mito}$ responses in IP₃R2-KO astrocytes (Fig. 5A). Similar to WT astrocytes, a subpopulation of IP₃R2-KO astrocytes showed no or very little responses. Compared with the minimal $[Ca^{2+}]_{Cyt}$ responses, $[Ca^{2+}]_{Mito}$ showed significant responses in IP₃R2-KO astrocytes upon NE stimulation. The $[Ca^{2+}]_{Mito}$ responses in IP₃R2-KO astrocytes were also abolished after treatment with CPA (Fig. 5B and C). These results indicate that IP₃R2-independent Ca^{2+} release efficiently induces an increase in $[Ca^{2+}]_{Mito}$.

Discussion

The results of the present study provide clear evidence that Ca^{2+} is released from the ER in $\text{IP}_3\text{R2-KO}$ astrocytes upon agonist stimulation. Although the $\text{IP}_3\text{R2}$ -independent Ca^{2+} release from the ER was ineffective in increasing the cytosolic Ca^{2+} concentration in $\text{IP}_3\text{R2-KO}$ astrocytes, it was effective to increase the Ca^{2+} concentration within mitochondria that make close contacts with the ER. These results provide new insights into the functional significance of ER Ca^{2+} release in astrocytes. The controversy derived from the assumption that ER Ca^{2+} release is abolished in $\text{IP}_3\text{R2-KO}$ astrocytes should be re-evaluated considering the presence of $\text{IP}_3\text{R2}$ -independent Ca^{2+} release.

Highly sensitive and specific detection of ER Ca^{2+} release using G-CEPIA1 er

Although $\text{IP}_3\text{R2}$ -independent Ca^{2+} release from the ER in $\text{IP}_3\text{R2-KO}$ astrocytes was detectable by G-CEPIA1 er responses, the resulting changes in $[\text{Ca}^{2+}]_{\text{Cyt}}$ were only barely detected by GCaMP6f responses. The difficulty in detecting the $[\text{Ca}^{2+}]_{\text{Cyt}}$ response is consistent with previous reports that failed to detect evoked $[\text{Ca}^{2+}]_{\text{Cyt}}$ responses in $\text{IP}_3\text{R2-KO}$ astrocytes (Petravicz, Fiacco, & McCarthy, 2008; Agulhon, Fiacco, & McCarthy, 2010; Agulhon et al., 2013; Nizar et al., 2013; Takata et al., 2013; Petravicz, Boyt, & McCarthy, 2014). These results suggest that the amount of Ca^{2+} release from the ER sufficient to change free $[\text{Ca}^{2+}]_{\text{ER}}$ might be insufficient to change free $[\text{Ca}^{2+}]_{\text{Cyt}}$. There are a couple of potential reasons for the puzzling observation. First, the volume occupied by the smooth ER (V_{ER}) is about 10% of that of the cytoplasm (V_{Cyt}) (Paumgartner, Losa, & Weibel, 1981). Therefore, Ca^{2+} released from the ER is 10-fold diluted in the cytosol. Another possible reason is the difference in the Ca^{2+} buffering capacity k (a change in the total Ca^{2+} concentration in a subcellular compartment divided by the corresponding change in free Ca^{2+} concentration in the compartment) between the ER and cytosol. In many cell types, cytosolic k (k_{Cyt}) is estimated to be 10–2,000 due

mainly to the presence of high concentrations of Ca^{2+} binding proteins in the cytosol (Allbritton, Meyer, & Stryer, 1992; Neher & Augustine, 1992; Neher, 1995; Mogami et al., 1999). However, k of the ER (k_{ER}) is estimated to be two orders of magnitude less than k_{Cyt} in pancreatic acinar cells (Mogami et al., 1999). If k_{ER} is similarly lower than k_{Cyt} in astrocytes, this would also reduce the Ca^{2+} release-induced change in $[\text{Ca}^{2+}]_{\text{Cyt}}$. We thus estimated the dilution and buffering effects in astrocytes. We previously measured both $[\text{Ca}^{2+}]_{\text{Cyt}}$ and $[\text{Ca}^{2+}]_{\text{ER}}$ in astrocytes stimulated with 30 μM ATP (Suzuki et al., 2014). Since the peak changes in $[\text{Ca}^{2+}]_{\text{Cyt}}$ and $[\text{Ca}^{2+}]_{\text{ER}}$ were observed almost simultaneously, we assumed that the change in the total Ca^{2+} concentration in the cytosol can be approximated by that of ER Ca^{2+} concentration as follows:

$$k_{\text{Cyt}} \cdot \Delta[\text{Ca}^{2+}]_{\text{Cyt}} \cdot V_{\text{Cyt}} \approx -k_{\text{ER}} \cdot \Delta[\text{Ca}^{2+}]_{\text{ER}} \cdot V_{\text{ER}}.$$

$\Delta[\text{Ca}^{2+}]_{\text{Cyt}}$ was similar to those of HeLa, HEK, and BHK cells, and most likely to be of the order of 1 μM , while $\Delta[\text{Ca}^{2+}]_{\text{ER}}$ estimated by the ratiometric ER Ca^{2+} indicator GEM-CEPIA1 $_{\text{er}}$ was about $-260 \mu\text{M}$ (Suzuki et al., 2014). Thus, the product of buffering and volume factors can be described as follows:

$$(k_{\text{ER}} / k_{\text{Cyt}}) \cdot (V_{\text{ER}} / V_{\text{Cyt}}) \approx -\Delta[\text{Ca}^{2+}]_{\text{Cyt}} / \Delta[\text{Ca}^{2+}]_{\text{ER}} \approx 1/260.$$

This indicates that Ca^{2+} release from the ER is highly buffered and diluted in the cytosol. In addition to the presence of Ca^{2+} binding proteins, effects of the efflux of Ca^{2+} to the extracellular space and influx to other intracellular organelles such as mitochondria would partly contribute to the estimated buffering/dilution factor. Thus, free Ca^{2+} concentration changes in the cytosol are only about 0.4% of those in the ER. Therefore, G-CEPIA1 $_{\text{er}}$ is expected to report ER Ca^{2+} release with much higher sensitivity than cytosolic Ca^{2+} indicators. Furthermore, G-CEPIA1 $_{\text{er}}$ can specifically detect ER Ca^{2+} release, whereas cytosolic Ca^{2+} indicators do not specify the source of Ca^{2+} .

Contribution of IP₃R2-independent Ca²⁺ release to Ca²⁺ signaling in IP₃R2-KO

astrocytes

When the extent of the NE-induced increase in Ca²⁺ concentration was compared between WT and IP₃R2-KO astrocytes, there was a much higher Ca²⁺ increase in mitochondria than in the cytoplasm of IP₃R2-KO astrocytes (Figs. 4 and 5). These results suggest an IP₃R2-independent mechanism that allows privileged transfer of Ca²⁺ from the ER to mitochondria. Indeed, it has been shown that the ER and mitochondria make close contacts (10–30 nm in distance), and that such contact sites are crucial for efficient Ca²⁺ transfer (Rizzuto et al., 1998; de Brito & Scorrano, 2010; Hirabayashi et al., 2017). This finding suggests that IP₃R2-independent Ca²⁺ release is capable of increasing [Ca²⁺]_{Cyt} within close proximity to the ER membrane, generating Ca²⁺ nanodomains around the ER (Rizzuto & Pozzan, 2006). These Ca²⁺ nanodomains generated within the ER-mitochondria contact sites should mediate efficient Ca²⁺ transfer between these two organelles (Rizzuto, De Stefani, Raffaello, & Mammucari, 2012). Other functional significances of IP₃R2-independent Ca²⁺ nanodomains around the ER require further clarification. Thus, in addition to extracellular and mitochondrial sources of Ca²⁺ (Srinivasan et al., 2015; Agarwal et al., 2017), IP₃R2-independent Ca²⁺ release from the ER may contribute to Ca²⁺ signaling in IP₃R2-KO astrocytes.

Ca²⁺ channels that mediate IP₃R2-independent Ca²⁺ release

Although IP₃R2 is the major ER Ca²⁺ release channel in astrocytes, expression of IP₃R1 and/or IP₃R3 in astrocytes has been indicated by transcriptome analyses (Cahoy et al., 2008; Zhang et al., 2014; Chai et al., 2017) and immunohistochemical studies (Yamamoto-Hino et al., 1995; Sharp et al., 1999). Furthermore, comparison between IP₃R2-KO mice and IP₃R2/IP₃R3-double KO mice indicate a functional contribution of IP₃R3 to astrocytic Ca²⁺

signaling, albeit small (Tamamushi et al., 2012; Sherwood et al., 2017). These previous studies suggest the presence of IP₃-induced Ca²⁺ release via IP₃R1 and/or IP₃R3 in IP₃R2-KO astrocytes. Indeed, crucial roles of IP₃R1 and IP₃R3 for the ER-mitochondria coupling have been reported. IP₃R1 has been shown to be one of the components that physically link the ER to mitochondria (Szabadkai et al., 2006). IP₃R3 is considered as a major subtype that mediates Ca²⁺ transfer from the ER to mitochondria (Mendes et al., 2005; Hayashi & Su, 2007; Cárdenas et al., 2010). Therefore, the involvement of IP₃R1 and/or IP₃R3 in mitochondrial Ca²⁺ signaling in IP₃R2-KO astrocytes is now being addressed.

In addition, expression of ryanodine receptor type 3 (RyR3) in astrocytes has been reported (Matyash et al., 2002; Chai et al., 2017). The functional significance of RyR3 was reported in both astrocytic Ca²⁺ signaling and motility (Matyash et al., 2002). Therefore, it also appears possible that Ca²⁺-induced Ca²⁺ release via RyR3 may enhance Ca²⁺ release from the ER via IP₃R1 and/or IP₃R3.

Reinterpretation of astrocytic functions in IP₃R2-KO mice

It has been assumed that Ca²⁺ release from the ER is absent in IP₃R2-KO astrocytes. Thus, the absence of functional deficits in IP₃R2-KO mice (Petravicz, Fiacco, & McCarthy, 2008; Agulhon, Fiacco, & McCarthy, 2010; Agulhon et al., 2013; Nizar et al., 2013; Takata et al., 2013; Bonder & McCarthy, 2014; Petravicz, Boyt, & McCarthy, 2014) has been considered to indicate the absence of functional roles of intracellular Ca²⁺ release in astrocytes. The presence of IP₃R2-independent Ca²⁺ release may necessitate reinterpretation of such results.

Ca²⁺ release from the ER may be inhibited by expressing the IP₃ hydrolyzing enzyme, IP₃ 5-phosphatase (5ppase) (Kanemaru, Okubo, Hirose, & Iino, 2007; Mashimo et al., 2010; Foley et al., 2017). Indeed, exogenous expression of 5ppase effectively suppresses IP₃ production following IP₃R-dependent Ca²⁺ release in astrocytes (Kanemaru, Okubo, Hirose,

& Iino, 2007; Mashimo et al., 2010). Because 5ppase is expected to suppress IP₃R-induced Ca²⁺ release regardless of IP₃R subtypes, the phenotypic difference between 5ppase-expressing mice and IP₃R2-KO mice may clarify the role of IP₃R2-independent Ca²⁺ release. In fact, a recent study has indicated that astrocytic expression of 5ppase disrupts sleep (Foley et al., 2017), whereas sleep disruption was not observed in IP₃R2-KO mice (Cao et al., 2013). Although there remains the possibility of developmental adaptation or compensatory mechanisms in these mice, the results suggest that residual Ca²⁺ signaling, including IP₃R2-independent Ca²⁺ release in IP₃R2-KO mice, accounts for the absence of functional deficits in IP₃R2-KO mice.

In conclusion, the presence of IP₃R2-independent Ca²⁺ release as well as Ca²⁺ signaling generated by sources other than the ER limit the use of IP₃R2-KO mice to assess the role of Ca²⁺ signaling in astrocytic functions when functional deficits are not observed in the mutant mice. This notion may help to resolve the long-lasting controversy arising from the assumption that Ca²⁺ signaling is abolished in IP₃R2-KO astrocytes. Furthermore, the present results facilitate the study of the functional roles of Ca²⁺ signaling in close proximity to the ER in astrocytes.

References

- Agarwal, A., Wu, P.-H., Hughes, E. G., Fukaya, M., Tischfield, M. A., Langseth, A. J., ... Bergles, D. E. (2017). Transient opening of the mitochondrial permeability transition pore induces microdomain calcium transients in astrocyte processes. *Neuron* 93, 587–605.
- Agulhon, C., Boyt, K. M., Xie, A. X., Friocourt, F., Roth, B. L., & McCarthy, K. D. (2013). Modulation of the autonomic nervous system and behaviour by acute glial cell G_q protein-coupled receptor activation in vivo. *The Journal of Physiology* 591, 5599–5609.
- Agulhon, C., Fiacco, T. A., & McCarthy, K. D. (2010). Hippocampal short- and long-term plasticity are not modulated by astrocyte Ca²⁺ signaling. *Science* 327, 1250–1254.
- Allbritton, N., Meyer, T., & Stryer, L. (1992). Range of messenger action of calcium ion and inositol 1,4,5-trisphosphate. *Science* 258, 1812–1815.
- Barres, B. A. (2008). The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron* 60, 430–440.
- Bazargani, N., & Attwell, D. (2016). Astrocyte calcium signaling: the third wave. *Nature Neuroscience* 19, 182–189.
- Bonder, D. E., & McCarthy, K. D. (2014). Astrocytic G_q-GPCR-linked IP₃R-dependent Ca²⁺ signaling does not mediate neurovascular coupling in mouse visual cortex *in vivo*. *Journal of Neuroscience* 34, 13139–13150.
- de Brito, O. M., & Scorrano, L. (2010). An intimate liaison: spatial organization of the endoplasmic reticulum-mitochondria relationship. *The EMBO journal* 29, 2715–23.
- Cahoy, J. D., Emery, B., Kaushal, A., Foo, L. C., Zamanian, J. L., Christopherson, K. S., ... Barres, B. A. (2008). A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function. *Journal of Neuroscience* 28, 264–278.
- Cao, X., Li, L.-P., Wang, Q., Wu, Q., Hu, H.-H., Zhang, M., ... Gao, T.-M. (2013). Astrocyte-derived ATP modulates depressive-like behaviors. *Nature Medicine* 19, 773–777.
- Cárdenas, C., Miller, R. A., Smith, I., Bui, T., Molgó, J., Müller, M., ... Foskett, J. K. (2010). Essential regulation of cell bioenergetics by constitutive InsP₃ receptor Ca²⁺ transfer to mitochondria. *Cell* 142, 270–283.
- Carreras-Sureda, A., Pihán, P., & Hetz, C. (2018). Calcium signaling at the endoplasmic reticulum: fine-tuning stress responses. *Cell Calcium* 70, 24–31.
- Chai, H., Diaz-Castro, B., Shigetomi, E., Monte, E., Oceau, J. C., Yu, X., ... Khakh, B. S. (2017). Neural circuit-specialized astrocytes: transcriptomic, proteomic, morphological,

- and functional evidence. *Neuron* 95, 531–549.
- Chen, N., Sugihara, H., Sharma, J., Perea, G., Petravicz, J., Le, C., & Sur, M. (2012). Nucleus basalis-enabled stimulus-specific plasticity in the visual cortex is mediated by astrocytes. *Proceedings of the National Academy of Sciences* 109, E2832–E2841.
- Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., ... Kim, D. S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499, 295–300.
- Ding, F., O'Donnell, J., Thrane, A. S., Zeppenfeld, D., Kang, H., Xie, L., ... Nedergaard, M. (2013). $\alpha 1$ -Adrenergic receptors mediate coordinated Ca^{2+} signaling of cortical astrocytes in awake, behaving mice. *Cell Calcium* 54, 387–394.
- Dong, Q., He, J., & Chai, Z. (2013). Astrocytic Ca^{2+} waves mediate activation of extrasynaptic NMDA receptors in hippocampal neurons to aggravate brain damage during ischemia. *Neurobiology of Disease* 58, 68–75.
- Edwards, F. A., Konnerth, A., Sakmann, B., & Takahashi, T. (1989). A thin slice preparation for patch clamp recordings from neurones of the mammalian central nervous system. *Pflügers Archiv : European journal of physiology* 414, 600–12.
- Fiacco, T. A., & McCarthy, K. D. (2018). Multiple lines of evidence indicate that gliotransmission does not occur under physiological conditions. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 38, 3–13.
- Foley, J., Blutstein, T., Lee, S., Erneux, C., Halassa, M. M., & Haydon, P. (2017). Astrocytic $\text{IP}_3/\text{Ca}^{2+}$ signaling modulates theta rhythm and REM sleep. *Frontiers in neural circuits* 11, 3.
- Hamilton, N. B., & Attwell, D. (2010). Do astrocytes really exocytose neurotransmitters? *Nature Reviews Neuroscience* 11, 227–238.
- Hayashi, T., & Su, T.-P. (2007). Sigma-1 receptor chaperones at the ER- mitochondrion interface regulate Ca^{2+} signaling and cell survival. *Cell* 131, 596–610.
- Hirabayashi, Y., Kwon, S.-K., Paek, H., Pernice, W. M., Paul, M. A., Lee, J., ... Polleux, F. (2017). ER-mitochondria tethering by PDZD8 regulates Ca^{2+} dynamics in mammalian neurons. *Science (New York, NY)* 358, 623–630.
- Holtzclaw, L. A., Pandhit, S., Bare, D. J., Mignery, G. A., & Russell, J. T. (2002). Astrocytes in adult rat brain express type 2 inositol 1,4,5-trisphosphate receptors. *Glia* 39, 69–84.
- Kanemaru, K., Kubota, J., Sekiya, H., Hirose, K., Okubo, Y., & Iino, M. (2013). Calcium-dependent N-cadherin up-regulation mediates reactive astrogliosis and neuroprotection after brain injury. *Proceedings of the National Academy of Sciences* 110, 11612–11617.

- Kanemaru, K., Okubo, Y., Hirose, K., & Iino, M. (2007). Regulation of neurite growth by spontaneous Ca^{2+} oscillations in astrocytes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 27, 8957–8966.
- Kanemaru, K., Sekiya, H., Xu, M., Satoh, K., Kitajima, N., Yoshida, K., ... Tanaka, K. F. (2014). *In vivo* visualization of subtle, transient, and local activity of astrocytes using an ultrasensitive Ca^{2+} indicator. *Cell Reports* 8, 311–318.
- Kim, S. K., Hayashi, H., Ishikawa, T., Shibata, K., Shigetomi, E., Shinozaki, Y., ... Nabekura, J. (2016). Cortical astrocytes rewire somatosensory cortical circuits for peripheral neuropathic pain. *The Journal of Clinical Investigation* 126, 1983–1997.
- Li, H., Xie, Y., Zhang, N., Yu, Y., Zhang, Q., & Ding, S. (2015). Disruption of $\text{IP}_3\text{R}2$ -mediated Ca^{2+} signaling pathway in astrocytes ameliorates neuronal death and brain damage while reducing behavioral deficits after focal ischemic stroke. *Cell Calcium* 58, 565–576.
- Li, X., Zima, A. V., Sheikh, F., Blatter, L. A., & Chen, J. (2005). Endothelin-1–induced arrhythmogenic Ca^{2+} signaling is abolished in atrial myocytes of inositol-1,4,5-trisphosphate (IP_3)–receptor type 2–deficient mice. *Circulation Research* 96, 1274–81.
- Mashimo, M., Okubo, Y., Yamazawa, T., Yamasaki, M., Watanabe, M., Murayama, T., & Iino, M. (2010). Inositol 1,4,5-trisphosphate signaling maintains the activity of glutamate uptake in Bergmann glia. *European Journal of Neuroscience* 32, 1668–1677.
- Matyash, M., Matyash, V., Nolte, C., Sorrentino, V., & Kettenmann, H. (2002). Requirement of functional ryanodine receptor type 3 for astrocyte migration. *The FASEB Journal* 16, 84–86.
- Mendes, C. C. P., Gomes, D. A., Thompson, M., Souto, N. C., Goes, T. S., Goes, A. M., ... Leite, M. F. (2005). The type III inositol 1,4,5-trisphosphate receptor preferentially transmits apoptotic Ca^{2+} signals into mitochondria. *The Journal of biological chemistry* 280, 40892–900.
- Mogami, H., Gardner, J., Gerasimenko, O., Camello, P., Petersen, O., & Tepikin, A. (1999). Calcium binding capacity of the cytosol and endoplasmic reticulum of mouse pancreatic acinar cells. *The Journal of physiology* 518, 463–7.
- Monai, H., Ohkura, M., Tanaka, M., Oe, Y., Konno, A., Hirai, H., ... Hirase, H. (2016). Calcium imaging reveals glial involvement in transcranial direct current stimulation-induced plasticity in mouse brain. *Nature Communications* 7, 11100.
- Navarrete, M., Perea, G., de Sevilla, D. F., Gómez-Gonzalo, M., Núñez, A., Martín, E. D., & Araque, A. (2012). Astrocytes mediate *in vivo* cholinergic-induced synaptic plasticity.

PLoS Biol 10, e1001259.

- Neher, E. (1995). The use of fura-2 for estimating Ca buffers and Ca fluxes. *Neuropharmacology* 34, 1423–1442.
- Neher, E., & Augustine, G. J. (1992). Calcium gradients and buffers in bovine chromaffin cells. *The Journal of physiology* 450, 273–301.
- Nizar, K., Uhlirova, H., Tian, P., Saisan, P. A., Cheng, Q., Reznichenko, L., ... Devor, A. (2013). *In vivo* stimulus-induced vasodilation occurs without IP₃ receptor activation and may precede astrocytic calcium increase. *Journal of Neuroscience* 33, 8411–8422.
- Okubo, Y., Suzuki, J., Kanemaru, K., Nakamura, N., Shibata, T., & Iino, M. (2015). Visualization of Ca²⁺ filling mechanisms upon synaptic inputs in the endoplasmic reticulum of cerebellar Purkinje cells. *Journal of Neuroscience* 35, 15837–15846.
- Padmashri, R., Suresh, A., Boska, M. D., & Dunaevsky, A. (2015). Motor-skill learning is dependent on astrocytic activity. *Neural Plasticity* 2015, 938023.
- Patrushev, I., Gavrilov, N., Turlapov, V., & Semyanov, A. (2013). Subcellular location of astrocytic calcium stores favors extrasynaptic neuron–astrocyte communication. *Cell Calcium* 54, 343–349.
- Paukert, M., Agarwal, A., Cha, J., Doze, V. A., Kang, J. U., & Bergles, D. E. (2014). Norepinephrine controls astroglial responsiveness to local circuit activity. *Neuron* 82, 1263–1270.
- Paumgartner, D., Losa, G., & Weibel, E. R. (1981). Resolution effect on the stereological estimation of surface and volume and its interpretation in terms of fractal dimensions. *Journal of Microscopy* 121, 51–63.
- Perez-Alvarez, A., Navarrete, M., Covelo, A., Martin, E. D., & Araque, A. (2014). Structural and functional plasticity of astrocyte processes and dendritic spine interactions. *Journal of Neuroscience* 34, 12738–12744.
- Petravicz, J., Boyt, K. M., & McCarthy, K. D. (2014). Astrocyte IP₃R2-dependent Ca²⁺ signaling is not a major modulator of neuronal pathways governing behavior. *Frontiers in Behavioral Neuroscience* 8, 384.
- Petravicz, J., Fiocco, T. A., & McCarthy, K. D. (2008). Loss of IP₃ receptor-dependent Ca²⁺ increases in hippocampal astrocytes does not affect baseline CA1 pyramidal neuron synaptic activity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 28, 4967–73.
- Rakers, C., & Petzold, G. C. (2016). Astrocytic calcium release mediates peri-infarct depolarizations in a rodent stroke model. *Journal of Clinical Investigation* 127, 511–516.
- Rizzuto, R., Pinton, P., Carrington, W., Fay, F. S., Fogarty, K. E., Lifshitz, L. M., ... Pozzan, R.

- T. (1998). Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca^{2+} responses. *Science (New York, NY)* 280, 1763–1766.
- Rizzuto, R., & Pozzan, T. (2006). Microdomains of intracellular Ca^{2+} : molecular determinants and functional consequences. *Physiological Reviews* 86, 369–408.
- Rizzuto, R., De Stefani, D., Raffaello, A., & Mammucari, C. (2012). Mitochondria as sensors and regulators of calcium signalling. *Nature Reviews Molecular Cell Biology* 13, 566–578.
- Rungta, R. L., Bernier, L.-P., Dissing-Olesen, L., Groten, C. J., LeDue, J. M., Ko, R., ... MacVicar, B. A. (2016). Ca^{2+} transients in astrocyte fine processes occur via Ca^{2+} influx in the adult mouse hippocampus. *Glia* 64, 2093–2103.
- Saito, K., Shigetomi, E., Yasuda, R., Sato, R., Nakano, M., Tashiro, K., ... Koizumi, S. (2018). Aberrant astrocyte Ca^{2+} signals “AxCa signals” exacerbate pathological alterations in an Alexander disease model. *Glia* 66, 1053–1067.
- Savtchouk, I., & Volterra, A. (2018). Gliotransmission: beyond black-and-white. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 38, 14–25.
- Sharp, A. H., Nucifora, F. C., Blondel, O., Sheppard, C. A., Zhang, C., Snyder, S. H., ... Ross, C. A. (1999). Differential cellular expression of isoforms of inositol 1,4,5-triphosphate receptors in neurons and glia in brain. *The Journal of comparative neurology* 406, 207–220.
- Sherwood, M. W., Arizono, M., Hisatsune, C., Bannai, H., Ebisui, E., Sherwood, J. L., ... Mikoshiba, K. (2017). Astrocytic IP_3Rs : contribution to Ca^{2+} signalling and hippocampal LTP. *Glia* 65, 502–513.
- Shigetomi, E., Bushong, E. A., Haustein, M. D., Tong, X., Jackson-Weaver, O., Kracun, S., ... Khakh, B. S. (2013). Imaging calcium microdomains within entire astrocyte territories and endfeet with GCaMPs expressed using adeno-associated viruses. *The Journal of General Physiology* 141, 633–647.
- Shigetomi, E., Tong, X., Kwan, K. Y., Corey, D. P., & Khakh, B. S. (2012). TRPA1 channels regulate astrocyte resting calcium and inhibitory synapse efficacy through GAT-3. *Nature Neuroscience* 15, 70–80.
- Srinivasan, R., Huang, B. S., Venugopal, S., Johnston, A. D., Chai, H., Zeng, H., ... Khakh, B. S. (2015). Ca^{2+} signaling in astrocytes from *Ip3r2^{-/-}* mice in brain slices and during startle responses *in vivo*. *Nature Neuroscience* 18, 708–717.
- Suzuki, J., Kanemaru, K., & Iino, M. (2016). Genetically encoded fluorescent indicators for organellar calcium imaging. *Biophysical Journal* 111, 1119–1131.
- Suzuki, J., Kanemaru, K., Ishii, K., Ohkura, M., Okubo, Y., & Iino, M. (2014). Imaging

- intraorganellar Ca^{2+} at subcellular resolution using CEPIA. *Nature communications* 5, 4153.
- Szabadkai, G., Bianchi, K., Várnai, P., De Stefani, D., Wieckowski, M. R., Cavagna, D., ... Rizzuto, R. (2006). Chaperone-mediated coupling of endoplasmic reticulum and mitochondrial Ca^{2+} channels. *The Journal of cell biology* 175, 901–11.
- Takata, N., Mishima, T., Hisatsune, C., Nagai, T., Ebisui, E., Mikoshiba, K., & Hirase, H. (2011). Astrocyte calcium signaling transforms cholinergic modulation to cortical plasticity *in vivo*. *Journal of Neuroscience* 31, 18155–18165.
- Takata, N., Nagai, T., Ozawa, K., Oe, Y., Mikoshiba, K., & Hirase, H. (2013). Cerebral blood flow modulation by basal forebrain or whisker stimulation can occur independently of large cytosolic Ca^{2+} signaling in astrocytes. *PLoS ONE* 8, e66525.
- Tamamushi, S., Nakamura, T., Inoue, T., Ebisui, E., Sugiura, K., Bannai, H., & Mikoshiba, K. (2012). Type 2 inositol 1,4,5-trisphosphate receptor is predominantly involved in agonist-induced Ca^{2+} signaling in Bergmann glia. *Neuroscience Research* 74, 32–41.
- Verkhratsky, A., Orkand, R. K., & Kettenmann, H. (1998). Glial calcium: homeostasis and signaling function. *Physiological Reviews* 78, 99–141.
- Volterra, A., & Meldolesi, J. (2005). Astrocytes, from brain glue to communication elements: the revolution continues. *Nature Reviews Neuroscience* 6, 626–640.
- Wang, F., Smith, N. A., Xu, Q., Fujita, T., Baba, A., Matsuda, T., ... Nedergaard, M. (2012)(a). Astrocytes modulate neural network activity by Ca^{2+} -dependent uptake of extracellular K^{+} . *Science Signaling* 5, ra26.
- Wang, F., Xu, Q., Wang, W., Takano, T., & Nedergaard, M. (2012)(b). Bergmann glia modulate cerebellar Purkinje cell bistability via Ca^{2+} -dependent K^{+} uptake. *Proceedings of the National Academy of Sciences of the United States of America* 109, 7911–7916.
- Yamamoto-Hino, M., Miyawaki, A., Kawano, H., Sugiyama, T., Furuichi, T., Hasegawa, M., & Mikoshiba, K. (1995). Immunohistochemical study of inositol 1,4,5-trisphosphate receptor type 3 in rat central nervous system. *Neuroreport* 6, 273–6.
- Yang, J., Yang, H., Liu, Y., Li, X., Qin, L., Lou, H., ... Wang, H. (2016). Astrocytes contribute to synapse elimination *via* type 2 inositol 1,4,5-trisphosphate receptor-dependent release of ATP. *eLife* 5, e15043.
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O’Keeffe, S., ... Wu, J. Q. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *Journal of Neuroscience* 34, 11929–11947.

Figure legends

Figure 1. Ca^{2+} imaging within the ER of cortical astrocytes using G-CEPIA1 er .

(A) G-CEPIA1 er -expressing astrocytes in the acute cortical slices. Maximum intensity projection of 22 (top) and 7 (bottom) images acquired by 1- μm interval. Scale bars, 100 μm (top) and 10 μm (bottom).

(B) SERCA inhibitor-induced Ca^{2+} depletion confirmed that G-CEPIA1 er reflected $[\text{Ca}^{2+}]_{\text{ER}}$. Representative $\Delta F/F_0$ traces of G-CEPIA1 er upon application of CPA (50 μM , gray bar) to WT (black traces) and IP₃R2-KO (magenta trace) astrocytes. Majority of WT astrocytes (17 cells among 23 cells) showed a stepwise decrease in G-CEPIA1 er fluorescence as indicated by arrows in the top right trace. The rest of WT astrocytes (6 cells among 23 cells, top left) and all IP₃R2-KO astrocytes (21 cells, bottom left) showed a gradual decrease in G-CEPIA1 er fluorescence. The region of interest to measure $\Delta F/F_0$ covered the entire cell. Bottom right: summary of the amplitude of G-CEPIA1 er fluorescence decrease upon application of CPA ($n = 23$ cells for WT and 21 cells for IP₃R2-KO, mean $\pm$ s.e.m.). No significant difference was detected between WT and IP₃R2-KO ($P = 0.447$, unpaired Student's t -test).

Figure 2. IP₃R2-independent Ca^{2+} release in cortical astrocytes.

(A and B) Representative G-CEPIA1 er responses upon application of NE (10 μM , cyan bars) to WT astrocytes (A) and IP₃R2-KO astrocytes (B) in cortical slices. The yellow circles in the leftmost images indicate the region of interest for the $\Delta F/F_0$ traces. Pseudocolor images indicate the average of five consecutive $\Delta F/F_0$ images within the time zones indicated by gray vertical bars. Scale bars, 10 μm .

(C) Left: G-CEPIA1 er and ER-YFP responses upon application of NE (10 μM , cyan bars) to WT (black) and IP₃R2-KO (magenta) astrocytes. Individual $\Delta F/F_0$ traces (pale color) and

average traces (deep color) are shown. Right: summary of the response amplitudes of $\Delta F/F_0$ traces (mean $\pm$ s.e.m.). For G-CEPIA1er, $n = 11$ cells for WT and 13 cells for IP₃R2-KO. For ER-YFP, $n = 12$ cells for WT and 10 cells for IP₃R2-KO. Asterisks indicate the statistical significance. ** $P < 0.01$, *** $P < 0.001$, unpaired Student's t -test.

Figure 3. Spontaneous ER Ca²⁺ release in WT cortical astrocytes.

(A and B) Representative spontaneous G-CEPIA1er responses that emerged in WT astrocytes either synchronously throughout the astrocyte (global response, A) or in confined regions of the processes (process-localized response, B). The yellow circles in the leftmost images indicate the region of interest for the $\Delta F/F_0$ traces. Pseudocolor images indicate the average of five consecutive $\Delta F/F_0$ images within the time zones indicated by gray vertical bars. Scale bar, 20 μ m.

(C) Representative G-CEPIA1er responses in IP₃R2-KO astrocytes without external stimulation. Data were shown as in A and B.

(D) Summary of frequencies (left) and amplitudes (right) of global and process-localized responses in WT astrocytes. $n = 25$ global responses and 27 process-localized responses in 15 cells (mean $\pm$ s.e.m.). No significant difference was detected between global and process-localized responses ($P = 0.905$ for frequency, $P = 0.831$ for amplitude, unpaired Student's t -test).

Figure 4. IP₃R2-independent Ca²⁺ release in hippocampal astrocytes.

(A) Representative G-CEPIA1er responses upon application of NE (10 μ M, cyan bars) to WT and IP₃R2-KO astrocytes in hippocampal slices. Yellow circles indicate the region of interest for the $\Delta F/F_0$ traces (black trace for WT and magenta trace for IP₃R2-KO). Pseudocolor images indicate the average of ten consecutive $\Delta F/F_0$ images within the time zones indicated

by gray vertical bars. Scale bar, 10 μm .

(B) Left: G-CEPIA1*er* responses upon application of NE (10 μM , cyan bars) to WT (black) and IP₃R2-KO (magenta) astrocytes. Individual $\Delta F/F_0$ traces (pale color) and average traces (deep color) are indicated. Right: Summary of the response amplitudes of $\Delta F/F_0$ traces ($n = 16$ cells for WT and 18 cells for IP₃R2-KO, mean $\pm$ s.e.m.).

(C and D) Representative spontaneous G-CEPIA1*er* responses that emerged in WT hippocampal astrocytes either synchronously throughout the astrocyte (global response, C) or in confined regions of the processes (process-localized response, D). The yellow circles in the image indicate the region of interest for the $\Delta F/F_0$ traces. Scale bar, 10 μm .

(E) Representative G-CEPIA1*er* responses in IP₃R2-KO hippocampal astrocytes without external stimulation. Data were shown as in C and D.

(F) Summary of frequencies and amplitudes of global and process-localized responses in hippocampal WT astrocytes. $n = 29$ global responses and 27 process-localized responses in 16 cells (mean $\pm$ s.e.m.). No significant difference was detected in frequency of global and process-localized responses ($P = 0.914$, unpaired Student's *t*-test). Amplitude of process-localized responses was significantly larger than global responses ($***P < 0.001$, unpaired Student's *t*-test).

Figure 5. Cytosolic Ca^{2+} transients induced by IP₃R2-independent Ca^{2+} release.

(A) Representative GCaMP6f responses upon application of NE (10 μM , cyan bars) to WT and IP₃R2-KO astrocytes. The yellow circles indicate the region of interest for the $\Delta F/F_0$ traces (black trace for WT and magenta trace for IP₃R2-KO). Pseudocolor images indicate $\Delta F/F_0$ at the time points indicated by gray vertical bars in the $\Delta F/F_0$ traces. Fluorescence increase in the soma of WT astrocytes was underestimated due to saturation of the dynamic range of photodetector. Scale bar, 10 μm .

(B) GCaMP6f responses upon application of NE (10 μ M, cyan bars) to WT (black) and IP₃R2-KO (magenta) astrocytes in the control condition [CPA (-)] or after treatment with CPA [CPA (+)]. A whole astrocyte was selected as the region of interest to measure $\Delta F/F_0$. Individual $\Delta F/F_0$ traces (pale color) and average traces (deep color) are indicated. $n = 13$ cells for WT [CPA (-)], 10 cells for WT [CPA (+)], 16 cells for IP₃R2-KO [CPA (-)], and 10 cells for IP₃R2-KO [CPA (+)].

(C) Summary of the response amplitudes of $\Delta F/F_0$ traces shown in (B) (mean $\pm$ s.e.m.). Asterisks indicate the statistical significance. * $P < 0.05$, *** $P < 0.001$, unpaired Student's t -test.

Figure 6. Mitochondrial Ca²⁺ transients induced by IP₃R2-independent Ca²⁺ release.

(A) CEPIA2mt-expressing astrocyte. The right image is the magnification of the region indicated by the yellow box in the left image. Scale bars, 10 μ m (left) and 2 μ m (right).

(B) Representative CEPIA2mt responses upon application of NE (10 μ M, cyan bars) to WT and IP₃R2-KO astrocytes. The yellow circles indicate the region of interest for the $\Delta F/F_0$ traces (black trace for WT and magenta trace for IP₃R2-KO). Pseudocolor images indicate the average of ten consecutive $\Delta F/F_0$ images within the time zones indicated by gray vertical bars. Scale bar, 10 μ m.

(C) CEPIA2mt responses upon application of NE (10 μ M, cyan bars) to WT (black) and IP₃R2-KO (magenta) astrocytes in the control condition [CPA (-)] or after treatment with CPA [CPA (+)]. A whole astrocyte was selected as the region of interest to measure $\Delta F/F_0$. Individual $\Delta F/F_0$ traces (pale color) and average traces (deep color) are shown. $n = 18$ cells for WT [CPA (-)], 10 cells for WT [CPA (+)], 16 cells for IP₃R2-KO [CPA (-)], and 12 cells for IP₃R2-KO [CPA (+)].

(D) Summary of the response amplitudes of $\Delta F/F_0$ traces shown in (B) (mean $\pm$ s.e.m.),

indicating significant CEPIA2 mt responses in IP₃R2-KO astrocytes. Absence of response after CPA indicates the requirement of ER Ca²⁺ release to CEPIA2 mt responses. Asterisks indicate the statistical significance of the responses in CPA (–) condition versus CPA (+) condition in each genotype. ** $P < 0.01$, *** $P < 0.001$, unpaired Student's t -test. Difference between WT [CPA (–)] and IP₃R2-KO [CPA (–)] responses was not statistically significant ($P = 0.0553$, unpaired Student's t -test).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6