1 Susceptibility of European freshwater fish to climate change: species profiling based on
2 life-history and environmental characteristics

3

4 Running head: European freshwater fish and climate change

5

6 Ivan Jarić^{1,2,3*}, Robert J. Lennox⁴, Gregor Kalinkat², Gorčin Cvijanović³ and Johannes
 7 Radinger^{2,5}

8

9 ¹ Biology Centre of the Czech Academy of Sciences, Institute of Hydrobiology, Ceske
 10 Budejovice, Czech Republic

11 ² Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany

12 ³ Institute for Multidisciplinary Research, University of Belgrade, Serbia

13 ⁴ Fish Ecology and Conservation Physiology Laboratory, Department of Biology, Carleton
 14 University, Ottawa, Ontario, Canada

15 ⁵ GRECO, Institute of Aquatic Ecology, University of Girona, Spain

16

17 * Corresponding author: Ivan Jarić, Biology Centre of the Czech Academy of Sciences,
 18 Institute of Hydrobiology, Na Sádkách 702/7, 370 05 České Budějovice, Czech Republic, E-
 19 mail: ivan.jaric@hbu.cas.cz, phone: +420 38 777 5855, fax: +420 385 310 24

20

21 Keywords: IUCN; Red List; extinction threat; global warming; climate change.

22

23

24

25 **Abstract**

26 Climate change is expected to strongly affect freshwater fish communities. Combined with
 27 other anthropogenic impacts, the impacts will alter species distributions and contribute to
 28 population declines and local extinctions. To provide timely management and conservation of
 29 fishes, it is relevant to identify species that will be most impacted by climate change and those
 30 that will be resilient. Species traits are considered a promising source of information on
 31 characteristics that influence resilience to various environmental conditions and impacts. We
 32 collated life history traits and climatic niches of 443 European freshwater fish species and
 33 compared those identified as susceptible to climate change to those that are considered to be
 34 resilient. Significant differences were observed between the two groups in their distribution,
 35 life-history and climatic niches, with climate-change susceptible species being distributed
 36 more southwardly within Europe, and being characterized by higher threat levels, lower
 37 commercial relevance, lower vulnerability to fishing, smaller body size and warmer thermal
 38 envelopes. We establish a list of species revealed to be of highest priority for further research
 39 and monitoring regarding climate change susceptibility within Europe. The presented
 40 approach represents a promising tool, to quickly assess large groups of species regarding their
 41 susceptibility to climate change and other threats, and to identify research and management
 42 priorities.

43

44 **Introduction**

45 As ectothermic organisms, fishes are intimately linked to local climatic conditions
 46 through physiological mechanisms that delimit tolerance or resilience (Comte & Olden,
 47 2017a). Zoogeography of fishes is therefore greatly influenced by the average and spread of

temperatures experienced in a given watershed (Pörtner & Farrell, 2008; Isaak & Rieman, 2013). Relative to seas and oceans, freshwater habitats are more drastically impacted by changes in climate, especially due to changes in temperature and flow, and climate change is projected to strongly affect freshwater fish communities (O'Reilly et al., 2003; Buisson et al., 2008; Graham & Harrod, 2009; Harrod, 2016; Radinger et al., 2017). Combined with other anthropogenic impacts (e.g. land use change and thermal pollution; Radinger et al., 2016; Raptis et al., 2017), climate change will restrict or redraw thermal envelopes, contribute to population declines and local extinctions, and overall shifts in the distribution of species. Riverine fish species losses due to climate change and reduced water discharge are predicted to reach 75% in some river basins (Xenopoulos et al., 2005). Phenological changes in fish behaviour (Otero et al., 2014; Dempson et al., 2017; Hovel et al., 2017) have been also detected and emphasize the powerful changes imposed by a changing climate. In Europe, there is a broad range of climatic conditions experienced across the landscape and a diverse ichthyofauna distributed throughout the lakes and rivers (Ficke et al., 2007). Within the IUCN (International Union for Conservation of Nature and Natural Resources) Red List, as much as 33% of European freshwater fish species are recognized as threatened by climate change (IUCN, 2017).

Efforts to preserve ecosystem integrity must focus on maintaining species richness and diversity to ensure that the services provided by freshwater ecosystems are maintained. Conservation is often limited by funding and therefore must undergo triage to identify priorities and allocate resources efficiently (McDonald-Madden et al., 2011). To provide timely management and conservation and allocate resources efficiently, it is important to identify those species that will be most impacted by climate change and those that might be rather resilient. Species traits are considered as a promising source of information on

characteristics that influence resilience to various environmental conditions and impacts (Jiguet et al., 2007; Comte & Olden, 2017b). Species traits represent any morphological, physiological or phenological feature that is measurable at the individual level of a species (Floeter et al., 2018). Trait-based evaluation has been demonstrated to be linked to the risk status of species and can be used to investigate mechanisms that contribute to imperilment, make predictions about unassessed species, or rank and prioritize species based on their relative risk (Olden et al., 2007; Bland & Böhm, 2016; Comte & Olden, 2018).

Here we assess various ecological and life-history characteristics of European freshwater fish species to identify traits that are characteristic for those that are susceptible to the effects of climate change. Automated scraping of an online trait database and calculation of climate envelopes using IUCN range maps overlaid on climate maps allowed us to collate species-specific data on life history, distribution, climatic niches, as well as data on threat and economic status. This allowed us to compare species identified as susceptible to climate change with those that are considered to be resilient. Results of the study will contribute to a better understanding of the expected climate change effects on European freshwater fish fauna. We also establish a list of European species of highest priority for further research and monitoring regarding climate change susceptibility. The method allows to extrapolate results and characterize rare and less studied species, with scarce autecological information.

Materials and methods

Dataset

Our analysis comprised comparisons of in total 443 European freshwater fishes between those that were identified as threatened by climate change (n=148) within the IUCN Red List Database (IUCN, 2017) and those without climate change listed as a threat (n=295).

96 A list of native European freshwater fish species belonging to 25 families, mainly to
 97 Cyprinidae (45%) and Salmonidae (20%), was obtained from the IUCN Red List database
 98 (IUCN, 2017). It included both exclusively freshwater species, as well as those that partly
 99 enter brackish and saltwater. Obtained data also comprised IUCN Red List classification and
 100 maps of their distributional range within Europe. In addition, we obtained information
 101 whether climate change was indicated as one of the threats for each species, which is based on
 102 threat analyses and expert judgement by IUCN species experts. Overall, the dataset comprised
 103 33% species which were categorized as susceptible to climate change.

104 In addition, for each species we collated trait information related to their life
 105 history, ecology, fishery and threat status, and spatial and bioclimatic data variables (Table 1).
 106 Life-history data were obtained from the FishBase database (Froese & Pauly, 2017) by using
 107 the rfishbase R package (Boettiger et al., 2012, 2017). Traits with low data coverage (i.e.
 108 those that were available for less than 1% of all species) were excluded from the analysis.
 109 Bioclimatic spatial data were obtained from the MERRAclim database (Vega et al., 2017) and
 110 included 19 variables related to temperature and humidity (Table 1). Global Human Footprint
 111 map (map of anthropogenic impacts on the environment) was obtained from WCS and
 112 CIESIN (2005) and the spatial elevation data were obtained from USGS (2010). Based on the
 113 distributional range maps of each species (IUCN, 2017), mean values within each species'
 114 range were estimated for each of the spatial variables using the intersect tool in ArcGIS
 115 (version 10.5) and the *extract* function in the R (version 3.4.3; R Core Team, 2017) package
 116 raster (version 2.6-7; Hijmans, 2017). Range maps were also used to estimate the number of
 117 watersheds covered by each species based on WRI (2006), as well as the area and coordinates
 118 of the range centroid for each species. General descriptive statistics and information on data
 119 sources of all variables used in the analysis is presented in Table 1 and Supplementary

material S1. The complete dataset is presented in Supplementary material S2.

Statistical Analysis

We calculated boosted regression trees (BRT) to evaluate the relationship between the membership of a species to the group of susceptible vs. non-susceptible species and the 45 explanatory variables. BRT are a statistical learning method that combines and averages (boosting) many simple single regression trees to form collective models of improved predictive performance (Elith et al., 2008). BRT can accommodate continuous and categorical variables, are not affected by missing values or transformation or outliers and are considered to effectively select relevant variables, identify variable interactions and avoid overfitting (Elith et al., 2008, Radinger et al., 2015).

Specifically, we first fitted an initial global BRT model (R package *dismo*, *gbm.step*, version 1.1-4; Hijmans et al., 2017) using the complete set of explanatory variables. An automatized stepwise backward selection of explanatory variables (*gbm.simplify*) was applied to eliminate non-informative variables based on model-internal cross-validation of changes in a models' predictive deviance (Hijmans et al., 2017). Thereafter, we calculated a final BRT model (*gbm.step*) based on the selected set of explanatory variables. For all BRT modeling steps, tree complexity and learning rate was set to 3 and 0.001, respectively, to achieve the recommended number of more than 1000 regression trees (Elith et al., 2008). All other model settings were set to default or were automatically adjusted by the boosting algorithm. We calculated a 10-fold cross validation of the BRT model as already implemented in the algorithm. In addition, we extracted the mean AUC (area under the receiver operating characteristic curve) as a measure of the model's predictive quality. The AUC is a threshold-independent rank-correlation coefficient with high values typically indicating a strong

agreement between the model prediction and the membership of species to the susceptible vs. non-susceptible group (Hijmans & Elith, 2017).

The relative importance (%) of each explanatory variable in the final BRT model was quantified based on the number of times each variable was used for splitting, weighted by the squared improvement at each split and averaged over all trees (Elith et al., 2008). For BRT models with Gaussian distribution the relative variable importance equals the reduction of squared error attributable to a given variable. Differences between groups were also assessed by bootstrapping, by sampling each group independently and estimating the difference based on confidence intervals (functions *two.boot* and *boot.ci*, R package *simpleboot*, version 1.1-3; Peng 2008). Differences were considered to be significant if 95% confidence intervals did not overlap with zero.

Subsequently, species were ranked based on the subset of variables selected by the BRT analysis (i.e., those with >1% variable importance score), and weighed by the importance of each variable, in order to estimate their position along the climate change susceptibility continuum. For each species, the value of each variable was standardized based on its position between the minimum ($t_{\min}$) and maximum values ($t_{\max}$) observed in the dataset, with 0-1 possible range, and multiplied by the importance score (I_x) of the given variable:

$$R_{tx} = \frac{t_x - t_{\min}}{t_{\max} - t_{\min}} \times I_x \quad (1)$$

where R_{tx} represents the rank value of variable t in species x , and t_x is the value of variable t for species x . For variables where the lower endpoint ($t_{\min}$) is associated with the climate change susceptibility, equation should be adjusted as follows:

$$R_{tx} = \frac{t_{\max} - t_x}{t_{\max} - t_{\min}} \times I_x \quad (2)$$

Summing of all ranking scores across all variables yielded the total species ranking score, which can range from 0 to 100, with higher values indicating stronger climate change susceptibility.

Results

Our analyses indicated substantial differences between the two groups, climate change susceptible versus non-susceptible species. The BRT analysis selected 35 most relevant variables, which were subsequently assessed for their relative importance to discriminate between the two groups (Figure 1 and Supplementary material S3). The BRT model with the selected set of explanatory variables was successfully modeled (Supplementary material S3) with a good cross-validated AUC of 0.87 (standard error = 0.014). Out of all explanatory variables, latitude of the species range centroid was selected as by far the most relevant variable (31% variable importance), followed by the IUCN Red List classification (8%), commercial relevance (6%) and vulnerability to fishing (6%). Climate susceptible species were characterized by more southwardly positioned distribution range centroids (41.6° vs. 47.8° N as a mean value in susceptible and non-susceptible species, respectively), smaller range sizes ($175 \times 10^3 \text{ km}^2$ vs. $1686 \times 10^3 \text{ km}^2$), and lower elevations within their ranges (717.7 m vs. 892.2 m a.s.l.), with a higher proportion of exclusively freshwater species (93% vs. 66%; Figure 2). Susceptible species were also characterized by a smaller maximum body length (23.4 cm vs. 41.0 cm), higher proportion of threatened species (63% vs. 31%), lower proportion of commercially relevant species (25% vs. 74% of highly commercial, commercial and minor commercial species), and lower vulnerability to overfishing (32.6 vs. 38.5 vulnerability index; Figure 3), as well as by higher temperature-related values (Supplementary material S4). Bootstrapping indicated significant differences

between the groups in each of the variables. Species that are susceptible to climate change are mainly distributed within the Mediterranean region, while the non-susceptible species distribution mainly covers central and northern European regions, as well as the Carpathian region (Figure 4).

Species ranking based on the association of their traits with climate change susceptibility characteristics is presented in Table 2 and Supplementary material S5. The five top-ranked climate susceptible species were Acheron spring goby (*Knipowitschia milleri*), Corfu toothcarp (*Valencia letourneuxi*), Iberochondrostoma almakai, Evia barbel (*Barbus euboicus*) and Malaga chub (*Squalius malacitanus*). Most of the species with the high climate-susceptibility ranks are also classified as highly threatened according to the IUCN classification (Table 2). Interestingly, the highest ranked species, *K. milleri*, was not classified within the IUCN Red List as threatened by climate change. Other high-ranking species that were not recognized as threatened by climate change were Malaga chub (*Squalius malacitanus*), Almiri toothcarp (*Aphanius almiriensis*), and Trichonis dwarf goby (*Economidichthys trichonis*). Species with the lowest ranking scores, i.e. with low climate change susceptibility, were humpback whitefish (*Coregonus pidschian*), Arctic flounder (*Liopsetta glacialis*), northern pike (*Esox lucius*), burbot (*Lota lota*), and European perch (*Perca fluviatilis*). A complete list of all species' rankings is presented in Supplementary material S5.

Discussion

In the present study, significant differences in life-history and climatic niches were observed between the European freshwater species susceptible to climate change and those that are not, such as species body size, range size, distribution and thermal envelopes. The

latitude of the species range centroid was by far the most influential trait. Overall, southern regions with the warmer, Mediterranean climate comprised a higher proportion of species susceptible to climate change (Figure 4). These results support recent findings that the species from lower latitudes and tropical, warm-water habitats, are at greater risk from climate change and warming (Payne & Smith, 2016; Payne et al., 2016; Comte & Olden, 2017b). In such species, evolved towards higher upper thermal tolerances, specialization to thermal extremes is accompanied by a reduced physiological flexibility and adaptation capacity to respond to changing environmental conditions (Payne & Smith, 2016; Payne et al., 2016; Comte & Olden, 2017b). Such heat-tolerant species are also adapted to temperatures close to their physiological thermal limits, with a narrow safety margin for further temperature increases (Sinclair et al., 2016; Comte & Olden, 2017b). Freshwater basins in Southern Europe are also of particular conservation concern due to an elevated pressure by a range of anthropogenic impacts that further exacerbate effects of climate change, such as pollution, water resource development and consumption, and biological invasions (Xenopoulos et al., 2005; Clavero & García-Berthou, 2005; Walther et al., 2009; Vörösmarty et al., 2010; Comte & Olden, 2017a).

Climate-susceptible species were also characterized by a smaller body and range size (Figures 2, 3). These traits, which are also related to a lower dispersal ability (Radinger & Wolter, 2014), are well recognized as predictors of climate change susceptibility in fish (e.g. Ficke et al., 2007; Isaak & Rieman, 2013; Chessman, 2013; Pearson et al., 2014; Radinger et al., 2017). Smaller-bodied fish are facing elevated overall extinction risk in freshwater habitat due to multiple threats, such as habitat loss and fragmentation (Olden et al., 2007; Kalinkat et al., 2017; Kopf et al., 2017), which explains higher threat level observed in climate-susceptible species in the present study. Observed lower commercial relevance and lower vulnerability to fishing of climate-susceptible species both stem from a lower body size and

related faster life history of such species.

It is important to acknowledge certain limitations of the data sources used in this study, such as species and trait coverage, reliability of methods applied for threat and extinction risk classification, and potential assessors' biases (Clavero & García-Berthou, 2005; Keith et al., 2014; Trull et al., 2018). Furthermore, species that are not classified within IUCN Red List as threatened by climate change can comprise also those that are not yet assessed for their major threats. Nevertheless, the focus of our study on a well-studied group such as European species ensured that the basic life history data and IUCN Red List assessments were mostly available (Kopf et al., 2017). IUCN Red List is sometimes considered to understate or improperly account for climate change as a threat, mostly due to ambiguous definitions and criteria (Trull et al., 2018). However, recent studies have indicated that the IUCN classification is more efficient in detecting species vulnerable to climate change than anticipated (Keith et al., 2014; Pearson et al., 2014). Notwithstanding all the caveats, the databases used in the present study still represent the most comprehensive sources of data and the best available knowledge (Olden et al., 2007; Vega et al., 2017).

Trait-based risk assessments are increasingly used for species profiling (Pacifi et al., 2015; Liu et al., 2017; MacLean & Beissinger, 2017). The approach presented in this study might be considered a valid and promising approach to be used as a screening tool, i.e. to quickly assess large groups of species regarding their susceptibility to climate change and other threats based on species traits, and to identify research and management priorities. Our results indicate that the European environmental policy related to climate change mitigation and adaptation (EEA, 2012, 2017) should be mainly focused on Mediterranean region. This is especially important since this region is also predicted to have the highest frequency of droughts and extreme high temperatures, strongest reduction in precipitation and river

discharges, the highest aggregate climate change impact and the lowest adaptation capacity (Milly et al., 2005; Dankers & Feyen, 2008; Fischer & Schär, 2010; ESPON Climate, 2011; Stagge et al., 2011; Rojas et al., 2012; Jacob et al., 2014; Russo et al., 2014). Moreover, Mediterranean region was also identified as a European priority area regarding climate change impacts for other species groups. A similar distributional pattern of species susceptible to climate change was previously also reported for aquatic insects such as Plecoptera, Ephemeroptera and Trichoptera (Hering et al., 2009; Conti et al., 2014), mammals (Levinsky et al., 2007), as well as for terrestrial species in general (Pacifici et al., 2015).

Species ranking conducted here indicated priority species for further research and monitoring regarding climate change (e.g. *V. letourneuxi*, *I. almacai* and *B. euboicus*; Table 2). Moreover, it also identified species whose IUCN Red List status potentially needs to be reconsidered or updated, such as highly ranked but apparently non-susceptible species (e.g. *K. milleri*), or highly ranked species without a proper threat category (e.g. *K. goerneri*, classified as Data Deficient species). As such, it has a potential to be used as a “Robin Hood Approach” (Punt et al., 2011), where assessments based on information-rich species are used to evaluate and categorize those that are information-poor. There is a need for climate change vulnerability assessments that would be based on quantitative approaches and consistent set of criteria, such as trait-based approaches advocated by IUCN (Foden et al., 2013; Trull et al., 2018). The approach presented here could be a good step in that direction.

Acknowledgements

IJ acknowledges the sponsorship provided by the The J. E. Purkyně Fellowship of the Academy of Sciences of the Czech Republic and funding from the European Union’s Horizon 2020 research and innovation programme through the project ClimeFish (grant No. 677039),

as well as funding provided by the Alexander von Humboldt Foundation and the German Federal Ministry of Education and Research (BMBF). GK acknowledges funding from the BMBF through the project “GLANCE” (Global Change Effects in River Ecosystems; 01 LN1320A). JR acknowledges funding through the 2015-2016 BiodivERsA COFUND call for research proposals and the Spanish Ministry of Economy, Industry and Competitiveness (project ODYSSEUS, BiodivERsA3-2015-26, PCIN-2016-168). GC acknowledges funding from the Ministry of Education, Science and Technological Development, Republic of Serbia (grant number 173045). Thanks are extended to Ryder Burt for assistance with geographic information processing.

References

- Bland, L. M., & Böhm, M. (2016). Overcoming data deficiency in reptiles. *Biological Conservation*, 204, 16-22.
- Boettiger, C., Lang, D. T., & Wainwright, P. (2012). rfishbase: exploring, manipulating and visualizing FishBase data from R. *Journal of Fish Biology*, 81, 2030–2039.
- Boettiger, C., Chamberlain, S., Lang, D. T., & Wainwright, P. (2017). Package ‘rfishbase’. R Interface to 'FishBase'. Version 2.1.2. <https://github.com/ropensci/rfishbase>. Accessed 15 January 2018
- Buisson, L., Thuiller, W., Lek, S., Lim, P., & Grenouillet, G. (2008). Climate change hastens the turnover of stream fish assemblages. *Global Change Biology*, 14, 2232–2248.

311 Chessman, B. C. (2013). Identifying species at risk from climate change: traits predict the
312 drought vulnerability of freshwater fishes. *Biological Conservation*, 160, 40-49.

313

314 Clavero, M., & García-Berthou, E. (2005). Invasive species are a leading cause of animal
315 extinctions. *Trends in Ecology & Evolution*, 20, 110.

316

317 Comte, L., & Olden, J. D. (2017a). Climatic vulnerability of the world's freshwater and
318 marine fishes. *Nature Climate Change*, 7, 718.

319

320 Comte, L., & Olden, J. D. (2017b). Evolutionary and environmental determinants of
321 freshwater fish thermal tolerance and plasticity. *Global Change Biology*, 23, 728-736.

322

323 Comte, L., & Olden, J. D. (2018). Evidence for dispersal syndromes in freshwater fishes.
324 *Proceedings of the Royal Society B*, 285, 20172214.

325

326 Conti, L., Schmidt-Kloiber, A., Grenouillet, G., & Graf, W. (2014). A trait-based approach to
327 assess the vulnerability of European aquatic insects to climate change. *Hydrobiologia*, 721,
328 297-315.

329

330 Dankers, R., & Feyen, L. (2008). Climate change impact on flood hazard in Europe: An
331 assessment based on high resolution climate simulations. *Journal of Geophysical Research*,
332 113, D19105.

333

334 Dempson, B., Schwarz, C. J., Bradbury, I. R., Robertson, M. J., Veinott, G., Poole, R., &

335 Colbourne, E. (2017). Influence of climate and abundance on migration timing of adult
336 Atlantic salmon (*Salmo salar*) among rivers in Newfoundland and Labrador. *Ecology of*
337 *Freshwater Fish*, 26, 247-259.

338

339 EEA (2012). *Climate Change, Impacts and Vulnerability in Europe 2012: An Indicator-Based*
340 *Report*. EEA Report No 12/2012. Copenhagen: European Environment Agency.

341

342 EEA (2017). *Climate Change, Impacts and Vulnerability in Europe 2016: An Indicator-Based*
343 *Report*. EEA Report No 1/2017. Copenhagen: European Environment Agency.

344

345 Elith, J., Leathwick, J. R., & Hastie, T. (2008). A working guide to boosted regression trees.
346 *Journal of Animal Ecology*, 77, 802–813.

347

348 ESPON Climate (2011). *Climate Change and Territorial Effects on Regions and Local*
349 *Economies (Scientific Report)*. Institute of Spatial Planning (IRPUD), TU Dortmund
350 University. http://www.espon.eu/main/Menu_Projects/Menu_AppliedResearch/climate.html.
351 Accessed 20 June 2018

352

353 Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate
354 change on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, 17, 581-613.

355

356 Fischer, E. M., & Schär, C. (2010). Consistent geographical patterns of changes in high-
357 impact European heatwaves. *Nature Geoscience*, 3, 398-403.

358

359 Floeter, S. R., Bender, M. G., Siqueira, A. C., & Cowman, P. F. (2018). Phylogenetic
360 perspectives on reef fish functional traits. *Biological Reviews*, 93, 131-151.
361

362 Foden, W. B., Butchart, S. H., Stuart, S. N., Vié, J. C., Akçakaya, H. R., Angulo, A., ... &
363 Donner, S. D. (2013). Identifying the world's most climate change vulnerable species: a
364 systematic trait-based assessment of all birds, amphibians and corals. *PloS One*, 8, e65427.
365

366 Froese, R., & Pauly, D. (Eds.) (2017). FishBase. World Wide Web electronic publication.
367 version 10/2017. <https://www.fishbase.org>. Accessed 15 January 2018
368

369 Graham, C. T., & Harrod, C. (2009). Implications of climate change for the fishes of the
370 British Isles. *Journal of Fish Biology*, 74, 1143-1205.
371

372 Harrod, C. (2016). Climate change and freshwater fisheries. In J. F. Craig (Ed.), *Freshwater*
373 *Fisheries Ecology* (pp. 641-694). John Wiley & Sons.
374

375 Hering, D., Schmidt-Kloiber, A., Murphy, J., Lücke, S., Zamora-Munoz, C., López-
376 Rodríguez, M. J., ... & Graf, W. (2009). Potential impact of climate change on aquatic insects:
377 a sensitivity analysis for European caddisflies (Trichoptera) based on distribution patterns and
378 ecological preferences. *Aquatic Sciences*, 71, 3-14.
379

380 Hijmans, R. J. (2017). Raster: Geographic Data Analysis and Modeling. R package version
381 2.6-7. <https://CRAN.R-project.org/package=raster>. Accessed 20 June 2018
382

383 Hijmans, R. J., & Elith, J. (2017). Species distribution modeling with R. R vignette.
384 <https://cran.r-project.org/web/packages/dismo/vignettes/sdm.pdf>. Accessed 20 June 2018
385
386 Hijmans, R. J., Phillips, S., Leathwick, J., & Elith, J. (2017). Dismo: Species distribution
387 modeling. <http://cran.r-project.org/package=dismo>. Accessed 20 June 2018
388
389 Hovel, R. A., Carlson, S. M., & Quinn, T. P. (2017). Climate change alters the reproductive
390 phenology and investment of a lacustrine fish, the three-spine stickleback. *Global Change*
391 *Biology*, 23, 2308-2320.
392
393 Isaak, D. J., & Rieman, B. E. (2013). Stream isotherm shifts from climate change and
394 implications for distributions of ectothermic organisms. *Global Change Biology*, 19, 742-751.
395
396 IUCN (2017). The IUCN Red List of Threatened Species. Version 2017-1.
397 <https://www.iucnredlist.org>. Accessed 15 January 2018
398
399 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., ... &
400 Georgopoulou, E. (2014). EURO-CORDEX: new high-resolution climate change projections
401 for European impact research. *Regional Environmental Change*, 14, 563-578.
402
403 Jiguet, F., Gadot, A. S., Julliard, R., Newson, S. E., & Couvet, D. (2007). Climate envelope,
404 life history traits and the resilience of birds facing global change. *Global Change Biology*, 13,
405 1672-1684.
406

- 407 Kalinkat, G., Jähnig, S. C., & Jeschke, J. M. (2017). Exceptional body size–extinction risk
408 relations shed new light on the freshwater biodiversity crisis. *Proceedings of the National*
409 *Academy of Sciences*, 201717087.
- 410
- 411 Keith, D. A., Mahony, M., Hines, H., Elith, J., Regan, T. J., Baumgartner, J. B., ... & Penman,
412 T. (2014). Detecting extinction risk from climate change by IUCN Red List criteria.
413 *Conservation Biology*, 28, 810-819.
- 414
- 415 Kopf, R. K., Shaw, C., & Humphries, P. (2017). Trait based prediction of extinction risk of
416 small bodied freshwater fishes. *Conservation Biology*, 31, 581-591.
- 417
- 418 Levinsky, I., Skov, F., Svenning, J. C., & Rahbek, C. (2007). Potential impacts of climate
419 change on the distributions and diversity patterns of European mammals. *Biodiversity and*
420 *Conservation*, 16, 3803-3816.
- 421
- 422 Liu, C., Comte, L., & Olden, J. D. (2017). Heads you win, tails you lose: Life history traits
423 predict invasion and extinction risk of the world's freshwater fishes. *Aquatic Conservation:*
424 *Marine and Freshwater Ecosystems*, 27, 773-779.
- 425
- 426 MacLean, S. A., & Beissinger, S. R. (2017). Species' traits as predictors of range shifts under
427 contemporary climate change: a review and meta analysis. *Global Change Biology*, 23,
428 4094-4105.
- 429
- 430 McDonald-Madden, E., Chadès, I., McCarthy, M. A., Linkie, M., & Possingham, H. P. (2011).

431 Allocating conservation resources between areas where persistence of a species is uncertain.
432 *Ecological Applications*, 21, 844-858.
433
434 Milly, P. C., Dunne, K. A., & Vecchia, A. V. (2005). Global pattern of trends in streamflow
435 and water availability in a changing climate. *Nature*, 438, 347-350.
436
437 Olden, J. D., Hogan, Z. S., & Zanden, M. J. V. (2007). Small fish, big fish, red fish, blue fish:
438 size-biased extinction risk of the world's freshwater and marine fishes. *Global Ecology and*
439 *Biogeography*, 16, 694-701.
440
441 O'Reilly, C. M., Alin, S. R., Plisnier, P. D., Cohen, A. S., & McKee, B. A. (2003). Climate
442 change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature*, 424,
443 766-768.
444
445 Otero, J., L'Abée Lund, J. H., Castro-Santos, T., Leonardsson, K., Storvik, G. O., Jonsson,
446 B., ... & Dionne, M. (2014). Basin scale phenology and effects of climate variability on
447 global timing of initial seaward migration of Atlantic salmon (*Salmo salar*). *Global Change*
448 *Biology*, 20, 61-75.
449
450 Pacifici, M., Foden, W. B., Visconti, P., Watson, J. E., Butchart, S. H., Kovacs, K. M., ... &
451 Corlett, R. T. (2015). Assessing species vulnerability to climate change. *Nature Climate*
452 *Change*, 5, 215-225.
453
454 Payne, N. L., & Smith, J. A. (2017). An alternative explanation for global trends in thermal

455 tolerance. *Ecology Letters*, 20, 70-77.

456

457 Payne, N. L., Smith, J. A., Meulen, D. E., Taylor, M. D., Watanabe, Y. Y., Takahashi, A., ... &

458 Suthers, I. M. (2016). Temperature dependence of fish performance in the wild: links with

459 species biogeography and physiological thermal tolerance. *Functional Ecology*, 30, 903-912.

460

461 Pearson, R. G., Stanton, J. C., Shoemaker, K. T., Aiello-Lammens, M. E., Ersts, P. J., Horning,

462 N., ... & Akçakaya, H. R. (2014). Life history and spatial traits predict extinction risk due to

463 climate change. *Nature Climate Change*, 4, 217-221.

464

465 Peng, R. D. (2008). Simple bootstrap routines. <https://cran.r-project.org/package=simpleboot>.

466 Accessed 20 June 2018

467

468 Pörtner, H. O., & Farrell, A. P. (2008). Physiology and climate change. *Science*, 322, 690-692.

469

470 Punt, A. E., Smith, D. C., & Smith, A. D. (2011). Among-stock comparisons for improving

471 stock assessments of data-poor stocks: the “Robin Hood” approach. *ICES Journal of Marine*

472 *Science*, 68, 972-981.

473

474 R Development Core Team (2017). R: A language and environment for statistical computing.

475 R Foundation for Statistical Computing, Vienna, Austria.

476

477 Radinger, J., & Wolter, C. (2014). Patterns and predictors of fish dispersal in rivers. *Fish and*

478 *Fisheries*, 15, 456-473.

479

480 Radinger, J., Wolter, C., & Kail, J. (2015). Spatial Scaling of Environmental Variables
481 Improves Species-Habitat Models of Fishes in a Small, Sand-Bed Lowland River. *PLoS One*,
482 10, e0142813.

483

484 Radinger, J., Hölker, F., Horký, P., Slavík, O., Dendoncker, N., & Wolter, C. (2016).
485 Synergistic and antagonistic interactions of future land use and climate change on river fish
486 assemblages. *Global Change Biology*, 22, 1505-1522.

487

488 Radinger, J., Essl, F., Hölker, F., Horký, P., Slavík, O., & Wolter, C. (2017). The future
489 distribution of river fish: the complex interplay of climate and land use changes, species
490 dispersal and movement barriers. *Global Change Biology*, 23, 4970-4986.

491

492 Raptis, C. E., Boucher, J. M., & Pfister, S. (2017). Assessing the environmental impacts of
493 freshwater thermal pollution from global power generation in LCA. *Science of the Total*
494 *Environment*, 580, 1014-1026.

495

496 Rojas, R., Feyen, L., Bianchi, A., & Dosio, A. (2012). Assessment of future flood hazard in
497 Europe using a large ensemble of bias corrected regional climate simulations. *Journal of*
498 *Geophysical Research*, 117, D17109.

499

500 Russo, S., Dosio, A., Graversen, R. G., Sillmann, J., Carrao, H., Dunbar, M. B., ... & Vogt, J.
501 V. (2014). Magnitude of extreme heat waves in present climate and their projection in a
502 warming world. *Journal of Geophysical Research: Atmospheres*, 119, 12500-12512.

503

504 Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., ... &

505 Huey, R. B. (2016). Can we predict ectotherm responses to climate change using thermal

506 performance curves and body temperatures?. *Ecology Letters*, 19, 1372-1385.

507

508 Stagge, J. H., Rizzi, J., Tallaksen, L. M., & Stahl, K. (2015). Future meteorological drought:

509 projections of regional climate models for Europe. In *EGU General Assembly Conference*

510 *Abstracts* (Vol. 17). Technical Report No. 25, DROUGHT-R&SPI, European Commission.

511

512 Trull, N., Böhm, M., & Carr, J. (2018). Patterns and biases of climate change threats in the

513 IUCN Red List. *Conservation Biology*, 32, 135-147.

514

515 USGS (2010). Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010). U.S.

516 Geological Survey (USGS). https://topotools.cr.usgs.gov/gmted_viewer. Accessed 15 January

517 2018

518

519 Vega, G. C., Pertierra, L. R., & Olalla-Tárraga, M. Á. (2017). MERRAclim, a high-resolution

520 global dataset of remotely sensed bioclimatic variables for ecological modelling. *Scientific*

521 *Data*, 4, 170078.

522

523 Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ...

524 & Davies, P. M. (2010). Global threats to human water security and river biodiversity. *Nature*,

525 467, 555-561.

526

527 Walther, G. R., Roques, A., Hulme, P. E., Sykes, M. T., Pyšek, P., Kühn, I., ... & Czucz, B.
528 (2009). Alien species in a warmer world: risks and opportunities. *Trends in Ecology and*
529 *Evolution*, 24, 686-693.

530

531 WCS, & CIESIN (2005). Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human
532 Footprint Dataset (IGHP). Wildlife Conservation Society (WCS) and and Center for
533 International Earth Science Information Network (CIESIN), Columbia University. Palisades,
534 NY: NASA Socioeconomic Data and Applications Center (SEDAC).
535 <http://dx.doi.org/10.7927/H4GF0RFQ>. Accessed 15 January 2018

536

537 WRI (2006). World Resources Institute (WRI) Major Watersheds of the World Delineation.
538 FAO GeoNetwork, FAO - Aquaculture Management and Conservation Service (FIMA).
539 <http://www.fao.org/geonetwork/srv/en/metadata.show?id=30914>. Accessed 15 January 2018

540

541 Xenopoulos, M. A., Lodge, D. M., Alcamo, J., Märker, M., Schulze, K., & Van Vuuren, D. P.
542 (2005). Scenarios of freshwater fish extinctions from climate change and water withdrawal.
543 *Global Change Biology*, 11, 1557-1564.

544

545

546

547

548

549

550

Table 1. Variables used in the analysis, with their data sources, general descriptive statistics and coverage (proportion of species with available data). See Supplementary material I for more information.

	Variable name	Data source	Median; mean $\pm$ SD (range); proportions (%) for categorical data	Coverage (%)
1	Game fish	FishBase	Yes: 14, No: 86	100
2	IUCN Red List status	FishBase/IUCN	EX: 3, CR: 12, EN: 11, VU: 16, NT: 4, LC: 48, DD: 6	100
3	Climate zone (Köppen climate classification)	FishBase	Subtropical: 26, temperate 73, polar: 1	100
4	Preferred habitat	FishBase	Pelagic: 3, pelagic-neritic: 3, benthopelagic: 64, demersal: 30	100
5	Minimum value of the water depth range (m)	FishBase	2; 11.2 $\pm$ 17.9 (0 - 100)	18
6	Maximum value of the water depth range (m)	FishBase	80; 106.7 $\pm$ 138.8 (1 - 700)	14
7	Freshwater preference	FishBase	Exclusively freshwater: 75, enters saltwater: 25	100
8	Maximum recorded body length (cm)	FishBase	21.7; 35.2 $\pm$ 60.5 (2.2 - 800)	98
9	Lateral body shape type	FishBase	Eel-like: 1, elongated: 38, fusiform/normal: 59, short and/or deep: 1	77
10	Aspect ratio of the caudal fin	FishBase	1.6; 1.6 $\pm$ 0.6 (0.4 - 3.4)	34
11	Trophic level	FishBase	3.3; 3.4 $\pm$ 0.6 (2.1 - 4.5)	13
12	Batch spawner	FishBase	Yes: 12, no: 88	55
13	Reproductive guild (first classification)	FishBase	Bearers: 1, guarders: 18, nonguarders: 81	53
14	Reproductive guild (second classification)	FishBase	Brood hiders: 10, clutch tenders: 5, external brooders: 10, internal live bearers: 2, nesters: 18, open water/substratum egg scatterers: 57	28
15	Maximum recorded longevity (years)	FishBase	9; 13.8 $\pm$ 17.4 (1 - 118)	30
16	Commercial importance	FishBase	Of no interest: 41, subsistence fisheries: 4, minor commercial: 15, commercial: 35, highly commercial: 5	40
17	Average global landings/production	FishBase	828.4; 10461.4 $\pm$ 23020.9 (0 - 104902.8)	12
18	Resilience to fishing pressure	FishBase	1: 3, 2: 12, 3: 58, 4: 27	100
19	Vulnerability to fishing	FishBase	32.9; 36.5 $\pm$ 16.6 (10 - 88.7)	100
20	Temperature tolerance (max - min reported T, °C)	FishBase	15; 14.4 $\pm$ 5.2 (1 - 32)	21
21	Number of inhabited freshwater basins	WRI (2006)/IUCN(range)	2; 8.5 $\pm$ 14.6 (0 - 82)	96
22	Global Human Footprint	WCS & CIESIN (2005)/IUCN(range)	6; 7.8 $\pm$ 7.8 (0 - 46)	96
23	Longitude of the centroid of species range	IUCN(range)	18.6; 17.2 $\pm$ 17.4 (-81.1 - 117.2)	96

24	Latitude of the centroid of species range	IUCN(range)	44.6; 45.8 ± 6.3 (34.4 - 70.1)	96
25	Range size (km ²)	IUCN(range)	32499; 1180481 ± 3799809 (13 - 35987250)	96
26	Mean elevation within the species range (m)	USGS (2010)/IUCN(range)	590.4; 833.8 ± 652.5 (3.9 - 2373.6)	96
27	Annual mean temperature (°C)	MERRAclim/IUCN(range)	13.5; 13.4 ± 4.1 (-3.7 - 22.7)	96
28	Mean diurnal temperature range (°C)	MERRAclim/IUCN(range)	19.9; 19.5 ± 2.9 (7.2 - 27.0)	96
29	Temperature isothermality (°C)	MERRAclim/IUCN(range)	44.9; 45.3 ± 3.9 (34.5 - 55.0)	96
30	Temperature seasonality (st. dev. x 100) (°C)	MERRAclim/IUCN(range)	780.6; 800.0 ± 180.6 (312.6 - 1515.3)	96
31	Maximum temperature of the warmest month (°C)	MERRAclim/IUCN(range)	35.7; 35.0 ± 4.6 (20.3 - 43.2)	96
32	Minimum temperature of the coldest month (°C)	MERRAclim/IUCN(range)	-8.6; -8.5 ± 7.6 (-36.4 - 8.8)	96
33	Temperature annual range (°C)	MERRAclim/IUCN(range)	43.3; 43.5 ± 7.5 (17.1 - 65.9)	96
34	Mean temperature of the most humid quarter (°C)	MERRAclim/IUCN(range)	22.9; 22.7 ± 3.5 (10.8 - 30.5)	96
35	Mean temperature of the least humid quarter (°C)	MERRAclim/IUCN(range)	4.2; 3.9 ± 5.8 (-19.8 - 16.5)	96
36	Mean temperature of the warmest quarter (°C)	MERRAclim/IUCN(range)	23.2; 23.1 ± 3.7 (11.0 - 31.0)	96
37	Mean temperature of the coldest quarter (°C)	MERRAclim/IUCN(range)	3.9; 3.6 ± 5.7 (-19.9 - 15.6)	96
38	Annual mean specific humidity (g of water / kg of air)	MERRAclim/IUCN(range)	7.2; 7.2 ± 0.9 (3.5 - 10.1)	96
39	Specific humidity of the most humid month (g of water / kg of air)	MERRAclim/IUCN(range)	11.2; 11.1 ± 1.0 (6.7 - 14.3)	96
40	Specific humidity of the least humid month (g water / kg air)	MERRAclim/IUCN(range)	4.0; 4.0 ± 1.1 (0.8 - 7.1)	96
41	Specific humidity seasonality (g water / kg air)	MERRAclim/IUCN(range)	255.4; 249.7 ± 52.3 (123.0 - 395.5)	96
42	Specific humidity of the most humid quarter (g water / kg air)	MERRAclim/IUCN(range)	10.4; 10.4 ± 1.0 (6.2 - 13.6)	96
43	Specific humidity of the least humid quarter (g water / kg air)	MERRAclim/IUCN(range)	4.5; 4.4 ± 1.1 (1.0 - 7.5)	96
44	Specific humidity of the warmest quarter (g water / kg air)	MERRAclim/IUCN(range)	10.3; 10.3 ± 1.0 (6.2 - 13.3)	96
45	Specific humidity of the coldest quarter (g water / kg air)	MERRAclim/IUCN(range)	4.5; 4.4 ± 1.2 (1.0 - 7.6)	96

Table 2. European freshwater fish species with the highest ranking scores, estimated based on the association of their traits with the climate change susceptibility characteristics, as indicated by the BRT model. Complete ranking list of all species is presented in Supplementary material 5.

Rank	Species	Climate change susceptibility	IUCN Red List category	Ranking score
1	<i>Knipowitschia milleri</i>	non-susceptible	CR	82.5
2	<i>Valencia letourneuxi</i>	susceptible	CR	82.1
3	<i>Iberochondrostoma almakai</i>	susceptible	CR	81.6
4	<i>Barbus euboicus</i>	susceptible	CR	81.4
5	<i>Squalius malacitanus</i>	non-susceptible	EN	80.9
6	<i>Aphanius baeticus</i>	susceptible	EN	80.7
7	<i>Knipowitschia goerneri</i>	susceptible	DD	80.5
8	<i>Squalius keadicus</i>	susceptible	EN	80.5
9	<i>Pelagus laconicus</i>	susceptible	CR	80.4
10	<i>Tropidophoxinellus spartiaticus</i>	susceptible	VU	80.3
11	<i>Iberocypris palaciosi</i>	susceptible	CR	80.1
12	<i>Pelagus epiroticus</i>	susceptible	CR	80.0
13	<i>Aphanius almiriensis</i>	non-susceptible	CR	79.8
14	<i>Anaocypris hispanica</i>	susceptible	EN	79.5
15	<i>Salapia economidisi</i>	susceptible	CR	79.5
16	<i>Squalius torgalensis</i>	susceptible	EN	79.4
17	<i>Cobitis trichonica</i>	susceptible	EN	78.9
18	<i>Valencia hispanica</i>	susceptible	CR	78.6
19	<i>Economidichthys trichonis</i>	non-susceptible	EN	78.6
20	<i>Knipowitschia thessala</i>	susceptible	EN	78.5

Figure captions:

Figure 1. Variables selected by the boosted regression tree (BRT) model as the most relevant descriptors of climate change susceptibility in European freshwater fish species; 20 most relevant variables are presented, which together account for 90% of the total relative variable influence.

Figure 2. Violin-boxplots and barplots of the most relevant spatial variables in European freshwater fish species indicated as either susceptible (n = 148) or non-susceptible (n = 295) to climate change. Habitat preference: blue - exclusively freshwater species, green - species that also enter saltwater.

Figure 3. Violin-boxplots and barplots of the most relevant life history traits and variables related to threat and commercial status in European freshwater fish species indicated as either susceptible (n = 148) or non-susceptible (n = 295) to climate change.

Figure 4. Richness of freshwater fish species across Europe indicated as either susceptible (middle panel) or non-susceptible (left panel) to climate change, and the relative share of susceptible species in the local total species richness (right panel).

Number of non-susceptible species

Number of susceptible species

Share (%) of susceptible species