Plos Genetics

Genetic heterogeneity in *Anopheles darlingi* related to biting behavior in western Amazon

Melina Campos^a

Diego Peres Alonso^a

Jan E. Conn^{b,c}

Joseph M. Vinetz^{d, e}

Kevin J. Emerson^f

Paulo Eduardo Martins Ribolla^{a *}

^aBiotechnology Institute (IBTEC) & Biosciences Institute at Botucatu (IBB), Sao Paulo State University (UNESP), Sao Paulo, Brazil; ^bDepartment of Biomedical Sciences, School of Public Health, University at Albany (State University of New York), Albany, NY; ^cWadsworth Center, New York State Department of Health, Albany, NY; ^dDivision of Infectious Diseases, Department of Medicine, University of California, San Diego, La Jolla, California, USA; ^eInstituto de Medicina Tropical “Alexander von Humboldt,” and Departamento de Ciencias Celulares y Moleculares, Laboratorio de Investigación y Desarrollo, Universidad Peruana Cayetano Heredia, Lima, Peru; and ^fBiology Department, St. Mary’s College of Maryland, St. Mary’s City, MD, USA.

* Corresponding author: PEMR Biotechnology Institute from Sao Paulo State University (UNESP), Alameda das Tecomarias, s/n, Chácara Capão Bonito, Botucatu, SP, Brazil 18607-440.

ABSTRACT

In the Amazon Basin, *Anopheles (Nyssorhynchus) darlingi* is the most aggressive and effective malaria vector. In endemic areas, behavioral aspects of anopheline species such as host preference, biting time and resting location after a blood meal have a key impact on malaria transmission dynamics and transmission control strategies. *An. darlingi* present a variety in behavior throughout its broad distribution including blood feeding related. To investigate the genetic basis of its biting behaviors, host-seeking *An. darlingi* were collected in two settlements (Granada and Remansinho) in Acre, Brazil. Mosquitoes were classified by captured location (indoors or outdoors) and time (dusk or dawn). Genome-wide SNPs were used to assess the degree of genetic diversity and structure in these groups. There was evidence of genetic component of biting behavior regarding both location and time in this species. This study supports that *An. darlingi* blood-feeding behavior has a genetic component. Additional ecological and genomic studies may help to understand the genetic basis of mosquito behavior and address appropriate surveillance and vector control.

Keywords: *Anopheles darlingi*, biting behavior, genetic population, malaria vector, SNPs, genetic heterogeneity

Author Summary

Malaria is a disease caused by parasite of the genus *Plasmodium* and is transmitted by mosquitoes of the genus *Anopheles*. In the Amazon Basin, the main malaria vector is *Anopheles darlingi*, which is present in high densities in this region. Egg development requires that females of this mosquito seek hosts for blood meals. *Anopheles* females blood feeding may occur indoor or outdoor the houses and typically from the sunset to dawn. *Anopheles darlingi* in particular present great variability regarding its behaviour, presenting variety of peak biting times and patterns. This work shows that there is a genetic component that partially explains these two behaviors: location of the blood meal (inside or outside the houses) and time of feeding. Single nucleotide polymorphisms (SNPs) scattered throughout the genome of *Anopheles darlingi* showed genetic diversity and structure in these groups. A comprehensive understanding of the genetic basis for mosquito behaviour may support innovative vector surveillance and control strategies.

Introduction

Anopheles (Nyssorhynchus) darlingi is the main neotropical malaria vector in the Amazon Basin due to its role in human *Plasmodium* transmission [1,2]. Following a continued decrease in the number of malaria cases in America from 2005 to 2014, an increase was observed in the last three years in which Brazil was the major contributor [3,4]. In Brazil, transmission remains entrenched in the Amazon Basin, which accounts for 99.5% of the country's malaria burden [5]. Human migration to this region over the past century has been accompanied by dramatic environmental modifications and the promotion of malaria transmission [6–8]. Deforestation and anthropogenic land changes are known to be associated with increases in *An. darlingi* presence and abundance and thus malaria risk [9–11].

Root (1926) first described *An. darlingi* based on morphological characters of the egg, fourth-instar larva, pupa, male and female adults. Thenceforth, successive studies reported morphological, behavioral, ecological and genetic heterogeneity throughout its broad distribution from Central to South America [12–15]. Despite of that, *An. darlingi* was considered as monophyletic species until Emerson and collaborators [16] proposed the occurrence of three putative species based on well-supported genetic clusters. In that study, Neotropical biogeographical events explained dispersal of *An. darlingi* population, in a continental scale, leading to diversification within this species. However, additional genetic diversity in a minor scale has been shown for this vector by ecological differences [17,18] and seasonality [19].

With regard to mosquito behavior, two aspects are important to determine pathogens transmission to human: anthropophily and endophily. The first is the predilection of the vector to blood-feed on humans instead of other animals and, the second is its preference for living close to human habitats. *An. darlingi* has presented persistently high anthropophily in many areas but substantial variation has been reported in behavior activity regarding endophily and endophagy i.e. blood-feed indoors [20,21]. This species has successfully colonized anthropogenic habitats such as fish

tanks, ditches, micro dams and drains besides being found in natural breeding sites such as lakes, rivers and streams [10,22]. Depending on the region, *An. darlingi* was considered essentially exophagic/endophagic or both. It also has presented resting behavior indoors (endophily) or outdoors (exophily) and a wide range of peak of biting times and patterns [23–26].

The drivers of behavioral shifting have been difficult to isolate and pinpoint in anophelines, although a general consensus is that the responses are mainly due to the distribution of long-lasting insecticidal nets (LLINs) and indoor residual spraying (IRS) [27–29]. Vector behavioral heterogeneity could be resulted of genetically differentiated subpopulations or behavioral plasticity i.e. individuals with same genotype adopt different behavior. The present article investigated genetic population diversity in *An. darlingi* at a micro-scale considering biting behavior. More specifically, we explored the genetic basis of this species by blood-feeding location (indoors or outdoors) and time (dusk or dawn) using genome-wide SNPs. Understanding the genetic contribution to mosquito biting behaviors could lead to the development of molecular methods towards vector surveillance and malaria control.

Results

SNP genotyping

A total of 128 individual mosquitoes were sequenced from both settlements, Granada and Remansinho. We considered four collection categories: indoor collection at dusk i.e., 18-21 h; indoors at dawn i.e., 03-06 h; outdoors at dusk and outdoors at dawn (Table 1). From 124,841,110 ddRAD tag sequences (NCBI SRA BioProject PRJNA298241), ~104 million sequences passed several levels of quality filtering and 34.9% (+/- 9.8 SD) of this set of reads was aligned to the *An. darlingi* genome [32]. An average of 10,107 (+/- 4,123 SD) ddRAD loci were genotyped per sample, and 25% presented SNPs (S1 Table). Analyses were conducted in two parts, considering locality of collection (Granada/Remansinho) and place (indoor/outdoor); and time (dusk/dawn) of collection. First, analyses included indoor and outdoor specimens collected at dusk from both settlements, whereas the second analyses included dusk and dawn collection only from Granada. Analysis presented a number of SNPs genotyped in at least 50% of individuals in that set without previous population assignment.

Heterogeneity genetic of *Anopheles darlingi* by biting behavior

Location: Indoor vs. Outdoor

Dataset analyses were conducted for indoor and outdoor samples collected in the dusk in Granada and Remansinho. After filtering, 944 SNPs were found in at least 50% of all individuals from the four groups. Principal component analysis (PCA) showed that indoor specimens were concisely grouped together regardless of the collection location, whereas outdoor individuals were found in two different areas in the plot (Fig 2A). Discriminant analysis of principal component (DAPC) showed $K = 3$ as the best number of genetic *clusters* i.e. the lowest Bayesian Information Criterion (BIC) (Fig 2C). Cluster “1” had only outdoor *An. darlingi* specimens from Granada and cluster “2” had nearly all outdoor individuals from Remansinho, whereas “3” included indoor individuals from both locations (Fig 2B). Indoor groups showed a higher percentage of polymorphism compared with outdoor ones (Table 2). Interestingly, the highest pairwise F_{ST} was between indoor Remansinho and outdoor Granada samples ($F_{ST} = 0.177$, $p <$

0.0001), and the lowest was between indoors samples ($F_{ST} = 0.094$, $p < 0.0001$) (Table 4).

Time: dusk vs. dawn

Four groups of samples, indoors and outdoors collected in the dusk and dawn, were analyzed in this set. A total of 589 SNPs was filtered in at least 50% in dusk/dawn and indoor/outdoor samples. The first axis in the PCA divided indoor and outdoor samples, whereas the second axis separated time of collection (Fig 3A). The lowest BIC for K -means was 4 in the DAPC where essentially the four groups were partitioned (Fig 3B-C). Samples collected in the dawn showed a higher percentage of polymorphic loci compared with dusk ones (Table 3). For this dataset, the highest pairwise F_{ST} was between outdoor dusk and indoor dawn samples ($F_{ST} = 0.259$, $p < 0.0001$), and the lowest was between outdoors samples ($F_{ST} = 0.081$, $p < 0.0001$) (Table 5).

Discussion

Reduced representation genome-sequencing methods, such as ddRAD, have proved powerful to assess a large number of SNPs on a genome-wide scale, with considerable lower library construction and sequencing effort. However, these approaches suffer from sampling biases i.e. allele dropout (ADO), due to the absence or polymorphism on a restriction site [35]. *In silico* simulations showed that pruned datasets with incomplete loci sampling i.e. missing data are more likely to accurately reflect population genetic [36]. Here, we applied two strict filters for all set of analysis: no prior information of population and loci were selected when presented in at least 50% of individuals.

In this study, we investigated the genetic basis of biting behavior regarding location (indoor/outdoor) and time (dusk/dawn) in females of *An. darlingi*. There was evidence of a genetic component for both local and time biting behaviors features using genome-wide SNPs. In the Peruvian Amazon, there was no evidence of genetic differentiation between exophagic and endophagic *An. darlingi* [29]. Hence, this is the first study to report a genetic component related to *An. darlingi* behavior. Previously, studies with other species such as *An. gambiae* s.l. and *An. funestus* showed genetic differentiation regarding behavior using chromosomal inversion frequencies [37,38]. In addition, a genetic component was detected for host choice for blood-feeding in *An. arabiensis* using whole-genome sequencing [39]. However, this species showed no evidence of genetic differentiation for biting and resting behavior [39,40].

Concerning location, females collected outdoor in two different rural settlements (Granada and Remansinho), about 60 km apart, comprised two genetic clusters. This finding corroborated prior reported of microgeographic genetic differentiation studies found in *An. darlingi* by habitat [18]. On the other hand, these two independent places presented a similar indoor population of *An. darlingi* by PCA, DAPC and presented the

lowest pairwise F_{ST} (Fig 2). This single indoor group could be explained by similar selection of *An. darlingi* female related to environmental conditions within houses such as temperature, sheltering and blood meal availability. In fact, Paaijmans and Thomas [41] showed that indoor environments are warmer than outdoor ones and usually presented less daily variation. Besides temperature, indoor residual spraying could be an important drive in the selection of mosquito females. Different reports already showed fast selection of mosquito resistance genes correlated to insecticide usage, either by insecticide-treated bed nets (ITNs) and indoors residual spraying [42–45]. Both rural settlement studied (Granada and Remansinho) have no long history in use of ITNs, but both places usually use indoors residual spraying.

In the matter of the time of biting behavior in *An. darlingi*, the present study showed non-random distribution of individuals in the PCA and clustering in DAPC. As shown in the first set of analysis, PC1 segregated indoors and outdoors individuals (Fig 3). In addition, PC2 segregated dusk and dawn individuals. In fact, DAPC presented four genetic clusters containing samples from each category. Across its broad distribution, *An. darlingi* populations present a range of peak biting times and patterns i.e. unimodal, bimodal, trimodal [21,24]. Biting time variation of *An. darlingi* showed associated with seasonality [19,46] and local and ecological factor [47]. The current study collected *An. darlingi* biting throughout the night with two peaks, one in the dusk and another in the dawn (data not published, supplementary). Prussing and collaborators [29] found no evidence of genetic differentiation between *An. darlingi* biting times during the night in Iquitos, Peru. However, *An. darlingi* have presented especially early-evening biting activity in that area [48,49], which disagreed with the pattern found in the present study. Additional sampling could help to determine whether the differentiation detected in this study is consistent between populations of *An. darlingi* showing similar behavior patterns.

It is widely accepted that behavioral aspects of anopheline species such as host preference, biting time and resting location after a blood meal have a major impact on

malaria transmission dynamics in endemic areas [50,51]. From an evolutionary point of view, it is tempting to assume that during *An. darlingi* colonization of the Amazon region, mosquito subpopulations began to evolve in order to occupy the myriad niches in this diverse environment. Together with recent events directly attributable to human presence such as deforestation and land use changes on a massive scale [9,21], we hypothesize that these processes have led to structured populations of *An. darlingi* that may at least partially explain the strong evidence for genetic and behavioral diversity. Besides deforestation, another important human modification in Amazon region is the construction of small ponds to supplement water all over the year. This permanent water collection near the houses could be an important breeding site to support anophelines populations with close access to human host. A better understanding of the genetic basis for diversity in *An. darlingi* may help to predict and improve vector control methods and effectiveness of malaria frontline interventions.

Materials & Methods

Samples

Collections of *Anopheles darlingi* adults were performed outdoors (peridomestic, within 10m of each house) and indoors in two rural settlements, Granada (9°45'S 67°04'W) and Remansinho (9°29'S 66°34'W) in Acre State in April 2011 (Fig 1). Collections were performed using human landing catch (HLC) by the authors MC and PEMR, between 18-06 h. All collected specimens were identified using the key of Consoli & Lourenço-Oliveira [30] based on morphological characters and conserved individually in eppendorf tubes at -20°C; only *An. darlingi* specimens were used for further analysis.

SNP genotyping

DNA was individually extracted using ReliaPrep™ Blood gDNA kit (Promega, Madison, USA) and DNA concentration was estimated using a Qubit fluorometer (Invitrogen, Carlsbad, USA). Double restriction digestion of 200 ng of high quality genomic DNA with *EcoRI*-*MspI* restriction enzymes was performed in a 40 µl reaction volume and then purified with AMPure XP beads following the manufacturer's protocol (Beckman Coulter, California, USA). Hybridized customized adapters P1 (0.3µM) and P2 (4.8µM) were ligated to the digested DNA (T4 DNA Ligase, Promega) as described in Campos et al. (2017). After another purification with AMPure XP beads, DNA was size selected on an agarose gel to 350-400 bp followed by another AMPure XP bead purification. PCR amplification for Nextera® indexing was carried out to generate Illumina sequencing libraries, according to these cycling conditions: an initial denaturation step at 72°C for 3 min and at 95°C for 30s, followed by 16 cycles of 95°C for 10s, annealing at 55°C for 30s, elongation at 72°C for 30s, and a final extension cycle at 72°C for 5 minutes, then each PCR product was purified one last time. Samples were individually quantified using the KAPA library quantification kit (KAPA Biosystems, Wilmington, USA) and equimolarly combined to compose a final library. Final libraries were quantified, normalized, denatured, and finally loaded on the Illumina reagent

cartridge 150-cycle of paired-end sequencing in a Hiseq 2500 (Genomics and Bioinformatics Core, State University of New York at Buffalo).

SNP data processing

Raw ddRAD reads were processed to identify SNP loci within and between individuals using scripts implemented in Stacks v1.31 [31]. First, sequences were quality filtered using the default parameters of *process_radtags* script. Then, each individual's sequence reads were aligned to the *An. darlingi* reference genome [32] using Bowtie2 with default parameters [33]. Aligned reads were input to *ref_map.pl* Perl script in Stacks, using a minimum of 5 reads (*-m*) to report a stack and join the catalog. The dataset was corrected using another Stacks script called *rxstacks* with the following parameters: prune out non-biological haplotypes unlikely to occur in the population (*--prune_haplo*), minimum log likelihood required was -10.0 (*--lnl_lim*). A new catalog was built by *cstacks* Stacks' script and each individual was matched against the catalog with *sstacks* script. We then used the *populations* script in Stacks to filter loci that were called in at least 50% (*-r*) of all samples i.e., the first step was performed without population information to avoid population bias in the SNP selection. This latter step was run with minimum coverage of 5 (*-m*) and a random single SNP for each RAD locus was selected (*write_random_snp*). The *populations* script produced genotype output in several formats (e.g. VCF, Genepop) and summary statistics such as nucleotide diversity, pairwise F_{ST} and private alleles.

SNP data analysis

Principal component analysis (PCA) and discriminant analysis of principal components (DAPC) were performed in the R package Adegenet [34]. The former described global diversity overlooking group information whereas DAPC maximize differences between groups and minimizes variation within clusters. The optimum number of genetic clusters in DAPC was the lowest Bayesian Information Criterion (BIC) associated with several *K*-means calculated.

Abbreviations

BIC: Bayesian information criterion; DAPC: Discriminant analysis of principal components; ddRADseq: Double digestion restriction-site associated DNA sequencing; PCA: Principal component analysis; SNP: single nucleotide polymorphism.

Author contributions

PEMR and MC designed the field and laboratory work; MC and DPA performed the laboratory research; PEMR, MC and KJE analyzed data. All authors actively contributed to the interpretation of the findings; PEMR, MC, DPA wrote, JEC and KJE revised the manuscript. All authors approved the final manuscript.

Funding

MC was supported by FAPESP. JMV received funding from International Centers for Excellence in Malaria Research (grant U19AI089681). KJE received funding St. Mary's College of Maryland. PEMR has a CNPq fellowship.

References

1. Deane L, Deane MP. Notas sobre a distribuição ea biologia dos anofelinos das regiões nordestina e amazônica do Brasil. Revista do Serviço Especial de Saúde Pública; 1948.
2. Tadei WP, Dutary Thatcher B. Malaria vectors in the Brazilian amazon: Anopheles of the subgenus Nyssorhynchus. Rev Inst Med Trop Sao Paulo. 2000;42: 87–94.
3. World Health Organization. World malaria report 2017 [Internet]. 2017. Available: <http://www.who.int/malaria/publications/world-malaria-report-2017/report/en/>
4. Pan American Health Organization / World Health Organization. Epidemiological Alert: Increase of malaria in the Americas [Internet]. 2018. Available: https://www.paho.org/hq/index.php?option=com_docman&task=doc_view&Itemid=270&gid=43434&lang=en
5. Ferreira MU, Castro MC. Challenges for malaria elimination in Brazil. Malar J. 2016;15: 284.
6. Cruz Marques A. Human migration and the spread of malaria in Brazil. Parasitol Today. 1987;3: 166–170.
7. Deane LM, Daniel Ribeiro C, Lourenço de Oliveira R, Oliveira-Ferreira J, Guimarães AE. Study on the natural history of malaria in areas of the Rondonia State-Brazil and problems related to its control. Rev Inst Med Trop Sao Paulo. 1988;30: 153–156.
8. de Castro MC, Monte-Mór RL, Sawyer DO, Singer BH. Malaria risk on the Amazon frontier. Proc Natl Acad Sci U S A. 2006;103: 2452–2457.
9. Vittor AY, Gilman RH, Tielsch J, Glass G, Shields T, Lozano WS, et al. The effect of deforestation on the human-biting rate of Anopheles darlingi, the primary vector of Falciparum malaria in the Peruvian Amazon. Am J Trop Med Hyg. 2006;74: 3–11.
10. Vittor AY, Pan W, Gilman RH, Tielsch J, Glass G, Shields T, et al. Linking deforestation to malaria in the Amazon: characterization of the breeding habitat of the principal malaria vector, Anopheles darlingi. Am J Trop Med Hyg. 2009;81: 5–12.
11. Hahn MB, Gangnon RE, Barcellos C, Asner GP, Patz JA. Influence of deforestation, logging, and fire on malaria in the Brazilian Amazon. PLoS One. 2014;9: e85725.
12. Lourenço-de-Oliveira R, Guimarães AE da G, Arlé M, Silva TF da, Castro MG, Motta MA, et al. Anopheline species, some of their habits and relation to malaria in endemic areas of Rondonia State, Amazon region of Brazil. Mem Inst Oswaldo Cruz. SciELO Brasil; 1989;84: 501–514.
13. Rosa-Freitas MG, Broomfield G, Priestman A, Milligan PJ, Momen H, Molyneux DH.

- Cuticular hydrocarbons, isoenzymes and behavior of three populations of *Anopheles darlingi* from Brazil. *J Am Mosq Control Assoc.* 1992;8: 357–366.
14. Sinka ME, Rubio-Palis Y, Manguin S, Patil AP, Temperley WH, Gething PW, et al. The dominant *Anopheles* vectors of human malaria in the Americas: occurrence data, distribution maps and bionomic précis. *Parasit Vectors.* 2010;3: 72.
 15. Motoki MT, Suesdek L, Bergo ES, Sallum MAM. Wing geometry of *Anopheles darlingi* Root (Diptera: Culicidae) in five major Brazilian ecoregions. *Infect Genet Evol.* 2012;12: 1246–1252.
 16. Emerson KJ, Conn JE, Bergo ES, Randel MA, Sallum MAM. Brazilian *Anopheles darlingi* Root (Diptera: Culicidae) Clusters by Major Biogeographical Region. *PLoS One.* 2015;10: e0130773.
 17. Lainhart W, Bickersmith SA, Moreno M, Rios CT, Vinetz JM, Conn JE. Changes in Genetic Diversity from Field to Laboratory During Colonization of *Anopheles darlingi* Root (Diptera: Culicidae). *Am J Trop Med Hyg.* 2015;93: 998–1001.
 18. Campos M, Conn JE, Alonso DP, Vinetz JM, Emerson KJ, Ribolla PEM. Microgeographical structure in the major Neotropical malaria vector *Anopheles darlingi* using microsatellites and SNP markers. *Parasit Vectors.* 2017;10: 76.
 19. Angêlla AF, Salgueiro P, Gil LHS, Vicente JL, Pinto J, Ribolla PEM. Seasonal genetic partitioning in the neotropical malaria vector, *Anopheles darlingi*. *Malar J.* 2014;13: 203.
 20. Dusfour I, Carinci R, Gaborit P, Issaly J, Girod R. Evaluation of four methods for collecting malaria vectors in French Guiana. *J Econ Entomol.* 2010;103: 973–976.
 21. Hiwat H, Bretas G. Ecology of *Anopheles darlingi* Root with respect to vector importance: a review. *Parasit Vectors.* 2011;4: 177.
 22. dos Reis IC, Codeço CT, Degener CM, Keppeler EC, Muniz MM, de Oliveira FGS, et al. Contribution of fish farming ponds to the production of immature *Anopheles* spp. in a malaria-endemic Amazonian town. *Malar J.* 2015;14: 452.
 23. Moutinho PR, Gil LHS, Cruz RB, Ribolla PEM. Population dynamics, structure and behavior of *Anopheles darlingi* in a rural settlement in the Amazon rainforest of Acre, Brazil. *Malar J.* 2011;10: 174.
 24. Zimmerman RH, Lounibos LP, Nishimura N, Galardo AKR, Galardo CD, Arruda ME. Nightly biting cycles of malaria vectors in a heterogeneous transmission area of eastern Amazonian Brazil. *Malar J.* 2013;12: 262.
 25. Ahumada ML, Pareja PX, Buitrago LS, Quiñones ML. [Biting behavior of *Anopheles darlingi* Root, 1926 (Diptera: Culicidae) and its association with malaria transmission in Villavicencio (Meta, Colombia)]. *Biomedica.* 2013;33: 241–250.

26. Barbosa LMC, Souto RNP, Dos Anjos Ferreira RM, Scarpassa VM. Behavioral patterns, parity rate and natural infection analysis in anopheline species involved in the transmission of malaria in the northeastern Brazilian Amazon region. *Acta Trop.* 2016;164: 216–225.
27. Elliott R. The influence of vector behavior on malaria transmission. *Am J Trop Med Hyg.* 1972;21: 755–763.
28. Russell TL, Beebe NW, Cooper RD, Lobo NF, Burkot TR. Successful malaria elimination strategies require interventions that target changing vector behaviours. *Malar J.* 2013;12: 56.
29. Prussing C, Moreno M, Saavedra MP, Bickersmith SA, Gamboa D, Alava F, et al. Decreasing proportion of *Anopheles darlingi* biting outdoors between long-lasting insecticidal net distributions in peri-Iquitos, Amazonian Peru. *Malar J.* 2018;17: 86.
30. Consoli RAGB, Rotraut A G, de Oliveira RL. Principais mosquitos de importância sanitária no Brasil. 1994.
31. Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. Stacks: an analysis tool set for population genomics. *Mol Ecol.* 2013;22: 3124–3140.
32. Marinotti O, Cerqueira GC, de Almeida LGP, Ferro MIT, Loreto EL da S, Zaha A, et al. The genome of *Anopheles darlingi*, the main neotropical malaria vector. *Nucleic Acids Res.* 2013;41: 7387–7400.
33. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012;9: 357–359.
34. Jombart T, Ahmed I. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics.* 2011;27: 3070–3071.
35. Gautier M, Gharbi K, Cezard T, Foucaud J, Kerdelhué C, Pudlo P, et al. The effect of RAD allele dropout on the estimation of genetic variation within and between populations. *Mol Ecol.* 2013;22: 3165–3178.
36. Arnold B, Corbett-Detig RB, Hartl D, Bomblies K. RADseq underestimates diversity and introduces genealogical biases due to nonrandom haplotype sampling. *Mol Ecol.* 2013;22: 3179–3190.
37. Coluzzi M, Sabatini A, Petrarca V, Di Deco MA. Chromosomal differentiation and adaptation to human environments in the *Anopheles gambiae* complex. *Trans R Soc Trop Med Hyg.* 1979;73: 483–497.
38. Guelbeogo WM, Sagnon N 'fale, Liu F, Besansky NJ, Costantini C. Behavioural divergence of sympatric *Anopheles funestus* populations in Burkina Faso. *Malar J.* 2014;13: 65.
39. Main BJ, Lee Y, Ferguson HM, Kreppel KS, Kihonda A, Govella NJ, et al. The

Genetic Basis of Host Preference and Resting Behavior in the Major African Malaria Vector, *Anopheles arabiensis*. PLoS Genet. 2016;12: e1006303.

40. Maliti DV, Marsden CD, Main BJ, Govella NJ, Yamasaki Y, Collier TC, et al. Investigating associations between biting time in the malaria vector *Anopheles arabiensis* Patton and single nucleotide polymorphisms in circadian clock genes: support for sub-structure among *An. arabiensis* in the Kilombero valley of Tanzania. Parasit Vectors. BioMed Central; 2016;9: 109.
41. Paaijmans KP, Thomas MB. The influence of mosquito resting behaviour and associated microclimate for malaria risk. Malar J. 2011;10: 183.
42. Matoke-Muhia D, Gimnig JE, Kamau L, Shililu J, Bayoh MN, Walker ED. Decline in frequency of the 2La chromosomal inversion in *Anopheles gambiae* (s.s.) in Western Kenya: correlation with increase in ownership of insecticide-treated bed nets. Parasit Vectors. 2016;9: 334.
43. Abeku TA, Helinski MEH, Kirby MJ, Ssekitooleko J, Bass C, Kyomuhangi I, et al. Insecticide resistance patterns in Uganda and the effect of indoor residual spraying with bendiocarb on *kdr* L1014S frequencies in *Anopheles gambiae* s.s. Malar J. 2017;16: 156.
44. Sougoufara S, Sokhna C, Diagne N, Doucouré S, Sembène PM, Harry M. The implementation of long-lasting insecticidal bed nets has differential effects on the genetic structure of the African malaria vectors in the *Anopheles gambiae* complex in Dielmo, Senegal. Malar J. 2017;16: 337.
45. Vontas J, Grigoraki L, Morgan J, Tsakireli D, Fuseini G, Segura L, et al. Rapid selection of a pyrethroid metabolic enzyme CYP9K1 by operational malaria control activities. Proc Natl Acad Sci U S A. 2018;115: 4619–4624.
46. Jiménez IP, Jiménez IP, Conn JE, Brochero H. Preliminary biological studies on larvae and adult *Anopheles* mosquitoes (Diptera: Culicidae) in Miraflores, a malaria endemic locality in Guaviare department, Amazonian Colombia. J Med Entomol. 2014;51: 1002–1009.
47. Charlwood JD. Biological variation in *Anopheles darlingi* Root. Mem Inst Oswaldo Cruz. 1996;91: 391–398.
48. Turell MJ, Sardelis MR, Jones JW, Watts DM, Fernandez R, Carbajal F, et al. Seasonal distribution, biology, and human attraction patterns of mosquitoes (Diptera: Culicidae) in a rural village and adjacent forested site near Iquitos, Peru. J Med Entomol. 2008;45: 1165–1172.
49. Moreno M, Saavedra MP, Bickersmith SA, Lainhart W, Tong C, Alava F, et al. Implications for changes in *Anopheles darlingi* biting behaviour in three communities in the peri-Iquitos region of Amazonian Peru. Malar J. 2015;14: 290.
50. Russell TL, Lwetoijera DW, Knols BGJ, Takken W, Killeen GF, Ferguson HM.

Linking individual phenotype to density-dependent population growth: the influence of body size on the population dynamics of malaria vectors. *Proc Biol Sci.* 2011;278: 3142–3151.

51. Gatton ML, Chitnis N, Churcher T, Donnelly MJ, Ghani AC, Godfray HCJ, et al. The importance of mosquito behavioural adaptations to malaria control in Africa. *Evolution.* 2013;67: 1218–1230.

Table 1. Number of *Anopheles darlingi* genotyped by settlement, location and time.

Location	Region	State	Latitude	Longitude	Dusk	Outdoor	Indoor	
						Dawn	Dusk	Dawn
Granada	Acrelandia	Acre	-9.752	-67.071	24	30	19	23
Remansinho	Labrea	Amazon	-9.497	-66.582	16	-	16	-

Table 2. Summary statistics for the indoor and outdoor *Anopheles darlingi*.

Location	Collection	Private	Variant	Polymorphic	% Polymorphic
----------	------------	---------	---------	-------------	---------------

	n	e	Sites	Sites	Loci
Granada	Indoor	111	920	469	0.337
	Outdoor	104	937	379	0.267
Remansinho	Indoor	134	938	532	0.375
	Outdoor	96	936	468	0.330

Table 3. Summary statistics for the indoor/outdoor and dusk/dawn *Anopheles darlingi*.

Location	Time	Private	Variant Sites	Polymorphic Sites	% Polymorphic Loci
Indoor	Dusk	55	556	243	0.289
	Dawn	85	557	315	0.374
Outdoor	Dusk	70	556	190	0.226
	Dawn	91	557	291	0.346

Table 4. Pairwise F_{ST} between indoor and outdoor *Anopheles darlingi*.

	Granada Outdoor	Granada Indoor	Remansinho Outdoor
Granada Indoor	0.140		
Remansinho Outdoor	0.146	0.140	

Remansinho	0.177	0.094	0.129
Indoor			

Table 5. Pairwise F_{ST} between indoor/outdoor and dusk/dawn *Anopheles darlingi*.

	Indoor Dawn	Indoor Dusk	Outdoor Dawn
Indoor Dusk	0.155		
Outdoor Dawn	0.134	0.102	
Outdoor Dusk	0.259	0.139	0.081

Figure legends

Figure 1. Collection points in Brazilian Amazon Basin. Map of Brazil showing Amazon domain. Square box: zoom in showing Acre (AC) and Amazonas (AM) states and the two collection localities, Granada and Remansinho.

Figure 2. Principal Component Analysis (PCA) and Discriminant analysis of principal components (DAPC) of outdoor and indoor *An. darlingi* genotypes. (A) PCA of outdoor samples (red and yellow dots) and indoor samples (blue and green dots) from Granada and Remansinho. (B) DAPC - ordination of all samples in the three genetic clusters (1-3) in two axes. (C) Bayesian Information Criterion (BIC) to estimate the appropriate number of genetic clusters, which is the lowest BIC value.

Figure 3. Principal Component Analysis (PCA) and Discriminant analysis of principal components (DAPC) of outdoor/indoor and dusk/dawn *An. darlingi* genotypes. (A) PCA of dusk samples (red and blue dots) and dawn samples (purple and orange dots) from Granada and Remansinho. (B) DAPC - ordination of all samples in the four genetic clusters (1-4) in two axes. (C) Bayesian Information Criterion (BIC) to estimate the appropriate number of genetic clusters, which is the lowest BIC value.

Supporting Information Legends

S1 Table. Per-individual *An. darlingi* detail of the number of sequence reads and unique stacks genotyped.

Figure 1. Collection points in Brazilian Amazon Basin. Map of Brazil showing Amazon domain. Square box: zoom in showing Acre (AC) and Amazonas (AM) states and the two collection localities, Granada and Remansinho.

Figure 2. Principal Component Analysis (PCA) and Discriminant analysis of principal components (DAPC) of outdoor and indoor *An. darlingi* genotypes. (A) PCA of outdoor samples (red and yellow dots) and indoor samples (blue and green dots) from Granada and Remansinho. (B) DAPC - ordination of all samples in the three genetic clusters (1-3) in two axes. (C) Bayesian Information Criterion (BIC) to estimate the appropriate number of genetic clusters, which is the lowest BIC value.

Figure 3. Principal Component Analysis (PCA) and Discriminant analysis of principal components (DAPC) of outdoor/indoor and dusk/dawn *An. darlingi* genotypes. (A) PCA of dusk samples (red and blue dots) and dawn samples (purple and orange dots) from Granada and Remansinho. (B) DAPC - ordination of all samples in the four genetic clusters (1-4) in two axes. (C) Bayesian Information Criterion (BIC) to estimate the appropriate number of genetic clusters, which is the lowest BIC value.