1 Transcription-dependent spreading of canonical yeast GATA factor 2 across the body of highly expressed genes

3 Short title: Transcription factor binding at gene bodies

4

5 Aria Ronsmans^{1,2¶}, Maxime Wery^{3¶}, Camille Gautier³, Marc Describes³, Evelyne Dubois^{1,2},
6 Antonin Morillon^{3*} and Isabelle Georis^{1*}

7

8 ¹ Metabolism of Model Microorganisms, Institut de Recherches Microbiologiques J.M.
9 Wiame - Labiris, Brussels, Belgium

10 ² Laboratoire de Microbiologie, Université Libre de Bruxelles, Brussels, Belgium

11 ³ ncRNA, epigenetic and genome fluidity, Institut Curie, PSL Research University, CNRS
12 UMR 3244, Université Pierre et Marie Curie, Paris, France

13

14 ^{*}Corresponding authors

15 E-mail: : igeoris@ulb.ac.be (IG), antonin.morillon@curie.fr (AM)

16

17 [¶]These authors contributed equally to this work.

18 **Abstract**

19 GATA transcription factors are highly conserved among eukaryotes and play roles in
 20 transcription of genes implicated in cancer progression and hematopoiesis. However, although
 21 their consensus binding sites have been well defined *in vitro*, the *in vivo* selectivity for
 22 recognition by GATA factors remains poorly characterized. Using ChIP-Seq, we identified
 23 the Dal80 GATA factor targets in yeast. Our data reveal Dal80 binding to a large set of
 24 promoters, sometimes independently of GATA sites. Strikingly, Dal80 was also detected
 25 across the body of promoter-bound genes, correlating with high, Dal80-sensitive expression.
 26 Mechanistic single-gene experiments showed that Dal80 spreading across gene bodies is
 27 independent of intragenic GATA sites but requires transcription elongation. Consistently,
 28 Dal80 co-purified with the post-initiation form of RNA Polymerase II. Our work suggests that
 29 GATA factors could play dual, synergistic roles during transcription initiation and post-
 30 initiation steps, promoting efficient remodeling of the gene expression program in response to
 31 environmental changes.

33 **Author Summary**

34 GATA transcription factors are highly conserved among eukaryotes and play key roles in
 35 cancer progression and hematopoiesis. In budding yeast, four GATA transcription factors are
 36 involved in the response to the quality of nitrogen supply. We have determined the whole
 37 genome binding profile of one of them, Dal80, and revealed that it also binds across the body
 38 or promoter-bound genes. Our observation that ORF binding correlated with elevated
 39 transcription levels and exquisite Dal80 sensitivity suggests that GATA factors could play
 40 other, unexpected roles at post-initiation stages in eukaryotes.

41

42 **Introduction**

43 In eukaryotes, gene transcription by RNA polymerase II (PolII) is initiated by the binding of
 44 specific transcription factors to double stranded DNA. The yeast transcription factors target
 45 regulatory regions called UAS or URS (for *Upstream Activating/Repressing Sequences*),
 46 generally directly adjacent to the core promoter. The generated regulatory signals converge at
 47 the core promoter where they permit the regulation of PolII recruitment via the ‘TATA box-
 48 binding protein’ and associated general transcription factors [1,2]. The transcription factor
 49 binding sites are usually short sequences ranging from 8 to 20 bp [3]. They are most often
 50 similar but generally not identical, differing by some nucleotides from one another [3],
 51 making it sometimes difficult to predict whether a given UAS will function as such *in vivo*.
 52 GATA factors constitute a family of transcription factors highly conserved among eukaryotes
 53 and characterized by the presence of a DNA binding domain which consists of four cysteines
 54 (fitting the consensus sequence CX₂CX₁₇₋₁₈CX₂C) coordinating a zinc ion followed by a basic
 55 carboxy-terminal tail [4]. *In vitro*, this conserved binding domain allows the recognition of
 56 consensus GATA DNA sequences (GATAA, GATAAG or GATTAG), found in the promoter
 57 of numerous genes, which was demonstrated in various organisms either with direct or
 58 indirect methods [4–8]. Structure elucidation of GATA factor binding domains in complex
 59 with their DNA target suggest an important role for the basic tail in sequence specificity
 60 determination [4,8].
 61 Since its discovery 40 years ago in chicken cells, the family of GATA factors was extended in
 62 human cells and represents master regulators of hematopoiesis and cancer [9]. However,
 63 although approximately 7 million GATA motifs can be found in the human genome, the
 64 GATA factors occupy only 0.1–1% of them. Conversely, other regions are occupied by
 65 GATA factors despite lacking the consensus motif [10,11]. In addition, GATA factors can

66 swap among them for the same motif and switch from active or repressive transcriptional
67 activity. All these observations raised the main paradigm on understanding the selective
68 recruitment and activity of GATA factors and what the determinants of their residency on the
69 chromatin are [12,13].

70 In yeast, the family of GATA transcription factors contains over 10 members [14]. Four of
71 them are implicated in the regulation of *Nitrogen Catabolite Repression*- (NCR-) sensitive
72 genes, the expression of which is repressed in the presence of a preferred nitrogen source
73 (glutamine, asparagine, ammonia) and derepressed when only poor nitrogen sources (e.g.
74 proline, leucine, urea) are available [15]. These key GATA factors involved in NCR signaling
75 are two activators (Gln3 and Gat1/Nil1) and two repressors (Gzf3/Nil2/Deh1 and
76 Dal80/Uga43) [16–21]. Moreover, the expression of *DAL80* and *GAT1* is also NCR-sensitive,
77 which implies cross- and autogenous regulations of the GATA factors in the NCR
78 mechanisms [21–24]. Under nitrogen limitation, expression of *DAL80* is highly induced [18],
79 and Dal80 enters the nucleus where it competes with the two GATA activators for the same
80 binding sites [5,22,25]. Although initially described as being active under nitrogen abundance
81 [20,21], the Gzf3 repressor also localizes to NCR-sensitive promoters in conditions of
82 activation [23].

83 *In vitro*, the Gln3 and Gat1 activators bind, presumably as monomers, to single GATA
84 sequences [26], like their orthologous vertebrate counterparts. On the other hand, the Dal80
85 repressor can form homo- and heterodimers with Gzf3, thanks to the presence of an N-
86 terminal leucine zipper [27] and, consistently, Dal80 binds *in vitro* to two GATA sequences
87 [5,22,27,28]. *In vivo*, GATA factor binding site recognition also appears to require repeats of
88 GATA motifs within promoters. Indeed, synthetic constructions containing a portion of the
89 NCR-sensitive *DAL5* promoter have shown in a β -galactosidase assay that at least two copies
90 of the consensus are required to support efficient gene activation. This led to the actual fuzzy

91 definition of UAS_{NTR} , consisting in two GATA sites located sufficiently close to one another
92 to present a binding platform for GATA factors [29–31].

93 *In vivo*, GATA ChIP analyses have been carried out genome-wide in yeast [32–35]. They
94 allowed gaining insights into the GATA factor gene network through the identification of
95 direct targets and characterization of the DNA elements required to render a gene NCR-
96 sensitive. However, these studies were not performed in activating conditions, when all
97 GATA factors are expressed, localized in the nucleus and active, so that the current list of
98 GATA factor targets are likely to be underestimated. On another hand, bioinformatic analyses
99 have shown that, since GATA sequences are short, they can be found almost everywhere
100 throughout the genome. Therefore, based on the sole criteria of the presence of repeated
101 GATA sequences in yeast promoters, a third of the yeast genes could hypothetically be NCR
102 regulator targets [36]. Such GATA motif repetitions have been found in the promoter of 91
103 genes, inducible by GATA activators in absence of a good nitrogen source, supposed to be
104 directly targeted by the GATA activators [37]. Nevertheless, the functionality of these
105 supposed UAS still needs to be directly demonstrated *in vivo* [1].

106 Here, we provide the first genome-wide identification of Dal80 targets in yeast, in
107 physiological conditions where Dal80 is fully expressed and active. Using a ChIP-Seq
108 approach combined to a bioinformatic peak-calling procedure, we defined the exhaustive set
109 of Dal80-bound promoters, which turned out to be much larger than anticipated. Our data
110 indicate that at some promoters, Dal80 recruitment occurs independently of GATA sites.
111 Strikingly, Dal80 was also detected across the body of a subset of genes bound at the
112 promoter, globally correlating with high and Dal80-sensitive expression. Mechanistic single-
113 gene experiments confirmed the Dal80 binding profiles, further indicating that Dal80
114 spreading across gene bodies occurs independently of intragenic GATA sites but requires
115 transcription elongation. Finally, co-immunoprecipitation experiments revealed that Dal80

physically interacts with both initiating and elongating form of PolII. Our work suggests that in addition to their well-established role at the (pre)initiation stage, GATA factors could be involved in transcription elongation, promoting efficient remodeling of the gene expression program in response to environmental changes.

Results

Genome-wide identification of Dal80-bound promoters

In order to determine the genome-wide occupancy of a GATA factor in yeast, our rationale was to choose Dal80 as it is known to be highly expressed in derepressing conditions and forms chromosome foci when tagged by GFP [38]. We thus grew yeast cells in proline-containing media and performed a ChIP-Seq analysis using a Dal80-Myc¹³ tagged strain and the isogenic untagged strain, as a control (Fig 1A). Dal80-bound regions were identified using a peak-calling algorithm (see Material & Methods). A promoter was defined as bound by Dal80 when >75% of the -350 to -100 region (relative to the downstream ORF start site) was overlapped by a peak (Fig 1B). We chose to use as the reference coordinate the translation initiation codon rather than the transcription start site (TSS) because the latter has not been accurately determined for all genes. Then, our arbitrary definition of the promoter as the -350 to -100 region relative to the ATG codon was based on the distribution of the TSS-ATG distance for genes with an annotated TSS (median and average distance = 58 and 107 bp, respectively; see S1 Fig).

Strikingly, Dal80 was found to bind to 1269 gene promoters (Fig 1C-D and S1 Table). Fig 1E shows a snapshot of ChIP-Seq signals at the promoter of the *FUR4* gene, identified as bound by Dal80. The number of Dal80-bound promoters, corresponding to 22% of all protein-coding genes promoters, is much higher than expected given the hundred target genes that are

generally cited for the GATA transcriptional activators [37,39], that are presumed to bind to the same sites as Dal80. However, we noted that some peaks (221) overlapped with several promoters (471), mainly of divergent genes (442). In such cases, it is therefore possible that only one of the two divergent promoters is indeed targeted by Dal80, so that the number of Dal80-bound promoters could actually be a little less important than 1269.

As expected, genes related to previous nitrogen regulation screens were present within our list [37,39–46](S2 Table). Among the genes showing Dal80-binding to the promoter, we also noticed a strong enrichment for cytoplasmic translation genes, as well as genes involved in small molecule biosyntheses, including amino acids (S3 Table).

Surprisingly, analysis of GATA site occurrence over Dal80-bound and unbound promoters revealed no major difference between the two classes (Fig 1F), 48.2% and 51.3% of Dal80-bound and unbound promoters containing at least two GATA sites, respectively (S4 Table). Intriguingly, 20% of Dal80-bound promoters do not contain any GATA site (Fig 1F and S4 Table), indicating that Dal80 recruitment can also occur independently of the presence of consensus GATA sites (see S1 Fig for visualization of Dal80 binding to a GATA-less promoter).

In summary, our ChIP-Seq analysis revealed that Dal80 binds to a set of promoters larger than previously expected, targeting biosynthetic functions and protein synthesis in addition to nitrogen catabolite repression.

Dal80 binds the intragenic region of a subset of genes

Metagene analysis revealed that the genes bound by Dal80 at the promoter also display a signal along the gene body, although this intragenic signal remains globally lower than in the promoter-proximal region (Fig 1D). This observation prompted us to investigate the possibility that Dal80 also occupies the gene body, at least for a subset of genes.

We identified 189 genes showing Dal80 intragenic occupancy, according to a >75% overlap of the ORF by a Dal80-Myc¹³ peak (Fig 2A-B). Among them, 144 (76%) were also bound at the promoter (Fig 2B). Conversely, only 11% of the Dal80-bound promoters (144/1269) showed downstream intragenic binding (Fig 2B). On the other hand, 45 genes showing Dal80 intragenic binding were not bound at the promoter (Fig 2B). Hence, we distinguished four classes of genes (S5 Table): (i) those bound by Dal80 at the promoter only (“P” class; Fig 2C), (ii) those showing both promoter and intragenic binding (“P & O” class; Fig 2D), (iii) those bound across the ORF only (“O” class; Fig 2E), (iv) the unbound genes (Fig 2F). Interestingly, we noted that the global Dal80-Myc¹³ signal at the promoter was higher for the “P & O” class in comparison to the “P” class (compare Fig 2C-D). In conclusion, a subset of genes displays intragenic Dal80 binding, and in most cases, this correlates with strong Dal80 recruitment at the promoter.

Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels

In order to correlate Dal80 binding with gene expression and Dal80-sensitivity in similar culture conditions, we performed total RNA-seq analysis in wild type and *dal80Δ* cells grown in medium with proline as the nitrogen source.

The expression of 546 genes was significantly altered in proline-grown *dal80Δ* cells compared to wild type (Fig 3A, S6 Table), among which 232 and 314 genes were negatively and positively regulated by Dal80, respectively. The first group was enriched for genes involved in small molecule catabolic processes (S7 Table), while the positively regulated genes were mostly involved in amino acid biosynthesis (S8 Table). Again, we noticed an overlap between Dal80-repressed genes and nitrogen regulated genes that were identified in other genome-wide screens (S4 Table).

Analysis of mRNA levels in proline-grown cells revealed a significant correlation between

Dal80 binding and regulation at the mRNA level: 39.6% (216/546) of Dal80-regulated genes were bound by Dal80 at their promoter and/or across the ORF, which is significantly more than expected if Dal80 binding and regulation were independent ($P < 0.00001$, Chi-square test; S2 Fig). More importantly, additional differences are emerging between the different classes of genes. First, gene expression was found to be globally similar between the unbound genes and those of the “O” class ($P = 0.15$, Wilcoxon rank-sum test; Fig 3B). Second, Dal80 binding at the promoter of genes correlated with significantly increased expression levels, in comparison to the unbound genes ($P < 2.2 \times 10^{-16}$, Wilcoxon rank-sum test; Fig 3B). Third, the genes bound by Dal80 both at the promoter and across the ORF (“P & O” class) were even more expressed than the genes showing promoter binding only ($P = 1.3 \times 10^{-14}$, Wilcoxon rank-sum test; Fig 3B). Together with the observation that genes of the “P & O” class globally showed higher Dal80-Myc¹³ ChIP-Seq signal at the promoter than those of the “P” class (Fig 2C-D), our results indicate that Dal80 spreading across gene bodies correlates with a strong recruitment at the promoter, higher sensitivity to Dal80 and high expression in proline-containing medium. Consistently, metagene analysis of Dal80-regulated genes is very similar to that of “P & O” class (S2 Fig). Noticeably, 15% (170) and 2% (75) of the genes belonging to the “P” and “unbound” classes respectively display expression levels higher than median levels of expression of the genes belonging to the “P&O” class (S2 Fig). These proportions raise even further when the first quartile (25%) of the “P&O” class is considered (S2 Fig), suggesting that elevated expression levels does not automatically imply binding at ORF.

In conclusion, our genome-wide analyses revealed a significant correlation between Dal80 recruitment to the chromatin and Dal80-sensitive expression. We also showed that Dal80 binding both at the promoter and across the gene body correlates with high levels of gene expression.

Dal80 binding across the ORF of well-characterized NCR-sensitive genes

In order to validate our genome-wide observations at the level of known NCR-sensitive genes, we characterized the binding profiles of Dal80-Myc¹³ along four NCR-sensitive genes of the “P & O” class (see snapshots of ChIP-Seq signals in Fig 4A-D). We chose the NAD⁺-dependent glutamate dehydrogenase gene (*GDH2*) and three permease-coding genes (*GAP1*, *DAL5* and *MEP2*), aware that their regulation could differ [37,39,47] and that GATA sites are differently represented throughout these genes (Fig 4A-D).

As expected from our genome-wide observations, Dal80 bound not only the promoter, but also the coding region of *DAL5*, *GAP1*, *GDH2* and *MEP2* in proline-grown cells (Fig 4E and S3 Fig). Binding of Dal80 to the *GDH2* ORF was reduced (Fig 4E), which is consistent with the ChIP-Seq profile (Fig 4D). We ruled out any erroneous interpretation of our data upon showing that, using the same experimental design, the Gal4-Myc¹³ protein only occupied the *GAL10* promoter, and not the coding region (S3 Fig). Moreover, intragenic GATA factor binding did not rely on an intermediate RNA molecule, since the Dal80-Myc¹³ signal across the *MEP2* gene body was unaffected upon RNase treatment of the chromatin extracts before the immunoprecipitation (Fig 4F and S3 Fig).

The results of the ChIP-qPCR show that the level of Dal80 binding also varied from one gene to another. As striking example, Dal80 was more present at the *GAP1* and *MEP2* ORFs, which are more highly expressed than *DAL5* in proline-grown cells (S6 Table). This is consistent with the correlation observed at the genomic level between Dal80 binding both to the promoter and across gene bodies and high expression levels (Fig 3B). Altogether, these observations prompted an important mechanistic question: how can Dal80 be localized to gene bodies ?

Dal80 binding within gene bodies occurs independently of intragenic GATA sites

Like most NCR-sensitive genes, the four genes described above are all characterized by

stretches of GATA sequences in their promoters (Fig 4A-D). A possible explanation for Dal80 binding across gene bodies could be that functional, intragenic GATA sequences, directly recruit Dal80. Given the low complexity of the GATA consensus, the probability that a gene is devoid of intragenic GATA site is very low. Indeed, only 10% of all yeast ORFs have no GATA site, and most genes contain an average of 5 intragenic GATA sites (S3 Fig). However, according to current models, GATA sites should be present in stretches of at least two in order to be functional *in vivo*, which is not the case within *DAL5*, *GDH2*, *GAP1* and *MEP2* ORFs, that contain some GATA sites, but relatively distant from each other. Interestingly, 23 of the 144 P&O-bound genes did not contain any GATA site in their ORF, hence showing that Dal80 spreading across the body of genes does not necessarily require intragenic GATA sites, at least for these genes.

In order to test if the presence of a NCR-sensitive promoter could confer intragenic Dal80 binding across the body of a non-NCR-sensitive gene, devoid of any GATA cluster, we placed the *URA3* ORF under the control of the *MEP2* promoter, at the *MEP2 locus* (P_{MEP2} -*URA3*; Fig 5A). Only two GATA sites are present in the *URA3* ORF, and separated by 300bp. In this strain, the expression of *URA3* became NCR-sensitive and followed wild type *MEP2* expression (S4 Fig), correlating with PolIII recruitment over the *URA3* ORF (S4 Fig). In these conditions, we observed Dal80-Myc¹³ binding at the promoter of *MEP2* and also across *URA3* (Fig 5B). Importantly, Dal80 was not detected across *URA3* when it was expressed from its native *locus*, under the control of its promoter (Fig 5B). These results show that significant recruitment of Dal80 at an ORF region must be preceded by promoter recruitment.

Finally, we analyzed Dal80 binding at the *UGA4 locus*, another well-characterized NCR-sensitive gene, which is devoid of intragenic GATA site clusters (Fig 5C). *UGA4* expression is induced by GABA (γ -aminobutyric acid) and is strongly repressed by Dal80 in the absence of the inducer [48]. Therefore, to derepress *UGA4* without inducer and to analyze Dal80

binding along *UGA4*, we deleted the repressive C-terminal leucine zipper domain of Dal80 (Dal80 Δ LZ-Myc¹³), known for its homodimerization and heterodimerization with the Gzf3 transcriptional repressor [17,27](Fig 5D). As a control experiment, *UGA4* expression was quantified in Dal80-Myc¹³ and Dal80 Δ LZ-Myc¹³ cells. As expected, *UGA4* mRNA level was very low in Dal80-Myc¹³ cells, whereas deletion of Dal80 leucine zipper led to a strong derepression of *UGA4* (S4 Fig). Analysis of PolII occupancy across *UGA4* using ChIP confirmed that elevated mRNA levels correlate with increased transcription (S4 Fig). In these conditions, full-length Dal80-Myc¹³ binding was restricted to the *UGA4* promoter, while Dal80 Δ LZ-Myc¹³ binding was detected at the promoter and across the body of *UGA4* (Fig 5E), correlating with the elevated transcript and PolII levels. Surprisingly, the leucine zipper of Dal80 and consequently, its dimerization, needed for *UGA4* repression, were not required for its spreading across *UGA4* gene body. Importantly, these results show that promoter binding is not sufficient to confer intragenic binding, but suggest that transcription activation is required.

Together, these data indicate that Dal80 binding at the promoter and across the body of NCR-sensitive genes correlated with expression levels and required transcription activation.

Transcription elongation is necessary for Dal80 binding across *MEP2*

We hypothesized that the elongating PolII complex could be responsible for Dal80 spreading beyond Dal80-bound promoters.

To assess whether the transcriptionally active PolII complex is necessary for Dal80 binding across the *MEP2* gene, we used an *rpb1-1* strain, known to rapidly involve a cessation of PolII activity when cells are shifted to a restrictive temperature of 37°C [49,50]. We analyzed Dal80-Myc¹³ binding along *MEP2* in WT and *rpb1-1* cells. When *rpb1-1* cells were shifted at 37°C for 1hr, *MEP2* mRNA and PolII levels showed a 2-fold (S5 Fig) and >10-fold decrease

(S5 Fig), respectively, reflecting the expected elongation defect when *rpb1-1* cells are shifted in non-permissive conditions. In the same conditions, we observed a significant >5-fold reduction of Dal80-Myc¹³ levels across the *MEP2* ORF, while the binding at the promoter was not affected (Fig 6A-B). This result suggests that Dal80 spreading across the body of NCR-sensitive genes would be a consequence of the PolII-driven elongation process.

To get insights into the mechanism by which Dal80 associates to gene bodies, we tested whether it physically interacts with the transcriptionnaly engaged form of PolII (Fig 6C). Total protein extracts from Dal80-Myc¹³ cells were immunoprecipitated with antibodies directed against the PolII CTD and its phospho-forms Ser2P and Ser5P, respectively characteristic of elongating and initiating PolII forms. All three antibodies enabled effective immunoprecipitation, whereas no antibody and nonspecific antibody controls generated a lower or no signal at all. Thus, Dal80 physically interacts with phosphoforms of the PolIII, suggesting a strong association with PolII engaged in active transcription from initiating to elongating polymerase.

Together, our data indicate that Dal80 spreading across the body of NCR-sensitive genes depends on active transcription and that Dal80 interacts with the transcriptioanlly active form of PolII, supporting a model where Dal80 spreading across the body of highly expressed, NCR-sensitive genes might be the result of Dal80-PolII association at post-initiation transcription phases.

Discussion

Eukaryotic GATA factors belong to a small but important family of DNA binding proteins involved in development and response to environmental changes in multicellular organisms in unicellular organisms, respectively. In yeast, four GATA factors are involved in Nitrogen

Catabolite Repression (NCR), controlling gene expression in response to nitrogen source availability. One of them, the Dal80 repressor, itself NCR-sensitive, acts to modulate the intensity of NCR responses.

Over the past decade, a number of studies have screened the genome aiming at gathering an inventory of genes regulated by the nitrogen source. Although >500 genes have been shown to be differentially expressed upon change of the nitrogen source [39,46], the list of NCR-sensitive genes was reduced to about 100, based on their sensitivity to GATA factors [37,39,42,45], suggesting that the number of Dal80 targets would be situated in that range.

Here, our ChIP-Seq analysis identified more than 1200 Dal80-bound promoters, which considerably extends the list of potential Dal80 targets. In fact, the number of Dal80-bound promoters could even have been more important. Indeed, the GATA consensus binding site is rather simple and short, so that in yeast, a total number of 10,000 putative binding sites can be found in all protein-coding gene promoters, 2930 promoters having at least 2 GATA sites, which is thought to be a prerequisite for *in vivo* binding and function of the GATA factors.

The difference between the number of promoters with ≥ 2 GATA sites and the number of Dal80-bound promoters suggests the existence of a selectivity for Dal80 recruitment. This selectivity could rely on promoter architecture and/or chromatin structure, conditioning the requirement for auxiliary DNA binding factors that would stabilize Dal80 at some promoters.

Moreover, although we observed a significant correlation between Dal80 binding and regulation, the expression of most of the Dal80-bound genes was not affected in a *dal80Δ* mutant strain. Again, Dal80-dependence for transcribing these genes, as well as their NCR sensitivity, could require the presence of cofactors which are not produced or inactive under the tested growth conditions. In mammals, GATA factors also display an extraordinary complexity in the relationships between binding and expression regulation. Like Dal80, GATA-1 and GATA-2 only occupy a small subset of their abundant binding motif throughout

the genome, and the presence of the conserved binding site is insufficient to cause GATA-dependent regulation in most instances [51]. GATA-1 binding kinetics, stoichiometry and heterogenous complex formations, conditioned by composite promoter architecture, influence its transcriptional activity and hence diversify gene expression profiles [51].

Given the high conservation at the amino acid level between the DNA binding domain of the four yeast NCR GATA factors, it is likely that they all recognize identical sequences (GATAA, GATAAG or GATTAG). This consensus has been largely validated in the past using gene reporter experiments, mutational analyses and *in vitro* binding experiments on naked DNA. Nonetheless, of the 1269 bound promoters, 48% contained at least two GATA sites, a proportion that is not different from that observed among unbound promoters, and the amount of GATA sites per promoter was not different between the two groups either. In addition, Dal80 recruitment was found to occur independently of the presence of GATA sites in 20% of Dal80-bound promoters, as also previously observed in mammalian cells [10,52]. Future experiments will be required to decipher the molecular mechanism by which Dal80 can be recruited to these GATA-less promoters.

Unexpectedly, although Dal80 has always been described as a repressor, we identified 314 genes that are positively regulated by Dal80 (their expression is significantly decreased upon Dal80 deletion). These genes are distinct from the NCR-sensitive genes : only 4 belong to the NCR-sensitive list of [39] and are significantly enriched in amino acid biosynthetic processes, resembling the amino acid starvation response mediated by the Gcn4 transcriptional activator. Interestingly, the promoter of 122/314 Dal80-activated genes contain Gcn4-binding sites (S3 Fig), and this group of 314 Dal80-activated genes is significantly enriched for genes regulated by the General Amino Acid Control (GAAC; YeastMine Gene List, Publication Enrichment, $p < 1.6e-13$), through the Gcn4 activator. Interconnections between NCR and GAAC have already been demonstrated, mostly at the level of nitrogen catabolism control: 1-a large

number of non-preferential nitrogen sources leads to increased transcription of GAAC targets [39]; and 2- Gcn4 contributes, with Gln3, to the expression of NCR-sensitive genes [53,54]. However, this is the first time that evidence are provided suggesting a positive function for the negative GATA factor at the level biosynthetic gene expression.

The most striking and unexpected finding of this work is the observation that for 189 genes, Dal80 occupied the gene body (“O” and “P & O” classes). The 45 genes displaying Dal80 intragenic occupancy with no recruitment at the promoter (“O” class) were globally neither Dal80-sensitive nor NCR-sensitive, similar to the unbound genes. Furthermore, we also noted that a substantial fraction of these genes correspond to small dubious ORFs, near or even overlapping the promoter, Dal80-bound, of the adjacent gene. In these cases, the limited resolution of the ChIP-Seq technique, combined to the small size of these genes, might have allowed them to pass through the filters of the procedure to identify the genes showing Dal80 intragenic binding. Overall, these observations suggest that the existence of the “O” class is likely to be physiologically irrelevant. Dal80 binding at the promoter and spreading across the body of the 144 genes belonging to the “P & O” class correlated with high expression levels and sensitivity to Dal80-dependent repression. The observation that a considerable fraction of “P” class genes have similar or higher expression levels than most of the “P&O” class genes suggest that the well-known artifact of hyper-chippable regions is not applicable here [55]. Our observations were also experimentally confirmed using ChIP experiments, at the level of well-characterized NCR-sensitive genes. Our data indicate that Dal80 spreading across genes bodies does not rely on intragenic GATA sites. Rather, promoter binding appears to be required but not sufficient: Dal80 spreading across genes depends on active transcription. Indeed, the observation that impaired PolII elongation correlates with decreased intragenic binding further indicates that Dal80 binding across gene bodies depends on ongoing transcription elongation. Consistently, we detected a physical interaction between

Dal80 and transcriptionally active forms of PolII. Together, our data leads us to propose a model where Dal80 could travel from the promoter of highly expressed, NCR-sensitive genes through the gene body by accompanying the elongating PolII complex (Fig 7). Additional investigations will be required to define which domain of Dal80 is responsible for the interaction with the transcription machinery, to determine whether there is any causal relationship between Dal80 intragenic binding and high expression levels, and to decipher the potential role of Dal80 during transcription elongation.

Whereas the binding of elongation factors across gene bodies has been thoroughly documented [56], it has also been described for few specific transcription factors. For example, Gal4 was reported to bind to its consensus DNA target within the *ACC1* ORF, but the authors concluded that the observed transcriptional repression of the *ACC1* gene was most likely resulting from random *GAL4* binding “noise” over the genome, thus having no physiological explanation for this ORF-bound transcription factor [57]. Likewise, Gcn4 was detected across the *PHO8* ORF, with concomitant recruitment of the SAGA complex, but without any impact on gene expression [58]. More recently, binding of the Gcn4 transcription factor to its consensus site at some ORFs, when located in proximity of the transcriptional start site, was found to play a consistent role in controlling embedded cryptic promoters in yeast, thereby affecting Gcn4-dependent transcription of some genes [59].

A recent study has identified CTD phosphorylation of polII as a hub that optimizes transcriptome changes to adequately balance optimal growth and stress tolerance responses [60]. The addition of nitrogen to nitrogen-limited cells rapidly results in the transient overproduction of transcripts required for protein translation (stimulated growth) whereas accelerated mRNA degradation favours rapid clearing of the most abundant transcripts, like those involved in high affinity permease production, that are highly expressed NCR-sensitive genes, for example [46]. The involvement of the Nrd1-Nab3-Sen1 (NNS) and TRAMP

complexes in these regulatory responses has been envisioned very recently [61,62]; deadenylation, decapping and exonuclease mutants display impaired *GAPI* mRNA clearance upon nitrogen upshift [63]. Thus, a possible role of Dal80 (and possibly of the other GATA factors) binding along highly expressed genes could be to transmit nutritional signals to the RNA decay machinery for rapid and efficient remodeling of gene expression in response to changing nitrogen supplies. Here, an attractive model (Fig 7) would hypothesize that Dal80 binding to polII could prevent the decay of the dedicated stress-response mRNAs to enhance their cellular life-cycle by regulating/repressing RNA decay factors that might associate otherwise on the polymerase. Other elongation-related processes, like histone modification and chromatin remodeling [64,65], mRNA export [66] or roadblock termination [67] can also be proposed.

In this respect, in human cells, GATA factors are also reported to occupy non-canonical sites within the genome, further reinforcing that they can be recruited to the chromatin independently of their motif [10,52]. In addition, 43% of the GATA1 peaks were collected among exon, introns and 3'UTR of coding genes in human erythroleukemia cells [52]. It is tempting to hypothesize that GATA factors could have a dual or synergistic role during transcription, i.e. recruiting/stabilizing the PIC complex as for any classical transcription factor in the promoter/enhancer regions and promoting elongation at a post initiation step interacting with the RNAPII. Interestingly, GATA1 in erythropoid cells interacts with the mediator [68], a complex also known to travel with PolII [69].

Materials and methods

Experimental model and subject details

Experiments were conducted using *S. cerevisiae* strains of the FY genetic background. The

strains used are listed in S9 Table. Dal80 was tagged with 13 copies of the *c-myc* epitope (Myc¹³) as described [70] using primers listed in S9 and S10 Tables. The *P_{MEP2}-URA3* allele in strains FV806-808 was created by amplification of the *URA3* gene using the same strategy, with primers listed in S9 and S10 Tables.

Cultures were grown at 29°C to mid-log phase ($A_{660\text{nm}} = 0.5$) in YNB (without amino acids or ammonia) minimal medium containing the indicated nitrogen source at a 0.1% final concentration, glucose (3%) and the appropriate supplements (20 µg/ml uracil, histidine and tryptophan) to cover auxotrophic requirements.

Chromatin immunoprecipitation

Cell extracts and immunoprecipitations were conducted as described [23] using primers listed in S10 Table. The cells (100-ml cultures grown to an absorbance ($A_{660\text{ nm}} = 0.6$) corresponding to 6×10^6 cells/ml) were treated with 1% formaldehyde for 30 min at 25 °C and mixed by orbital shaking. Glycine was then added to a final concentration of 500 mM and incubation continued for 5 min. The cells were collected, washed once with cold 10 mM Tris-HCl, pH 8, washed once with cold FA-SDS buffer (50 mM HEPES-KOH, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS, 1 mM phenylmethylsulfonyl fluoride), and resuspended in 1 ml of cold FA-SDS buffer. An equal volume of glass beads (0.5 mm in diameter) was added, and the cells were disrupted by vortexing for 30 min in a cold room. The lysate was diluted into 4 ml of FA-SDS buffer, and the glass beads were discarded. The cross-linked chromatin was then pelleted by centrifugation ($17,000 \times g$ for 35 min), washed for 60 min with FA-SDS buffer, resuspended in 1.6 ml of FA-SDS buffer for 15 min at 4 °C, and sonicated three times for 30 s. each (Bioruptor, Diagenode) to yield an average DNA fragment size of 700 base pairs. Finally, the sample was clarified by centrifugation at $14,000 \times g$ for 30 min and diluted 4-fold in FA-SDS

buffer, and aliquots of the resultant chromatin containing solution were stored at -80°C . Myc¹³-tagged proteins were immunoprecipitated by incubating 100 μl of the chromatin containing solution for 180 min at 4°C with 2 μl of mouse anti-Myc antibodies (SC-40, Santa Cruz) prebound to 10 μl of Dynabeads Pan Mouse IgG (Dyna) according to the manufacturer's instructions. Immune complexes were washed six times in FA-SDS buffer and recovered by treating with 50 μl of Pronase Buffer (25 mM Tris, pH 7.5, 5 mM EDTA, 0.5% SDS) at 65°C with agitation. Input (IN) and immunoprecipitated (IP) fractions were then subjected to Pronase treatment (0.5 mg/ml; Roche Applied Science) for 60 min at 37°C , and formaldehyde cross-links were reversed by incubating the eluates overnight at 65°C . Finally, the samples were treated with RNase (50 $\mu\text{g}/\text{ml}$) for 60 min at 37°C . DNA from the IP fractions was purified using the High Pure PCR Product Purification Kit (Roche Applied Science) and eluted in 50 μl of 20 mM Tris buffer, pH 8. IN fractions were boiled 10 min and diluted 500-fold with no further purification prior to quantitative PCR analysis.

Quantitative RT-PCR

Quantitative RT-PCR was performed as described previously [23] using primers listed in S10 Table. Total RNA was extracted from 4-ml cultures and cDNA was generated from 100 to 500 ng of total RNA using a RevertAid H Minus first-strand cDNA synthesis kit with oligo(dT)₁₈ primers from Fermentas using the manufacturer's recommended protocol. cDNAs were subsequently quantified by RT-PCR using the Maxima SYBR green qPCR master mix from Fermentas.

Co-immunoprecipitation

Cultures (100 ml) were harvested, washed once in 50 mM Tris, pH 8, and resuspended in 1 ml of buffer (50 mM Tris, pH 8, 150 mM NaCl, 5 mM EDTA, 0.05% NP-40, 1 mM phenylmethylsulfonyl fluoride, and complete protease inhibitor cocktail tablets [Roche]).

Lysis was performed by shaking with 425-600 μ m acid-washed glass beads (Sigma) on an IKA Vibrax VXR orbital shaker at maximum speed for 30 min at 4°C. Cell debris and glass beads were removed by centrifugation. Immunoprecipitation was performed by incubating 200 μ l of total cell extracts with 20 μ l of Dynabeads PAN mouse immunoglobulin G (Invitrogen) that were preincubated with anti-HA (SCBT, SC-7392), anti-CTD (SCBT, CTD4H8), anti-Ser2P (BioLegend, H5) or anti-Ser5P (BioLegend, H14) antibodies and 20 μ l of 1% phosphate-buffered saline-bovine serum albumin for 2 h under orbital shaking (800 rpm) at 30°C. Immune complexes were washed three times in lysis buffer, eluted by boiling in sodium dodecyl sulfate (SDS) sample buffer, and loaded on SDS-polyacrylamide gel for anti-Myc Western blotting.

ChIP-Seq analysis and peak-calling

ChIP-Seq analysis was performed from two biological replicates of proline-grown 25T0b (no tag) and FV078 (Dal80-Myc¹³) cells. For each condition, libraries were prepared from 10 ng of “input” or “IP” DNA using the TruSeq ChIP Sample Preparation Kit (Illumina). Single-read sequencing (50 nt) of the libraries was performed on a HiSeq 2500 sequencer.

Reads were uniquely mapped to the *S. cerevisiae* S288C reference genome using Bowtie2 v2.1.0 [71], with a tolerance of 1 mismatch in seed alignment. Tags densities were normalized on the total number of uniquely reads mapped.

Dal80-bound regions were identified through a peak-calling procedure using version 2.0.9 of MACS [72], with a minimum false discovery rate (FDR) of 0.001.

Total RNA-Seq

For each strain and condition, total RNA was extracted from two biological replicates using standard hot phenol procedure, ethanol-precipitated, resuspended in nuclease-free H₂O (Ambion) and quantified using a NanoDrop 2000c spectrophotometer. Ribosomal RNAs were

depleted from 1 µg of total RNA using the RiboMinus™ Eukaryote v2 Kit (Life Technologies). After concentration using the Ribominus™ Concentration Module (Life Technologies), rRNA-depleted RNA was quantified using the Qubit RNA HS Assay kit (Life Technologies). In parallel, rRNA depletion efficiency and integrity of both total and rRNA-depleted RNA were checked by analysis in a RNA 6000 Pico chip, in a 2100 bioanalyzer (Agilent). Strand-specific total RNA-Seq libraries were prepared from 125 ng of rRNA-depleted RNA using the TruSeq® Stranded Total RNA Sample Preparation Kit (Illumina), following manufacturer's instructions. Paired-end sequencing (2 x 50 nt) of the libraries was performed on a HiSeq 2500 sequencer. Sequenced reads were mapped to the reference genome using version 2.0.6 of TopHat [73], as described [74]. Tags densities were normalized on the total number of reads uniquely mapped on ORFs. Differential expression analysis was performed using DESeq [75].

Quantification and Statistical Analysis

Statistical details are to be found in the figure legends. Error bars correspond to standard error. Statistical significance tests were carried out using the Student's t test (Microsoft Excel) when indicated.

Availability of data and materials

Sequence data can be accessed at the NCBI Gene Expression Omnibus using accession numbers GSE86307 and GSE86325. Genome browsers for visualization of processed ChIP-Seq and RNA-Seq data are accessible at <http://vm-gb.curie.fr/dal80>.

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Isabelle Georis (igeoris@ulb.ac.be). Bioinformatics and genome wide dataset requests could also be addressed to antonin.morillon@curie.fr for rapid processing.

530

531 **Acknowledgements**

532 This work has benefited from the facilities and expertise of the NGS platform of Institut
533 Curie.

534 We are grateful to all members of our labs for discussions and critical reading of the
535 manuscript. We thank F. Vierendeels for excellent technical assistance.

536

537 **References**

- 538 1. Hahn S, Young ET. Transcriptional regulation in *Saccharomyces cerevisiae*:
539 Transcription factor regulation and function, mechanisms of initiation, and roles of
540 activators and coactivators. *Genetics*. 2011;189: 705–736.
541 doi:10.1534/genetics.111.127019.
- 542 2. Levine M, Tjian R. Transcription regulation and animal diversity. *Nature*. 2003;424:
543 147–151. doi:10.1038/nature01763.
- 544 3. Zambelli F, Pesole G, Pavesi G. Motif discovery and transcription factor binding sites
545 before and after the next-generation sequencing era. *Brief Bioinform*. 2013;14: 225–
546 237. doi:10.1093/bib/bbs016.
- 547 4. Omichinski JG, Clore GM, Schaad O, Felsenfeld G, Trainor C, Appella E, et al. NMR
548 structure of a specific DNA complex of Zn-containing DNA binding domain of
549 GATA-1. *Science*. 1993;261: 438–46. doi:10.1126/science.8332909.
- 550 5. Cunningham TS, Dorrington RA, Cooper TG. The UGA4 UAS(NTR) site required for
551 GLN3-dependent transcriptional activation also mediates DAL80-responsive regulation
552 and DAL80 protein binding in *Saccharomyces cerevisiae*. *J Bacteriol*. 1994;176: 4718–

4725.

6. Horak CE, Mahajan MC, Luscombe NM, Gerstein M, Weissman SM, Snyder M. GATA-1 binding sites mapped in the beta-globin locus by using mammalian chIp-chip analysis. *Proc Natl Acad Sci U S A*. 2002;99: 2924–2929. doi:10.1073/pnas.0527069999.
7. Stanbrough M, Magasanik B. Two transcription factors, Gln3p and Nil1p, use the same GATAAG sites to activate the expression of GAP1 of *Saccharomyces cerevisiae*. *J Bacteriol*. 1996;178: 2465–2468.
8. Starich MR, Wikström M, Arst HN, Clore GM, Gronenborn AM. The solution structure of a fungal AREA protein-DNA complex: an alternative binding mode for the basic carboxyl tail of GATA factors. *J Mol Biol*. 1998;277: 605–20. doi:10.1006/jmbi.1998.1625.
9. Bresnick EH, Katsumura KR, Lee HY, Johnson KD, Perkins AS. Master regulatory GATA transcription factors: Mechanistic principles and emerging links to hematologic malignancies. *Nucleic Acids Res*. 2012;40: 5819–5831. doi:10.1093/nar/gks281.
10. Cheng Y, Wu W, Kumar SA, Yu D, Deng W, Tripic T, et al. Erythroid GATA1 function revealed by genome-wide analysis of transcription factor occupancy, histone modifications, and mRNA expression. *Genome Res*. 2009;19: 2172–2184. doi:10.1101/gr.098921.109.
11. Tripic T, Deng W, Cheng Y, Zhang Y, Vakoc CR, Gregory GD, et al. SCL and associated proteins distinguish active from repressive GATA transcription factor complexes. *Blood*. 2009;113: 2191–2201. doi:10.1182/blood-2008-07-169417.
12. Bresnick EH, Lee HY, Fujiwara T, Johnson KD, Keles S. GATA switches as developmental drivers. *J Biol Chem*. 2010;285: 31087–31093. doi:10.1074/jbc.R110.159079.

13. Grass JA, Boyer ME, Pal S, Wu J, Weiss MJ, Bresnick EH. GATA-1-dependent transcriptional repression of GATA-2 via disruption of positive autoregulation and domain-wide chromatin remodeling. *Proc Natl Acad Sci U S A*. 2003;100: 8811–8816. doi:10.1073/pnas.1432147100.
14. Lowry JA, Atchley WR. Molecular Evolution of the GATA Family of Transcription Factors: Conservation Within the DNA-Binding Domain. *J Mol Evol*. 2000;50: 103–115. doi:10.1007/s002399910012.
15. Magasanik B, Kaiser CA. Nitrogen regulation in *Saccharomyces cerevisiae*. *Gene*. 2002;290: 1–18. doi:10.1016/S0378-1119(02)00558-9.
16. Coffman JA, Rai R, Cunningham T, Svetlov V, Cooper TG. Gat1p, a GATA family protein whose production is sensitive to nitrogen catabolite repression, participates in transcriptional activation of nitrogen-catabolic genes in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1996;16: 847–858.
17. Coornaert D, Vissers S, Andre B, Grenson M. The UGA43 negative regulatory gene of *Saccharomyces cerevisiae* contains both a GATA-1 type zinc finger and a putative leucine zipper. *Curr Genet*. 1992;21: 301–307.
18. Cunningham TS, Cooper TG. Expression of the DAL80 gene, whose product is homologous to the GATA factors and is a negative regulator of multiple nitrogen catabolic genes in *Saccharomyces cerevisiae*, is sensitive to nitrogen catabolite repression. *Mol Cell Biol*. 1991;11: 6205–15. doi:10.1128/MCB.11.12.6205.
19. Minehart PL, Magasanik B. Sequence and expression of GLN3, a positive nitrogen regulatory gene of *Saccharomyces cerevisiae* encoding a protein with a putative zinc finger DNA-binding domain. *Mol Cell Biol*. 1991;11: 6216–6228. doi:10.1128/MCB.11.12.6216.
20. Rowen DW, Esiobu N, Magasanik B. Role of GATA factor Nil2p in nitrogen

regulation of gene expression in *Saccharomyces cerevisiae*. *J Bacteriol.* 1997;179: 3761–3766.

21. Soussi-Boudekou S, Vissers S, Urrestarazu A, Jauniaux JC, André B. Gzf3P, a fourth GATA factor involved in nitrogen-regulated transcription in *Saccharomyces cerevisiae*. *Mol Microbiol.* 1997;23: 1157–1168. doi:10.1046/j.1365-2958.1997.3021665.x
22. Coffman JA, Rai R, Loprete DM, Cunningham T, Svetlov V, Cooper TG. Cross regulation of four GATA factors that control nitrogen catabolic gene expression in *Saccharomyces cerevisiae*. *J Bacteriol.* 1997;179: 3416–3429.
23. Georis I, Feller A, Vierendeels F, Dubois E. The Yeast GATA Factor Gat1 Occupies a Central Position in Nitrogen Catabolite Repression-Sensitive Gene Activation. *Mol Cell Biol.* 2009;29: 3803–3815. doi:10.1128/MCB.00399-09.
24. Georis I, Tate JJ, Vierendeels F, Cooper TG, Dubois E. Premature termination of GAT1 transcription explains paradoxical negative correlation between nitrogen-responsive mRNA, but constitutive low-level protein production. *RNA Biol.* 2015;12: 824–837. doi:10.1080/15476286.2015.1058476.
25. André B, Talibi D, Boudekou SS, Hein C, Vissers S, Coornaert D. Two mutually exclusive regulatory systems inhibit UASGATA, a cluster of 5'-GAT(A/T)A-3' upstream from the UGA4 gene of *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 1995;23: 558–564. doi:10.1093/nar/23.4.558.
26. Cunningham TS, Svetlov V V., Rai R, Smart W, Cooper TG. Gln3p Is capable of binding to UASNTR elements and activating transcription in *Saccharomyces cerevisiae*. *J Bacteriol.* 1996;178: 3470–3479.
27. Svetlov V V., Cooper TG. The *Saccharomyces cerevisiae* GATA factors Dal80p and Deh1p can form homo- and heterodimeric complexes. *J Bacteriol.* 1998;180: 5682–5688.

28. Cunningham TS, Cooper TG. The *Saccharomyces cerevisiae* DAL80 repressor protein binds to multiple copies of GATAA-containing sequences (URS(GATA)). *J Bacteriol.* 1993;175: 5851–5861. doi:10.1128/JB.175.18.5851-5861.1993.
29. Bysani N, Daugherty JR, Cooper TG. Saturation mutagenesis of the UAS(NTR) (GATAA) responsible for nitrogen catabolite repression-sensitive transcriptional activation of the allantoin pathway genes in *Saccharomyces cerevisiae*. *J Bacteriol.* 1991;173: 4977–4982.
30. Cooper TG, Rai R, Yoo HS. Requirement of upstream activation sequences for nitrogen catabolite repression of the allantoin system genes in *Saccharomyces cerevisiae*. *Mol Cell Biol.* 1989;9: 5440–5444.
31. Rai R, Genbauffe FS, Sumrada RA, Cooper TG. Identification of sequences responsible for transcriptional activation of the allantoate permease gene in *Saccharomyces cerevisiae*. *Mol Cell Biol.* 1989;9: 602–8.
32. Harbison CT, Gordon DB, Lee TI, Rinaldi NJ, Macisaac KD, Danford TW, et al. Transcriptional regulatory code of a eukaryotic genome. *Nature.* 2004;431: 99–104. doi:10.1038/nature02800.
33. Lee TI. Transcriptional Regulatory Networks in *Saccharomyces cerevisiae*. *Science* (80-). 2002;298: 799–804. doi:10.1126/science.1075090.
34. MacIsaac KD, Wang T, Gordon DB, Gifford DK, Stormo GD, Fraenkel E. An improved map of conserved regulatory sites for *Saccharomyces cerevisiae*. *BMC Bioinformatics.* 2006;7: 113. doi:10.1186/1471-2105-7-113.
35. Venters BJ, Pugh BF. A canonical promoter organization of the transcription machinery and its regulators in the *Saccharomyces* genome. *Genome Res.* 2009;19: 360–371. doi:10.1101/gr.084970.108.
36. van Helden J, André B, Collado-Vides J. Extracting regulatory sites from the upstream

- 653 region of yeast genes by computational analysis of oligonucleotide frequencies. *J Mol*
654 *Biol.* 1998;281: 827–842. doi:10.1006/jmbi.1998.1947.
- 655 37. Scherens B, Feller A, Vierendeels F, Messenguy F, Dubois E. Identification of direct
656 and indirect targets of the Gln3 and Gat1 activators by transcriptional profiling in
657 response to nitrogen availability in the short and long term. *FEMS Yeast Res.* 2006;6:
658 777–791. doi:10.1111/j.1567-1364.2006.00060.x.
- 659 38. Distler M, Kulkarni A, Rai R, Cooper TG. Green fluorescent protein-Dal80p
660 illuminates up to 16 distinct foci that colocalize with and exhibit the same behavior as
661 chromosomal DNA proceeding through the cell cycle of *saccharomyces cerevisiae*. *J*
662 *Bacteriol.* 2001;183: 4636–4642. doi:10.1128/JB.183.15.4636-4642.2001.
- 663 39. Godard P, Urrestarazu A, Vissers S, Kontos K, Bontempi G, van Helden J, et al. Effect
664 of 21 Different Nitrogen Sources on Global Gene Expression in the Yeast
665 *Saccharomyces cerevisiae*. *Mol Cell Biol.* 2007;27: 3065–3086.
666 doi:10.1128/MCB.01084-06.
- 667 40. Daugherty JR, Rai R, El Berry HM, Cooper TG. Regulatory circuit for responses of
668 nitrogen catabolic gene expression to the GLN3 and DAL80 proteins and nitrogen
669 catabolite repression in *Saccharomyces cerevisiae*. *J Bacteriol.* 1993;175: 64–73.
- 670 41. Cox KH, Pinchak AB, Cooper TG. Genome-wide transcriptional analysis in *S.*
671 *cerevisiae* by mini-array membrane hybridization. *Yeast.* 1999;15: 703–713.
672 doi:10.1002/(SICI)1097-0061(19990615)15:8<703::AID-YEA413>3.0.CO;2-Z.
- 673 42. Boer VM, De Winde JH, Pronk JT, Piper MDW. The genome-wide transcriptional
674 responses of *Saccharomyces cerevisiae* grown on glucose in aerobic chemostat cultures
675 limited for carbon, nitrogen, phosphorus, or sulfur. *J Biol Chem.* 2003;278: 3265–
676 3274. doi:10.1074/jbc.M209759200.
- 677 43. Tai SL, Boer VM, Daran-Lapujade P, Walsh MC, de Winde JH, Daran JM, et al. Two-
28

- dimensional transcriptome analysis in chemostat cultures. *J Biol Chem.* 2005;280: 437–447.
44. Wu J, Zhang N, Hayes A, Panoutsopoulou K, Oliver SG. Global analysis of nutrient control of gene expression in *Saccharomyces cerevisiae* during growth and starvation. *Proc Natl Acad Sci USA.* 2004;101: 3148–3153.
45. Boer VM, Tai SL, Vuralhan Z, Arifin Y, Walsh MC, Piper MDW, et al. Transcriptional responses of *Saccharomyces cerevisiae* to preferred and nonpreferred nitrogen sources in glucose-limited chemostat cultures. *FEMS Yeast Res.* 2007;7: 604–620. doi:10.1111/j.1567-1364.2007.00220.x.
46. Airoidi EM, Miller D, Athanasiadou R, Brandt N, Abdul-Rahman F, Neymotin B, et al. Steady-state and dynamic gene expression programs in *Saccharomyces cerevisiae* in response to variation in environmental nitrogen. *Mol Biol Cell.* 2016;27: 1–38. doi:10.1091/mbc.E14-05-1013.
47. Georis I, Tate JJ, Cooper TG, Dubois E. Nitrogen-responsive regulation of GATA protein family activators Gln3 and Gat1 occurs by two distinct pathways, one inhibited by rapamycin and the other by methionine sulfoximine. *J Biol Chem.* 2011;286: 44897–44912. doi:10.1074/jbc.M111.290577.
48. André B, Hein C, Grenson M, Jauniaux JC. Cloning and expression of the UGA4 gene coding for the inducible GABA-specific transport protein of *Saccharomyces cerevisiae*. *Mol Gen Genet.* 1993;237: 17–25. doi:10.1007/BF00282779.
49. Kim TS, Liu CL, Yassour M, Holik J, Friedman N, Buratowski S, et al. RNA polymerase mapping during stress responses reveals widespread nonproductive transcription in yeast. *Genome Biol.* 2010;11: R75. doi:10.1186/gb-2010-11-7-r75.
50. Nonet M, Scafe C, Sexton J, Young R. Eucaryotic RNA polymerase conditional mutant that rapidly ceases mRNA synthesis. *Mol Cell Biol.* 1987;7: 1602–1611.

doi:10.1128/MCB.7.5.1602.Updated.

51. Hasegawa A, Kaneko H, Ishihara D, Nakamura M, Watanabe A, Yamamoto M, et al. GATA1 binding kinetics on conformation-specific binding sites elicit differential transcriptional regulation. *Mol Cell Biol.* 2016;36: 2151–67. doi:10.1128/MCB.00017-16.
52. Fujiwara T, O’Geen H, Keles S, Blahnik K, Linnemann AK, Kang YA, et al. Discovering Hematopoietic Mechanisms through Genome-wide Analysis of GATA Factor Chromatin Occupancy. *Mol Cell.* 2009;36: 667–681. doi:10.1016/j.molcel.2009.11.001.
53. Staschke KA, Dey S, Zaborske JM, Palam LR, McClintick JN, Pan T, et al. Integration of general amino acid control and Target of Rapamycin (TOR) regulatory pathways in nitrogen assimilation in yeast. *J Biol Chem.* 2010;285: 16893–16911. doi:10.1074/jbc.M110.121947.
54. Valenzuela L, Aranda C, González A. TOR modulates GCN4-dependent expression of genes turned on by nitrogen limitation. *J Bacteriol.* 2001;183: 2331–2334. doi:10.1128/JB.183.7.2331-2334.2001.
55. Teytelman L, Thurtle DM, Rine J, van Oudenaarden A. Highly expressed loci are vulnerable to misleading ChIP localization of multiple unrelated proteins. *Proc Natl Acad Sci.* 2013;110: 18602–18607. doi:10.1073/pnas.1316064110.
56. Sims RJ, Belotserkovskaya R, Reinberg D. Elongation by RNA polymerase II: The short and long of it. *Genes Dev.* 2004;18: 2437–2468. doi:10.1101/gad.1235904.
57. Li Q, Johnston SA. Are All DNA Binding and Transcription Regulation by an Activator Physiologically Relevant? *Mol Cell Biol.* 2001;21: 2467–2474. doi:10.1128/MCB.21.7.2467-2474.2001.
58. Topalidou I, Thireos G. Gcn4 occupancy of open reading frame regions results in the

- recruitment of chromatin-modifying complexes but not the mediator complex. *EMBO Rep.* 2003;4: 872–876. doi:10.1038/sj.embor.embor931.
59. Rawal Y, Chereji R, Valabhoju V, Qiu H, Ocampo J, Clark D, et al. Gcn4 Binding in Coding Regions Can Activate Internal and Canonical 5' Promoters in Yeast. *Mol Cell.* 2018;70: 297–311. doi:10.1016/j.molcel.2018.03.007.
60. Chasman D, Ho Y-H, Berry DB, Nemec CM, MacGilvray ME, Hose J, et al. Pathway connectivity and signaling coordination in the yeast stress-activated signaling network. *Mol Syst Biol.* 2014;10: 759. doi:10.15252/msb.20145120.
61. Bresson S, Tuck A, Staneva D, Tollervey D. Nuclear RNA Decay Pathways Aid Rapid Remodeling of Gene Expression in Yeast. *Mol Cell.* 2017;65: 787–800. doi:10.1016/j.molcel.2017.01.005.
62. Merran J, Corden JL. Yeast RNA-Binding Protein Nab3 Regulates Genes Involved in Nitrogen Metabolism. *Mol Cell Biol.* 2017;37: 1–11.
63. Miller D, Brandt N, Gresham D. Systematic identification of factors mediating accelerated mRNA degradation in response to changes in environmental nitrogen. *PLoS Genet.* 2018;14: 1–27. doi:10.1371/journal.pgen.1007406.
64. Henikoff S, Shilatifard A. Histone modification: Cause or cog? *Trends Genet.* 2011;27: 389–396. doi:10.1016/j.tig.2011.06.006.
65. Tessarz P, Kouzarides T. Histone core modifications regulating nucleosome structure and dynamics. *Nat Rev Mol Cell Biol.* 2014;15: 703–708. doi:10.1038/nrm3890.
66. Meinel DM, Burkert-Kautzsch C, Kieser A, O'Duibhir E, Siebert M, Mayer A, et al. Recruitment of TREX to the Transcription Machinery by Its Direct Binding to the Phospho-CTD of RNA Polymerase II. *PLoS Genet.* 2013;9: e1003914. doi:10.1371/journal.pgen.1003914.
67. Candelli T, Challal D, Briand J, Boulay J, Porrua O, Colin J, et al. High-resolution

- transcription maps reveal the widespread impact of roadblock termination in yeast.
EMBO J. 2018;37: e97490. doi:10.15252/emboj.201797490.
68. Stumpf M, Waskow C, Krötschel M, van Essen D, Rodriguez P, Zhang X, et al. The mediator complex functions as a coactivator for GATA-1 in erythropoiesis via subunit Med1/TRAP220. Proc Natl Acad Sci U S A. 2006;103: 18504–9. doi:10.1073/pnas.0604494103.
69. Allen BL, Taatjes DJ. The Mediator complex: a central integrator of transcription. Nat Rev Mol Cell Biol. 2015;16: 155–166. doi:10.1038/nrm3951.
70. Longtine MS, McKenzie A, Demarini DJ, Shah NG, Wach A, Brachet A, et al. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. Yeast. 1998;14: 953–961. doi:10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U.
71. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9: 357–9. doi:10.1038/nmeth.1923.
72. Zhang Y, Liu T, Meyer CA, Eeckhoutte J, Johnson DS, Bernstein BE, et al. Model-based Analysis of ChIP-Seq (MACS). Genome Biol. 2008;9: R137. doi:10.1186/gb-2008-9-9-r137.
73. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. 2013;14: R36. doi:10.1186/gb-2013-14-4-r36.
74. Wery M, Describes M, Vogt N, Dallongeville AS, Gautheret D, Morillon A. Nonsense-Mediated Decay Restricts LncRNA Levels in Yeast Unless Blocked by Double-Stranded RNA Structure. Mol Cell. 2016;61: 379–392. doi:10.1016/j.molcel.2015.12.020.
75. Anders S, Huber W. Differential expression analysis for sequence count data. Genome

Biol. 2010;11: R106. doi:10.1186/gb-2010-11-10-r106.

76. Describes M, Zouari Y Ben, Wery M, Legendre R, Gautheret D, Morillon A. VING: a software for visualization of deep sequencing signals. BMC Res Notes. 2015;8: 419. doi:10.1186/s13104-015-1404-5.

Supporting information

S1 Fig. Genome-wide identification of Dal80-bound promoters. (A) Box-plot of the distance between the annotated TSS and ORF start site (translation initiation codon, ATG) for protein-coding genes. (B) Snapshot of ChIP-Seq signals along a GATA-less *locus* (*ALD6*). Densities (tag/nt) are shown for the untagged (black line) and Dal80-Myc¹³ (blue line) strains. ORFs and the Dal80-Myc¹³ ChIP-Seq peak are represented by grey arrows and the red dashed line, respectively. The snapshot was produced using the VING software [76].

S1 Table. Genome-wide identification of Dal80-bound promoters. List of 1269 gene promoters bound by Dal80.

S2 Table. Genome-wide identification of Dal80-bound promoters. Overlap of genes identified in our screens and in previous genome-wide expression screens.

S3 Table. Genome-wide identification of Dal80-bound promoters. GO term analysis of 1269 Dal80-bound promoters.

S4 Table. Genome-wide identification of Dal80-bound promoters. Whole genome GATA site contents of promoters and ORFs, correlated with Dal80 binding, regulation, NCR status and presence of Gcn4-binding sites.

S5 Table. Dal80 binds the intragenic region of a subset of genes. Inventory of gene promoters and ORFs bound by Dal80 and their intersection.

S6 Table. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels. List of 546 Dal80-regulated genes on proline.

S7 Table. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels. Gene ontology term analysis of the 232 genes repressed by Dal80 on proline.

S8 Table. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels. Gene ontology term analysis of the 314 genes activated by Dal80 on proline.

S2 Fig. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels. (A) Contingency table showing the number of Dal80-regulated or unaffected genes (defined upon RNA-Seq analysis in WT and *dal80Δ* cells) among the Dal80-bound and unbound genes (identified upon ChIP-Seq analysis in untagged and Dal80-Myc¹³ cells). Results that were experimentally observed or that were expected in case of independence are indicated in bold and in brackets, respectively. (B) Metagene view of the ChIP-Seq signal along the 216 Dal80-regulated and -bound genes (solid lines), and the 330 Dal80-regulated and -unbound genes (dashed lines), in the untagged (black) and Dal80-Myc¹³ (blue) strains. The shading surrounding each line denotes the 95% confidence interval. (C) Density-plot of RNA-Seq signal (tag/nt, log₂ scale) in the WT strain grown in proline-containing , for genes of the “unbound” (blue, n=4484), “P” (red, n=1125) and P&O” (black, n=144) classes. Y-axis: proportion of genes for each class. The highlighted areas correspond to the 75 (2%) and 170 (15%) genes of the “unbound” and “P” classes, respectively, showing a signal higher than

the median of the “P&O” class. A box-plot representation of the same RNA-Seq signals is shown on the top of the density-plot. (D) Same as c., highlighting the 949 (21%) and 632 (56%) genes of the “unbound” and “P” classes, respectively, showing a signal higher than the first quartile value for the “P&O” class.

S3 Fig. Dal80 binding across the ORF of well-characterized NCR-sensitive genes –

Control experiments. (A) Cells of FV078 (wt *DAL80-MYC*¹³) strain were grown in glutamine (Gln) or proline (Pro) as unique nitrogen sources. Anti-Myc ChIP was performed as described in Experimental procedures. Each histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of at least 3 independent cultures on which two IPs were performed. The associated error bars correspond to the standard error (DAL5_{P1-P2}, n=6; DAL5_{O3-O4}, n=5; GAP1_{P1-P2}, n=4; GAP1_{O9-O10}, n=4; GDH2_{P1-P2}, n=4; GDH2_{O5-O6}, n=4; MEP2_{P9-P10}, n=11; MEP2_{O1-O2}, n=7). Black histograms represent qPCR performed on promoter regions whereas grey ones indicate qPCR data obtained within genes. Primers are described in S10 Table: DAL5_{P1-P2}, DAL5_{O3-O4}, GAP1_{P1-P2}, GAP1_{O9-O10}, GDH2_{P1-P2}, GDH2_{O5-O6}, MEP2_{P9-P10}, MEP2_{O1-O2}. (B) Gal4-Myc¹³ binding at the *GAL1-10* promoter. Schematic representation of the *GAL1-10* locus. Position of the Gal4-binding sites (vertical segments across the locus line) and qPCR probes (black and grey lettered segments) are indicated. Cells of FV437 (wt *GAL4-MYC*¹³) strain were grown in YP with glucose or galactose as unique carbon sources. Anti-Myc ChIP was performed as described in Experimental procedures. Each histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of 2 independent cultures. The associated error bars correspond to the standard error (n=2). Black histograms represent qPCR performed on promoter regions whereas grey ones indicate qPCR data obtained within genes. Primers are described in S10 Table. (C) Binding of Dal80-Myc¹³ at the *MEP2* locus, in untreated or RNase-treated cells. Schematic representation of the *MEP2* locus. Position of the GATA sites (vertical segments above -sense GATA sites- or

below -antisense GATA sites- the *locus* line) and qPCR probes (black and grey lettered segments) are indicated. Cells of untagged (25T0b) or FV078 (*DAL80-MYC^{I3}*; wt) strains were grown at 29°C in the presence of proline (Pro) as unique nitrogen source. ChIP analysis was conducted with or without RNase treatment before immunoprecipitation. The first 3 black columns represent qPCR with 3 primer pairs within the *MEP2* promoter (*MEP2*_{P5-P6}; *MEP2*_{P13-P14}; *MEP2*_{P9-P10}) and the 3 following grey ones correspond to qPCR data obtained with 3 primer pairs within the *MEP2* ORF (*MEP2*_{O1-O2}; *MEP2*_{O11-O12} *MEP2*_{O9-O10}). Each histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of independent cultures. The associated error bars correspond to the standard error (n=3).

S4 Fig. Dal80 binding within gene bodies occurs independently of intragenic GATA sites

– **Control experiments.** (A) Expression of the non-NCR-sensitive gene *URA3* under the control of *P_{MEP2}*. Cells of wild type (25T0b) and *P_{MEP2}-URA3* (FV806) strains were cultivated at 29°C in the presence of proline (Pro) as unique nitrogen source. The expression of *URA3* is normalized to the expression of the *TBP1* gene. Each column corresponds to the mean value of data obtained for at least 2 independent cultures and the error bars correspond to the standard error (n=2). (B) PolIII occupancy at the *URA3* and *P_{MEP2}-URA3* loci. Cells of *DAL80-MYC^{I3}* wild type (FV078) and *P_{MEP2}-URA3* (FV808) strains were cultivated at 29°C in the presence of glutamine (Gln) or proline (Pro) as unique nitrogen sources. Anti-polIII (CTD4H8) ChIP was performed as described in Experimental procedures. Each histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of at least 2 independent cultures on which two IPs were performed. Histograms represent qPCR performed with *URA3*_{O1-O2} primers (n=2). (C) *UGA4* expression in a *DAL80ΔLZ* mutant strain. Cells of FV078 (*DAL80-MYC^{I3}*; wt) and FV136 (*DAL80ΔLZ-MYC^{I3}*; ΔLZ) strains were cultivated at 29°C in the presence of proline (Pro) as unique nitrogen source. Total RNA was isolated and *TBP1*-normalized *UGA4* mRNA levels were quantified by qRT-PCR using *UGA4*_{O1-O2}

primers. Histograms represent the averages of at least 2 independent experiments and the associated error bars correspond to the standard error (n=3). (D) Occupancy of the *UGA4* locus by PolII. Cells of FV078 (*DAL80-MYC¹³*; wt) and FV136 (*DAL80ΔLZ-MYC¹³*; ΔLZ) strains were grown at 29°C in the presence of proline (Pro) as unique nitrogen source. Anti-polII (CTD4H8) ChIP analysis was conducted as in Panel B using UGA4_{O1-O2} primers (n=2).

S5 Fig. Transcription elongation is necessary for Dal80 binding across *MEP2* – Control

experiments. (A) Effect of PolII elongation defects on *MEP2* expression. Wild type (FV673) or *rpb1-1* (FV675) *DAL80-MYC¹³* cells were grown at 29°C in the presence of glutamine (Gln) or proline (Pro) as unique nitrogen sources, and shifted at 37°C for one hour at mid-log phase. Total RNA was isolated and *TBPI*-normalized *MEP2* mRNA levels were quantified by qRT-PCR using MEP2_{O9-O10} primers. Histograms represent the average of at least 2 independent experiments and the associated error bars correspond to the standard error (n=2). (B) PolII occupancy at the *MEP2* locus. Wild type (FV673) or *rpb1-1* (FV675) *DAL80-MYC¹³* cells were grown at 29°C in the presence of glutamine (Gln) or proline (Pro) as unique nitrogen sources, and shifted at 37°C for one hour at mid-log phase. ChIP analysis was conducted as in S4 Fig using MEP2_{O9-O10} primers (n=4).

S9 Table. Strains used in this study. [50]

S10 Table. Primers used in this study.

Figure captions

Fig 1. Genome-wide identification of Dal80-bound promoters.

(A) Overview of the ChIP-Seq analysis. Biological duplicates of FV78 (Dal80-Myc¹³) and 25T0b (no tag) cells were grown to mid-log phase in proline-containing medium, and then

harvested. After chromatin extraction and sonication, Dal80-Myc¹³ was immunoprecipitated using α -Myc antibody. Co-precipitated DNA fragments were purified and used to construct ChIP-Seq libraries. After sequencing of the libraries, signals were computed using uniquely mapped reads. Dal80-bound regions were identified using a peak-calling procedure using MACS2. (B) Identification of Dal80-bound promoters. After peak-calling, Dal80-bound promoters were identified on the basis of a >75% overlap of the -100 to -350 regions (relative to the downstream ORF start site) by the peak (represented in as a red dashed line). (C) Heatmap view of the ChIP-Seq signal in the ATG +/- 500 bp region for the 1269 genes identified as bound by Dal80 at the promoter, in untagged (left) and Dal80-Myc¹³ cells (right). The colour goes from yellow to red as the signal increases. (D) Metagene view of the ChIP-Seq signal along the ATG +/- 600 bp region for the 1269 genes identified as bound by Dal80 at the promoter (solid lines) and for the unbound genes (dashed lines), in untagged (black) and Dal80-Myc¹³ cells (blue). For each group of genes, normalized coverage (tag/nt) for each gene was piled up, and the average signal was plotted. The shading surrounding each line denotes the 95% confidence interval. (E) Snapshot of ChIP-Seq signals along the *FUR4* locus. Densities (tag/nt) are shown for the untagged (black line) and Dal80-Myc¹³ (blue line) strains. ORFs and the Dal80-Myc¹³ ChIP-Seq peak are represented by grey arrows and the red dashed line, respectively. The position (and orientation) of each GATA site is represented by vertical segments above (sense GATA sites) or below (antisense GATA sites) the *locus* line. The snapshot was produced using the VING software (Descrimes *et al.*, 2015). (F) Number of GATA sites in the promoter of Dal80-unbound (4522) and promoter-bound (1268) genes. The analysis was performed using RSAT (Medina-Rivera *et al.*, 2015), across the -500 to -1 region (relative to the ATG codon of the downstream ORF).

Fig 2. Dal80 binds the intragenic region of a subset of genes.

(A) Identification of genes showing Dal80 intragenic binding. (B) Venn diagram showing the number of genes that are bound by Dal80 at the promoter and/or across the ORF. (C) Metagene view of the ChIP-Seq signal along the ATG +/- 600 bp region for the 1125 genes identified as bound by Dal80 at the promoter only ("P"), in untagged (black) and Dal80-Myc¹³ cells (blue). MetaSignal computation was as described in Fig 1d. The shading surrounding each line denotes the 95% confidence interval. (D) Same as above for the 144 genes bound by Dal80 at the promoter and across the ORF ("P & O"). (E) Same as above for the 45 genes bound by Dal80 across the ORF only ("O"). (F) Same as above for the 4484 genes that are not bound by Dal80.

Fig 3. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels.

(A) Scatter plot of densities (tag/nt, log₂ scale) for genes in WT and *dal80Δ* cells grown in proline-containing medium. Total RNA was extracted from exponentially growing biological replicates of 25T0b (WT) and FV080 (*dal80Δ*). After rRNA depletion, strand-specific RNA-Seq libraries were constructed and then sequenced. Tag densities were computed using uniquely mapped reads. Dal80-regulated genes were identified on the basis of a min. 2-fold change (up or down) in the *dal80Δ* mutant relative to the WT control, with a *P* < 0.05 upon differential expression analysis using DESeq [75]. Unaffected (n=5252), Dal80-repressed (n=232) and Dal80-activated (n=314) genes are shown as grey, red and blue dots, respectively. (B) Box-plot of densities (tag/nt, log₂ scale) in the WT strain grown in proline-containing medium, for genes of the unbound, "O", "P" and "P & O" classes. *P*-values obtained upon Wilcoxon rank-sum test are indicated.

Fig 4. Dal80 binding across the ORF of well-characterized NCR-sensitive genes.

(A-D) Position of GATA sites (vertical segments above -sense GATA sites- or below -

antisense GATA sites- the *locus* line) and qPCR probes (black and grey lettered segments) along 4 NCR-sensitive loci: *DAL5* (A), *GAP1* (B), *GDH2* (C) and *MEP2* (D). (E) Cells of FV078 (wt *DAL80-MYC*¹³) strain were grown in proline as unique nitrogen source. Anti-Myc ChIP was performed as described in Experimental procedures. Each histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of at least 3 independent cultures on which two IPs were performed. The associated error bars correspond to the standard error (*DAL5*_{P1-P2}, n=6; *DAL5*_{O3-O4}, n=5; *GAP1*_{P1-P2}, n=4; *GAP1*_{O9-O10}, n=4; *GDH2*_{P1-P2}, n=4; *GDH2*_{O5-O6}, n=4; *MEP2*_{P9-P10}, n=11; *MEP2*_{O1-O2}, n=7). Black histograms represent qPCR performed on promoter regions whereas grey ones indicate qPCR data obtained within genes. Dotted line represents the highest no tag control levels. Primers are described in S10 Table: a, *DAL5*_{P1-P2}; b, *DAL5*_{O3-O4}; c, *GAP1*_{P1-P2}; d, *GAP1*_{O9-O10}; e, *GDH2*_{P1-P2}; f, *GDH2*_{O5-O6}; g, *MEP2*_{P9-P10}; h, *MEP2*_{O1-O2}. (F) ChIP analysis was conducted as in Panel E, with or without RNase treatment before immunoprecipitation (n=3).

Fig 5. Dal80 binding within gene bodies occurs independently of intragenic GATA sites.

(A) Schematic representation of the *P_{MEP2}-URA3 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, *MEP2* promoter; grey segments, *URA3* ORF). (B) Occupancy of Dal80-Myc¹³ at the *MEP2* promoter and *URA3* ORF at the *MEP2* and *URA3 loci* in wild type and *P_{MEP2}-URA3* strains. Cells of *DAL80-MYC*¹³ wild type (FV078) and *P_{MEP2}-URA3* (FV808) strains were cultivated at 29°C in the presence of proline (Pro) as unique nitrogen source. ChIP analysis was conducted as in Fig 4. The first histogram summarizes the data obtained by qPCR probing the promoter of *MEP2* (a, *MEP2*_{P9-P10}), whereas the two following histograms are generated by qPCR performed with oligos positioned in the ORF of *URA3* (b, *URA3*_{O1-O2}; c, *URA3*_{O3-O4}), either at the *URA3 locus* (in wild type strain), or both at the *URA3* and *MEP2*

loci. Dotted line represents the highest no tag control levels. The associated error bars correspond to the standard error (n=6). (C) Schematic representation of the *UGA4 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, promoter; grey segments, ORF). (D) Wild type and truncated (Δ LZ) versions of Dal80-Myc¹³ used for the ChIP experiment at the *UGA4 locus*. (E) Occupancy of the *UGA4 locus* by Dal80-Myc¹³. Cells of FV078 (*DAL80-MYC*¹³; wt) and FV136 (*DAL80* Δ LZ-*MYC*¹³; Δ LZ) strains were grown at 29°C in the presence of proline (Pro) as unique nitrogen source. ChIP analysis was conducted as described in Fig 4. Black histograms represent qPCR performed with promoter-specific primers (a, UGA4_{P1-P2}; b, UGA4_{P9-P10}) whereas grey ones indicate qPCR data obtained with ORF-specific primers (c, UGA4_{O1-O2}; d, UGA4_{O3-O4}) at the *UGA4 locus*. The associated error bars correspond to the standard error (n=6).

Fig 6. Transcription elongation is necessary for Dal80 binding across *MEP2*.

(A) Schematic representation of the *MEP2 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line), and qPCR probe positions (black segments, promoter; grey segments, ORF). (B) Occupancy of Dal80-Myc¹³ at the *MEP2 locus* when transcription elongation is impaired. Wild type (FV673) or *rpb1-1* (FV675) *DAL80-MYC*¹³ cells were grown at 29°C in the presence of proline (Pro) as unique nitrogen source, and shifted at 37°C for one hour at mid-log phase. ChIP analysis was conducted as in Fig 4. The first 2 black columns represent qPCR with 2 primer pairs within the *MEP2* promoter (a, MEP2_{P5-P6}; b, MEP2_{P13-P14}) and the 2 following grey ones correspond to qPCR data obtained with 2 primer pairs within the *MEP2* ORF (c, MEP2_{O11-O12} d, MEP2_{O9-O10}). The associated error bars correspond to the standard error (n=6). (C) Coimmunoprecipitation of Dal80-Myc¹³ with different phosphoforms of PolII. Total proteins were extracted,

991 immunoprecipitated with the indicated antibodies, and subjected to anti-Myc western blot
992 analysis.

993 **Fig 7. Model summarizing the correlations observed between Dal80 binding, Dal80**
994 **dependency and expression levels.**

Fig 1. Genome-wide identification of Dal80-bound promoters.

Fig 2. Dal80 binds the intragenic region of a subset of genes.

(A)

(B)

Fig 3. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels.

Fig 4. Dal80 binding across the ORF of well-characterized NCR-sensitive genes.

(A)

MEP2 locus

URA3 locus

(B)

(C)

(D)

(E)

Fig 5. Dal80 binding within gene bodies occurs independently of intragenic GATA sites.

(A)

(B)

(C)

Dal80 co-immunoprecipitation

Fig 6. Transcription elongation is necessary for Dal80 binding across *MEP2*.

Fig 7. Model summarizing the correlations observed between Dal80 binding, Dal80 dependency and expression levels.