1 Classification: Biological Sciences

2 Title (135 characters incl. spaces)

3 Temperature but not the circadian clock determines nocturnal carbohydrate availability for
4 growth in cereals

5 Authors: Lukas M. Müller^{1,2*}, Leonard Gol^{1,2}, Jong-Seong Jeon³, Andreas P.M. Weber^{4,5}, Seth
6 J. Davis^{2,6}, Maria von Korff^{1,2,5*}

7 ¹ Institute for Plant Genetics, Heinrich-Heine University Düsseldorf, c/o Max-Planck-Institute
8 for Plant Breeding Research, Carl-von-Linné Weg 10, 50829 Köln/Cologne, Germany

9 ² Max-Planck-Institute for Plant Breeding Research, Carl-von-Linné Weg 10, 50829
10 Köln/Cologne, Germany

11 ³ Graduate School of Biotechnology, Kyung Hee University, Yongin, 446-701, Korea

12 ⁴ Institute of Plant Biochemistry, Heinrich-Heine University Düsseldorf, Universitätsstraße 1,
13 40225 Düsseldorf, Germany

14 ⁵ Cluster of Excellence on Plant Sciences (CEPLAS), "From Complex Traits towards Synthetic
15 Modules"

16 ⁶ Department of Plant Biology, University of York, Heslington, York, YO10 5DD, United
17 Kingdom

18 *Corresponding authors: Lukas M. Müller (lmueeller@mpipz.mpg.de), Maria von Korff
19 (korff@mpipz.mpg.de)

20 Other authors' email addresses: Leonard Gol (gol@mpipz.mpg.de), Jong-Seong Jeon
21 (jjjeon@khu.ac.kr), Andreas PM Weber (andreas.weber@uni-duesseldorf.de), Seth J Davis
22 (seth.davis@york.ac.uk)

23 ORCID ID: 0000-0002-6816-586X (M.v.K)

24 ORCID ID: 0000-0002-2089-0800 (L.M.M)

25 Keywords: circadian clock, plant growth, sucrose metabolism, starch degradation, cool
26 temperature, barley (*Hordeum vulgare*), *Arabidopsis thaliana*, rice (*Oryza sativa*), SUCROSE
27 TRANSPORTER 2 (SUT2), SUCROSE TRANSPORTER 4 (SUT4/SUC4)

28 Abstract (max 250 words):

29 The circadian clock is considered a key target for crop improvement because it controls
30 metabolism and growth in *Arabidopsis*. Here, we show that the clock gene *EARLY*
31 *FLOWERING 3* (*ELF3*) controls vegetative growth in *Arabidopsis* but not in the cereal crop
32 barley. Growth in *Arabidopsis* is determined by the degradation of leaf starch reserves at
33 night, which is controlled by *ELF3*. The vegetative growth of barley, however, is determined
34 by the depletion of leaf sucrose stores through an exponential kinetics, presumably catalyzed
35 by the vacuolar sucrose exporter *SUCROSE TRANSPORTER 2* (*SUT2*). This process depends
36 on the sucrose content and the nighttime temperature but not on *ELF3*. The regulation of
37 starch degradation and sucrose depletion in barley ensures efficient growth at favorable
38 temperature as stores become exhausted at dawn. On cool nights, however, only the starch
39 degradation rate is compensated against low nighttime temperatures, whereas the sucrose
40 depletion rate is reduced. This coincides with reduced biomass in barley but not in
41 *Arabidopsis* after growth in consecutive cool nights. The sucrose depletion metabolism
42 determines growth in the cereal crops barley, wheat, and rice but is not generally conserved
43 in monocot species and is not a domestication-related trait. Therefore, the control of growth
44 by endogenous (clock) versus external factors (temperature) is species-specific and depends
45 on the predominant carbohydrate store. Our results give new insights into the physiology of
46 growth in cereals and provide a basis for studying the genetics and evolution of different
47 carbohydrate stores and their contribution to plant productivity and adaptation.

48 Significance Statement (120 words max):

49 The circadian clock controls growth in the model plant *Arabidopsis thaliana* by regulating the
50 starch degradation rate so that reserves last until dawn. This prevents nocturnal starvation
51 until photosynthesis resumes. The cereal crops barley, wheat and rice, however,
52 predominantly consume sucrose instead of starch as carbohydrate source. We find that
53 carbohydrate supply from sucrose at night is regulated by enzyme kinetics and night-time
54 temperature, but not the circadian clock. We postulate that the regulation of growth depends
55 on the predominant carbohydrate store, where starch degradation is controlled by
56 endogenous cues (clock) and sucrose depletion by external factors (temperature). These
57 differences in the regulation of carbohydrate availability at night may have important
58 implications for adapting crops yields to climate change.

59 \body

60 **Introduction**

61 Plants, as photoautotrophic organisms, harvest light energy through photosynthesis during
 62 the day to drive growth, development, and reproduction. To supply the energy demand
 63 during the night, plants partition photosynthetic assimilates for transitory storage in leaves
 64 during the day and deplete them at night (1–3). Arabidopsis accumulates almost exclusively
 65 starch in its chloroplast during the day and uses the circadian clock to exhaust the starch
 66 stores in coincidence with the end of the night (4). This circadian control of starch
 67 degradation provides a benefit for growth at night because starch is neither unproductively
 68 sequestered nor prematurely depleted until photosynthesis resumes at dawn (1–3). The
 69 clock control of starch degradation is maintained in a wide range of conditions, including
 70 fluctuating and interrupted photoperiods (4), varying light intensity (5) and low night
 71 temperature (6). As a consequence, the circadian control of starch degradation in the
 72 chloroplast during the night is a key trait for optimal growth in Arabidopsis (4). However, the
 73 cereal crops barley, wheat and rice predominantly accumulate soluble carbohydrates,
 74 particularly sucrose, to high amounts in the leaf vacuole during the day and only little
 75 amounts of starch in the chloroplast (7–14). Furthermore, barley and wheat, but not rice,
 76 can transform abundant sucrose in the vacuole into the storage intermediate fructan during
 77 the day to mitigate the osmotic consequences of sucrose accumulation in the leaf cell (7).
 78 The conversion between sucrose and fructans is interrelated and determined by the
 79 concentration of vacuolar sucrose (7). Due to its molecular size and complex structure,
 80 fructans require to be converted back to sucrose before they can be released from the
 81 vacuole (7). Therefore, the export of sucrose from the vacuole but not the degradation of
 82 crystalline starch in the chloroplast, is the deciding pathway for carbohydrate supply in cereal
 83 crops during the night (8, 10, 11, 14, 15). Work from Gordon *et al.* and Farrar *et al.* and
 84 others in the 1980s have described carbohydrate fluxes in the barley leaf during the
 85 day/night cycle, but it remained unknown if the circadian clock controls depletion of sucrose
 86 and starch from the leaf (1, 10, 15–18) and what consequences result for overall growth
 87 when sucrose export from the vacuole but not starch degradation in the chloroplast
 88 dominates carbohydrate supply during the night (1, 11, 17). The circadian clock is
 89 structurally and functionally conserved between Arabidopsis and the cereal crops barley,
 90 wheat and rice (19–21), suggesting that the circadian regulation of metabolism and growth
 91 as observed in Arabidopsis offers a target for crop improvement (20, 21). However, the link
 92 between the circadian clock, carbohydrate metabolism and growth is not well understood in
 93 crops. Here, we investigate the link between temporal control of the carbohydrate supply
 94 during the night and vegetative growth in the cereals in comparison to Arabidopsis.

95 **Results and discussion**

96 The clock component ELF3 determines metabolism and growth in Arabidopsis but not in 97 barley

98 We measured the accumulation and the turnover of carbohydrates in short photoperiods (8 h
99 light/16 h dark) to compare the nocturnal carbohydrate metabolism in barley and Arabidopsis
100 leaves. Barley primarily accumulated sucrose during the day, followed by small amounts of
101 fructans and starch but did not store fructose and glucose (Figure 1a). At the end of the
102 night, all the sucrose, fructan and starch was consumed (Figure 1a). This confirmed earlier
103 reports stating that soluble storage carbohydrates from the vacuole, respectively sucrose,
104 but not crystalline starch from the chloroplast dominate the carbohydrate supply in the
105 barley leaf (10, 16, 22, 23). As a consequence, barley consumed around five times more
106 carbohydrates from vacuolar sucrose than from starch during the night (Figure 1a). If the
107 conversion of fructans into sucrose before export from the vacuole was considered (7), even
108 seven times more carbohydrates were exported in the form of sucrose from the vacuole than
109 provided from starch degradation in the chloroplast (Figure 1a). In contrast, Arabidopsis
110 accumulated starch in the leaf during the day but no sucrose or glucose and consumed
111 nearly all starch reserves during the night (Figure 1b). As a consequence, Arabidopsis
112 depended almost exclusively on the degradation of chloroplastic starch for carbohydrate
113 supply in the dark (Figure 1b). We then tested if the carbohydrate supply during the night is
114 under control of the circadian clock in barley and Arabidopsis. We measured the depletion of
115 sucrose and the degradation of starch in the leaf of wild-type and *early flowering 3 (elf3)*-
116 mutant plants of barley (*hvelf3*) (24) and Arabidopsis (*atelf3*). The oscillator in the *elf3*-
117 mutant of both species is severely disturbed as it is arrhythmic in constant light and partly
118 arrested in day/night cycles (24, 25). Depletion of sucrose from the barley leaf did not differ
119 between *hvelf3* and wild-type plants during the night (Figure 1c). However, starch degraded
120 significantly faster in the last half of the night in *hvelf3*-plants compared to wild-type plants
121 (Figure 1d). In Arabidopsis, sucrose levels remained low throughout the night and did not
122 differ between *atelf3* and wild-type plants (Figure 1e). By contrast, the degradation of
123 transitory starch terminated prematurely 2 h before the end of the night in *atelf3* while it
124 continued its linear trend until the end of the night in wild-type plants (Figure 1f). These
125 findings demonstrated that a functional ELF3 was required for the temporal control of starch
126 breakdown at the end of the night in both barley and Arabidopsis, although species-specific
127 differences in the degradation pattern existed. In contrast, the depletion of sucrose from the
128 vacuole in barley was not under the control of ELF3. Sucrose depletion was also not affected
129 by the shortfall of starch degradation in the barley *hvelf3*-mutant at the end of the night
130 (Figure 1c, d). The Arabidopsis *atelf3*-mutant, however, was impaired in starch degradation
131 which represented the primary carbohydrate source during the night. To demonstrate that

atelf3 experienced carbohydrate starvation after the termination of starch degradation before dawn (Figure 1f), we analyzed the expression of the sugar-repressed genes At3g59940 and BRANCHED CHAIN AMINO ACID TRANSFERASE 2 (BCAT2) as starvation reporters (4, 26) (Figure 1g, h). Both transcripts were upregulated 2 h before the end of the 24 h cycle in Arabidopsis *atelf3*-mutant plants but not in wild-type plants (Figure 1g, h). Wild-type plants only increased expression of the starvation markers when they were kept in extended darkness beyond 24 h after dawn (Figure 1g, h). As even small but repetitive reductions of carbohydrate supply at night reduce growth in plants (2–4), we tested if the *elf3*-mutants of barley and Arabidopsis were impaired in biomass accumulation. We found that the *elf3*-mutant of Arabidopsis, but not that of barley, showed a significantly reduced fresh and dry weight in comparison to wild-type plants in the growth chamber and in the greenhouse at different stages of vegetative growth (Figure 2a-c, Supplementary Figure 1). Therefore, our data demonstrates that nocturnal carbohydrate supply for metabolism and growth is determined by the circadian clock component ELF3 in Arabidopsis but not in barley. This explains why only *atelf3*, but not *hvel3*, reduces biomass accumulation during vegetative growth.

Sucrose depletion from the leaf is controlled by the sucrose content and the night temperature through an exponential kinetics

Sucrose levels in the barley *elf3*-mutant were almost exhausted at the end of the night 24 h after dawn although sucrose depletion was not controlled by ELF3 (Figure 1c). We, therefore, investigated if nocturnal sucrose depletion from the leaf is controlled by a clock-independent mechanism to adjust carbohydrate supply to the length of the night. We grew wild-type plants under cycles of 8 h light/16 h dark and measured the sucrose content over the whole diel cycle. During the night, sucrose depletion was exponential and sucrose levels 24 h after dawn were identical to those 24 h before (Figure 3a). This cycle demonstrated that sucrose depletion is under temporal control. To analyze the adaptability of sucrose depletion to the length of the night, we exposed plants to an unexpected extension of the light period from 8 h to 12 h. This extension increased the sucrose content in the leaf at the end of the light period and enhanced the sucrose depletion rate at night so that levels were comparable between entrained and extended photoperiods at the end of the night 24 h after dawn (Figure 3a). This demonstrated that the sucrose content at the end of the light period relates to the rate of sucrose depletion during the night so that sucrose is almost completely exhausted in the leaf 24 h after dawn, even after unexpected variation of the light period. We modeled the sucrose export from the leaf during the night as a function of the sucrose concentration and fitted the equation of first-order chemical kinetics $[suc]_t = [suc]_0 e^{-kt}$ to the measured data. Chemical kinetics of the first order describes enzyme catalyzed reactions that depend, at constant temperature, only on the substrate concentration. We were able to

approximate the sucrose content in the leaf over time based on the sucrose content at the beginning of the night and extracted the depletion constant k to quantify the rate of sucrose depletion. When the light period was unexpectedly extended from 8 h to 12 h, the rate of sucrose depletion was 0.03 units higher than under entrained conditions ($k \approx 0.18$ vs. $k \approx 0.15$), reflecting the increased sucrose depletion rate that was necessary to deplete a higher amount of sucrose in a shorter night (Figure 3a). To further investigate the adaptability of the sucrose depletion kinetics, we analyzed barley plants shifted from 12 h of light to an unexpected 16, 20, and 24 h period of light and released them to constant darkness (Figure 3b). An unexpected extension of the light period from 12 h to 16 h increased the sucrose content in the leaf and increased the sucrose export rate in the dark for 0.08 units from $k \approx 0.22$ to $k \approx 0.3$ (Figure 3b). However, an unexpected extension of the light period from 12 h to 20 h or even 24 h saturated the sucrose content in the leaf and the depletion rate in the dark did not increase much further in comparison to 16 h of light ($k \approx 0.3$ vs. $k \approx 0.31$) (Figure 3b). Together, this demonstrated a close relationship between the sucrose content in the leaf at dusk and the depletion rate during the following night.

As the activity of metabolic enzymes is temperature-dependent (27), we tested if the night temperature influences the kinetics of sucrose depletion from the leaf. After an unexpected reduction of the night temperature from entrained 18°C to 10°C at the onset of darkness, the sucrose content in the leaf was significantly higher throughout the night at 10°C and the rate of sucrose depletion during the night was 0.02 units lower at 10°C compared to 18°C ($k \approx 0.17$ and $k \approx 0.19$, respectively) (Figure 3c). Therefore, less sucrose was turned over in the leaf during the night at 10°C compared to 18°C. The initial delay in sucrose depletion in the night at 10°C in Figure 3c was probably due to a temperature shock that plants experienced when they were transferred from 20°C to the dark pre-cooled growth chamber of 10°C at the end of the light period. However, even after this initial delay, sucrose depletion rates were lower under 10°C compared to 18°C. This reduction in the sucrose depletion rate in cool nights was comparable to the change observed after an unexpected extension of the day by 4h (change in k after temperature drop in Figure 3c: $\Delta k \approx 0.2$, change in k after day extension in Figure 3a: $\Delta k \approx 0.3$). In contrast, the starch content was, after a similar initial delay, not significantly different throughout the night between 10°C and 18°C. Consequently, the rate of starch degradation was identical between the cool and the entrained night conditions (Figure 3d). This demonstrated that an internal mechanism was capable to compensate the rate of starch degradation, but not that of sucrose depletion, against a sudden reduction in night temperature. We then investigated how the sucrose depletion and the starch degradation adapted to consecutive cool nights. We grew 3 week old barley wild-type plants for one further week at 20°C during the day and either the entrained 18°C or the cool 10°C during the night. The light intensity during the day was identical for both treatments. At the seventh day, plants grown in the 10°C cool night

contained around three times more sucrose in the leaf at the end of the night than those from 18°C (Figure 3e). In contrast, the starch content was not significantly different between both temperature treatments at the end of the night and similar to the onset of the experiment at day 0 (Figure 3e). This demonstrated that the nocturnal sucrose depletion in barley does not adapt to consecutive cool nights and is under environmental and not endogenous control. It also indicated that the effects of consecutive cool nights amplify the incomplete turnover of sucrose in barley as observed for a single cool night in Figure 3c. When taken together, our findings demonstrate that the sucrose content and the night temperature control the depletion of sucrose from the barley leaf in the dark following first-order chemical kinetics. Variation in the sucrose content at the onset of the night alters the depletion rate so that sucrose levels exhaust at dawn, even under unexpected variation of the photoperiod. On the contrary, low night temperatures reduce the sucrose depletion rate, resulting in an incomplete turnover of sucrose during the night. This effect is magnified by consecutive cool nights. Starch degradation, on the other hand, is controlled by an endogenous mechanism in barley that involves the circadian clock component ELF3 and is capable of compensating the degradation rate against low night temperatures.

Cool night temperatures reduce growth (biomass accumulation) in species predominantly consuming sucrose instead of starch during the night

Pyl *et al.* (2012) (6) identified unused capacity in biosynthetic pathways and processes for growth in Arabidopsis plants in cool nights. They concluded that the carbohydrate supply but not the biochemical pathways involved in growth limits biomass accumulation during nights at low temperature (6). They demonstrated in Arabidopsis that the systemic control of starch degradation through the circadian clock adjusts enzyme activity for starch breakdown to low temperatures so that the rate of starch degradation and, as a consequence, the carbohydrate supply and, subsequently, growth was compensated against cool nights (6). As barley consumes both sucrose and starch during the night but only the degradation of starch was compensated against low night temperature (Figure 3 c-e), we tested if the reduction in sucrose depletion from the barley leaf during cool nights affects growth. We confirmed that growth in Arabidopsis plants is compensated against cool nights of 10°C as fresh weight biomass did not differ between plants grown for one week in cool versus control nights (Figure 4a). However, the fresh weight of barley plants grown in 10°C nights for one week was significantly lower than that of plants grown under control night temperature of 18°C (Figure 4b). Therefore, the reduced sucrose turnover in the barley leaf at low night temperatures correlated with reduced growth. This suggested that low night temperatures reduced carbohydrate supply and, subsequently, nocturnal growth in barley. We then investigated if the sucrose depletion metabolism from barley and its associated growth reduction in cool nights is a domestication-related trait and if it is present in further monocot

species. Wild barley, wheat and rice metabolized approximately three times more sucrose than starch during the night (Figure 5a, Supplementary Figure 2). By contrast, *Brachypodium* consumed 1.4-fold more starch than sucrose during the night (sucrose-to-starch ratio: 0.7, Figure 5a, Supplementary Figure 2). Wild barley, rice, and wheat, but not *Brachypodium*, significantly reduced fresh weight after one week of growth in cool nights (Figure 5b). These findings showed that the sucrose depletion metabolism and its association with reduced growth in cool nights is prevalent in several agriculturally important cereal crops. This includes rice, barley and wheat but not all monocot species as nocturnal growth in *Brachypodium* primarily depends on starch instead of sucrose and was compensated against low night temperature. Moreover, the sucrose depletion metabolism is apparently not a domestication trait because it is present in both wild and cultivated barley (Figure 5a, b). When taken together, we observed that the primary carbohydrate store correlated with differences in the growth response of monocot species to cool nights. It is likely that the reduced carbohydrate supply from sucrose depletion in cool nights is responsible for the growth reduction, but the causal link between the major carbohydrate store at night and growth responses to external versus internal cues awaits further investigations.

Sucrose export from the vacuole catalyzed by SUCROSE TRANSPORTER 2 regulates the sucrose depletion metabolism and affects growth and yield

Finally, we investigated the molecular basis of the sucrose depletion metabolism in cereal crops. Previous studies in barley, wheat and rice demonstrated that the SUCROSE TRANSPORTER 2 (SUT2) is located in the vacuolar membrane and catalyzes the export of sucrose through a proton coupled co-transport (13, 14, 28, 29). Isolation of the *sut2*-mutant in rice provided evidence *in planta* that SUT2 is necessary for the sucrose export from the vacuole into the cytoplasm as *sut2*-mutant plants over-accumulated sucrose in the leaf vacuole at dawn (14). As rice, like barley, depends on the sucrose depletion metabolism for carbohydrate supply at night (Figure 5a, b) and the barley *sut2*-mutant was not available to us, we investigated in rice if SUT2 is involved in the regulation of the nocturnal sucrose depletion metabolism. The sucrose depletion metabolism is based on a positive relationship between the sucrose content in the leaf at dusk and the amount of sucrose depleted in an exponential fashion during the following night (Figure 3a, b). While sucrose depletion was exponential with a constant of $k \approx 0.12$ in wild-type plants, sucrose depletion in the *sut2*-mutant was rather linear than exponential due to the low depletion constant of $k \approx 0.03$ (Figure 6a). In addition, despite the nearly two-fold higher sucrose content at dusk, the *sut2*-mutant depleted only around half of the amount of sucrose per Gram fresh weight (mg/gFW) until dawn than wild-type plants (13mg and 22mg sucrose/gFW, respectively) (Figure 6a). This demonstrated that SUT2 function is required for the exponential depletion of sucrose from the leaf based on the sucrose content. This disturbance of the sucrose

depletion metabolism correlated with a strong reduction of growth, biomass and 1000 kernel weight in the *sut2*-mutant (Figure 6b-d). This confirmed that the sucrose export from the leaf vacuole during the night affects growth and yield. The functional redundancy of the SUT2 transporter in rice, barley and wheat (13, 14, 28, 29) and the predominant consumption of sucrose instead of starch in these species during the night (Figure 5a) together with the growth reduction in cool nights (Figure 5b) suggests that the vacuolar sucrose exporter SUT2 is also a key regulator of the sucrose depletion metabolism and growth in barley and wheat. By contrast, Arabidopsis plants, which accumulate almost exclusively starch in the chloroplast during the day but almost no sucrose in the vacuole (Figure 1b), do not express the homologue of OsSUT2, AtSUT4/SUC4, in the leaf (30). This highlights the different regulatory principles that exist in plants for the depletion of sucrose and the degradation of starch in the leaf during the night and their specific effects on growth.

Conclusion

Our results suggest that metabolism and growth in plants at night is determined by the predominant nocturnal carbohydrate stores of starch and sucrose. The nocturnal starch turnover is controlled by the circadian clock, whereas sucrose depletion is determined by vacuolar sucrose concentrations at the end of the day and night-time temperatures. This endogenous versus external regulation of carbohydrate supply correlates with variation in biomass accumulation between sucrose and starch storing species. In this context, it remains to be understood why different species either use starch or sucrose or a combination of both as transient storage carbohydrates for the night. The starch storing mechanism might buffer growth against environmental fluctuations, but the sucrose depletion metabolism may offer a growth advantage over the starch degradation metabolism because the carbohydrate supply from low molecular sucrose is energetically advantageous compared to the synthesis and the degradation of high molecular starch. However, the nocturnal sucrose export and the subsequent growth are susceptible to cool nights. A "hybrid" system with sucrose and starch as seen in *Brachypodium* may combine the benefits from sucrose-driven growth under favorable and starch-driven growth under unfavorable conditions. Barley, wheat and rice might accumulate little amounts of starch during the day to mitigate a growth reduction in cool nights.

We demonstrate that the physiological control of growth in important cereal crops cannot be resolved in the current model species Arabidopsis and *Brachypodium*, despite the conservation of clock components in crop plants (17, 20, 21, 31). We propose that the sucrose depletion metabolism gives new insight into the physiology of growth and yield in cereal crops. In addition, the monocot lineage with variation in carbohydrate storage

317 provides the opportunity to study the genetics, evolution and adaptive significance of
318 carbohydrate use and growth in plants. A mechanistic understanding of growth as controlled
319 by sucrose versus starch is important for the adaptation of crops to climate change.

320

Acknowledgments:

This work was supported by the Max Planck Society through an International Max Planck Research School fellowship to L.M.M., by an ERASysApp project ("CropClock"), a trilateral DFG grant ("Circadian clock and stress adaptation in barley") and by the Excellence Cluster EXC1028.

Author contributions:

L.M.M. conceived the study and conducted the experiments; L.M.M. and M.v.K. designed the experiments; L.G. contributed experimental data; L.M.M., M.v.K., A.P.M.W., and S.J.D. analyzed the data; J-S.J. contributed new material; L.M.M. and M.v.K. prepared the manuscript.

The authors declare no competing financial interests.

Correspondence and requests for materials should be addressed to L.M.M. or M.v.K.

Materials and Methods

Plant material and growth conditions:

The spring cultivar Bowman (BW) was used as the wild-type barley. The introgression line BW290, which carries an introgression of the *eam8.k* allele in the background of Bowman, was denoted by *hvelf3*. The *eam8.k* allele is characterized by a base-pair mutation leading to a premature stop codon in *HvELF3*, which is orthologous to *ELF3* in Arabidopsis (24). For Arabidopsis, the *elf3-4* mutant in the Wassilewskija (WS) background was used (32). Both barley and Arabidopsis were grown in growth chambers at 450 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux density (PPFD) at 20°C during the day and 18°C during the night as standard growth conditions. For the cool night treatment, standard growth conditions were applied during the day and only the nighttime temperature was altered to 10°C. The significant difference in biomass between *hvelf3* and wild-type plants was confirmed in a temperature-controlled greenhouse (20°C during the day, 18°C during the night). Both barley and Arabidopsis seeds were directly sown into soil (Einheitserde) and stratified for 4 days at 4°C in the dark.

The rice mutant *ossut2* carried a T-DNA insertion in the fifth exon of the *SUT2* gene in the Hwayong background (14). The rice seeds were directly sown into potting soil (Einheitserde) and cultured in growth chambers under 12 h light at 28°C and 350-450 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux density (PPFD) and 12 h dark at 25°C.

Diel/circadian sampling:

Barley and rice plants were harvested during vegetative growth and grown to the three-leaf stage. The whole second leaf of two plants was cut at the base and placed into a 2 ml Eppendorf tube (200-300 mg) as one biological replicate. For Arabidopsis, plants were grown for 4 weeks before complete rosettes from at least 4 different plants were pooled to one biological replicate (100-200 mg). At least 4 different biological replicates were sampled per time point, genotype and plant species and snap frozen in liquid nitrogen. The time points of sampling were applied as given in the respective diagrams of the individual experiments.

Biomass measurements:

Biomass was measured as the fresh weight of the shoot on an accuracy balance directly after cutting from the rootstock. Individual shoots were collected, oven dried and measured for dry weight. At least 20 biological replicates were measured per genotype for the biomass analysis. Arabidopsis plants were harvested before the first bud became visible, while the development of the monocot species was established by dissecting the shoot apical meristem and scoring the development of the main shoot apex as described in (33). This ensured that biomass measurements captured the vegetative and not the reproductive growth.

Measurement of sugars and starch:

Frozen samples were ground to powder using milling beads. The soluble sugar fraction was separated from the insoluble starch fraction by boiling the samples twice in 1 ml of 80% (vol/vol) ethanol at 85°C for 30 min. Chlorophyll was precipitated from the soluble fraction by adding 0.5 ml of chloroform and 1 ml of water. The clear alcoholic extract was concentrated in a vacuum concentrator and resuspended in sterile water. The content of sucrose, glucose, fructose, and maltose was measured using a photometer at 340 nm wavelength as NADPH absorbance in a glucose-6-phosphate dehydrogenase (G6PDH)-dependent assay and quantified against a glucose standard curve using glucose/sucrose enzyme kits from r-biopharm (Cat. No. 11 113 950 035).

To measure fructan content, the resuspended concentrate was hydrolyzed with 37% HCl for 20 min at 80°C, then neutralized with 7.67 M NaOH and measured as fructose against the fructose background using an enzyme kit to measure fructose (r-biopharm, Cat. No. 10 139 106 35).

For the measurement of starch, the insoluble fraction was boiled in KOH at 95°C for 45 min, neutralized with acetic acid (HAc), digested to glucose by α -amylase and amyloglucosidase at room temperature overnight, and quantified as glucose.

Equation of first-order enzyme kinetics:

The equation $[\text{suc}]_t = [\text{suc}]_0 e^{-kt}$ was fitted to the sucrose content at night to describe the depletion kinetics. $[\text{suc}]_t$ is the sucrose concentration in the leaf at any given time t during the night, $[\text{suc}]_0$ is the initial sucrose concentration at the beginning of the night, and k is the depletion constant, which is a measure of the depletion rate.

Differences between treatments or genotypes were analyzed via a t-test ($p \leq 0.01$). The number of biological replicates was at least 4 for one time point in the time series data and at least 20 for biomass measurements.

Transcript Analysis:

Five whole rosettes were sampled and pooled per biological replicate, for a total of 4 biological replicates per time point. Extraction of RNA, reverse transcription and qRT-PCR were then performed as previously described (Digel et al., 2015). Briefly, RNA was extracted using the TRIzol reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. The DNase treated RNA was transcribed into first strand cDNA using Superscript II reverse transcriptase (Thermo Fisher Scientific) in accordance to the manufacturer's instructions. Two technical replicates were performed for each sample and absolute quantification was based on a titration curve of cloned fragments for each target gene. All qRT-PCR reactions were performed on a LightCycler 480 (Roche; Software version 1.5) using the following primers: AtBCAT2 FW 5'-GTACAACGCACAAGCTGCAT-3', RV 5'-ACCCGAGATTATCCCAATCC-3'; At3g59940 FW 5'-TGGAACGATGATGGTGAAGA-3', RV 5'-AACCAGAGGGAGTGTGACG-3'. The geometric mean of AtPP2A (FW 5'-CAGCAACGAATTGTGTTTGG-3', RV 5'-AAATACGCCCAACGAACAAA-3') and AtPEX4 (FW 5'-TTACGAAGGCGGTGTTTTTC-3', RV 5'-GGCGAGGCGTGTATACATTT-3') absolute expression was used to calculate relative gene expression values.

References

1. Graf A, Smith AM (2011) Starch and the clock: the dark side of plant productivity. *Trends Plant Sci* 16(3):169–175.
2. Stitt M, Zeeman SC (2012) Starch turnover: pathways, regulation and role in growth. *Curr Opin Plant Biol* 15(3):282–292.
3. Smith A, Stitt M (2007) Coordination of carbon supply and plant growth. *Plant Cell Environ* 30(9):1126–1149.
4. Graf A, Schlereth A, Stitt M, Smith A (2010) Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. *Proc Natl Acad Sci U S A* 107(20):9458–9463.
5. Scialdone A, et al. (2013) Arabidopsis plants perform arithmetic division to prevent starvation at night. *Elife* 2. Available at: <http://elife.elifesciences.org/content/2/e00669.abstract>.
6. Pyl E-T, et al. (2012) Metabolism and Growth in Arabidopsis Depend on the Daytime Temperature but Are Temperature-Compensated against Cool Nights. *Plant Cell Online*. doi:10.1105/tpc.112.097188.
7. Pollock CJ, Cairns AJ (1991) Fructan Metabolism in Grasses and Cereals. *Annu Rev Plant Physiol Plant Mol Biol* 42(1):77–101.
8. Hirose T, et al. (2013) Disruption of a rice gene for α -glucan water dikinase, OsGWD1, leads to hyperaccumulation of starch in leaves but exhibits limited effects on growth. *Front Plant Sci* 4(147). doi:10.3389/fpls.2013.00147.
9. Nie G, Hendrix DL, Webber AN, Kimball BA, Long SP (1995) Increased accumulation of carbohydrates and decreased photosynthetic gene transcript levels in wheat grown at an elevated CO₂ concentration in the field. *Plant Physiol* 108(3):975–983.
10. Gordon AJ, Ryle GJA, Mitchell DF, Powell CE (1982) The dynamics of carbon supply from leaves of barley plants grown in long or short days. *J Exp Bot* 33(2):241–250.
11. Smith A, Zeeman S, Smith S (2005) Starch degradation. *Annu Rev Plant Biol* 56:73–98.
12. Kaiser G, Heber U (1984) Sucrose transport into vacuoles isolated from barley mesophyll protoplasts. *Planta* 161(6):562–568.
13. Endler A, et al. (2006) Identification of a Vacuolar Sucrose Transporter in Barley and Arabidopsis Mesophyll Cells by a Tonoplast Proteomic Approach. *Plant Physiol* 141(1):196–207.
14. Eom J-S, et al. (2011) Impaired Function of the Tonoplast-Localized Sucrose Transporter in Rice, OsSUT2, Limits the Transport of Vacuolar Reserve Sucrose and Affects Plant Growth. *Plant Physiol* 157(1):109–119.
15. Sicher RC, Kremer DF, Harris WG (1984) Diurnal Carbohydrate Metabolism of Barley Primary Leaves. *Plant Physiol* 76(1):165–169.
16. FARRAR SC, FARRAR JF (1985) CARBON FLUXES IN LEAF BLADES OF BARLEY. *New Phytol* 100(3):271–283.
17. Müller LM, von Korff M, Davis SJ (2014) Connections between circadian clocks and

- 451 carbon metabolism reveal species-specific effects on growth control. *J Exp Bot*.
452 Available at:
453 <http://jxb.oxfordjournals.org/content/early/2014/04/03/jxb.eru117.abstract>.
- 454 18. Webb AAR, Satake A (2015) Understanding circadian regulation of carbohydrate
455 metabolism in arabidopsis using mathematical models. *Plant Cell Physiol* 56(4):586–
456 593.
- 457 19. Campoli C, Shtaya M, Davis S, von Korff M (2012) Expression conservation within the
458 circadian clock of a monocot: natural variation at barley Ppd-H1 affects circadian
459 expression of flowering time genes, but not clock orthologs. *BMC Plant Biol* 12(1):97.
- 460 20. Hsu PY, Harmer SL (2013) Wheels within wheels: the plant circadian system. *Trends*
461 *Plant Sci* (0).
- 462 21. Bendix C, Marshall CM, Harmon FG (2015) Circadian clock genes universally control
463 key agricultural traits. *Mol Plant* (0).
- 464 22. GORDON AJ, RYLE GJA, POWELL CE, MITCHELL D (1980) Export, Mobilization, and
465 Respiration of Assimilates in Uniculm Barley during Light and Darkness. *J Exp Bot*
466 31(2):461–473.
- 467 23. GORDON AJ, RYLE GJA, WEBB G (1980) The Relationship between Sucrose and
468 Starch during `Dark` Export from Leaves of Uniculm Barley. *J Exp Bot* 31(3):845–850.
- 469 24. Faure S, et al. (2012) Mutation at the circadian clock gene EARLY MATURITY 8 adapts
470 domesticated barley (*Hordeum vulgare*) to short growing seasons. *Proc Natl Acad Sci*.
471 doi:10.1073/pnas.1120496109.
- 472 25. Hicks KA, et al. (1996) Conditional Circadian Dysfunction of the Arabidopsis early-
473 flowering 3 Mutant. *Science* (80-) 274(5288):790–792.
- 474 26. Usadel B, et al. (2008) Global transcript levels respond to small changes of the carbon
475 status during progressive exhaustion of carbohydrates in Arabidopsis rosettes. *Plant*
476 *Physiol* 146(4):1834–1861.
- 477 27. Atkin OK, Tjoelker MG (2003) Thermal acclimation and the dynamic response of plant
478 respiration to temperature. *Trends Plant Sci* 8(7):343–351.
- 479 28. Weschke W, et al. (2000) Sucrose transport into barley seeds: molecular
480 characterization of two transporters and implications for seed development and starch
481 accumulation. *Plant J* 21(5):455–467.
- 482 29. Deol KK, et al. (2013) Identification and characterization of the three homeologues of
483 a new sucrose transporter in hexaploid wheat (*Triticum aestivum* L.). *BMC Plant Biol*
484 13(1). doi:10.1186/1471-2229-13-181.
- 485 30. Schneider S, et al. (2012) Vacuoles release sucrose via tonoplast-localised SUC4-type
486 transporters. *Plant Biol* 14(2):325–336.
- 487 31. Filichkin SA, et al. (2011) Global Profiling of Rice and Poplar Transcriptomes
488 Highlights Key Conserved Circadian-Controlled Pathways and *cis*-
489 Regulatory Modules. *PLoS One* 6(6):e16907 EP-.
- 490 32. Hicks KA, Albertson TM, Wagner DR (2001) EARLY FLOWERING3 Encodes a Novel
491 Protein That Regulates Circadian Clock Function and Flowering in Arabidopsis. *Plant*
492 *Cell Online* 13(6):1281–1292.
- 493 33. Digel B, Pankin A, von Korff M (2015) Global Transcriptome Profiling of Developing

494 Leaf and Shoot Apices Reveals Distinct Genetic and Environmental Control of Floral
495 Transition and Inflorescence Development in Barley. *Plant Cell* 27(9):2318–2334.
496
497

498 Figures:

499

500 **Figure 1: Sucrose depletion dominates carbohydrate supply from the barley leaf**
 501 **during the night and is not controlled by ELF3 while starch degradation**
 502 **dominates carbohydrate supply from the Arabidopsis leaf and requires ELF3 to**
 503 **prevent starvation before dawn.**

504 **a), b)** Content of carbohydrates in the leaf of a) barley and b) Arabidopsis after the end of
 505 the day and the end of the night in 8 h light/16 h dark photoperiods. *N*=4 per time point. **c),**

d) Content of c) sucrose and d) starch in the leaf of barley *elf3*-mutant and wild-type plants during the night in 12 h light/12 h dark photoperiods. *N=4 per time point*. **e), f)** Content of e) sucrose and f) starch in the leaf of Arabidopsis *elf3*-mutant and wild-type plants during the night in 8 h light/16 h dark photoperiods. *N=4 per time point*. **g), h)** Expression of the starvation reporters g) At3g59940 and h) AtBCAT2 in Arabidopsis *elf3*-mutant and wild-type plants before dawn (light grey) and in extended darkness (dark grey). *N=4 per time point*.

*Significant differences per time point (t-test, $p \leq 0.01$) are marked with *. Error bars are the standard deviation of the mean throughout.*

515

516

517 **Figure 2: The clock component ELF3 controls biomass accumulation during**
518 **vegetative growth in Arabidopsis but not in barley.**

519 **a)** Fresh and dry weight of *elf3*-mutant and wild-type plants from barley and Arabidopsis
520 during vegetative growth before transition to reproductive growth. $N \geq 20$. **b), c)** Phenotypes
521 of b) barley and c) Arabidopsis *elf3*-mutant and wild-type plants.

522 Boxes show the 25 and 75% percentile, whiskers denote the 5-95% percentile, the line
523 denotes the median and the square denotes the mean. Significant differences were tested by
524 *t*-test ($p \leq 0.01$). White scale bars represent 1cm.

Figure 3: The sucrose content and the temperature control the depletion of sucrose from the barley leaf at night while starch degradation is compensated against low temperature.

a) Leaf sucrose content in wild-type barley plants entrained to a photoperiod of 8 h light/16 h dark and then kept in 8 h light/16 h dark (8/16 LD, blue) or unexpectedly shifted to 12 h light/12 h dark (12/12 LD, red). $N=4$ per time point. **b)** Leaf sucrose content in wild-type barley plants entrained to 12 h light/12 h dark and then kept in 12 h light (12 L, green) and released to constant darkness (DD) or unexpectedly shifted to an extended light period of 16 h (16 L, red), 20 h (20 L, blue) or 24 h (24 L, orange) before being released to constant darkness (DD). $N=3$ per time point. **c), d)** Content of c) sucrose and d) starch in the leaf of barley wild-type plants entrained to 12 h light at 20°C and 12 h dark at 18°C before plants were exposed to either an unexpected cool night at 10°C (blue) or kept in the entrained night temperature of 18°C (red). $N=4$ per time point. **e)** Sucrose and starch content at dawn in three week old barley plants grown in 12 h light at 20°C and 12 h dark at 18°C (Day 0) and after seven days of growth at 20°C during the day and either 18°C or 10°C during the night (Day 7).

Significant differences per time point (t -test, $p \leq 0.01$) are marked with *. Error bars are the standard deviation of the mean throughout. Boxes show the 25 and 75% percentile, whiskers denote the 5-95% percentile, the line denotes the median and the square denotes the mean.

Figure 4: Barley, but not Arabidopsis, reduces biomass accumulation during one week of growth in cool nights.

a), b) Fresh weight of a) Arabidopsis and b) barley wild-type plants after one week of growth in the entrained night temperature of 18°C or cool nights of 10°C. Plants were grown at 20°C during the day in 12h light/ 12 h dark photocycles and the light intensity was identical in both treatments.

*Significant differences per time point (t-test, $p \leq 0.0001$) are marked with ***. Boxes show the 10, 25, 75 and 90% percentile, whiskers denote the 5-95% percentile, the line denotes the median and the square denotes the mean.*

Figure 5: Biomass is reduced during growth in cool nights in several monocot species when they predominantly consume sucrose instead of starch during the night.

a) Ratio of sucrose to starch consumed during the night in 12 h light/12 h dark photoperiods in several monocot species. A ratio of 1 indicates equal consumption of sucrose and starch during the night. **b)** Fresh weight of species depicted in a) after one week of growth at a low night time temperature compared to the entrained favorable night time temperature for the particular specie. $N=50$.

Significant differences per time point (t -test, $p \leq 0.01$) are marked with *. Boxes show the 10, 25, 75 and 90% percentile, whiskers denote the 5-95% percentile, the line denotes the median and the square denotes the mean.

Figure 6: Sucrose export from the vacuole catalyzed by SUCROSE TRANSPORTER 2 (SUT2) in rice plays a key role in the regulation of the sucrose depletion metabolism.

a) Sucrose depletion from the leaf during a 12 h night of rice *sut2*-mutant and wild-type plants. **b), c), d)** Loss of SUT2-function reduces b) growth, c) biomass and d) 1000 kernel weight in rice.

Significant differences per time point (*t*-test, $p \leq 0.01$) are marked with *. Boxes show the 10, 25, 75 and 90% percentile, whiskers denote the 5-95% percentile, the line denotes the median and the square denotes the mean.

Supplementary Figures:

Supplementary Figure 1: Fresh weight and dry weight of barley *elf3*-mutant and wild-type plants grown in the greenhouse in early summer one week after emergence from the soil. *N*=30.

Supplementary Figure 2: Sucrose and starch content at the end of a 12 h day and a 12 h night in several monocot species. *N*=5 per time point.