1 **The association between alcohol consumption and** 2 **telomere length: A meta-analysis focusing on** 3 **observational studies**

4

5

6 Jianqiang Li^{1,2¶}, Yu Guan^{1¶}, Faheem Akhtar^{1,3}, Xi Xu¹, Ji-Jiang Yang^{4*}, Shi Chen⁵,
7 Qing Wang⁴, Hui Pan⁵, Weiliang Qiu⁶

8

9 ¹ Faculty of Information Technology, Beijing University of Technology, Beijing,
10 China.

11 ² Beijing Engineering Research Center for IoT Software and Systems, Beijing, China.

12 ³ Department of Computer Science, Sukkur IBA University, Sukkur, Sindh, Pakistan.

13 ⁴ Tsinghua National Laboratory for Information Science and Technology, Tsinghua
14 University, Beijing, China.

15 ⁵ Department of Endocrinology, Peking Union Medical College Hospital, Chinese
16 Academy of Medical Sciences & Peking Union Medical College, Beijing, China.

17 ⁶ Channing Division of Network Medicine, Brigham and Women's Hospital/Harvard
18 Medical School, Boston, USA.

19

20 ¶These authors contributed equally to this work.

21

22 *Corresponding author:

23 E-mail: Jijiang_Yang@126.com (JY)

25 **Abstract**

26 **Background**

27 Both telomere length and alcohol consumption play important roles in carcinogenesis
 28 and biological age. Many efforts have been made to investigate the association
 29 between alcohol consumption and telomere length. However, no consensus has been
 30 reached yet.

31 **Methods**

32 In this article, we performed a meta-analysis to integrate the investigation results in
 33 the literature about the association between alcohol consumption and telomere length.
 34 After searching articles published between 2000 and 2016, 21 articles (including 27
 35 analyses, total sample size 35,891) met our eligibility criteria.

36 **Results**

37 We found a significant association between alcohol consumption and telomere length
 38 (Fisher's combined p-value = 3.52E-8 and Liptak's weighted p-value = 8.24E-3). We
 39 also found that the significance of the association between alcohol consumption and
 40 telomere length varies with study type (cohort, case-control, or cross-sectional) and
 41 study population (Europe, Asia, American, or Australia).

42 **Conclusions**

43 Combined evidence showed that alcohol consumption is associated with telomere
 44 length. The consistent quantifications of alcohol consumption and telomere length
 45 would benefit the future aggregation of the evidence from different studies.

46

47

48 **Introduction**

49 A telomere is a segment of repetitive nucleotide sequences at both tails of a
 50 chromosome, which protects the tail of the chromosome from deterioration or fusion
 51 with neighboring chromosomes[1]. Over time, the end of the chromosome becomes
 52 shorter during cell replications[1]. Once the telomere length (TL) is shortened to a
 53 critical length, the chromosomes would become unstable and the cells will
 54 immediately activate the apoptotic mechanism and lose viability[2]. So TL reflects the
 55 cell copy history and replication potential, known as the "mitotic clock" of the cell's
 56 lifespan[1]. Defects in telomere length have been linked to several age-related
 57 diseases, premature aging syndromes, and cancers[3]. Telomere shortening may result
 58 in genomic instability during the initial stage of tumorigenesis[4].

59
 60 Although telomere shortening occurs as a natural part of aging, it is known to be
 61 affected by several factors including age, gender, ethnicity, paternal age at birth,
 62 genetic mutations of telomerase, and telomere maintenance genes[5]. Telomere
 63 shortening also could be accelerated by mechanisms like oxidative stress and
 64 inflammation[5]. Furthermore, several studies suggest that psychosocial,
 65 environmental, and behavioral exposures can impact TL as well[6]. Since alcohol
 66 exposure is one of the behavioral exposures as well as a source of oxidation[7]. It
 67 would be interesting to investigate the effect of alcohol consumption (AC) on
 68 telomere length.

69
 70 Alcohol is one of the most widely used recreational drugs in the world. Alcoholic
 71 drinks are classified by the International Agency for Research on Cancer (IARC) as a
 72 Group 1 carcinogen (carcinogenic to humans)[8]. IARC classifies alcoholic drink

consumption as a cause of colorectal, larynx, liver, esophagus, oral cavity, pharynx, and female breast cancers; and as a probable cause of pancreatic cancer[8-10]. The World Health Organization estimates that as of 2010 there were 208 million people with alcoholism worldwide[11]. AC is the world's third-largest risk factor for public health; in middle-income countries, which constitute almost half of the world's population, it is the greatest risk factor for public health[12]. Also, it could reduce a person's life expectancy by around ten years[13].

Many efforts have been made to investigate the association between AC and TL. However, no consensus has been reached yet about the AC-TL association. Some studies showed significant inverse associations between TL and alcohol use[14-16]. In other words, the more alcohol consumption, the shorter TL. However, a few studies reported significant positive associations between AC and TL. For example, Liu et al. (2009)[17] observed significantly longer TL in ever drinkers than that in never drinkers for controls of gastric cancer. Liu et al. (2011) observed significantly longer TL in ever drinkers than that in never drinkers in hepatocellular carcinoma (HCC) patients[18]. Also, several studies reported no association between AC and TL[19-30]. To facilitate the investigation of the AC-TL association, we conducted a meta-analysis.

Methods

Search strategy and selection criteria.

This meta-analysis is based on a comprehensive and systematic search of ten research

97 databases: ProQuest, Science Online, Wiley-Blackwell, PubMed, Science Direct,
 98 Google Scholar, Nature, Baidu scholar (<http://xueshu.baidu.com/>), Chinese National
 99 Knowledge Infrastructure (CNKI, <https://en.wikipedia.org/wiki/CNKI>,
 100 <http://www.cnki.net/>), and Chongqing VIP (<http://lib.cqvip.com/>), the last three of
 101 which are Chinese research databases. Full-text available articles in English or
 102 Chinese, published or in press between January 2000 and December 2016 were
 103 considered. Since the effect of alcohol on health is usually manifested after a long
 104 time period and clinical trials (RCTs) were usually completed in a relatively
 105 short-time period, we only focused on observational studies in this meta-analysis.

107 **First round of selection.**

108 The first round of selection was based on title and abstract according to the search
 109 terms (“telomere”, “alcohol”, and “ethanol”). If either the set (“telomere” and
 110 “alcohol”) or the set (“telomere” and “ethanol”) were found in title or abstract, this
 111 article would be included for further screening.

113 **Second round of selection.**

114 When it came to the second round of selection, only papers with full text available
 115 would be kept. The papers that did not mention any association between AC and TL
 116 were also excluded. We then grouped the remaining papers based on the types of the
 117 original studies: random clinical trial or observational study.

119 **Third round of selection.**

120 Since the effect of alcohol on health is usually manifested after a long time period and

121 clinical trials (RCTs) were usually completed in a relatively short-time period, we
 122 excluded RCT studies in the third round. In the third round, we further excluded
 123 papers with low document quality.
 124
 125 Document quality evaluations were indispensable before any data processing, but
 126 criteria vary depending on the type of article. We applied NEWCASTLE-OTTAWA
 127 QUALITY ASSESSMENT (NOS) SCALE[31] to evaluate the quality of cohort
 128 studies and case-control studies. Meanwhile, we applied the 11-item checklist that
 129 was recommended by Agency for Healthcare Research and Quality (AHRQ) to
 130 evaluate the quality of cross-sectional studies. Based on NOS Scale, a study can be
 131 awarded a maximum of one star for each numbered item within the Selection and
 132 Outcome (for cohort studies)/Exposure (for case-control studies) categories, a
 133 maximum of two stars can be given for Comparability. The NOS ranges from zero up
 134 to nine stars[31]. According to AHRQ checklist, an item would be scored '0' if it was
 135 answered 'NO' or 'UNCLEAR'; if it was answered 'YES', then the item scored '1'.
 136 Article quality was assessed as follows: low quality = 0–3; moderate quality = 4–7;
 137 high quality = 8–11[32].
 138
 139 To pass the third round screening, we require that a cohort study or a case-control
 140 study needs to have NOS Scale ≥ 6 , and a cross-sectional study need to have AHRQ $\geq$
 141 7.
 142
 143 Study selection was performed by YG and double-checked by JL and WQ. All
 144 disagreements were resolved through consensus by the three authors. The procedure
 145 of study selection is illustrated in **Fig 1**.

146 **Fig 1. The flowchart of study selection**

147

148

149 **Extraction of relevant information.**

150 After the three rounds of screening, 21 articles met the requirements and were

151 selected for further analysis. We extracted the following information relevant to the

152 present meta-analysis: study type (case-control, cohort, or cross-sectional),

153 significance of the association between AC and TL (test statistics and p-values),

154 source of the population, gender, sample size, age, ethnicity, country of the

155 participants, statistical models, quantification of TL, and quantification of AC.

156

157 **Classification of analyses.**

158 We regarded a study as a case-control study if it compared TL between two AC

159 groups (e.g., alcohol abusers versus social drinkers). We regarded a study as a cohort

160 study if it tested the association of AC measured at baseline to TL measured at the end

161 of follow-up or to the change of TL from baseline to the end of follow-up. We

162 regarded a study as a cross-sectional study if it evaluated the association of

163 continuous-type AC to TL based on data measured at the same time period (e.g., both

164 measured at the baseline or both measured at the end of follow-up).

165

166 **Statistical Analysis.**

167 The standard method to combine evidence from independent studies is the

168 meta-analysis[33], in which the weighted average of the test statistics from individual

169 studies is used to pool evidence. However, it is challenging to get appropriate test

170 statistics for the present meta-analysis. Several different statistical models were used
171 in the related articles to test for the association of AC to TL, such as t-test and general
172 linear regression used in case-control studies, correlation analysis and general linear
173 regression used in cross-sectional and cohort studies. In some studies, AC was treated
174 as continuous variable. In other studies, AC was treated as categorical variables with
175 different categorizations (e.g., two-category, three-category, or four-category). Even
176 for the same number of categories, the definitions of categories might also be different.
177 For example, Pavanello et al. (2011)[14] defined three categories as 0 – 1
178 drink-units/day; 2-4 drink-units/day; > 4 drink-units/day, while Houben et al.
179 (2010)[23] defined three categories as 0 gram /day; 1-19 gram/day; >=20 gram/day.
180 Similarly, the measurements of TL also varied among the 21 articles. Most of the
181 studies regarded TL as a continuous variable. Houben et al. (2010)[23] and Kozlitina
182 et al. (2012)[24] categorized TL to tertiles, Weischer et al. (2014)[34] categorized TL
183 to quartiles, while Cassidy et al. (2010)[19] categorized TL to quintiles. Hence, the
184 interpretations of the test statistics in different studies would be different, indicating
185 that it is not appropriate to combine test statistics. Moreover, some studies did not
186 provide the values of test statistics. Fortunately, almost all of the studies provided
187 p-values and sample sizes. Therefore, we performed a meta-analysis by calculating
188 the combined p-value as pooled evidence about the association of AC to TL. If the
189 combined p-value < 0.05, we claim that AC is significantly associated with TL.
190 Several methods for combining p-values have been proposed. In the present study, we
191 used Fisher's method. In addition, to utilize the information of the sample sizes, we
192 used Liptak's method[35].
193
194 Since different studies used different statistical models to investigate the association

of AC to TL and did not report test statistics and their standard errors of the effect sizes, we could not directly access the potential publication bias and heterogeneity among studies. In this article, to roughly assess the heterogeneity, we used $-\log_{10}(\text{p-value})$ to surrogate the effect size of each study and used the inverse of sample size to surrogate the variance of the effect size. We then used $-\log_{10}(\text{Fisher's combined p-value})$ and $-\log_{10}(\text{Liptak's combined p-value})$, respectively, as the pooled effect size to calculate I^2 . To roughly assess the publication bias, we drew funnel plot by using signed $-\log_{10}(\text{p-value})$ to surrogate the effect size and using the square root of the inverse of sample size as the standard error. The sign of the effect size for a study is the same as the sign of the test statistic. If test statistic is missing, we assumed negative sign (i.e., the higher AC, the shorter TL). R package *metafor* was used to draw the funnel plot.

In this meta-analysis, we also applied Fisher's exact test to check if the following factors affect the significance of the association between AC and TL: (1) study type (cohort study, case-control study, or cross-sectional study), (2) article goal (if testing for AC-TL association is the primary goal of the article), (3) sex-specificity (men only study, female only study, or both sex study), (4) categorization of AC (whether containing never-drinker category), and (5) study population (American, Asian, Australian, or European). For binary factors (e.g., article goal), p-values of Fisher's exact test are obtained by directly using the hypergeometric distribution. Otherwise, the network developed by Mehta and Patel (1983, 1986)[36] and improved by Clarkson, Fan, and Joe (1993)[37] was used to calculate the p-values of Fisher's exact test. We also applied two-sample Wilcoxon rank sum tests to check if sample size and mean age affect the AC-TL association.

220

221 A test is claimed as significant if its two-sided p-value < 0.05. All analyses were

222 performed by using IBM SPSS Statistics Version 22.0 or R Version 3.4.2.

223

224

225 **Results**

226 After the three rounds of screening, we obtained 21 articles. The information

227 (including quality scores) of these 21 articles is listed in **Table 1**. Six of the 21 articles

228 performed more than one analyses. There were 44 analyses in total that evaluated the

229 association between AC and TL in the 21 articles. The information (including quality

230 scores) of these 44 analyses is listed in **S1 Table** online. We evaluated the 27

231 independent analyses and discussed the remaining 17 analyses in the Discussion

232 section. Among the 27 analyses (**Table 1**), there are four case-control studies[14, 15,

233 17, 18], three cohort studies[16, 34, 38], and 20 cross-sectional studies[14, 16, 19-30,

234 34, 38-40]. Among the remaining 17 analyses, there are ten case-control studies, three

235 cohort studies, and four cross-sectional studies.

236

237 Since different studies used different statistical models to investigate the association

238 of AC to TL and did not report test statistics and their standard errors of the effect

239 sizes, we could not directly access the potential publication bias and heterogeneity

240 among studies. In this article, we used surrogate test statistics and surrogate standard

241 errors. If we used $-\log_{10}(\text{Fisher's combined p-value})$ as the pooled effect size, the

242 value of I^2 is equal to 34.42, indicating small to medium heterogeneity. If we used

243 $-\log_{10}(\text{Liptak's combined p-value})$ as the pooled effect size, the value of I^2 is equal to

244 0, indicating small heterogeneity. **Fig 2** shows roughly symmetric funnel plot,
 245 indicating no publication bias.

246 **Fig 2. Funnel plot of surrogate effect size and surrogate standard error**

247

248

Table 1 The information of the 27 independent studies

Study type	First Author	publish Year	studyDesign Original	primary Analysis	sigAssoc	both Sex	Pvalue	testStat	nTotal	mean Age	sepNon Drinker	continent
crossSectional	Harris [21]	2006	cross-sectional	No	No	both	0.962	NA	185	79.1	N	Europe
crossSectional	Bekaert [30]	2007	Longitudinal study	No	No	women only	0.682	-7.453	1,291	45.9	N	Europe
crossSectional	Bekaert[30]	2007	Longitudinal study	No	No	men only	0.63	-8.033	1,218	46.1	N	Europe
crossSectional	Hou [22]	2009	case-control study	No	No	both	0.06	NA	416	65.5	Y	Europe
crossSectional	Mirabello [27]	2009	nested case-control study	No	No	men only	0.799	0.006	1,661	64	N	America
crossSectional	Houben [23]	2010	Longitudinal study	No	No	men only	0.28	NA	203	78.47	Y	Europe
crossSectional	Mather [39]	2010	cross-sectional	No	Yes	both	0.008	-0.155	646	56.75	N	Australia
crossSectional	Cassidy [19]	2010	cross-sectional	Yes	No	women only	0.59	NA	2,284	58.86	N	America
crossSectional	Fyhrquist [20]	2011	Longitudinal study	No	No	women only	0.962	NA	668	65	NA	Europe
crossSectional	Fyhrquist [20]	2011	Longitudinal study	No	No	men only	0.962	NA	603	63	NA	Europe
crossSectional	Pavanello [14]	2011	case-control study	Yes	Yes	men only	0.003	NA	457	41	Y	Europe
crossSectional	Strandberg [16]	2012	Longitudinal study	Yes	No	men only	0.11	-0.08	499	75.7	N	Europe
crossSectional	Kozlitina [24]	2012	Longitudinal study	No	No	both	0.526	NA	3,157	50	N	America
crossSectional	Sun [29]	2012	cross-sectional	Yes	No	women only	0.93	0.001	5,862	58.7	N	America
crossSectional	Marcon [26]	2012	cross-sectional	No	No	both	0.962	-0.006	56	56	N	Europe

(Continued)

Table 1 (Continued)

Study type	First Author	publish Year	studyDesign Original	primary Analysis	sigAssoc	both Sex	Pvalue	testStat	nTotal	mean Age	sepNon Drinker	continent
crossSectional	Bendix [38]	2014	Longitudinal study	No	No	both	0.078	-0.001	2,214	55	N	Europe
crossSectional	Weischer [34]	2014	Longitudinal study	Yes	Yes	both	0.01	NA	4,576	54.25	N	Europe
crossSectional	Latifovic [25]	2015	cross-sectional	Yes	No	both	0.57	(coef=-0.014)+/-(se=0.046) for abstainer; -0.055+/-0.039 for moderate AC; -0.024+/-0.050 for high AC	477	35	Y	America
crossSectional	Starnino [28]	2016	Longitudinal study	No	No	both	0.23	-1.205	132	45.34	N	America
crossSectional cohort	Shin [40]	2016	cross-sectional	Yes	Yes	both	0.04	NA	1,771	57.23	Y	Asia
cohort	Strandberg [16]	2012	Longitudinal study	Yes	Yes	men only	0.004	-0.13	499	46.7	N	Europe
cohort	Bendix [38]	2014	Longitudinal study	No	Yes	both	0.032	-0.001	1,356	44.7	N	Europe
cohort	Weischer [34]	2014	Longitudinal study	Yes	No	both	0.61	-0.154	4,535	54.25	N	Europe
case-control	Liu [17]	2009	case-control study	No	Yes	both	0.016	meanDiff=0.07	378	53	Y	Asia
case-control	Pavanello [14]	2011	case-control study	Yes	Yes	men only	0.00009	NA	457	38	Y	Europe
case-control	Aida [15]	2011	case-control study	Yes	Yes	both	0.002	NA	50	67.25	Y	Asia
case-control	Liu [18]	2011	case-control study	No	Yes	both	0.043	meanDiff=0.04	240	50.5	Y	Asia

N, No; Y=Yes, NA=Not Available (i.e., missing value); studytype, indicating the study type of the AC-TL association; firstAuthor, indicating the last name of the first author; publishYear, indicating the article publication year; studyDesignOriginal, indicating the original study design of

the article; primaryAnalysis, indicating if the primary goal of the study is to investigate the AC-TL association; sigAssoc, indicating if the AC-TL association is significant (* $p < 0.05$); bothSex, indicating if the AC-TL association study is based on women-only data, men-only data, or based on both women and men data; pvalue, the p-value for testing the AC-TL association; testStat, test statistic; nTotal, the number of total sample size in the AC-TL association study; meanAge, mean age of subjects in the AC-TL association study; sepNonDrinker, indicating if the AC-TL association study includes a non-drinker category; continent, indicating which continent the subjects were from.

249 Among the 27 studies, 11 studies have primary goals to detect the AC-TL association.
 250 Among the 11 studies, two were cohort studies[16, 34], two were case-control
 251 studies[14, 15], and seven were cross-sectional studies[14, 16, 19, 25, 29, 34, 40]. The
 252 other 16 studies had various primary goals, from the association between TL and red
 253 blood cell size[24] to the association between TL and mortality in humans[38].

254
 255 As for sex-specificity, eight studies are men only studies[14, 16, 20, 23, 27, 30], four
 256 studies are women only studies[19, 20, 29, 30], and 15 studies contain both men and
 257 women[15, 17, 18, 21, 22, 24-26, 28, 34, 38-40]. As for the categorization of AC, nine
 258 studies have the category of non-alcohol drinkers[14, 15, 17, 18, 22, 23, 25, 40], while
 259 two studies did not provide this information[20]. As for study population, 16 studies
 260 were based on populations in Europe[14, 16, 20-23, 26, 30, 34, 38], six studies were
 261 based on populations in America[19, 24, 25, 27-29], four studies were based on
 262 Asian[15, 17, 18, 40], and one study was based on Australian[39]. The range of total
 263 number of samples among the 27 articles is from 50[15] to 5862[29]. The mean age is
 264 ranged from 35 year old[25] to 79 year old[21]. The p-value is ranged from 9.00E-5 to
 265 0.962.

266
 267 Ten out of the 27 studies reported significant associations between AC and TL (i.e.,
 268 p-value < 0.05). The Fisher's combined p-value of the 27 studies is 5.75E-8 and the
 269 Liptak's combined p-value is 8.76E-3.

270
 271 Two out of the three cohort studies and all four case-control studies reported
 272 significant AC-TL associations, while only four out of the 20 (20%) cross-sectional
 273 studies reported significant AC-TL associations. For the association between study

274 type and AC-TL association, the cross-table is in **S2 Table** online and the parallel pie
275 chart is in **Fig 3**. Fisher's exact tests showed that study type is significantly associated
276 with the significance of AC-TL association (p-value=1.86E-3).

277 **Fig 3. Parallel pie chart: The association between study type and AC-TL**
278 **association**

279

280

281 All six studies from America reported non-significance of the AC-TL association,
282 while all four studies from Asia and the only one study from Australia reported
283 significant AC-TL association. Five out of 16 studies from Europe reported
284 significant AC-TL association. For the association between continent and AC-TL
285 association, the cross-table is in **S3 Table** online and the parallel pie chart is in **Fig 4**.
286 Fisher's exact test showed that study population (p-value = 2.67E-3) is significantly
287 associated with the significance of the AC-TL association.

288 **Fig 4. Parallel pie chart: The association between continent and AC-TL**
289 **association**

290

291

292 The significance of AC-TL association is not related to whether a study's primary
293 goal is to test for AC-TL association (p-value=0.22), whether a study is based only on
294 men, only on women, or on both men and women (p-value=0.39), or whether a study
295 has an AC category: never drinker (p-value=0.09). Cross-tables of the association
296 between these factors and AC-TL association are in **S4-S6 Tables** online,
297 respectively.

298

299 Wilcoxon rank sum tests showed that both the total number of sample size
300 (p-value=0.51) and mean age (p-value=0.11) are not associated with the significance
301 of AC-TL association. Parallel boxplots of the association between these factors and
302 AC-TL association are in **S1** and **S2 Figs** online.

303

304

305 **Discussion**

306 In this article, we conducted a meta-analysis to summarize the pieces of evidence in
307 the literature of observational studies about the association between alcohol
308 consumption (AC) and telomere length (TL), and to identify factors that might affect
309 the significance of the AC-TL association. There are 21 eligible articles included in
310 this meta-analysis, containing 44 studies of the AC-TL association. The pooled
311 evidence from the 27 independent studies showed that AC is significantly associated
312 with TL (Fisher's combined p-value=3.52E-8; Liptak's combined p-value=8.24E-3).
313 Study type (p-value=1.86E-3) and study population (p-value=2.67E-3) are
314 significantly associated with the significance of the AC-TL association.

315

316 Using all 44 studies and ignoring the dependency among the 44 studies, The AC-TL
317 association still remained significant (Fisher's combined p-value=1.11E-16 and
318 Liptak's combined p-value=3.57E-5).

319

320 If we considered only the four case-control studies and the 20 cross-sectional studies,
321 whether or not having an AC category "never drinker" is significantly associated with
322 the significance of the AC-TL association (p-value=0.026). All four case-control
323 studies reported significant AC-TL associations, while only four out of 20

324 cross-sectional studies reported significant AC-TL association. By observing that the
325 controls in all four case-control studies are never drinkers and that only five out of the
326 20 cross-sectional studies had the never-drinker category, we guess that the effect of
327 study type on the significance of the AC-TL association is probably due to the
328 comparison of ever drinkers to never drinkers. Parallel pie chart and the cross-table of
329 the association between whether a study has an AC category as never-drinker and the
330 AC-TL association are in **S3 Fig** and **S7 Table** online.

331
332 Since there is only one study from Australia, we regroup study population to four Asia
333 studies and 23 non-Asia studies; the study population is still significantly associated
334 with the significance of the AC-TL association (p-value=0.012). Parallel pie chart and
335 the cross-table of the association between whether the population of a study was
336 Asian and the AC-TL association are in **S4 Fig** and **S8 Table** online. It would be
337 interesting to investigate why all four studies from Asia reported significant AC-TL
338 association.

339
340 It is interesting to observe that all four women-only studies reported non-significant
341 associations between AC-TL, although sex-specificity is not significantly associated
342 with the significance of the AC-TL association (p-value=0.3948). More women-only
343 studies are needed to confirm if this observation is by chance or not.

344
345 Theoretically, longitudinal study (cohort study) would be better than cross-sectional
346 study or case-control study in that cohort study could infer if heavy alcohol
347 consumption causes shorter telomere length. However, there are several challenges
348 for cohort studies[38]. First, it would be hard to make sure the procedure to measure

349 TL at the end of long follow-up is the same as that at the baseline. Second, the blood
350 storage method would also be different between baseline and the end of follow-up.
351 Third, it would be difficult to make sure TL technically behaves in the same manner
352 at baseline as that at the end of follow-up.

353

354 Genetics may play an important role in the AC-TL association. Pavanello et al. (2011)
355 showed that carriers of the common ADH1B*1/*1 (rs1229984) genotype were more
356 likely to be alcohol abusers, while exhibiting shorter TL[14]. Shin et al. (2016)
357 showed that heavy alcohol consumption was inversely associated with leukocyte TL
358 only among carriers of the mutant alleles (CT and TT) of rs2074356 (ALDH2)[40].
359 These studies indicate genetics could help identify subtypes of subjects that have
360 significant AC-TL associations.

361

362 Several studies only provided lower or upper boundaries of p-values. For instance,
363 when evaluating the effect of interaction between AC and ADH1B genotypes on TL
364 among 255 controls, Pavanello et al. (2011) showed that the p-value < 0.001 [14].
365 Pavanello et al. (2011) showed p-values < 0.0001 for another four studies[14]. Harris
366 et al. (2006) showed p-value > 0.05 [21]. When we calculated Fisher's combined
367 p-value and Liptak's weighted p-value, we set p-value = 0.0009 for the study with
368 p-value < 0.001 , set p-value = 0.00009 for the four studies with p-value < 0.0001 , and
369 set p-value = 0.962 (the maximum p-value among the 44 studies) for the study with
370 p-value > 0.05 .

371

372 Several studies provided mean age only for each group of subjects (e.g., for each TL
373 tertile in Houben et al. (2010)[23]). We took the average of the means of these groups

374 when we investigated if age would affect the significance of the AC-TL association.

375

376 Six of the 21 articles included more than one study that investigated the association

377 between AC and TL. Among the 27 studies, two studies (one is men only study and

378 the other is women-only study) were from Bekaert et al. (2007)[30]. The same is for

379 Fyhrquist et al. (2011)[20]. Pavanello et al. (2011) included two studies: one is

380 case-control study, and the other is cross-sectional study[14]. Stranberg et al. (2012)

381 and Weischer (2014) both included two studies: one is cross-sectional study, and the

382 other is cohort study[16, 34]. We assumed that the significance of the AC-TL

383 association in different types of studies within the same article is independent.

384

385 The quantifications of TL are quite different among the 27 analyses. Sixteen analyses

386 quantified TL as proportion or ratio. Among the 16 analyses, one study used

387 normalized telomere-to-centromere ratio (NTCR)[15] and fifteen studies used the

388 relative telomere to single copy gene (T/S) ratio[14, 17-19, 22, 25, 27, 28, 34, 38-40].

389 One study assessed TL with Z-score of T/S ratio[29]. One study assessed TL by

390 terminal restriction fragment (TRF)[26]. Nine analyses used kilobase (kb) as the unit

391 of TL. Three analyses divided TL into three[23, 24] or four categories[34]. Three

392 analyses took the logarithm of TL[20, 40]. The difference in the TL quantification

393 makes it difficult to aggregate data and interpret results. In future, it would be

394 beneficial to have guidance on how to standardize the quantification of TL.

395

396 The quantifications of AC are also quite different among the 27 analyses.

397 Twenty-three analyses measured AC on a continuous scale as alcoholic beverages per

398 week[17, 21, 25, 28, 30, 38, 39], beverages per day[14, 29, 34], grams (g) per

399 week[16], and g per day[18, 19, 23, 24, 26, 27, 40]. Among them, a standard alcoholic
 400 beverage widely varied from 12 oz. beer (1 bottle), 5 oz. wine (1 glass), 1.5 oz. liquor,
 401 to 12 g alcohol. One analysis measured AC as total drink-years (yearly frequency *
 402 total years of alcohol use)[22]. Other studies used categorized AC. One defined an
 403 ever-drinker as a person drinking at least one serving of beer (12 oz.), wine (4 oz.), or
 404 liquor (1.5 oz.) per month for ≥ 6 months, while a study used a four-category AC
 405 categorization: nondrinker, <10, 10-29, and >29 years of drinking alcohol[22]. Two
 406 analyses took the natural logarithm of AC[30]. Three analyses did not report the
 407 quantifications of AC[15, 20]. In future, it would be beneficial to have guidance on
 408 how to standardize the quantification of AC.

409
 410 Among the 27 main studies investigated in the present study, 15 studies reported test
 411 statistic values, four of which reported positive associations between AC and TL[17,
 412 18, 27, 29]. That is, the more alcohol consumption is, the longer telomere length is.
 413 Two of the four positive associations are statistically significant[17, 18]. Liu et al.
 414 (2009)[17] gave a potential explanation the significant positive association:
 415 telomerase activation by alcohol drinking in target tissues. They also mentioned that
 416 they could not rule out the possibility of chance findings because of the limited size in
 417 each subgroup[17]. We hypothesize that the positive AC-TL association is due to
 418 moderate drinking, which is beneficial to human health. However, Liu et al. (2009)
 419 did not mention how they defined “ever drinkers”[17]. Further investigation is
 420 warranted to study the differences of the AC-TL association among never drinkers,
 421 moderate drinkers, and heavy drinkers.

422
 423 In summary, the present meta-analysis aggregated the p-values of 27 related studies

(total sample size 35,891) and showed a significant association between alcohol consumption and telomere length based on Fisher's combined p-value and Liptak's weighted p-value, although the different quantification methods of AC and TL hindered the use of meta-analysis to aggregate the evidence for the AC-TL association. Factors, such as study type (case-control, cross-sectional, or cohort study), study population (American, Asian, or European), and subject gender, might affect the AC-TL association. In future, it would help us uncover the true relationship between alcohol consumption and telomere length if we standardize the quantifications of AC and TL and if we take account for both beneficial and deteriorating effects of AC to human health.

434

435

436 **Supporting information**

437 **S1 Fig. Parallel boxplots of the association between total number of sample size**
438 **and AC-TL association.**

439 (PDF)

440 **S2 Fig. Parallel boxplots of the association between mean age and AC-TL**
441 **association.**

442 (PDF)

443 **S3 Fig. Parallel pie chart: The association between whether a study has an AC**
444 **category as never-drinker and the AC-TL association (no cohort).**

445 (PDF)

446 **S4 Fig. Parallel pie chart: The association between whether the population of a**
447 **study was Asian and the AC-TL association**

448 (PDF)

449 **S1 Checklist. PRISMA checklist.**

450 (DOC)

451 **S1 Table. The information of the 44 analyses in the 21 articles.**

452 (xlsx)

453 **S2 Table. Cross table: The relationship between study type and the significance**

454 **of the AC-TL association. Note: ratio: the ratio of number of studies with**

455 **significant AC-TL association to the number of total studies**

456 (xlsx)

457 **S3 Table. Cross table: The relationship between continent and the significance of**

458 **the AC-TL association.**

459 (xlsx)

460 **S4 Table. Cross table: The relationship between a study's primary goal and the**

461 **significance of the AC-TL association.**

462 (xlsx)

463 **S5 Table. Cross table: The relationship between gender of a study's participants**

464 **and the significance of the AC-TL association.**

465 (xlsx)

466 **S6 Table. Cross table: The relationship between whether a study has an AC**

467 **category as never-drinker and the significance of the AC-TL association.**

468 (xlsx)

469 **S7 Table. Cross table: The relationship between whether a study has an AC**

470 **category as never-drinker and the significance of the AC-TL association (cohort**

471 **study excluded).**

472 (xlsx)

473 **S8 Table. Cross table: The relationship between whether the population of a**
474 **study was Asian and the significance of the AC-TL association.**

475 (xlsx)

476

477 **Author Contributions**

478 **Conceptualization:** Jianqiang Li, Ji-Jiang Yang, Weiliang Qiu.

479 **Data curation:** Yu Guan, Faheem Akhtar, Xi Xu.

480 **Investigation:** Yu Guan, Jianqiang Li, Weiliang Qiu.

481 **Methodology:** Jianqiang Li, Ji-Jiang Yang, Weiliang Qiu.

482 **Project administration:** Jianqiang Li, Ji-Jiang Yang.

483 **Software:** Yu Guan, Faheem Akhtar, Xi Xu.

484 **Supervision:** Shi Chen, Qing Wang, Hui Pan.

485 **Validation:** Yu Guan, Weiliang Qiu.

486 **Writing - original draft:** Faheem Akhtar, Xi Xu, Shi Chen, Qing Wang, Hui Pan.

487 **Writing - review & editing:** Shi Chen, Qing Wang, Hui Pan.

488

489

490

References

1. Wong J, Collins K. Telomere maintenance and disease. *Lancet*. 2003;362(9388):983–8.
2. Hayashi MT, Cesare AJ, Rivera T, Karlseder J. Cell death during crisis is mediated by mitotic telomere deprotection. *Nature*. 2015;522(7557):492–6.
3. Blasco MA. Telomeres and human disease: ageing, cancer and beyond. *Nat Rev Genet*. 2005;6:611–22.
4. Londoño-Vallejo JA. Telomere instability and cancer. *Biochimie*. 2008;90(1):73–82.
5. Zglinicki Tv. Replicative senescence and the art of counting. *Exp Gerontol*. 2003;38(11–12):1259–64.
6. Ornish D, Lin J, Chan JM, Epel E, Kemp C, Weidner G, et al. Effect of comprehensive lifestyle changes on telomerase activity and telomere length in men with biopsy-proven low-risk prostate cancer: 5-year follow-up of a descriptive pilot study. *Lancet Oncol*. 2013;14(11):1112–20.
7. Wu D, Cederbaum AI. Alcohol, Oxidative Stress, and Free Radical Damage. *Alcohol Res Health*. 2003;27(4):277–84.
8. Secretan B, Straif K, Baan R, Grosse Y, Ghissassi FE, Bouvard V, et al. A review of human carcinogens--Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol*. 2009;10(11):1033–4.
9. Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, et al. Alcohol consumption and site-specific cancer risk: a comprehensive dose–response meta-analysis. *Br J Cancer*. 2014;112(3):580–93.
10. Smyth A, Teo KK, Rangarajan S, O'Donnell M, Zhang X, Rana P, et al. Alcohol consumption and cardiovascular disease, cancer, injury, admission to hospital, and mortality: a prospective cohort study. *Lancet*. 2015;386(10007):1945–54.
11. DESA UN. Global Population Estimates by Age, 1950–2050. United Nations, Department of Economic and Social Affairs, 2013.
12. World Health Organization. Global Status Report on Alcohol and Health. Geneva, Switzerland: World Health Organization, 2011.
13. Schuckit MA. Recognition and management of withdrawal delirium (delirium tremens). *N Engl J Med*. 2014;371(22):2109–13.
14. Pavanello S, Hoxha M, Dioni L, Bertazzi PA, Snenghi R, Nalesso A, et al. Shortened telomeres in individuals with abuse in alcohol consumption. *Int J Cancer*. 2011;129(4):983–92.
15. Aida J, Yokoyama A, Izumiyama N, Nakamura K, Ishikawa N, Poon SS, et al. Alcoholics show reduced telomere length in the oesophagus. *J Pathol*. 2011;223(3):410–6.
16. Strandberg TE, Strandberg AY, Saijonmaa O, Tilvis RS, Pitkala KH, Fyhrquist F. Association between alcohol consumption in healthy midlife and telomere length in older men. The Helsinki Businessmen Study. *Eur J Epidemiol*. 2012;27(10):815–22.
17. Liu X, Bao G, Huo T, Wang Z, He X, Dong G. Constitutive telomere length and gastric cancer risk: case-control analysis in Chinese Han population. *Cancer Sci*. 2009;100(7):1300–5.
18. Liu J, Yang Y, Zhang H, Zhao S, Liu H, Ge N, et al. Longer leukocyte telomere length predicts increased risk of hepatitis B virus-related hepatocellular carcinoma: a case-control analysis. *Cancer*. 2011;117(18):4247–56.
19. Cassidy A, De Vivo I, Liu Y, Han J, Prescott J, Hunter DJ, et al. Associations between diet, lifestyle factors, and telomere length in women. *Am J Clin Nutr*. 2010;91(5):1273–80.
20. Fyhrquist F, Silventoinen K, Saijonmaa O, Kontula K, Devereux RB, de Faire U, et al. Telomere length and cardiovascular risk in hypertensive patients with left ventricular hypertrophy: the LIFE study. *J Hum Hypertens*. 2011;25(12):711–8.
21. Harris SE, Deary IJ, MacIntyre A, Lamb KJ, Radhakrishnan K, Starr JM, et al. The association between telomere length, physical health, cognitive ageing, and mortality in non-demented older people. *Neurosci Lett*. 2006;406(3):260–4.
22. Hou L, Savage SA, Blaser MJ, Perez-Perez G, Hoxha M, Dioni L, et al. Telomere Length in Peripheral Leukocyte DNA and Gastric Cancer Risk. *Cancer Epidemiol Biomarkers Prevent*. 2009;18(11):3103–9.
23. Houben JM, Giltay EJ, Rius-Ottenheim N, Hageman GJ, Kromhout D. Telomere length and mortality in elderly men: the Zutphen Elderly Study. *J Gerontol A Biol Sci Med Sci*. 2011;66(1):38–44.

24. Kozlitina J, Garcia CK. Red blood cell size is inversely associated with leukocyte telomere length in a large multi-ethnic population. *PloS one*. 2012;7(12):e51046.
25. Latifovic L, Peacock SD, Massey TE, King WD. The Influence of Alcohol Consumption, Cigarette Smoking, and Physical Activity on Leukocyte Telomere Length. *Cancer Epidemiol Biomarkers Prevent*. 2016;25(2):374–80.
26. Marcon F, Siniscalchi E, Crebelli R, Saieva C, Sera F, Fortini P, et al. Diet-related telomere shortening and chromosome stability. *Mutagenesis*. 2012;27(1):49–57.
27. Mirabello L, Huang WY, Wong JY, Chatterjee N, Reding D, Crawford ED, et al. The association between leukocyte telomere length and cigarette smoking, dietary and physical variables, and risk of prostate cancer. *Aging cell*. 2009;8(4):405–13.
28. Starnino L, Busque L, Tardif JC, D'Antono B. Psychological Profiles in the Prediction of Leukocyte Telomere Length in Healthy Individuals. *PloS one*. 2016;11(10):e0165482.
29. Sun Q, Shi L, Prescott J, Chiuve SE, Hu FB, De Vivo I, et al. Healthy lifestyle and leukocyte telomere length in U.S. women. *PloS one*. 2012;7(5):e38374.
30. Bekaert S, Meyer TD, Rietzschel ER, Buyzere MLD, Bacquer DD, Langlois M, et al. Telomere length and cardiovascular risk factors in a middle-aged population free of overt cardiovascular disease. *Aging cell*. 2007;6(5):639–47.
31. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–5.
32. Hu J, Dong Y, Chen X, Liu Y, Ma D, Liu X, et al. Prevalence of suicide attempts among Chinese adolescents: A meta-analysis of cross-sectional studies. *Compr Psychiatry*. 2015;61:78–89.
33. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clin Trials*. 1986;7(3):177–88.
34. Weischer M, Bojesen SE, Nordestgaard BG. Telomere shortening unrelated to smoking, body weight, physical activity, and alcohol intake: 4,576 general population individuals with repeat measurements 10 years apart. *Plos Genetics*. 2014;10(3):e1004191.
35. Won S, Morris N, Lu Q, Elston RC. Choosing an optimal method to combine P-values. *Stat Med*. 2009;28(11):1537–53.
36. Mehta CR, Patel NR. A network algorithm for performing Fisher's exact test in $r \times c$ contingency tables. *J Am Stat Assoc*. 1983; 78:427–34.
37. Clarkson DB, Fan Y-a, Joe H. A Remark on Algorithm 643: FEXACT: An Algorithm for Performing Fisher's Exact Test in $r \times c$ Contingency Tables. *ACM Trans Math Softw*. 1993;19:484–8.
38. Bendix L, Thinggaard M, Fenger M, Kolvraa S, Avlund K, Linneberg A, et al. Longitudinal changes in leukocyte telomere length and mortality in humans. *J Gerontol A Biol Sci Med Sci*. 2014;69(2):231–9.
39. Mather KA, Jorm AF, Milburn PJ, Tan X, Easteal S, Christensen H. No associations between telomere length and age-sensitive indicators of physical function in mid and later life. *J Gerontol A Biol Sci Med Sci*. 2010;65(8):792–9.
40. Shin C, Baik I. Associations Between Alcohol Consumption and Leukocyte Telomere Length Modified by a Common Polymorphism of ALDH2. *Alcohol Clin Exp Res*. 2016;40(4):765–71.

PRISMA 2009 Flow Diagram

Standard Error

AC-TL association is significant

case-control

cohort

cross-sectional

AC-TL association is not significant

6%

94%

AC-TL association is significant

AC-TL association is not significant