Exogenous DCPTA ameliorates the soil drought effect on nitrogen metabolism in maize during the pre-female inflorescence emergence stage

Tenglong Xie¹, Wanrong Gu^{1*}, Liguang Zhang², Congfeng Li³, Wenhua Li², Shi Wei^{1*}

¹ College of Agriculture, Northeast Agricultural University, Harbin 150030, P.R. China;

² Maize Research Institute, Heilongjiang Academy of Agricultural Sciences, Harbin 150086, P.R. China; ³ Institute of Crop Science, Chinese Academy of Agricultural Sciences, Beijing 100086, P.R. China.

E-mail of Tenglong Xie: tenglongxie@126.com

E-mail of Wanrong Gu: wanronggu@163.com

E-mail of Liguang Zhang: zlgncy@163.com

E-mail of Congfeng Li: lcfaas@163.com

E-mail of Wenhua Li: lwhnky@163.com

E-mail of Shi Wei: weishi5608@126.com

*Correspondence:

Shi Wei and Wanrong Gu

E-mail: weishi5608@126.com and wanronggu@163.com

Tel.: +86-451-55190472

Tables number: 1

Figures number: 18, and Fig. 2, Fig. 4 and Fig. 6 should be in colour in print.

Word count: 7260

29 **Highlights:**

30 This is the first article that explores the effects of DCPTA on nitrogen metabolism and
31 the first article that explores the effects of DCPTA on crops under soil drought
32 conditions.

33 **Abstract**

34• 2-(3,4-Dichlorophenoxy) triethylamine (DCPTA) regulates many aspects of plant
35 development; however, its effects on soil drought tolerance are unknown. We
36 pre-treated maize (*Zea mays* L.) by foliar application of DCPTA and subsequently
37 exposed the plants to soil drought and rewatering conditions during the pre-female
38 inflorescence emergence stage. Exogenous DCPTA significantly alleviated
39 drought-induced decreases in maize yield, shoot and root relative growth rate (RGR),
40 leaf relative water content (RLWC), net photosynthetic rate (Pn), stomatal
41 conductance (Gs) and transpiration rate (Tr), nitrate (NO₃⁻), nitrite (NO₂⁻), and soluble
42 protein contents, and nitrate reductase (NR), nitrite reductase (NiR), isocitrate
43 dehydrogenase (ICDH), alanine aminotransferase (AlaAT) and aspartate
44 aminotransferase (AspAT) activities; increases in the intercellular CO₂ concentration
45 (Ci), the ammonium (NH₄⁺) and free amino acid contents, and the glutamate
46 dehydrogenase (GDH) and protease activities. Simultaneously, exogenous DCPTA
47 improved the spatial and temporal distribution of roots and increased the root
48 hydraulic conductivity (Lp), flow rate of root-bleeding sap and NO₃⁻ delivery rates.
49 Moreover, Exogenous DCPTA protected the chloroplast structure from drought injury.
50 Taken together, our results suggest that exogenous DCPTA mitigates the repressive
51 effects of drought on N metabolism and subsequently enhances drought tolerance
52 during the pre-female inflorescence emergence stage of maize.

53 **Key words:** DCPTA; maize (*Zea mays* L.); nitrogen metabolism; pre-female
54 inflorescence emergence stage; soil drought; yield

55

56

57 Introduction

58 Crops are frequently exposed to drought during the growth period because of
 59 limited and erratic rainfall patterns due to global climate change, which leads to
 60 restrictions on agricultural productivity worldwide (Trenberth *et al.*, 2014). Maize
 61 (*Zea mays* L.), an essential component of global food security, is widely cultivated
 62 around the world. The majority of the cultivated area of maize is almost wholly
 63 rain-fed and experiences sporadic drought and rewetting cycles (Nuccio *et al.*, 2015).
 64 However, maize is considered to be a drought-sensitive crop and loses approximately
 65 1/4 potential yield annually due to drought (Ziyomo and Bernardo, 2013). By 2050,
 66 the world population will reach 9 billion people, resulting in a high demand for maize
 67 (projected to double); furthermore, at that time, drought will severely restrict crop
 68 growth for more than 50% of the cultivated land (Schiermeier, 2014).

69 To stabilize and increase global crop production to satisfy the demand of the
 70 globally burgeoning population, it is imperative to design agronomic research to
 71 improve maize performance under drought stress (Zhao *et al.*, 2018). The application
 72 of plant growth regulators has been considered an effective way to enhance crop
 73 drought resistance (Ali *et al.*, 2017). Multiple investigations have indicated that a
 74 tertiary amine bioregulator known as 2-(3,4-dichlorophenoxy) triethylamine (DCPTA)
 75 regulates many aspects of plant development; for example, DCPTA promotes plant
 76 growth (Keithly *et al.*, 1990a), enlarges chloroplast volume (Keithly *et al.*, 1990b),
 77 enhances photosynthetic enzyme activity (Wan and Mendoza, 1992), accelerates CO₂
 78 fixation (Gausman *et al.*, 1985), and stimulates carotenoid biosynthesis (Benedict *et*
 79 *al.*, 1985). As far as we know, very few studies of DCPTA have focused on crops, and
 80 the effect of DCPTA on crops exposed to soil drought are still unclear.

81 Nitrogen (N) metabolism is a fundamental process in determining the growth and
 82 productivity of plants (Kusano *et al.*, 2011). After being taken up by root systems,
 83 nitrate (NO₃⁻) is converted to nitrite (NO₂⁻) by nitrate reductase (NR), the first step of
 84 N uptake and utilization. Subsequently, nitrite (NO₂⁻) is converted to NH₄⁺ by nitrite
 85 reductase (NiR) with reduction-ferredoxin (Fd_{red}) as an electron donor (Rajasekhar *et*

al., 2010). Afterward, the ammonium (NH_4^+), derived from NO_3^- reduction, photorespiration and/or other metabolic processes is assimilated into glutamine by the glutamine synthase/glutamine oxoglutarate aminotransferase (GS/GOGAT) cycle or the alternative glutamate dehydrogenase (GDH) pathway with 2-oxoglutarate (2-OG) and reducing equivalents provided by photosynthesis (Chardon *et al.*, 2012). Subsequently, glutamate serving as a donor of the amino group is used for the synthesis of other amino acids, which are used for the synthesis of various organic molecules such as chlorophyll, proteins and nucleic acids. The reactions are catalysed by aminotransferases such as alanine aminotransferase (AlaAT) and aspartate aminotransferase (AspAT) (Slattery *et al.*, 2017).

Drought disrupts N metabolism mainly via inhibiting the uptake and/or long-distance transportation of NO_3^- (Nacer *et al.*, 2013), altering the activities of enzymes involved in N metabolism (Robredo *et al.*, 2011), inhibiting amino acid synthesis, and promoting protein hydrolysis (Fresneau *et al.*, 2007). At present, the study of plant growth regulators mainly concentrates on the improvement of photosynthesis and antioxidant systems, and there have been only a limited number of publications related to N metabolism.

Our previous hydroponic trial found that exogenous DCPTA drastically promoted growth under non-stress conditions and mitigated the PEG-simulated drought-induced growth inhibition of maize at the seedling stage (Xie *et al.*, 2017). The present study was conducted to explore whether DCPTA can alleviate soil drought injuries to maize and whether the effects are associated with the modulation of nitrogen metabolism.

Materials and Methods

Plant material, growth conditions, design and sampling

Seeds of the maize cultivar ZhengDan 958 and DCPTA were obtained from the Henan Academy of Agricultural Sciences in China and the China Zhengzhou Zhengshi Chemical Limited Company, respectively.

These experiments were performed in 2016 and 2017 at the Experimental Station

of Northeast Agricultural University, Harbin (126°73'E, 45°73'N), Heilongjiang province, China. The research field area has a temperate continental monsoon climate. The rainfall and mean temperature data during the study period (2016 and 2017, May–October) are listed in Fig. 1. Pits (inner length, 10 m; width, 7 m; and height, 1.2 m) in the field were used as experiment containers (Fig. 2). Plastic sheets were used to cover the inner sides of the pits, and a rain-proof shed was used to ensure the crops were solely dependent on soil moisture and irrigation over the course of the experiment to maintain the soil water conditions. The soil used was Chernozem and was sieved (pore size, 1 cm) and diluted with vermiculite (particle diameter, 4-8 mm; soil to vermiculite, v/v, 2:1). Before planting, soil chemical analysis was conducted according to Cottenie *et al.* (1982), and the results are presented in Table 1. Fertilization was carried out by adding ammonium nitrate (33.5% N), calcium superphosphate (15.5% P₂O₅), and potassium sulfate (48% K₂O) at the rates of 8.0, 8.0 and 20 kg pit⁻¹, respectively, before planting. No fertilizer was applied after planting. All containers were watered to 85% before planting. The seeds were manually sown on 2nd May 2016 and 4th May 2017 and were harvested on 7th October 2016 and 3rd October 2017, respectively. Three seeds were sowed per hole to ensure germination, and only the healthiest seedling within 20 days was kept at each site. Each container consisted of 10 rows, and the plant-to-plant and row-to-row distances were 20 cm and 65 cm, respectively. In addition, the ground around the containers was manually sown with the same plant-to-plant and row-to-row distances. The control of plant diseases and insect pests was conducted by managers.

The maize at the nine-leaf stage (during the pre-female inflorescence emergence stage) were treated as follows:

- (1) plants were irrigated continuously and sprayed with either 10 mL water (well-watered) or DCPTA (well-watered+DCPTA) per plant;
- (2) irrigation was stopped to form the drought conditions and sprayed with 10 mL of either water (drought) or DCPTA (drought +DCPTA) per plant; plants were rehydrated after 20 days of drought treatment.

144 The concentrations of DCPTA (25 mg/L) were based on the results of
145 previous screening experiments, and Tween-20 (0.03%) was added as a surfactant to
146 the solution for spraying. Each treatment had five replicates, and experiments were
147 performed in a completely randomized design. The dynamic changes in soil water
148 contents during the experimental stage are exhibited in Fig. 3.

149 Random plants from each treatment were sampled on days 0, 5, 10, 15, 20, 25
150 and 30. For leaf sampling, the middle part of the 9th leaf (numbered basipetally) was
151 sampled for analysis of leaf gas exchange, and the same part of the leaf was stored at
152 -80°C after immersion in liquid nitrogen for 30 min for determination of physiological
153 parameters. For root sampling, a hand-held soil auger (inner diameter of 20 cm) was
154 used to obtain soil cores from 0 to 100 cm depth of the soil profile at 10 cm increments.
155 The soil cores were soaked in a plastic container overnight, and roots were stirred and
156 sieved through a mesh (400 holes cm⁻²). The soil cores were carefully washed by
157 swirling water through the cores. The soil material and old dead roots debris were
158 manually separated from the live roots.

159 **Plant measurement and analysis**

160 **Relative growth rate (RGR) and plant productivity**

161 The shoots and roots of maize were oven dried at 105°C for 45 min and then
162 held at 80°C for 48 h; the shoot and root dry weights plant⁻¹ were determined soon
163 afterwards. The RGR was determined as follows: $RGR \text{ (fresh weight)} = [\ln(\text{final dry weight}) - \ln(\text{initial dry weight})] / (\text{duration of treatment days})$ (Kingsbury *et al.*, 1984).
164 A leaf area metre (Li-COR 3100; Li-COR, Lincoln, NE, USA) was used to estimate
165 the leaf area; number of grains plant⁻¹ (GN) and grain yield plant⁻¹ (GY) were
166 recorded at the maize physiological maturity stage.

168 **Leaf relative water content of (RLWC) and soil water content (SWC)**

169 The RLWC was determined on fresh leaf disks (2×2 cm) from the middle part of

the 8th leaves (numbered basipetally). After they were weighed (FW), the disks were immersed in distilled water at 25 °C overnight to obtain the turgid weight (TW). The leaves were dried at 80°C for 48 h and then weighed a third time (DW). RLWC was calculated as follows:

$$\text{RLWC (\%)} = [(FW - DW) / (TW - DW)] \times 100.$$

SWC was determined in the soil from the internal area of each container. After being weighed (FW), the soil portion was dried at 85°C for 96 h and then weighed (DW). SWC was calculated as follows:

$$\text{SWC (\%)} = [(FW - DW) / DW] \times 100.$$

Gas exchange

The photosynthetic rate (Pn), transpiration rate (Tr), stomatal conductance (Gs), and intercellular CO₂ concentration (Ci) values were determined with a portable photosynthesis system (LI-6400XT; LI-COR Biosciences, Lincoln, NE, USA) between at 13:00~14:00 h. The 6-cm² leaf chamber was used, and the photo flux density was 1000 $\mu\text{mol m}^{-2}\text{s}^{-1}$.

Transmission electron microscopy of chloroplasts

Observations were performed according to the description of Hu *et al.* (2014), and the chloroplast ultrastructure was observed under a H-7650 transmission electron microscope (manufacture: Hitachi, Japan).

Root morphological traits, root hydraulic conductivity, and the collection of root-bleeding sap

Roots from each soil core were scanned using a digital scanner (Epson V700, Indonesia). The root images were analysed using the WinRHIZO Image Analysis system (Version 2013e) (Regent Instruments Inc., Canada). The root length density (RLD, cm root cm⁻³ soil) and root square area density (RSD, cm² root cm⁻³ soil) were calculated according to the method described by Mosaddeghi *et al.* (2009).

196 The hydrostatic root hydraulic conductivity (L_p) was measured with a
197 Scholander pressure chamber according to the method described by López-Pérez *et al.*
198 (2007).

199 The plants were cut by scissors at 10–12 cm above the soil surface at 18:00-19:00.
200 Centrifuge tubes (inner diameter 40 mm) with cotton were placed on the upper end of
201 the stalks, and the stalk joints and centrifuge tubes were wrapped by plastic film to
202 keep impurities and insects out (Fig. 4). The bleeding sap was collected for 12 h; then,
203 the cotton was extracted from each centrifuge tube and placed into a glass syringe
204 (100 ml), and the root-bleeding sap was squeezed out for volume measurement. The
205 NO_3^- content in the root-bleeding sap was determined by AA3 Continuous Flow
206 Analytical System (Seal, Germany) according to Guan *et al.* (2014). The flow rate of
207 the root-bleeding sap and the NO_3^- delivery rate were expressed as $\text{ml h}^{-1} \text{root}^{-1}$ and
208 $\mu\text{g h}^{-1} \text{root}^{-1}$, respectively.

209 **Foliar NO_3^- , NO_2^- and NH_4^+ contents**

210 The foliar NO_3^- content determination by the reduction of NO_3^- to NO_2^- followed
211 the salicylic acid methods of Cataldo *et al.* (1975), the absorbance was monitored at
212 410 nm. The foliar NO_2^- content was determined using the method described by Barro
213 *et al.* (1991). The NO_2^- content was calculated according to the standard curve
214 obtained by known concentrations of KNO_3 . The NH_4^+ content was determined by
215 measuring the absorbance changes at 620 nm, as described by Weber (2007). The
216 NH_4^+ content was calculated according to the standard curve obtained by known
217 concentrations of $(\text{NH}_4)_2\text{SO}_4$.

218 **Enzyme activities involved in nitrogen metabolism**

219 The activities of foliar NR and NiR were measured based on the conversion of
220 NO_3^- to NO_2^- and NO_3^- to NH_4^+ , following the methods of Barro *et al.* (1991), and Ida
221 and Morita (1973), respectively. One NR and NiR unit was expressed as the enzyme
222 amount required for the conversion of 1 mmol of NO_3^- to NO_2^- per hour and the

enzyme amount required for the conversion of 1 mmol of NO_3^- to NH_4^+ min^{-1} , respectively. GS activity was measured according to the methods of O'Neal and Joy (1973). One unit of GS was expressed as the enzyme amount required to catalyse the formation of 1 mmol of glutamylmonohydroxamate per min. The results were expressed as μmol NADH used per minute per milligram of Pro. The GOGAT activity was measured based on the conversion of 2-ketoglutarate to glutamate, following the methods of Groat and Vance (1981). One unit of GOGAT was expressed as the number of enzymes catalysing the oxidation of 1 mmol of NADH per min. The deaminating GDH activity (NAD-GDH) and aminating GDH activity (NADH-GDH) were determined by recording the reduction of NAD and the oxidation of NADH, respectively, as described by Groat and Vance (1981). One unit of GDH was calculated in units of mmol of NADH oxidized/NAD reduced per minute. Isocitrate dehydrogenase (ICDH) was assayed according to the method of Lòpez-Millàn *et al.* (2000). The activity was expressed as U per minute per milligram of protein.

Activity of AlaAT and AspAT

The AlaAT and AspAT activities were determined according to the methods of Jia *et al.* (2015). Enzyme activity was expressed as $\mu\text{mol g}^{-1}\text{min}^{-1}$.

Contents of free amino acids and soluble proteins and protease activity

The free amino acid contents were assayed with the ninhydrin reagent method according to Yemm and Cocking (1955), followed by absorbance readings at 570 nm using glycine as the standard. The soluble protein contents were determined using the Coomassie Brilliant Blue G-250 reagent following the description of Bradford (1976), followed by absorbance readings at 595 nm using bovine serum albumin as the standard. Protease activity was determined by the casein digestion assay described by Drapeau (1974). By this method, one unit is the number of enzymes required to release acid-soluble fragments equivalent to 0.001 A280 per minute at 37°C and pH 7.8.

250 **Statistical analysis**

251 The data were analysed using SPSS 17.0 and all the values are presented as the mean
252 $\pm$ SE. The means were separated using the least significant difference (LSD) test at
253 the 5% probability level.

254 **Results**

255 **Yield**

256 Drought stress significantly inhibited the maize yield (Fig. 5). Compared with
257 the well-watered treatment, in the drought treatment, the grain number decreased by
258 29.23% in 2016 and 33.24% in 2017, and the grain yield decreased by 34.06% in
259 2016 and by 38.22% in 2017. However, the decrease in maize yield was partially
260 recovered by DCPTA. Compared with the well-watered treatment, in the
261 drought+DCPTA treatment, the grain number decreased by 14.01% in 2016 and by
262 16.55% in 2017, and the grain yield decreased by 17.98% in 2016 and by 20.54% in
263 2017. Moreover, the application of DCPTA improved maize yield under well-watered
264 conditions. Compared with the well-watered treatment, in the DCPTA treatment, the
265 grain number increased by 5.97% in 2016 and by 6.50% in 2017, and the grain yield
266 increased by 7.31% in 2016 and by 8.02% in 2017.

267 **Relative growth rate (RGR) of shoot and root**

268 The shoot growth rate was inhibited during the drought period, the root growth
269 rate was promoted over days 0-10 and subsequently decreased, and the growth rates
270 of shoots and roots recovered during rehydration (Fig. 6 and 7). In the drought
271 treatment compared with the control in 2016, the RGR of shoots and roots decreased
272 by 52.26% and by 48.59%, respectively, between day 15 and 20 and by 39.76% and
273 37.70%, respectively, between day 25 and 30 in 2016. In the drought treatment
274 compared with the control in 2017, the RGR of shoots and roots decreased by 65.18%
275 and 66.52%, respectively, between day 15 and 20 and by 51.97% and 43.74%,
276 respectively, between day 25 and 30 in 2017. However, the decrease in growth was

277 partially recovered by DCPTA. In the drought+DCPTA treatment compared with the
278 control in 2016, the RGR of shoots and roots decreased by 29.96% and 26.31%,
279 respectively, between day 15 and 20, and by 10.11% and 11.49%, respectively,
280 between day 25 and 30. In the drought+DCPTA treatment compared with the control
281 in 2017, the RGR of shoots and roots decreased by 34.20% and 42.23%, respectively,
282 between day 15 and 20, and by 12.41% and 14.75%, respectively, between day 25 and
283 30. Moreover, the application of DCPTA improved maize growth under well-watered
284 conditions. The shoot RGR difference between the well-watered+DCPTA treatment
285 and well-watered treatment was significant at days 16-20 and 26-30 in 2016 and days
286 11-30 in 2017, respectively. Similarly, the root RGR difference between the DCPTA
287 treatment and well-watered treatment was significant at days 6-10 and 16-25 in 2016
288 and days 11-30 in 2017, respectively.

289 **Root length density (RLD) and root surface area density (RSD)**

290 Drought dramatically inhibited RLD and RSD in the 0–60 cm soil profile in
291 2016 and 2017 (Fig. 8). At day 20, there was a significant difference in RLD for the
292 drought treatment and drought+DCPTA treatment in 20–40 cm in 2016 and in 20–50
293 cm in 2017, respectively. On day 30, the RLD declines were partially reversed by
294 rehydration. On day 30, there was a significant difference in RLD for the drought
295 treatment and drought+DCPTA treatment in 0–50 cm in both 2016 and 2017. On day
296 30, there was a significant ($P > 0.05$) difference in RLD for the well-watered
297 treatment and well-watered+DCPTA treatment in 0–20 cm in both 2016 and 2017. On
298 day 20, there was a significant difference in RLD between the drought treatment and
299 drought+DCPTA treatment in 10–40 cm in 2016 and 10–50 cm in 2017. On day 30,
300 the declines in RLD were partially reversed by rehydration. At day 30, there was a
301 significant ($P > 0.05$) difference in RLD between the drought treatment and
302 drought+DCPTA treatment in 0–40 cm in 2016 and 0–50 cm in 2017. On day 30,
303 there was a significant difference between RLD for the well-watered treatment and
304 well-watered+DCPTA treatment in 0–20 cm in 2016 and in 0–30 cm in 2017.

305 **Root hydraulic conductivity, flow rate of root-bleeding sap and NO₃⁻ delivery** 306 **rates**

307 The root hydraulic conductivity, flow rate of root-bleeding sap and NO₃⁻
308 concentrations in the root-bleeding sap declined continuously during the drought
309 period and recovered during rehydration (Fig. 9). In the drought treatment compared
310 with the control, the root hydraulic conductivity, flow rate of root-bleeding sap and
311 NO₃⁻ delivery rates decreased by 34.21%, 75.69% and 76.58%, respectively, on day
312 20 and by 19.34%, 35.96% and 57.37%, respectively, on day 30 in 2016; these values
313 decreased by 47.01%, 78.80% and 61.34%, respectively, on day 20 and by 29.03%,
314 33.79%, and 50.47%, respectively, on day 30 in 2017. However, the DCPTA
315 application partially reversed the decline in root hydraulic conductivity, root-bleeding
316 sap flow and NO₃⁻ delivery rates caused by drought and resulted in a faster recovery
317 after rehydration. In the drought+DCPTA treatment compared with the control, the
318 root hydraulic conductivity, root-bleeding sap flow and NO₃⁻ delivery rates decreased
319 by 17.91%, 46.73% and 41.90%, respectively, on day 20 and by 9.87%, 18.80% and
320 23.51%, respectively, on day 30 in 2016; these values decreased by 26.60%, 47.66%
321 and 37.06%, respectively, on day 20 and by 16.94%, 14.68% and 25.23%,
322 respectively, on day 30.

323 Under well-watered conditions, DCPTA significantly increased root hydraulic
324 conductivity on the 15th, 20th, 25th and 30th days in 2016 and on the 20th and 25th days
325 in 2017; DCPTA significantly increased the flow rate of root-bleeding sap on the 10th,
326 15th, 20th and 25th days in both 2016 and 2017; DCPTA significantly increased the
327 NO₃⁻ concentration in root-bleeding sap on the 10th, 15th, 20th, 25th and 30th days in
328 2016 and on the 10th, 15th, 20th and 25th days in 2017.

329 **Leaf water status (RLWC)**

330 The RLWC declined continuously over the drought period and recovered
331 during rehydration (Fig. 10). In the drought treatment compared with the control,
332 RLWC decreased by 30.88% on day 20 and by 13.08% on day 30 in 2016 and

333 decreased by 37.48% on day 20 and by 21.89% on day 30 in 2017. However, the
334 DCPTA application partially reversed the decline in RLWC caused by drought and
335 resulted in a faster recovery of the foliar RLWC contents after rehydration. In the
336 drought+DCPTA treatment compared with the control, the RLWC decreased by 11.75%
337 on day 20 and by 3.84% on day 30 in 2016 and decreased by 21.15% on day 20 and
338 by 10.73% on day 30 in 2017. Under well-watered conditions, DCPTA application
339 had no significant effect on the RLWC.

340 **Gas exchange parameters**

341 Pn, Gs and Tr declined continuously over the drought period and recovered
342 during rehydration (Fig. 11). In the drought treatment compared with the control, Pn,
343 Gs and Tr decreased by 46.20%, 68.97% and 51.35%, respectively, on day 20 and by
344 35.06%, 46.30% and 38.17%, respectively, on day 30 in 2016 and decreased by
345 56.00%, 71.10% and 62.10%, respectively, on day 20 and by 45.89%, 44.02% and
346 43.79%, respectively, on day 30 in 2017. However, the DCPTA application partially
347 reversed the decline in Pn, Gs and Tr caused by drought and resulted in a faster
348 recovery of Pn, Gs and Tr rehydration. In the drought+DCPTA treatment compared
349 with the control, Pn, Gs and Tr decreased by 23.41%, 37.39% and 20.95%,
350 respectively, on day 20 and by 8.77%, 20.89% and 10.61%, respectively, on day 30 in
351 2016 and decreased by 35.31%, 32.03% and 34.11%, respectively, on day 20 and by
352 14.16%, 15.33% and 8.62%, respectively, on day 30. Under well-watered conditions,
353 DCPTA significantly increased Pn on the 10th, 15th, 25th and 30th days in 2016 and on
354 the 15th, 20th and 25th days in 2017. In addition, under well-watered conditions,
355 DCPTA significantly increased Gs on the 15th, 20th, 25th and 30th days in 2016 and on
356 the 10th, 15th, 20th, 25th and 30th days in 2017 and significantly increased Tr on the 10th,
357 15th, 20th, 25th and 30th days in both 2016 and 2017.

358 In 2016 and 2017, Ci showed the same variation tendency during the drought
359 period. In the drought treatment, Ci declined over days 0-10, subsequently increased
360 over days 10-20, and then decreased after rehydration. In the drought+DCPTA

361 treatment, Ci declined over days 0-15 day and subsequently increased over days
362 15-30. Under well-watered conditions, DCPTA application had no significant effect
363 on Ci.

364 **Chloroplast ultrastructure**

365 Regardless of whether DCPTA was applied, the photosynthetic mesophyll cells
366 of the non-stressed seedlings included a delimited cell wall containing chloroplasts.
367 These chloroplasts had intact membranes and a regular arrangement of granal and
368 stromal thylakoids, which were attached to the cell wall and exhibited typical
369 ellipsoidal shapes (Fig. 12). However, in the stressed seedlings, the cell wall structure
370 was incomplete and exhibited indistinct gradation, a lower density, and loose edges.
371 Plasmolysis and degradation were also evident in part of the cell membrane.
372 Moreover, the chloroplasts, which separated from the plasma membrane, were nearly
373 round and swelled asymmetrically, the thylakoids were overly disorganized, and the
374 thylakoid membranes were loose and showed an increased number of plastoglobules.
375 In the PEG-6000+DCPTA treatment, the complete membrane structures of the
376 chloroplasts were present, and the shapes of the chloroplasts changed slightly from
377 elongated ellipses to ellipses close to the cell walls. A well-aligned internal lamellar
378 system and fewer plastoglobules were observed in the leaves of the
379 PEG-6000+DCPTA treatment compared with the leaves of the PEG-6000 treatment.

380 **ICDH activity**

381 ICDH activity declined continuously over the drought period and recovered
382 during rehydration (Fig. 13). In the drought treatment compared with the control,
383 ICDH activity decreased by 40.75% on day 20 and by 36.33% on day 30 in 2016 and
384 decreased by 37.77% on day 20 and by 33.07% on day 30 in 2017. However, the
385 DCPTA application partially reversed the decline in ICDH activity caused by drought
386 and resulted in a faster recovery of ICDH activity after rehydration. In the
387 drought+DCPTA treatment compared with the control, ICDH decreased by 24.92% on

day 20 and by 19.94% on day 30 in 2016 and decreased by 19.88% on day 20 and by 13.63% on day 30 in 2017. Under well-watered conditions, the application of DCPTA significantly increased the foliar ICDH activity on the 10th, 15th, 20th and 25th days in 2016 and on the 15th, 20th, and 25th days in 2017.

Contents of NO₃⁻, NO₂⁻ and NH₄⁺

The foliar NO₃⁻ and NO₂⁻ contents declined continuously during the drought period and recovered during rehydration (Fig. 14). In the drought treatment compared with the control, the foliar NO₃⁻ and NO₂⁻ contents decreased by 39.82% and 38.27%, respectively, on day 20 and by 33.62% and 35.11%, respectively, on day 30 in 2016 and decreased by 57.97% and 32.27%, respectively, on day 20 and by 42.38% and 25.21%, respectively, on day 30 in 2017. However, the DCPTA application partially reversed the decline in the foliar NO₃⁻ and NO₂⁻ contents caused by drought and resulted in a faster recovery of the foliar NO₃⁻ and NO₂⁻ contents after rehydration. In the drought+DCPTA treatment compared with the control, the foliar NO₃⁻ and NO₂⁻ contents decreased by 25.94% and 23.31%, respectively, on day 20 and by 14.25% and 18.55%, respectively, on day 30 in 2016 and decreased by 33.86% and 18.26%, respectively, on day 20 and by 17.26% and 8.60%, respectively, on day 30 in 2017. In contrast, drought led to a marked elevation in the foliar NH₄⁺ content. In the drought treatment compared with the control, the foliar NH₄⁺ increased by 52.48% on day 20 and by 29.18% on day 30 in 2016 and increased by 98.68% on day 20 and by 72.21% on day 30 in 2017. In contrast, the DCPTA application suppressed the increase in the foliar NH₄⁺ content induced by drought. In the drought+DCPTA treatment compared with the control, the foliar NH₄⁺ increased by 20.83% on day 20 and by 13.33% on day 30 in 2016 and increased by 45.37% on day 20 and by 33.29% on day 30 in 2017.

Under well-watered conditions, DCPTA significantly increased the foliar NO₃⁻ content on the 20th and 25th days in 2016 and on the 15th, 20th, 25th and 30th days in 2017. However, the DCPTA application had no significant effect on the foliar contents of NO₂⁻ and NH₄⁺.

416 **Activities of NR and NiR**

417 The activities of foliar NR and NiR declined continuously during the drought
418 period and recovered during rehydration (Fig. 15). In the drought treatment compared
419 with the control treatment, the activities of foliar NR and NiR decreased by 40.37%
420 and 36.91%, respectively, on day 20 and by 34.42% and 29.82%, respectively, on day
421 30 in 2016 and decreased by 52.80% and 32.46%, respectively, on day 20 and by
422 37.36% and 21.28%, respectively, on day 30 in 2017. However, the DCPTA
423 application partially reversed the declines in the activities of foliar NR and NiR
424 caused by drought and resulted in a faster recovery of the foliar NR and NiR activities
425 after rehydration. In the drought+DCPTA treatment compared with the control, the
426 activities of foliar NR and NiR decreased by 15.54% and 14.21%, respectively, on
427 day 20 and by 10.79% and 7.79%, respectively, on day 30 in 2016 and decreased by
428 25.10% and 12.39%, respectively, on day 20 and by 13.55% and 4.96%, respectively,
429 on day 30 in 2017. Under well-watered conditions, DCPTA significantly increased
430 the foliar NR activity on the 10th, 15th and 25th days in 2016 and on the 10th, 15th, 20th
431 and 25th days in 2017. Similarly, under well-watered conditions, DCPTA significantly
432 increased the foliar NiR activity on the 10th and 25th days in 2016 and on the 15th and
433 20th days in 2017.

434 **Activities of GS, GOGAT and GDH**

435 The activities of foliar GS and GOGAT first increased and then decreased
436 during the drought period and recovered during rehydration (Fig. 16). In the drought
437 treatment compared with the control, the activities of foliar GS and GOGAT
438 decreased by 40.69% and 60.62%, respectively, on day 20 and by 33.65% and 51.39%,
439 respectively, on day 30 in 2016 and decreased by 47.57% and 66.84%, respectively,
440 on day 20 and by 37.96% and 56.77%, respectively, on day 30 in 2017. However, the
441 DCPTA application partially reversed the decline in the foliar GS and GOGAT
442 activities caused by drought and resulted in a faster recovery of the foliar GS and
443 GOGAT activities after rehydration. In the drought+DCPTA treatment compared with

the control, the foliar GS and GOGAT activities decreased by 19.87% and 33.37%, respectively, on day 20 and by 10.99% and 15.17%, respectively, on day 30 in 2016 and decreased by 23.56% and 33.02%, respectively, on day 20 and by 14.93% and 18.20%, respectively, on day 30 in 2017. Likewise, the DCPTA application alone caused an increase in the foliar GS and GOGAT activities. In contrast, drought led to marked increases in the activities of foliar NAD-GDH and NADH-GDH. In the drought treatment compared with the control, the activities of foliar NAD-GDH and NADH-GDH increased by 87.16% and 150.92%, respectively, on day 20 and by 84.01% and 134.71%, respectively, on day 30 in 2016 and increased by 103.99% and 137.36%, respectively, on day 20 and by 96.13% and 111.20%, respectively, on day 30 in 2017. However, the DCPTA application partially reversed the increases in the activities of foliar NAD-GDH and NADH-GDH caused by drought and resulted in a faster recovery of the foliar NAD-GDH and NADH-GDH activities after rehydration. In the drought+DCPTA treatment compared with the control, the activities of foliar NAD-GDH and NADH-GDH increased by 49.55% and 92.59%, respectively, on day 20 and by 36.52% and 49.51%, respectively, on day 30 in 2016 and increased by 46.90% and 80.94%, respectively, on day 20 and by 35.43% and 45.60%, respectively, on day 30 in 2017.

Under well-watered conditions, DCPTA significantly increased the foliar GS activity on the 20th and 25th days in 2016 and on the 10th 20th and 25th days in 2017. Similarly, under well-watered conditions, DCPTA significantly increased the foliar GOGAT activity on day 15 in 2017 and significantly increased the foliar NAD-GDH activity on day 15 in 2016

Activities of AlaAT and AspAT

The activities of foliar AlaAT and AspAT first increased and then continuously decreased during the drought period and recovered during rehydration (Fig. 17). In the drought treatment compared with the control, the activities of foliar AlaAT and AspAT decreased by 44.18% and 65.51%, respectively, on day 20 and by 36.43% and 38.52%,

472 respectively, on day 30 in 2016 and decreased by 56.80% and 54.92%, respectively,
473 on day 20 and by 41.43% and 41.29%, respectively, on day 30 in 2017. However, the
474 DCPTA application partially reversed the decline in the activities of foliar AlaAT and
475 AspAT caused by drought and resulted in a faster recovery of the foliar AlaAT and
476 AspAT activities after rehydration. In the drought+DCPTA treatment compared with
477 the control, the activities of foliar AlaAT and AspAT decreased by 17.45% and
478 35.39%, respectively, on day 20 and by 15.84% and 17.17%, respectively, on day 30
479 in 2016 and decreased by 32.49% and 37.82%, respectively, on day 20 and by 17.46%
480 and 18.21%, respectively, on day 30 in 2017.

481 Under well-watered conditions, DCPTA significantly increased the foliar
482 AlaAT activity on the 10th, 15th, and 20th days in 2016 and on the 10th, 15th, 20th and
483 25th days in 2017. Similarly, under well-watered conditions, DCPTA significantly
484 increased the foliar AspAT activity on day 15 in 2016 and on the 5th, 10th, 15th and 25th
485 days in 2017.

486 **Protease activity and contents of proteins and free amino acids**

487 The protease activity and free amino acid contents increased continuously
488 during the drought period and decreased during rehydration (Fig. 18). In the drought
489 treatment compared with the control, the protease activity and free amino acid
490 contents increased by 122.48% and 88.92%, respectively, on day 20 and by 55.03%
491 and 34.51%, respectively, on day 30 in 2016 and increased by 145.15% and 78.58%,
492 respectively, on day 20 and by 57.08% and 43.29%, respectively, on day 30 in 2017.
493 However, the DCPTA application partially reversed the increases in the protease
494 activity and free amino acid contents caused by drought.

495 In the drought+DCPTA treatment compared with the control, the protease
496 activity and free amino acid contents increased by 78.47% and 24.77%, respectively,
497 on day 20 and by 16.46% and 6.62%, respectively, on day 30 in 2016 and increased
498 by 65.41% and 30.24%, respectively, on day 20 and 14.50% and 13.57%, respectively,
499 on day 30 in 2017.

In contrast, drought led to a marked decrease in the foliar protein content. In the drought treatment compared with the control, the foliar protein content decreased by 35.51% on day 20 and by 18.32% on day 30 in 2016 and decreased by 44.81% on day 20 and by 22.50% on day 30 in 2017. In contrast, the DCPTA application suppressed the increase in the foliar protein content induced by drought. In the drought+DCPTA treatment compared with the control, the foliar protein content increased by 19.54% on day 20 and by 7.25% on day 30 in 2016 and increased by 25.46% on day 20 and by 10.30% on day 30 in 2017.

Significant differences between the foliar protein contents of the well-watered treatment and well-watered+DCPTA treatment were observed on day 10 in 2017 and on day 15 in 2017, and significant differences between the free amino acid contents of these treatments were observed on day 10 in 2017.

Discussion

One of the most clear and consistent effects of drought on crops is the inhibition of growth and yield (Sinclair, 2011). In this experiment, exogenous DCPTA partly mitigated the reductions in plant growth and yield induced by drought, as expressed by the stable shoot and root RGR, GN and GY (Fig. 5, Fig. 6 and Fig. 7). Moreover, exogenous DCPTA promoted growth and yield under well-watered conditions.

N is a necessary macro-nutrient element for plants and a main limiting factor in plant growth and development (Kusano *et al.*, 2011). The pre-female inflorescence emergence stage is the important stage determining maize yield (Talaat *et al.*, 2015). The inhibition of maize growth induced by drought could be partly attributed to nitrogen metabolism during the pre-female inflorescence emergence stage. NO_3^- is the main nitrogen source assimilated by higher plants in agricultural soils (Robredo *et al.*, 2011). Similar to previous reports for tomato (Sánchez-Rodríguez *et al.*, 2011) and wheat (Fresneau *et al.*, 2007), drought significantly diminished the NO_3^- content in maize leaves in both the DCPTA-treated and non-treated leaves (Fig. 14). This decrease may be explained by drought-induced inhibitions in nitrate uptake from the

528 roots and/or nitrate transport. However, the reduction in the non-treated leaves was
529 greater than that in the DCPTA-treated leaves in this study.

530 Maintaining the supply of nutrients and water from the soil to crop depends on
531 root morphology, which can be characterized by RLD and RSD (Kamran *et al.*, 2018).
532 Previous studies have reported that DCPTA promotes root development in tomatoes
533 (Keithly *et al.*, 1990a). In this study, there was a significant difference between the
534 RLD and RSD of the well-watered treatment and DCPTA treatment in 0–20 cm and
535 0–30 cm in 2016 and 2017 on day 30 (Fig. 8). This result suggests that DCPTA
536 promoted root development in maize under well-watered conditions. Interestingly,
537 under drought conditions, the DCPTA application significantly increased RLD, RSD
538 and the root RGR. These results indicate that the DCPTA application also promoted
539 maize root growth and improved the spatial and temporal distribution of roots, which
540 was beneficial to NO_3^- uptake under drought conditions.

541 The xylem sap transports water and nutrients from the roots throughout the plant
542 and depends on transpiration intensity and root pressure. The increased root hydraulic
543 conductivity and flow rate of root-bleeding sap induced by the DCPTA application
544 may, due to the enhanced root pressure, which depend on physiological activity of the
545 whole root system (Fig. 9) (Noguchi *et al.*, 2005; Lui *et al.*, 2014). In addition, the
546 NO_3^- delivery rate in the presence of DCPTA was significantly higher than that
547 without DCPTA under drought conditions, which may partly result from the improved
548 NO_3^- absorption and enhanced root pressure induced by DCPTA. The stable RLWC
549 in the drought+DCPTA treatment suggests an abundant supply water to the
550 aboveground parts, and balanced transpirational loss and water uptake under drought
551 conditions; as a result, the DCPTA-treated plants maintained G_s , which reduced the
552 leaf epidermal resistance and promoted the mass flow of water to the leaf surface and
553 the transportation of the NO_3^- required for N metabolism in leaves (Fig. 10 and Fig.
554 11). Under drought conditions, increases in the foliar NO_3^- were observed in the
555 DCPTA-treated plants (Fig. 14).

556 Whether stomatal or non-stomatal factors are the main cause of a reduced P_n

may be determined by changes in G_s and C_i (Bethke and Drew, 1992). During the early period of drought, the change of C_i were accompanied by continuously declined G_s , then G_s decreases but C_i shows an increase (Fig. 11). Thus, the decrease of the P_n in drought-treated plants was mainly attributed to stomatal limitations firstly, and then, non-stomatal limitations induced by the damage of photochemical mechanism, partly reflected by damaged chloroplast (Fig. 12). However, DCPTA application maintains relatively high G_s , ensuring the availability of CO_2 for the carbon reduction cycle. Simultaneously, the DCPTA application delayed the increase in C_i and protected the chloroplast ultrastructure against drought-induced oxidative damage, which suggests that DCPTA can protect the photochemical mechanism and, as a result, ensures a more efficient photosynthesis process after rehydration. Moreover, similar to previous studies on spruce (Keithly *et al.*, 1990a), sugar beets (Keithly *et al.*, 1990b) and guayule (Wan and Mendoza, 1992), DCPTA application also promoted photosynthesis under well-watered conditions.

In most plants, nitrate reduction occurs in leaves. NO_3^- , after being taken up into the leaf cell, is converted to NH_4^+ by two successive steps catalysed by NR and NiR. NR, the rate-limiting enzyme of nitrogen assimilation, is highly sensitive to stress (Plett *et al.*, 2016). Similar to previous studies on wheat and barley, the NR activity continuously declined in response to drought (Fig. 15) (Robredo *et al.*, 2011; Fresneau *et al.*, 2007). As a typical nitrate-induced enzyme, NR activity is primarily regulated by the NO_3^- concentration in the leaves (Chamizo-Ampudia *et al.*, 2017). The up-regulation of foliar NR activity in the drought+DCPTA treatment may result from the increase in the foliar NO_3^- content (Fig. 14). Moreover, the reduction in the foliar NiR activity under drought conditions was significantly reversed by the application of DCPTA, which may be because the DCPTA-stabilized photosynthesis resulted in a sufficient supply of Fd_{red} (Fig. 11 and Fig. 15), thus promoting the conversion of NO_2^- to NH_4^+ . The present results indicate that DCPTA pretreatment could maintain a high NO_3^- assimilation ability in maize under drought conditions.

Although the foliar NR and NiR activities declined during the drought period, the

foliar NH_4^+ content exhibited an increasing tendency in our experiment (Fig. 14). This increase may be associated with the glycine oxidation in activated photorespiration, which is induced by decreases in *Ci* levels under drought conditions (Wang *et al.*, 2012). The increased *Gs* induced by DCPTA was beneficial to the increase in CO_2 in the cellular spaces of the leaf, implying that photorespiration was partly alleviated (Fig. 11).

In plants cells, excessive levels of NH_4^+ are destructive, and the major NH_4^+ assimilation pathway is the GS/GOGAT cycle in higher plants. When the GS/GOGAT cycle is suppressed and the NH_4^+ content rises continuously under stress, NH_4^+ could serve as a substrate to form glutamate via the reversible amination of 2-OG by GDH catalyse, although the enzyme has a lower affinity for NH_4^+ (Fontaine *et al.*, 2012). In general, drought inhibited NH_4^+ assimilation (Naya *et al.*, 2007). During the early period of drought, GDH activity increased sharply, GS activity increased slightly, and GOGAT remained stable (Fig. 16). These results suggest that accelerated NH_4^+ assimilation in maize may be an adaptive mechanism to produce more glutamate and eliminate the accumulation of excess foliar NH_4^+ . Subsequently, the GS and GOGAT activities decreased, which may have resulted from an inadequate supply of energy and 2-OG because of photosynthetic inhibition and decreased ICDH activity (Fig. 13). GDH activity decreased with drought, which may be due to the oxidative degradation of GS caused by the overproduction of reactive oxygen species (ROS) induced by drought conditions (Xia *et al.*, 2015).

The DCPTA application altered the major NH_4^+ assimilation pathway, maintained the GOGAT/GS cycle and suppressed the GDH pathway, which may have contributed to maintaining the conversion of NH_4^+ to glutamine and the subsequent formation of glutamate from glutamine. This result may occur because the photosynthetic stability and ICDH activity induced by the DCPTA application promoted 2-OG synthesis and the reducing power (i.e., NADPH, ATP, or Fd_{red}) in plants during the drought period, thus providing the GS/GOGAT cycle with relatively sufficient substrates and energy and favouring the enhancement of foliar

615 GOGAT and GS activities (Du *et al.*, 2016). As a result, with the application of
616 DCPTA, drought had less of an effect on the activities of GS and GOGAT.

617 Although DCPTA promoted NO_3^- assimilation, as expressed by the increased NR
618 and NiR activities (Fig. 15), this treatment compared to the drought treatment caused
619 significant decreases in the NH_4^+ content, which means that exogenous DCPTA
620 resulted in the integration of NH_4^+ into the structure of organic compounds, thereby
621 contributing to the reduction in the NH_4^+ content. Therefore, the DCPTA application
622 effectively modulated the activities of ICDH, GS, GOGAT and GDH and accelerated
623 the conversion of NH_4^+ to glutamate, which is the precursor of other amino acids.

624 Transamination is a key step in the biosynthesis of various amino acids from
625 glutamate, with the availability of C skeletons from the Krebs cycle (Hodges, 2002).
626 In our studies, both the aminotransferases studied, AlaAT and AspAT, showed
627 increased activity in maize during the early drought period (Fig. 17). Such increases
628 in aminotransferases activities under drought conditions might help in the synthesis of
629 increased amounts of amino acids that act as compatible cytoplasmic solutes and
630 protect cell organelles and biomolecules, thus reducing the adverse effects of drought
631 on maize (Munns and Tester, 2008). Subsequently, the AlaAT and AspAT activities
632 decreased, which may be attributable to the weakened GS/NADH-GOGAT pathway
633 (Fig. 16) (Gangwar and Singh, 2011). Moreover, stable aminotransferase activities
634 were observed in DCPTA-treated plants. This finding may be associated with
635 increased GS/GOGAT activities, which can generate more glutamate to serve as a
636 substrate for transamination reactions in maize treated with DCPTA under drought
637 conditions.

638 Most soluble proteins are enzymes that participate in various metabolic
639 pathways in plants; therefore, the soluble protein content is considered one of the
640 most important indices reflecting the overall metabolic level in plants. Protein
641 synthesis in plants is very sensitive to abiotic stresses and is positively correlated with
642 stress tolerance (Fresneau *et al.*, 2007). Free amino acids are the building blocks of
643 proteins. Drought increased the free amino acid contents, which may mainly be

644 attributed to the increased AlaAT and AspAT activities in the early drought period
 645 and the subsequent promotion of protein degradation (Fig. 18) (Yang *et al.*, 2013).
 646 However, DCPTA-treated seedlings maintained higher soluble protein levels and
 647 lower free amino acid levels than did non-DCPTA-treated seedlings in response to
 648 drought. This result may occur because DCPTA inhibited protein degradation by
 649 stable protease activities and maintained protein stability, ensuring the series of
 650 physiological and biochemical processes that occur normally under stress conditions.
 651 Additionally, the DCPTA application increased the amino acid contents under
 652 well-watered conditions, which may be attributable to the promoted biosynthesis and
 653 accumulation of amino acids, which ultimately improved plant growth and
 654 development (Talaat and Shawky, 2016).

655 **Conclusions**

656 The present study suggested that DCPTA treatment increased NO_3^- uptake and
 657 the long-distance transportation of NO_3^- from the roots to the leaves via the
 658 production of excess roots and maintained a stabilized transpiration rate. The
 659 increased foliar NO_3^- content up-regulated NR activity and maintained a high N
 660 assimilation ability that was restrained by drought. Exogenous DCPTA effectively
 661 regulated the ICDH, GS, GOGAT and GDH activities to speed up the conversion of
 662 NH_4^+ to Glu, reduced the toxicity of excess NH_4^+ to the plant, and accelerated the
 663 synthesis of proteins and amino acids. Moreover, DCPTA treatment maintained
 664 increased the photosynthetic capacity, supply nitrogen metabolism of energy and
 665 carbon skeleton thus alleviating the inhibition of growth by drought in maize.

666
 667 **Acknowledgements** This work was supported by the National Key Research
 668 and Development Program of China (2016YFD0300103), the National Key Research
 669 and Development Program of China (2017YFD0300506), and the “Academic
 670 Backbone” Project of Northeast Agricultural University (17XG23).

671

672 **References**

- 673 **Ali F, Bano A, Fazal A.** 2017. Recent methods of drought stress tolerance in
674 plants. *Plant Growth Regulation* **82**, 1–13. doi: 10.1007/s10725-017-0267-2
- 675 **Bethke PC, Drew MC.** 1992. Stomatal and nonstomatal components to inhibition of
676 photosynthesis in leaves of capsicum annuum during progressive exposure to
677 NaCl salinity. *Plant Physiology* **99**, 219–26. doi: 10.1104/pp.99.1.219
- 678 **Barro F, Fontes AG, Maldonado JM.** 1991. Organic nitrogen content and nitrate and
679 nitrite reductase activities in tritordeum and wheat grown under nitrate or
680 ammonium. *Plant and Soil* **135**, 251–256. doi: 10.1007/BF00010913
- 681 **Benedict CR, Rosenfield CL, Mahan JR, Madhavan S, Yokoyama H.** 1985. The
682 chemical regulation of carotenoid biosynthesis in citrus. *Plant*
683 *Science* **41**, 169–173. doi: 10.1016/0168-9452(85)90084-6
- 684 **Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram
685 quantities of protein utilizing the principle of protein-dye binding. *Analytical*
686 *Biochemistry* **72**, 248–254. doi: 10.1006/abio.1976.9999
- 687 **Cataldo DA, Maroon M, Schrader LE, Youngs VL.** 1975. Rapid colorimetric
688 determination of nitrate in plant tissue by nitration of salicylic
689 acid. *Communications in Soil Science and Plant Analysis* **6**, 71–80. doi:
690 10.1080/00103627509366547
- 691 **Chamizo-Ampudia A, Sanz-Luque E, Llamas A, Galvan A, Fernandez E.** 2017.
692 Nitrate reductase regulates plant nitric oxide homeostasis. *Trends in Plant*
693 *Science* **22**, 163. doi: 10.1016/j.tplants.2016.12.001
- 694 **Chardon F, Noël V, Masclaux-Daubresse C.** 2012. Exploring new in crops and in
695 arabidopsis ideotypes to improve yield and seed quality. *Journal of Experimental*
696 *Botany* **63**, 3401. doi: 10.1093/jxb/err353
- 697 **Cottenie A, Verloo M, Kiekens L, Velghe G, Camerlynck R.** 1982. Chemical
698 analysis of plants and soils. Laboratory of analytical and agrochemistry. State
699 University, Ghent, 14–24.
- 700 **Drapeau G.** 1974. Protease from *Staphylococcus aureus*. In: Lorand L (ed) *Method of*

enzymology. Academic Press., New York, 469.

Du J, Shu S, An Y, Zhou H, Guo S, Sun J. 2016. Influence of exogenous spermidine on carbon–nitrogen metabolism under $\text{Ca}(\text{NO}_3)_2$ stress in cucumber root. *Plant Growth Regulation* **81**, 1–13. doi: 10.1007/s10725-016-0193-8

Fontaine JX, Tercé-laforge T, Armengaud P, Clément G, Renou JP, Pelletier S, Catterou M, Azzopardi M, Gibon Y, Lea PJ, Hirel B, Dubois F. 2012. Characterization of a NADH-dependent glutamate dehydrogenase mutant of *Arabidopsis* demonstrates the key role of this enzyme in root carbon and nitrogen metabolism. *Plant Cell* **24**, 4044–4065. doi: 10.1105/tpc.112.103689

Fresneau C, Ghashghaie J, Cornic G. 2007. Drought effect on nitrate reductase and sucrose-phosphate synthase activities in wheat (*Triticum durum* L.): role of leaf internal CO_2 . *Journal of Experimental Botany* **58**, 2983. doi: 10.1093/jxb/erm150

Gangwar S, Singh VP. 2011. Indole acetic acid differently changes growth and nitrogen metabolism in *Pisum sativum* L. seedlings under chromium (VI) phytotoxicity: implication of oxidative stress. *Scientia Horticulturae* **129**, 321–328. doi: 10.1016/j.scienta.2011.03.026

Gausman HW, Burd JD, Quisenberry J, Yokoyama H, Dilbeck R, Chan RB. 1985. Effect of 2-diethylaminoethyl-3,4-dichlorophenylether (DCPTA) on cotton plant (*Gossypium hirsutum*) growth and phenology. *Nature Biotechnology* **3**, 255–257. doi: 10.1038/nbt0385-255

Groat RG, Vance CP. 1982. Root and nodule enzymes of ammonia assimilation in two plant-conditioned symbiotically ineffective genotypes of alfalfa (*Medicago sativa* L.). *Plant Physiology* **69**, 614–618. doi: 10.1104/pp.69.3.614

Guan D, Al-Kaisi MM, Zhang Y, Duan L, Tan W, Zhang M, Li Z. 2014. Tillage practices affect biomass and grain yield through regulating root growth, root-bleeding sap and nutrients uptake in summer maize. *Field Crop Research* **157**, 89–97. doi: 10.1016/j.fcr.2013.12.015

Hodges M. 2000. Enzyme redundancy and the importance of 2-oxoglutarate in plant ammonium assimilation. *Journal of Experimental Botany* **53**, 905. doi:

10.1093/jexbot/53.370.905

Hu L, Xiang L, Li Z, Zhou X, Zou Z, Hu X. 2014. The photoprotective role of spermidine in tomato seedlings under salinity-alkalinity stress. *Plos One* **9**, e110855. doi: 10.1371/journal.pone.0110855

Ida S, Morita Y. 1973. Purification and general properties of spinach leaf nitrite reductase. *Plant and Cell Physiology* **14**, 661-671. doi: 10.1093/oxfordjournals.pcp.a074901

Jia Y, Zou D, Wang J, Liu H, Inayat MA, Sha H, Zheng H, Sun J, Zhao H. 2015. Effect of low water temperature at reproductive stage on yield and glutamate metabolism of rice (*Oryza sativa*, L.) in China. *Field Crop Research* **175**, 16–25. doi: 10.1016/j.fcr.2015.01.004

Kamran M, Su W, Ahmad I, Meng X, Cui W, Zhang X, Mou S, Aaqil K, Han Q, Liu T. 2018. Application of paclobutrazol affect maize grain yield by regulating root morphological and physiological characteristics under a semi-arid region. *Scientific Reports* **8**. doi: 10.1038/s41598-018-23166-z

Keithly JH, Yokoyama H, Gausman H. 1990. Enhanced yield of tomato in response to 2-(3,4-dichlorophenoxy) triethylamine (DCPTA). *Plant Growth Regulation* **9**, 127–136. doi: 10.1007/BF00027440

Keithly JH, Yokoyama H, Gausman HW. 1990. Effect of 2-(3,4-dichlorophenoxy)triethylamine (DCPTA) on the growth and development of sugarbeet. *Plant Science* **68**, 57–64. doi: 10.1016/0168-9452(90)90152-E

Kingsbury RW, Epstein E, Percy RW. 1984. Physiological responses to salinity in selected lines of wheat. *Plant Physiology* **74**, 417–23. doi: 10.1104/pp.74.2.417

Kusano M, Fukushima A, Redestig H, Saito K. 2011. Metabolomic approaches toward understanding nitrogen metabolism in plants. *Journal of Experimental Botany* **62**, 1439. doi: 10.1093/jxb/erq417

Liu P, Yin L, Deng X, Wang S, Tanaka K, Zhang S. 2014. Aquaporin-mediated increase in root hydraulic conductance is involved in silicon-induced improved root water uptake under osmotic stress in *Sorghum bicolor* L. *Journal of*

759 Experimental Botany **65**, 4747. doi: 10.1093/jxb/eru220

760 **López-Millán AF, Morales F, Andaluz S, Gogorcena Y, Abadía A, DE LRJ, Abadía J.**

761 2000. Responses of sugar beet roots to iron deficiency. Changes in carbon

762 assimilation and oxygen use. Plant Physiology **124**, 885. doi:

763 10.1104/pp.124.2.885

764 **López-Pérez L, Fernández-García N, Olmos E, Carvajal M.** 2007. The phi

765 thickening in roots of broccoli plants: an acclimation mechanism to

766 salinity? International Journal of Plant Sciences **168**, 1141–1149. doi:

767 10.1086/520722

768 **Mosaddeghi MR, Mahboubi AA, Safadoust A.** 2009. Short-term effects of tillage

769 and manure on some soil physical properties and maize root growth in a sandy

770 loam soil in western Iran. Soil and Tillage Research **104**, 173–179. doi:

771 10.1016/j.still.2008.10.011

772 **Munns R, Tester M.** 2008. Mechanisms of salinity tolerance. Annual Review of Plant

773 Biology **59**, 651–681. doi: 10.1146/annurev.arplant.59.032607.092911

774 **Naya L, Ladrera R, Ramos J, González EM, Arrese-Igor C, Minchin FR, Becana**

775 **M.** 2007. The response of carbon metabolism and antioxidant defenses of alfalfa

776 nodules to drought stress and to the subsequent recovery of plants. Plant

777 Physiology **144**, 1104. doi: 10.1104/pp.107.099648

778 **Noguchi A, Kageyama M, Shinmachi F, Schmidhalter U, Hasegawa I.** 2005.

779 Potential for using plant xylem sap to evaluate inorganic nutrient availability in

780 soil. Soil Science and Plant Nutrition **51**, 343–350. doi:

781 10.1111/j.1747-0765.2005.tb00038.x

782 **Nuccio ML, Wu J, Mowers R, Zhou H P, Meghji M, Primavesi LF, Paul**

783 **MJ, Chen X, Gao Y, Haque E, Basu SS, Lagrimini LM.** 2015. Expression of

784 trehalose-6-phosphate phosphatase in maize ears improves yield in well-watered

785 and drought conditions. Nature Biotechnology **33**, 862–869. doi:

786 10.1038/nbt.3277

787 **O'Neal D, Joy KW.** 1973. Glutamine synthetase of pea leaves. I. Purification,

788 stabilization, and PH optima. Archives of Biochemistry and Biophysics **159**,
789 113–22. doi: 10.1016/0003-9861(73)90435-9

790 **Plett D, Holtham L, Baumann U, Kalashyan E, Francis K, Enju A, Toubia**
791 **J, Roessner U, Bacic A, Rafalski A, Dhugga KS, Tester M, Garnett T, Kaiser**
792 **BN.** 2016. Nitrogen assimilation system in maize is regulated by developmental
793 and tissue-specific mechanisms. Plant Molecular Biology **92**, 293–312. doi:
794 10.1007/s11103-016-0512-5

795 **Sinclair TR.** 2011. Challenges in breeding for yield increase for drought. Trends in
796 Plant Science **16**, 289–293. doi: 10.1016/j.tplants.2011.02.008

797 **Rajasekhar VK, Oelmüller R.** 2010. Regulation of induction of nitrate reductase and
798 nitrite reductase in higher plants. Physiologia Plantarum **71**, 517–521. doi:
799 10.1111/j.1399-3054.1987.tb02893.x

800 **Nacer B, Gillen AM, Alemu M, Hirut K, Fisher DK, Smith JR, Reddy KN.** 2013.
801 Responses of nitrogen metabolism and seed nutrition to drought stress in
802 soybean genotypes differing in slow-wilting phenotype. Frontiers in Plant
803 Science **4**, 498. doi: 10.3389/fpls.2013.00498

804 **Robredo A, Pérez-López U, Miranda-Apodaca J, Lacuesta M, Mena-Petite A,**
805 **Muñoz-Rueda A.** 2011. Elevated CO₂ reduces the drought effect on nitrogen
806 metabolism in barley plants during drought and subsequent
807 recovery. Environmental and Experimental Botany **71**, 399–408. doi:
808 10.1016/j.envexpbot.2011.02.011

809 **Sánchez-Rodríguez E, Rubio-Wilhelmi MM, Ríos JJ, Blasco B, Rosales MÁ,**
810 **Melgarejo R, Romero L, Ruiz JM.** 2011. Ammonia production and
811 assimilation: its importance as a tolerance mechanism during moderate water
812 deficit in tomato plants. Journal of Plant Physiology **168**, 816. doi:
813 10.1016/j.jplph.2010.11.018

814 **Schiermeier Q.** 2014. The parched planet: water on tap. Nature **510**, 326–328. doi:
815 10.1016/j.jplph.2010.11.018

816 **Slaterry RA, Walker BJ, Weber A, Ort DR.** 2017. The impacts of fluctuating light

on crop performance. *Plant Physiology* **176**, 990–1003. doi:
10.1104/pp.17.01234

Talaat NB, Shawky BT. 2016. Dual application of 24-epibrassinolide and spermine confers drought stress tolerance in maize (*Zea mays*, L.) by modulating polyamine and protein metabolism. *Journal of Plant Growth Regulation* **35**, 518–533. doi: 10.1007/s00344-015-9557-y

Talaat NB, Shawky BT, Ibrahim AS. 2015. Alleviation of drought-induced oxidative stress in maize (*Zea mays*, L.) plants by dual application of 24-epibrassinolide and spermine. *Environmental and Experimental Botany* **113**, 47–58. doi: 10.1016/j.envexpbot.2015.01.006

Trenberth KE, Dai A, Schrier GVD, Jones PD, Barichivich J, Briffa KR, Sheffield J. 2014. Global warming and changes in drought. *Nature Climate Change* **4**, 17–22. doi: 10.1038/nclimate2067

Wan JH, Mendoza K. 1992. Effect of 2-(3,4-dichlorophenoxy) triethylamine on guayule (*Parthenium argentatum*) cell suspension cultures. *Journal of Agricultural and Food Chemistry* **40**, 924–926. doi: 10.1021/jf00017a046

Wang H, Wu Z, Han J, Zheng W, Yang C. 2012. Comparison of ion balance and nitrogen metabolism in old and young leaves of alkali-stressed rice plants. *Plos One* **7**, e37817. doi: 10.1371/journal.pone.0037817

Weber AP. 2007. High-throughput colorimetric method for the parallel assay of glyoxylic acid and ammonium in a single extract. *Analytical Biochemistry* **362**, 151–153. doi: 10.1016/j.ab.2006.12.033

Xia XJ, Zhou YH, Shi K, Zhou J, Foyer CH, Yu JQ. 2015. Interplay between reactive oxygen species and hormones in the control of plant development and stress tolerance. *Journal of Experimental Botany* **66**, 2839–2856. doi: 10.1093/jxb/erv089

Xie T, Gu W, Meng Y, Li J, Li L, Wang Y, Qu D, Wei S. 2017. Exogenous DCPTA ameliorates simulated drought conditions by improving the growth and photosynthetic capacity of maize seedlings. *Scientific Reports* **7**. doi:

846 10.1038/s41598-017-12977-1

847 **Yang Y, Lu X, Bei Y, Li B, Jin S, Guo S, Tezuka T.** 2013. Bottle gourd
848 rootstock-grafting affects nitrogen metabolism in nacl-stressed watermelon
849 leaves and enhances short-term salt tolerance. *Journal of Plant Physiology* **170**,
850 653. doi: 10.1016/j.jplph.2012.12.013

851 **Yemm EW, Cocking EC.** 1955. The determination of amino-acids with
852 ninhydrin. *Analyst* **80**, 209–214. doi: 10.1039/an9558000209

853 **Zhao J, Xue Q, Jessup KE, Hao B, Hou X, Marek TH, Xu W, Evett SR,**
854 **O’Shaughnessy SA, Brauer DK.** 2018. Yield and water use of drought-tolerant
855 maize hybrids in a semiarid environment. *Field Crops Research* **216**. doi:
856 10.1016/j.fcr.2017.11.001

857 **Ziyomo C, Bernardo R.** 2013. Drought tolerance in maize: indirect selection through
858 secondary traits versus genomewide selection. *Crop Science* **53**, 1269–1275. doi:
859 10.2135/cropsci2012.11.0651

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875 **Tables**

876 **Table 1** Chemical properties of the used soil.

Year	pH	HCO ₃ ⁻ + CO ₃ ²⁻ (mg kg ⁻¹)	Cl ⁻ (mg kg ⁻¹)	SO ₄ ²⁻ (mg kg ⁻¹)	Ca ₂ ⁺ (mg kg ⁻¹)	Mg ₂ ⁺ (mg kg ⁻¹)	Na ⁺ (mg kg ⁻¹)	K ⁺ (mg kg ⁻¹)	N (mg kg ⁻¹)	P (mg kg ⁻¹)
2016	7.2	204.3	297.6	463.8	87.4	37.6	4.1	28.9	17.3	3.9
2017	7.1	201.7	302.5	446.2	90.3	40.9	4.0	31.4	16.1	3.8

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897 **Figure legends:**

898 **Fig. 1** The rainfall (bar) and mean temperature (line) data during the study period
899 (2016 and 2017, May–October).

900 **Fig. 2** The pits (inner length, 10 m; width, 7 m; and height, 1.2 m) used for this study
901 and the plastic sheets used to cover inner sides of the pits.

902 **Fig. 3** Changes in the soil water content (SWC) in 2016 and 2017. The data represent
903 the means of independent measurements with five replicates, and the standard
904 deviations are indicated by the vertical error bars. Values with the same letters on the
905 bars are not significantly different at $P < 0.05$ (LSD test).

906 **Fig. 4** The tools used for the collection of bleeding sap. (A) cotton, (B) centrifuge
907 tube, (C) scissors, (D) plastic film, and (E) deionized water.

908 **Fig. 5** Changes in the grain number plant⁻¹ and grain yield plant⁻¹ (g) of maize in 2016
909 and 2017. The data represent the means of independent measurements with five
910 replicates, and the standard deviations are indicated by the vertical error bars. Values
911 with the same letters on the bars are not significantly different at $P < 0.05$ (LSD test).

912 **Fig. 6** Leaf phenotypic features of the maize seedlings after 20 days of treatment with
913 drought and/or DCPTA in 2016 and 2017.

914 **Fig. 7** Effect of drought and/or DCPTA treatment on the relative growth rate (RGR)
915 of the shoots and roots of the maize in 2016 and 2017. The data represent the means
916 of independent measurements with five replicates, and the standard deviations are
917 indicated by the vertical error bars. Values with the same letters on the bars are not
918 significantly different at $P < 0.05$ (LSD test).

919 **Fig. 8** Effect of drought and/or DCPTA treatment on root length density (RLD) and
920 root surface area density (RSD) for different soil depths on the 20th day and 30th day
921 after treatment in 2016 and 2017.

922 **Fig. 9** Effect of drought and/or DCPTA treatment on root hydraulic conductivity, the
923 flow rate of root-bleeding sap and the NO₃⁻ delivery rate in 2016 and 2017. The data
924 represent the means of independent measurements with five replicates, and the
925 standard deviations are indicated by the vertical error bars. Values with the same

926 letters on the bars are not significantly different at $P < 0.05$ (LSD test).

927 **Fig. 10** Effect of drought and/or DCPTA treatment on relative water content of the
928 leaves (RLWC) in 2016 and 2017. The data represent the means of independent
929 measurements with five replicates, and the standard deviations are indicated by the
930 vertical error bars. Values with the same letters on the bars are not significantly
931 different at $P < 0.05$ (LSD test).

932 **Fig. 11** Effect of drought and/or DCPTA treatment on the gas exchange parameters in
933 the maize leaves in 2016 and 2017. Pn, net photosynthetic rate; Gs, stomatal
934 conductance; Tr, transpiration rate; Ci, intercellular CO₂ concentration. The data
935 represent the means of independent measurements with five replicates, and the
936 standard deviations are indicated by the vertical error bars. Values with the same
937 letters on the bars are not significantly different at $P < 0.05$ (LSD test).

938 **Fig. 12** Ultrastructure of the photosynthetic apparatus in the maize leaves after 20
939 days of treatment with drought and/or DCPTA in 2016 and 2017. SL, stroma lamella;
940 GL, grana lamellae; CW, cell wall; and P, plastoglobule. The scale bars for the
941 photosynthetic apparatus represent 2000 nm.

942 **Fig. 13** Effects of drought and/or DCPTA treatment on isocitrate dehydrogenase
943 (ICDH) activity in 2016 and 2017. The data represent the means of independent
944 measurements with five replicates, and the standard deviations are indicated by the
945 vertical error bars. Values with the same letters on the bars are not significantly
946 different at $P < 0.05$ (LSD test).

947 **Fig. 14** Effects of drought and/or DCPTA treatment on the nitrate (NO₃⁻), nitrite
948 (NO₂⁻) and ammonium (NH₄⁺) contents in the leaves of maize in 2016 and 2017. The
949 data represent the means of independent measurements with five replicates, and the
950 standard deviations are indicated by the vertical error bars. Values with the same
951 letters on the bars are not significantly different at $P < 0.05$ (LSD test).

952 **Fig. 15** Effects of drought and/or DCPTA treatment on nitrate reductase (NR) and
953 nitrite reductase (NiR) activities in the leaves of maize in 2016 and 2017. The data
954 represent the means of independent measurements with five replicates, and the

955 standard deviations are indicated by the vertical error bars. Values with the same
956 letters on the bars are not significantly different at $P < 0.05$ (LSD test).

957 **Fig. 16** Effects of drought and/or DCPTA treatment on the glutamine synthetase (GS),
958 glutamate synthase (GOGAT) and glutamate dehydrogenase (GDH) activities in the
959 leaves of maize in 2016 and 2017. The data represent the means of independent
960 measurements with five replicates, and the standard deviations are indicated by the
961 vertical error bars. Values with the same letters on the bars are not significantly
962 different at $P < 0.05$ (LSD test).

963 **Fig. 17** Effects of drought and/or DCPTA treatment on the alanine aminotransferase
964 (AlaAT) and aspartate aminotransferase (AspAT) activities in the leaves of maize in
965 2016 and 2017. The data represent the means of independent measurements with five
966 replicates, and the standard deviations are indicated by the vertical error bars. Values
967 with the same letters on the bars are not significantly different at $P < 0.05$ (LSD test).

968 **Fig. 18** Effects of drought and/or DCPTA treatment on the protease activity and
969 protein and free amino acid contents in the leaves of maize in 2016 and 2017. The
970 data represent the means of independent measurements with five replicates, and the
971 standard deviations are indicated by the vertical error bars. Values with the same
972 letters on the bars are not significantly different at $P < 0.05$ (LSD test).

- Well-watered
- Well-watered+DCPTA
- Drought
- Drought +DCPTA

2016

bioRxiv preprint doi: <https://doi.org/10.1101/385203>; this version posted August 4, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

2017

Well-watered

Well-watered
+DCPTA

Drought

Drought
+DCPTA

Well-watered Well-watered+DCPTA Drought Drought +DCPTA

2016

2017

2016

2017

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

bioRxiv preprint doi: <https://doi.org/10.1101/385203>; this version posted August 4, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Days after treatment

Days after treatment

2016

2017

Well-watered

Well-watered
+DCPTA

Drought

Drought
+DCPTA

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Days after treatment

Days after treatment

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Days after treatment

Days after treatment

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Well-watered Well-watered+DCPTA Drought Drought+DCPTA

2016

2017

Drought

Rehydration

Drought

Rehydration

Days after treatment

Days after treatment