Title: Determinants of Minimum Acceptable Diet among 6–23 Months Age Children in Ethiopia: A Multilevel Analysis of The Ethiopian Demographic Health Survey

Short title: Determinants of Minimum Acceptable Diet

Aberash abay¹ ¶*, Dejen Yemane² ¶, Abate Bekele³ ¶, Beyene Meressa²

¹School of Public Health, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia

²Department of Environmental Health and Behavioral Sciences, School of Public Health, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia

³Department of Biostatistics, School of Public Health, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia

* Corresponding Author

E-mail: beyene2006m@gmail.com[BM]

Abstract

Background: Though infant and young children should be fed according to a minimum acceptable diet to ensure appropriate growth and development, only 7% of Ethiopian 6-23 months age children meet the minimum acceptable dietary standards, which is lower than the national target of 11% set for 2016. Therefore, this study aims to assess the individual and community level determinants of minimum acceptable diet among 6–23 months age children in Ethiopia.

Methods: This study analyzed retrospectively a cross-sectional data on a weighted sample of 2919 children aged 6-23 months nested within 617 clusters after extracting from Ethiopian Demographic and Health Survey 2016 via the link www.measuredhs.com. By employing bivariate multilevel logistic regression model, variables which were significant at the p-value < 0.25 were included in multivariable multilevel logistic regression analysis. Finally, variables with p-value < 0.05 were considered as significant predictors of minimum acceptable diet.

Results: Only 6.1% of 6-23 months age children feed minimum acceptable diet in Ethiopia. Children 18-23 months age (AOR=3.7, 95%CI 1.9, 7.2), father's with secondary or higher education (AOR=2.1, 95%CI 1.2, 3.6), Employed mothers (AOR=1.7, 95%CI 1.2, 2.5), mothers have access to drinking water (AOR=1.9, 95%CI 1.2, 2.9), mothers with media exposure (AOR=2.1 95%CI 1.1, 2.7) were positive individual level predictors. Urban mothers (AOR=4.8, 95%CI 1.7, 13.2)) and agrarian dominant region (AOR=5.6, 95%CI 2.2, 14.5) were community level factors that significantly associated with minimum acceptable diet of 6–23 months age children.

Conclusion: Both individual and community level factors were significantly associated with minimum acceptable diet of 6-23 months age children in Ethiopia, suggesting that nutritional

39 interventions designed to improve child health should not only be implemented at the individual
40 level but tailored to community context as well.

41 **Keywords:** Determinants, Ethiopia Demographic Health Survey, Minimum Acceptable Diet,
42 Multilevel Analysis

Introduction

After 6 months, breast milk is no longer adequate to meet the nutritional needs and increasing demand of nutritional requirements of infants and children[1]. Complementary feeding is the process of transition from exclusive breastfeeding to other foods besides breast milk[2]. During this period timely introduction of complementary feeding with a variety of foods should be added to the child's diet to ensure their nutritional requirement[3]. World Health Organization (WHO) has established guidelines for infant and young child feeding (IYCF) practices for 6–23 months age children by considering minimum acceptable diet (MAD) as one of the eight core indicators of complementary feeding [4]. It is the combination of minimum dietary diversity and minimum meal frequency. Minimum dietary diversity for breastfed and non breast feed children 6-23 months is defined as receiving four or more food groups out of the seven food groups[5] [7].

Minimum meal frequency for breastfed and non breast feed 6-23 month children is defined as two or more feedings of solid, semi-solid, or soft food for 6-8 months, three or more feedings for 9-23 months breast feed and four times for non breast feed children[5].

Low dietary diversity and meal frequency practices are determinant for health and growth in children less than 2 years of age. They increase the risk of under-nutrition, illness, and mortality in infants and young children [6]. Even with optimum breastfeeding, children will become stunted if they do not receive sufficient dietary diversity and frequency over 6 months of age [7].

Supplementing breast feeding with nutritious complementary foods can reduce stunting among children of this age by 20% [8].

According to the recent demographic and health survey reports of 10 Asian and African countries, including Ethiopia feeding a child minimum acceptable diet ranges from 7% in

Ethiopia to 36% in Nepal. This indicates that, feed the minimum acceptable diet is a major problem both globally and in developing countries[1,3, 9-16].

The Ethiopian government has been implementing the national nutrition program, IYCF and has recently developed a multi-sectoral plan of nutrition intervention (Sekota Declaration), which aims to address the immediate, basic and underlying causes of malnutrition to end child under nutrition in Ethiopia by 2030 [17]. These interventions are meant to tackle the nutrition problems in children, including those with inappropriate feeding. However, the progress was not satisfactory, particularly; the MAD has increased from 3% to 7% in a decade (2005-2016).

Factors associated with the minimum acceptable diet are complex, ranging from the individual to the community level factors. Though some studies conducted about determinants of complementary feeding in Ethiopia, no information was documented about minimum acceptable diet independently. However, these small scale studies were limited in scope as the data used were not nationally representative. In addition, they are also limited in their methodology in which they have applied traditional logistic regression to identify aforementioned predictors of minimum acceptable diet, using a traditional level logistic regression analysis to analyze a data that has a hierarchical structure (i.e. Children are nested within the communities) violates the independence assumptions of regression and gives an incomplete picture to understand the true association of minimum acceptable diet and its determinants. Likewise, findings from such studies could not be generalized to the entire Ethiopian children.

Hence, to address these limitations and to further document the significant effect of individual and community level factors on minimum acceptable diet, this study utilized a multilevel logistic regression model. Therefore, the purpose of this study will be to determine the individual and community level factors that are associated with minimum acceptable diet in Ethiopia.

Methods and materials

Data source and study subjects

For this study, the EDHS 2016 child recode data were accessed from www.measuredhs.com. It is the recent and the fourth nationally representative survey data from 9 regions and two administrative cities. The survey used a two-stage cluster sampling design with rural-urban and regions as strata yielding 21 sampling strata. In the first stage, a total of 645 enumeration areas (EAs) was selected. An EA is a geographic area consisting of 200-300 household, which served as a counting unit for the census. In the second stage, a fixed number of 28 households per cluster were selected randomly from the household listing. A total of 16,650 eligible households and 15,683 women (15-49 years) were interviewed, making up response rates of 98% and 95%, respectively. All women who had at least one child living with them who was last born two-years preceding the survey were asked questions about the types of food the child had consumed during the day or night before the interview. Mothers who had more than one child within the two years preceding the survey were asked questions about the youngest child living with them [1].

In our study, a total of 2919 children of 6-23 months age who had measurement on their feeding behavior and who live with their mother during the survey were included. Accordingly, children who did not live with their mother during the survey (n=26), and those children's who had incomplete information about MAD (n=98) were excluded from our study. The child recode data used in this study comprised all the data related with children and their parents. Further, the details of sampling design, data collection, and data quality are available in the EDHS 2016 reports [1].

Study variables

The DHS recode-6 manual and questionnaire at the end the EDHS 2016 report along with other relevant literatures were used to select appropriate variables for current analysis. The outcome variable was minimum acceptable diet (MAD). It is a binary outcome variable and coded as 0 if the child didn't feed minimum acceptable diet and coded as 1 if the child feed a minimum acceptable diet. The child is said to be fed with MAD if he/she had both minimum meal frequency and dietary diversity in both BF and non-BF children

(39). The children who received solid, semi-solid or soft foods, two times for breastfed infants 6–8 months, three times for breastfed children 9–23 months and four times for non-breastfed children is said to have minimum meal frequency. The children with 6–23 months of age who received foods from four or more food groups of the seven food groups (Cereals, Legumes and nuts, Dairy products, Eggs, Flesh foods, Vitamin A rich fruits and dark green leafy vegetables and other fruits) are said to have minimum dietary diversity [2].

In this study, the effect of two-level explanatory factors i.e. the individual and community level factors was examined. The relationship between the explanatory variables and the minimum acceptable diet is depicted using the socio-ecologic model [18]. According to this model, individual dietary or health behavior is not only determined by the individual's characteristics, but also the community level attributes (**Fig 1**). The individual level variables are further categorized as paternal and household characteristics, and child related variables. While, at community level, place of residence (either urban vs. rural or by agro-ecologic regions) and community level aggregate variables such as community poverty, women's education, media exposure and ANC utilization were included. The EDHS did not capture data that can directly describe the characteristics of the community/clusters. Hence, we created community variables by aggregating the individual mothers' characteristics within their clusters. The aggregates were computed using the average values of the proportions of women in each category of a given variable. Likewise, based on national median values the aggregate values were categorized into groups.

Statistical analysis

The data were analyzed using STATA version 12. A sampling weight was used for computing all descriptive statistics to adjust for the non-proportional allocation of the sample to different regions and their urban and rural areas and for the possible differences in response rates. Hence, the actual representativeness of the survey results at both the national and regional levels is ensured. The 'Svy' command was used to allow for adjustments for the cluster sampling design. Tables were used for data presentation. Frequency and percentage were used to report categorical variables.

To examine the effect of multilevel factors on the individual dietary behavior, multilevel modeling approach was used. The nested nature of EDHS data makes the use of traditional regression methods inappropriate because of the assumption of independence among individuals within the same group, assumption of equal variance across groups which are inherent in traditional regression methods are violated. Therefore, in this study, a two-level mixed effect logistic regression analysis was employed in order to estimate both independent (fixed) effects of the explanatory variables and community-level random effects on feeding MAD. The first level represents the individual (children) and the second level is the cluster (community). Hence, the log of the probability of feeding MAD was modeled using two-level multilevel model as follows:

$$\text{Log}\left(\frac{\pi_{ij}}{1 - \pi_{ij}}\right) = Y_{ij} = \beta_0 + \beta_1 X_{ij} + \beta_2 Z_{ij} + u_j$$

Where, Y_{ij} Is the feeding of MAD for the individual i in j cluster/community; X_{ij} and Z_{ij} are individual and community level variables for the i^{th} individual in group j , respectively. The β 's are fixed coefficients indicating a unit increase in X can cause a β unit increase in probability feeding MAD. While, the β_0 is intercept that is the effect on feeding MAD when the effect of all explanatory variables absent (38). The u_j shows random effect (effect of the community on mother's decision to provide MAD) for the j^{th} community. By assuming each community has different intercept (β_0) and fixed coefficient (β), the clustered data nature and the within and between community variations were taken in to account. The measures of community variation (random-effects) was estimated as intra class correlation coefficient (ICC) which is proportion of the total variance in feeding MAD due to variables operating at a community level. Hence, the ICC was estimated as: $\rho = (\sigma_u^2 / (\sigma_u^2 + \pi^2 / 3))$, Where, ρ is the ICC, σ_u^2 is the variance at the community level; $\pi^2 / 3 = 3.29$ represent the fixed individual level variance (37). Given the above assumptions, four models were developed, namely: Null-model – is a model with no explanatory variables; Model I – is a model with only individual level factors are controlled; Model II – a model with only community level factors are controlled and Model III – is also called combined model which controls the effect of both individual and community level explanatory variables on mothers decision to provide MAD to their child (**Table 1**).

The null model – the model with no covariates – was fitted to examine the between community variation and to justify the use multilevel analysis. Likewise, about 40% of the total variance

(ICC) in the odds of feeding a child with MAD is due to community level factors ($P < 0.001$), indicating using multilevel modeling is better to get valid estimates than using ordinary logistic regression. The Proportional Change in Variance (PCV) was computed for each model with respect to the empty model to understand the relative contributions of both individual and community level variables to the community level variance on feeding MAD to a child. It was calculated as $PCV = (V_e - V_{mi}) / V_e$; where, V_e is variance in the empty model and V_{mi} is variance in successive models (**Table 1**).

The between-community variability declined in successive models, from 39.6% in null model, to 28.3% in individual-level only model, 27.5% in community-level only model and 26.6% in combined model. Akaike information criterion (AIC) was used to select a best model that explains the variation in feeding MAD well. The model with the lowest AIC is the preferred, accordingly the combined model showed lowest AIC than others. The combined model implied that 44.9% of the total explained variance in the odds of feeding MAD to a child could be attributed to both individual and community level characteristics in the model. However, the variance in the combined model remained significant ($p < 0.001$), indicating the presence of other community level factors – which were not included in our model – that can explain the feeding of MAD to a child. In other words, even though the unexplained community level variance was reduced in the mixed model, the remaining community-level variance still remains significant (**Table 1**).

Table 1: Individual and community-level variances for multilevel random intercept Logit models predicting feeding 6-23 months age children with minimum acceptable diet in Ethiopia 2016.

Random effect	Null model	Model I	Model II	Model III (combined)
Variance	2.16*	1.30*	1.25*	1.19*
ICC (%)	39.6	28.3	27.5	26.6
PCV (%)	Reference	39.8	42.1	44.9
AIC	1296.4	1153.3	1238.7	1142.7

Note: Null model – a model with no covariates; Model I – only individual level explanatory variables were included; Model II – only community level explanatory variables were included; Model III (Combined) – both individual and community level variables were included; *significant at $p < 0.001$; AIC: Akaike Information Criterion.

207 The measures of association (fixed-effects) between the odds of feeding a child with MAD and
 208 various explanatory variables were expressed as adjusted odds ratio (AOR) at their 95 %
 209 confidence intervals (CI). The statistical significance was set at p-value of 0.05. The
 210 multicollinearity among independent variables was checked using variance inflation factor
 211 (VIF). The VIF value of all predictor variables was less than 10 indicating the absence of a
 212 significant correlation among the explanatory variables. The presence of interaction among the
 213 explanatory variables was checked and there was no significant interaction between them.

214 This study was approved by the Ethical Review Committee of College of Health Sciences,
 215 Mekelle University . Then, a written inquiry to access data from the measure DHS website was
 216 submitted to the ORC macro INC, Chicago and permission was obtained accordingly. This study
 217 was based on the EDHS 2016 dataset with all participants' identifiers removed, no further effort
 218 was made to trace back the subjects and the data was kept confidential as per the agreement
 219 made with the ORC macro. Otherwise, the DHS was approved by the Ethiopian Health Nutrition
 220 and Research Institute (EHNRI) Review Board and the National Research Ethics Review
 221 Committee (NRERC) at the Ministry of Science and Technology, Ethiopia.

Results

Individual level characteristics of the study participants

Five hundred fifty-six (19%) children were in the age group 6–8 months, 492 (17%) in age 9-11 months, 1071 (37%) in 12–17 months and the rest 799 (27%) found in 18–23 months. More than half of the children (52.9%) were females. More than two-third 1942 (67%) children were born within more than 24 months of pregnancy interval. About three-fourth mothers 2110 (72%) were in age between 20-34 years. Sixty-two percent of the mothers and 1207 (44%) of fathers had no education. About forty percent of participants were Muslims followed by Orthodox 998 (34.2%). Regarding the employment status, 1685 (58 %) of the mothers were not working at the time of EDHS 2016 survey. More than three-fourth 2521 (86.4%) of households were headed by males. More than two-third 2009 (69%) of the households have at least five family sizes and 1387 (47.5%) were with two under five children. More than half of households 1519 (53%), travel 30 minutes or longer round trip to fetch drinking water. On the other hand, only 90 (3%) were used efficient cooking fuel (electricity) but most 2373 (81.9%) of households use wood as cooking fuel. The proportions of mothers with no ANC visit and at least 4 visits during their last pregnancy were nearly the same, 1019 (35%) and 1002 (34.5%), respectively (**Table 2**).

Table 2: The individual level characteristics of 6-23 months age children in Ethiopia, EDHS 2016(n=2919).

Individual level variables	Frequency (n)	Percentage (%)
Child age(months)		
6-8 month	556	19.0
9-11 month	492	16.9
12-17 month	1071	36.7
18-23 month	799	27.4
Child Sex		
Female	1,543	52.9
Male	1376	47.1
Birth Order		
First	550	18.9
Second –fourth	1256	43.0
Fifth and above	1113	38.1

Birth Interval

No previous birth	552	18.9
< 24 months	422	14.5
>= 24 months	1942	66.6

Mother age

<20	361	12.4
20-34	2110	72.3
35-49	448	15.3

Mother Education

No education	1,797	61.6
Primary education	897	30.7
Secondary & Higher	225	7.7

Mother Occupation

Unemployed	1,685	57.7
Employed	1,234	42.3

Father Education

No education	1,207	44.1
Primary education	1,156	42.1
Secondary & Higher	378	13.8

Father Occupation

Unemployed	223	7.7
Employed	2,696	92.3

240

241 **Table 2:** continued

Individual level variables	Frequency (n)	Percentage (%)
ANC visit		
No	1019	35.1
1-3	884	30.4
>=4	1002	34.5
Media Exposure		
Yes	978	33.5

No	1941	66.5
Religion		
Protestant	642	22.0
Orthodox	998	34.2
Muslim	1,176	40.3
Others	103	3.5
Household wealth		
Poorest	681	23.3
Poor	623	21.4
Middle	644	22.1
Richer	531	18.2
Richest	440	15.0
Drug use Status		
Yes	24	0.8
No	2895	99.2
Time to get drinking water		
< 30 minutes	1,370	47.4
≥ 30 minutes	1519	52.6
Type of cooking fuel		
Charcoal	156	5.4
Electric	91	3.1
Wood	2373	82.0
Animal dug	173	6.0
Others	102	3.5
Family size		
< 5 family	910	31.0
≥ 5 family	2009	69.0
Under 5 Children		
One	1128	38.6
Two	1387	47.5
Three and above	404	13.9

Household head

Male	2,521	86.3
Female	398	13.7

Note: Birth interval, father education, ANC visits, time to get drinking water and types of cooking fuel have 3, 178, 14, 30 and 24 missing values, respectively.

Community level characteristics of the study subjects

More than nine in ten mothers 2679 (91.8%) were from the Agrarian dominant region. More than half 1937 (66.4%) of the mothers were from the community with high poverty. More than half mothers 1506 (52%) were from the low media access community. Most of the mothers 2576 (88.3%) were from rural areas and 2008 (69%) of the mothers were from the community with high ANC utilization (**Table 3**).

Table 3: The Community level characteristics of 6-23 months age children in Ethiopian, EDHS 2016 (n=2919)

Community level	Frequency (n)	Percentage (%)
Residence		
Urban	343	11.7
Rural	2576	88.3
Region		
Pastoralist dominant	144	4.9
Agrarian dominant	2679	91.8
City dwellers	96	3.3
Community poverty		
High	1937	66.4
Low	982	33.6
Community women education		
Low	2053	70.3
High	866	29.7
Community media exposure		
Low	1506	51.6
High	1413	48.4
Community ANC utilization		
Low	911	31.0
High	2008	69.0

Minimum acceptable diet feeding practice

In this study, of the total of 2919 children 6-23 months of age, only 6.1% were fed MAD. On the other hand, the proportion of children with minimum dietary diversity and minimum meal frequency were 11% and 42%, respectively (**Table 4**).

Table 4: Minimum meal frequency, Dietary diversity and minimum acceptable practice among children 6-13 months of age in Ethiopia, 2016 (n=2919).

Diets	Frequency (n)	Percentage (%)
Minimum acceptable diet		
No	2742	93.9
Yes	177	6.1
Meal frequency		
No	1697	58.1
Yes	1222	41.9
Dietary diversity		
No	2588	88.7
Yes	331	11.3

Individual and community level determinants of feeding MAD

The individual and community level determinants of feeding MAD are presented in Table 5. At individual level: child age, father's education, mother's occupation, time to get drinking water, media exposure; and at community level: region of residence were independent predictors of feeding the child MAD.

The odds of feed the child MAD was nearly 4 times (AOR=3.7; 95%CI 1.9, 7.3) higher among children in age between 18-23 months than children at age 6-8 months. The odds of feeding the child with a minimum acceptable diet were 70% (AOR=1.7; 95%CI 1.2, 2.5) higher among employed mothers compared with unemployed mothers.

Mothers who have media exposure had nearly 2 times (AOR=1.95; 95%CI 1.3, 2.9) higher odds of feeding a MAD for their children than mothers with no media access. The odds of feeding MAD to a child was twice (AOR=2.0; 95%CI 1.3, 3.1) higher among mothers who travel less than 30 minutes to fetch drinking water. Father's education was significantly positively associated with feeding minimum acceptable diet to the child. Fathers who educated secondary or higher had 2.4 times (AOR=2.4; 95%CI 1.4, 4.0) higher odds to feed their children MAD.

The region was a significant predictor of feeding the child MAD. Mothers from agrarian region had 5 times (AOR=5.1; 95% CI 2.0, 13.1) and city dwellers had nearly 5 times (AOR=5.4; 95%CI 1.9, 14.8) higher odds to feed their children MAD respectively compared with mothers from pastoralist (**Table 5**).

Table 5: Individual and community level determinants of feeding minimum acceptable diet for 6-23 months age children in Ethiopia, 2016(n=2919).

Individual and community-level	COR(95%CI)	AOR(95%CI)	p-value
Child age in months			
6-8 month	1:00	1:00	
9-11 month	3.2(1.6, 6.4)	3.1(1.5,6.4)	0.002
12-17 month	3.2(1.7, 6.1)	3.5(1.8,6.7)	0.0001
18-23 month	3.6(1.9, 6.8)	3.7(1.9,7.3)	0.0001
Time to get drinking water			
≥ 30 minutes	1:00	1:00	
< 30 minutes	3.2(2.1, 4.7)	2(1.3,3.1)	0.001
Mother's Occupation			
Unemployed	1:00	1:00	
Employed	1.7(1.4, 2.8)	1.7(1.2,2.5)	0.003
Media Exposure			
No exposure	1:00	1:00	
Has media exp	2(1.3, 3.3)	1.7(1.3,2.9)	0.002
Father Education			
No education	1:00	1:00	
Primary	2.1(1.3, 3.3)	1.4(0.9,2.3)	0.152
Secondary or higher	4.6(2.9, 7.3)	2.4(1.4,4.0)	0.001
Region			
Pastoralist dominant	1:00	1:00	
Agrarian dominant	8.9(3.5, 22.4)	5.1(2, 13.1)	0.001
City dwellers	16.0(5.9,42.9)	5.4(1.9,14.8)	0.001

Discussion

This study used multilevel logistic regression analysis to address these two level factors. Accordingly the individual level factors such as the child's age, mother's occupation, father's education, access to drinking water and media exposure showed a significant association with feeding minimum acceptable diet.

The age of the child was positively associated with feeding minimum acceptable diet for 6-23 months age children. Children with age 18-23 months were found to be significantly higher odds to be feed minimum acceptable diet than 6-8 months. This finding is consistent with evidences from Indonesia, Pakistan, Ghana and Uganda [7, 8, 19, 20]. This might be due to late introduction of complementary feeding and when they start complimentary feeding on time; they included only milk or cereal products. Other possibility could be mothers may perceive that younger the children, the poor ability of intestine to digest certain foods like banana, egg, pumpkin, carrot, green vegetables and Meat [21]. This could be further justified by traditional beliefs and practices, during introducing complementary food to infants in rural community, infants may develop diarrhea due to poor hygienic condition, but mothers could associate this problem with taking different food items and eventually she might not permit the child to taste unfamiliar foods. It can also be attributed to the feeding interest of the child too.

Father's education level was significantly and positively associated with feeding the child minimum acceptable diet. Other studies in South Asia, Bangladesh and Nepal had reported a similar finding indicating that children whose fathers had secondary or higher level of education were more likely to be provided with the recommended acceptable diet as compared to the children whose fathers did not have any education [22, 23, 24]. This could be due to educated fathers were more likely to have information (media exposure), understand educational messages about child feeding easily, might have received lessons on child feeding in the curricula at school that would increase their knowledge about the importance of child feeding. However, different finding was reported from study done in Serilanka, in which paternal education was not associated with any of complementary feeding indicators [25]. This difference could be due to study area and sample size. The current study included large population from different geographic regions with various culture, beliefs, and traditions.

Mother's occupational status was also significant predictor of feeding minimum acceptable diet to the child. Children who have employed mothers were more likely to be feed minimum acceptable diet. This result is consistent with study conducted in Indonesia and Serilanka [19, 25]. It could give us an indication that mother's earning ability is an important factor in feeding the child a minimum acceptable diet in Ethiopia. Enhanced access to resources, wider social networks and growing understanding of their social environment could increases opportunities to feed the child minimum acceptable diet than unemployed mothers.

This study also revealed that time to get drinking water showed a positive effect on feeding the children a minimum acceptable diet. Mothers who travel less than 30 minutes round trip to fetch drinking water had higher odds to feed their children a minimum acceptable diet than those who travel 30 or more minutes. This could be attributed to that, mothers who travel less than 30 minutes to fetch drinking water could have time to take care of their children, including feeding them properly and adequately which therefore could lead to meeting their MAD. In addition, if there is access to drinking water source this could also encourage households to use water for on-plot gardening, which can improve dietary diversity. Furthermore, if there is access to drinking water, time spent to fetching water could be invested in additional productive activities that can augment economic and educational empowerment of mothers.

In this study exposure to media was also a significant predictor of feeding minimum acceptable diet to the child. Families who have exposure to media had increased odds of feeding minimum acceptable diet to their child than those who have no media exposure. This is supported by studies conducted in south Asia and India which reported low feeding practice of minimum acceptable diet for the children in the families who have no media exposure [24, 26]. This might be pointing to the influence of media on behavioral change to improve the minimum acceptable diet through enhancing mother's knowledge on feeding a minimum acceptable diet to their children.

From the community level factors geographic region of the study participants was found to be statistically significant predictor of feeding minimum acceptable diet in Ethiopia. Children from city dwellers and agrarian dominant had more odds to be fed minimum acceptable diet than children from pastoralist dominant region. This might be due to Ethiopia is large country with

diverse cultures, religions reflected by different food habits and traditional practices [21]. This could also be linked with less food production and higher levels of poverty.

Furthermore, the current study found that; even though most variation on feeding the child minimum acceptable diet was explained by individual-level factors, some of the variation was also explained by unmeasured community-level factors. The random effects of the community-level model were significant in explaining feeding the child minimum acceptable diet, even it reduced, from 39.7% in null model to 27% in the full model. This indicates feeding the child minimum acceptable diet was explained by both individual and community-level factors. The efforts of improving the practice of MAD in Ethiopian should address both factors operating at individual and community level.

Conclusion

The proportion of young children aged between 6–23 months receiving minimum acceptable diet was very low (6.1%). This study also showed that this low proportion of minimum acceptable diet among 6-23 months age children is determined by number of individual and community level factors such as age of the child, mother's occupation, father's education level, and media exposure, access of household drinking water and region of residence. Therefore, interventions to improve MAD practice should not only be implemented at the individual level but also be tailored to the community context. Utilizing media to promote feeding young children with minimum acceptable diet through enhancing mother's knowledge on child feeding practices and advocacy for appropriate complementary feeding, particularly on meeting MAD and nutrition education and social and behavior change interventions should be strengthened targeting the infants and children aged 6–8 months and unemployed mothers.

Acknowledgments

Our sincere and deepest gratitude goes to Mekelle University, College of Health Sciences and MU- NMBU for financial support of this study. We are also very grateful to Measure DHS for making the data freely available and we would like to extend our heartfelt thanks and appreciation to family members of Mrs.Aberash Abay (Haset Beyene, Nobel Beyene, Lidiya Beyene and Beyene Meressa) and friends for their endless support and providing the required courage during the research work.

Author Contributions

Conceptualization: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa

Formal analysis: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa

Funding acquisition: Aberash Abay

Investigation: Aberash Abay, Dejen Yemane, Abate Bekele

Methodology: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa

Project administration: Aberash Abay, Dejen Yemane, Abate Bekele

Supervision: Dejen Yemane, Abate Bekele, Beyene Meressa

Validation: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa.

Writing original draft: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa.

Writing review & editing: Aberash Abay, Dejen Yemane, Abate Bekele, Beyene Meressa

References

1. Central Statistical Agency. Ethiopia Demographic and Health Survey [Internet]. Addis Ababa; 2016. Available from: www.DHSprogram.com
2. World Health Organization. Guiding Principles for Complementary Feeding of the Breastfed (PAHO and WHO) [Internet]. Washington, D.C; 2001. Available from: <http://www.paho.com>
3. Central Statistics of Afghanistan. Afghanistan Demographic and Health Survey [Internet]. Kabul; 2015. Available from: www.DHSprogram.com.
4. Republic FD. National Nutrition Programme Republic of Ethiopia. published [Internet]. 2013;(June 2013). Available from: www.unicef.org/ethiopia/National_Nutrition_Programme.pdf
5. USAID. Nutrition Indicator Reference Sheets : External Source Data [Internet]. 2010. Available from: http://pdf.usaid.gov/pdf_docs/PNADT413.pdf
6. Beyene M, Worku AG, Wassie MM. Dietary diversity, meal frequency and associated factors among infant and young children in Northwest Ethiopia: a cross- sectional study. BMC Public Health [Internet]. 2015;15(1):1007. Available from: <http://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-015-2333-x>
7. Issaka AI, Agho KE, Burns P, Page A, Dibley MJ. Determinants of inadequate complementary feeding practices among children aged 6 – 23 months in Ghana. Public Health Nutr [Internet]. 2014;18(4):669–78. Available from: <https://www.cambridge.org/>
8. Mokori A, Schonfeldt H, Hendriks SL. Child factors associated with complementary feeding practices in Uganda Child factors associated with complementary feeding practices in Uganda. South African J Clin Nutr [Internet]. 2017;30(1):7–14. Available from: <http://dx.doi.org/10.1080/16070658.2016.1225887>
9. National Statistical Office. Malawi Demographic and Health Survey [Internet]. Zombia; 2015. Available from: www.DHSprogram.com.
10. Zimbabwe National Statistics Agency. Zimbabwe Demographic and Health Survey [Internet]. Harare; 2015. Available from: www.DHSprogram.com.
11. International Institute for Population Sciences. India National Family Health Survey [Internet]. Mumbai; 2015. Available from: <https://dhsprogram.com/pubs/pdf/frind3/frind3-vollandvol2.pdf>
12. Ministry of Health, Community Development, Gender E and C. Tanzania [Internet]. daresalam; 2015. Available from: <http://www.dhsprogram.com>.

- 415 13. National Institute of Statistics of Rwanda. Rwanda Demographic and Health Survey
416 [Internet]. Kigali; 2014. Available from: www.DHSprogram.com
- 417 14. Pakistan National Institute of Population Studies. Pakistan Demographic and Health Survey
418 [Internet]. Islamabad, Pakistan; 2012. Available from: <http://www.nips.org.pk>
- 419 15. Armenia National Statistical Service. Armenia Demographic and Health Survey [Internet].
420 Yerevan; 2015. Available from: www.DHSprogram.com
- 421 16. Nepal Ministry of Health. Nepal Demographic and Health Survey [Internet]. Ramshah Path;
422 2016. Available from: info@DHSprogram.com
- 423 17. Federal Democratic Republic of Ethiopia. Implementation Plan (2016 – 2030) [Internet].
424 2016. Available from: <https://www.imf.org/external/pubs/ft/scr/2004/cr0437.pdf>
- 425 18. UNICEF. C4D Communication for Development (C4D) Capability Development Framework
426 [Internet]. UNICEF; 2016. Available from: [http://intranet.unicef.org/pd/cbsc.nsf/Site](http://intranet.unicef.org/pd/cbsc.nsf/SitePages/Page01?OpenDocument&TableRow=3.2.3#3.2)
427 [Pages/Page01?OpenDocument&TableRow=3.2.3#3.2](http://intranet.unicef.org/pd/cbsc.nsf/SitePages/Page01?OpenDocument&TableRow=3.2.3#3.2)
- 428 19. Ng CS, Dibley MJ, Agho KE. Complementary feeding indicators and determinants of poor
429 feeding practices in Indonesia: a secondary analysis of 2007 Demographic and Health Survey
430 data. Public Health Nutr [Internet]. 2012;15(5). Available from:
431 http://www.journals.cambridge.org/abstract_S1368980011002485
- 432 20. Na M, Arimond M, Stewart CP. Risk factors of poor complementary feeding practices in
433 Pakistani children aged 6 – 23 months : A multilevel analysis of the Demographic and Health
434 Survey 2012 – 2013. Matern Child Nutr [Internet]. 2017;13(November 2016):1–12. Available
435 from: <https://www.ncbi.nlm.nih.gov/pubmed/29032630>
- 436 21. Ethiopia Health and Nutrition Research Institute and U, Suggested. IYCF practices, beliefs
437 and influences in Tigray region, Ethiopia. Alive & Thrive [Internet]. 2010;(September).
438 Available from: aliveandthrive@aed.org www.aliveandthrive.org September
- 439 22. Khanal V, Sauer K, Zhao Y, Black R, Cousens S, Johnson H, et al. Determinants of
440 complementary feeding practices among Nepalese children aged 6–23 months: findings from
441 demographic and health survey 2011. BMC Pediatr [Internet]. 2013;13(1):131.
- 442 23. Kabir I, Khanam M, Agho KE, Mihrshahi S, Dibley MJ, Roy SK. Determinants of
443 inappropriate complementary feeding practices in infant and young children in Bangladesh :
444 secondary data analysis of Demographic Health Survey 2007. Matern Child Nutr [Internet].
445 2012;8:11–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22168516>
- 446 24. Department of Community Medicine, Faculty of Medicine, University of Colombo, 25 K.
447 Complementary feeding practices in South Asia : Matern Child Nutr [Internet]. 2012;8(Who
448 2002):5–10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22168515>

- 449 25. Senarath U, Godakandage SSP, Jayawickrama H, Siriwardena I, Dibley MJ. Determinants of
450 inappropriate complementary feeding practices in young children in Sri Lanka : secondary data
451 analysis of Demographic and Health Survey 2006 – 2007. 2012;8:60–77. Available from:
452 <https://www.ncbi.nlm.nih.gov/pubmed/22168519>
- 453 26. Patel A, Pusdekar Y, Badhoniya N, Borkar J, Agho KE, Dibley MJ. Determinants of
454 inappropriate complementary feeding practices in young children in India : secondary analysis of
455 National Family Health Survey 2005 – 2006. 2012;8:28–44. Available from:
456 <https://www.ncbi.nlm.nih.gov/pubmed/22168517>
457

459

460