Experimental and mathematical approaches to quantify recirculation kinetics of lymphocytes

Vitaly V. Ganusov^{1*} and Michio Tomura²

¹Department of Microbiology, University of Tennessee, Knoxville, TN 37996, USA

²Immunology laboratory, Pharmaceutical Division, Osaka Ohtani University, Osaka, Japan

*Corresponding author: vitaly.ganusov@gmail.com

August 17, 2018

Abstract

One of the properties of the immune system that makes it different from nervous and endocrine systems of mammals is the ability of immune cells to migrate between different tissues. Lymphocytes such as T and B cells have the ability to migrate from the blood to secondary lymphoid tissues such as spleen, lymph nodes, and Peyer's patches, and then migrate back to the blood, i.e., they can recirculate. Recirculation of lymphocytes has been a subject of intensive investigation decades ago with wealth of data on the kinetics of lymphocyte recirculation available. However, these data have not been widely used to estimate the kinetics of recirculation of different lymphocyte subsets in naive and immunized animals. In this paper we review pioneering studies addressing the question of lymphocyte recirculation, overview quantitative approaches that have been used to estimate the kinetics of lymphocyte recirculation, and provide currently published estimates of the residence times of resting lymphocytes in secondary lymphoid tissues of mammals.

Keywords: lymphocyte migration, recirculation, T cells, B cells, dwell time, residence time, mathematical modeling

Abbreviations: LNs: lymph nodes, iLNs: inguinal lymph nodes, mLNs: mesenteric lymph nodes, TDLs: thoracic duct lymphocytes, EEL: efferent lymph lymphocytes, LCMV: lymphocytic choriomeningitis virus, PPs: Peyer's patches.

Introduction

Adaptive immune system of mammals includes two major subsets of lymphocytes, B and T lymphocytes [1, 2]. One of the major functions of the adaptive immune system is to protect its host from invading microorganisms such as viruses and bacteria. Because microbes can enter the host from multiple areas such as via skin, lung or gut mucosa, there is a need to have lymphocyte being present near these tissues. In addition, because every lymphocyte is in general specific to only one microbial determinant (epitope) and there are many microbial determinants, only a small percent of lymphocytes would be able to recognize any specific microbe. For example, recent estimates suggest that only 1 in $10^5 - 10^6$ of T lymphocytes would recognize a given epitope; that is in a mouse that has $\sim 2 \times 10^8$ lymphocytes only about 200-2000 T cells would be specific to a given epitope [3]. While pathogens in general have multiple epitopes, only few of those can be recognized by lymphocytes. For example, in B6 mice lymphocytic choriomeningitis virus (LCMV) has about 30 epitopes that are recognized by CD8 T cells at all and only a few recognized strongly [4]. It is probably impossible to put a few thousands of T cells in all potential places of entry of a pathogen. Instead, lymphocytes have the ability to recirculate between different tissues in the body, thus increasing chances of encountering antigen they are specific for. This ability to migrate from the blood into several specific tissues and then back to the blood (i.e., to recirculate) makes adaptive immune system different from several other major systems of mammals such as nervous and endocrine systems. Because lymphocyte recirculation is a fundamental property of the mammalian immune system, our knowledge of how immune system works would be greatly incomplete if do not have solid understanding of the kinetics of lymphocytes recirculation, i.e., how quickly lymphocytes migrate to peripheral tissues, how long they spend in the tissues and return back to circulation. Understanding lymphocyte recirculation kinetics may be not just an academic exercise as blocking lymphocyte migration by anti-VLA4 antibodies – VLA4 is an integrin regulating lymphocyte entry into several tissues – has been shown to be effective in reducing symptoms of multiple sclerosis [5]. However, such treatment has serious side effects suggesting that deeper understanding how lymphocyte migration is regulated is needed [6, 7].

Ability of lymphocytes to recirculate between blood and tissues depends strongly on the type of lymphocyte, type of the tissue, and conditions of the host [8–18]. Specifically, naive T cells — cells that have not yet encountered their cognate antigen — are able to recirculate between blood and secondary lymphoid organs such as lymph nodes, spleen, and Peyer’s patches [9, 14, 19]. Activated T cells have the ability to migrate to nonlymphoid tissues [14, 20]; however, whether activated T cells in nonlymphoid tissues can migrate back to the blood remains poorly understood [21]. Inflammation may also change the pattern of lymphocyte migration; for example, intravenous (i.v.) infection may lead to trapping of recirculating lymphocytes in the spleen [22]. In this review we will focus on aspects of recirculation of resting (naive and memory) T lymphocytes with the major focus on migration of these cells via secondary lymphoid tissues, and how mathematical modeling has helped so far to quantify kinetics of this recirculation. Our main focus on recirculation of resting T cells is due to lack of good quantitative data and mathematical models on recirculation of activated T and B cells. However, we will provide a novel analysis of older data on recirculation kinetics of activated T cells in mice.

Because this review is about lymphocyte recirculation it is important to outline some basic anatomical features of the mammalian immune system. Since mice are the smallest mammalian animal model used to study lymphocyte recirculation, we focus our description specifically on murine secondary lymphoid organs. The major secondary lymphoid organs of mice are lymph nodes (LNs), spleen, and Peyer’s patches (PPs). Fluids that leak out of blood vessels are collected by the lymph

phatic vessels which bring this interstitial fluid via afferent lymphatics to tissue-draining lymph nodes. Each lymph node drains fluids from specific tissues and fluids (lymph) exit lymph nodes via efferent lymphatics often into another lymph node [23, 24]. Lymph from the final lymph nodes is collected into two big vessels, left and right lymphatic ducts, which are connected to the blood and which return collected interstitial fluids and cells back to circulation [2]. In mice and humans, the right lymphatic duct collects lymph from the upper right part of the body (about 1/4 of all interstitial fluid) and left lymphatic duct (also called thoracic duct) collects lymph from the rest (about 3/4) of the body. Lymph nodes can be roughly divided into several groups such as skin-draining lymph nodes, lung-draining lymph nodes, and gut-draining lymph nodes. A typical laboratory mice strain has about 30 lymph nodes [23] while humans have hundreds (perhaps over a thousand) lymph nodes [24–26]. Peyer’s patches are lymph node-like structures found in the gut. Peyer’s patches do not have afferent lymphatics and efferent lymph from Peyer’s patches flows into mesenteric (gut-draining) lymph nodes. Finally, spleen is probably the largest single secondary lymphoid organ in mice and humans [25, 27–31]. Spleen is not connected directly to the lymphatic system and lymphocytes enter the spleen from the blood and exit the spleen into the blood. In contrast, lymphocytes may enter lymph nodes or Peyer’s patches from the blood via high endothelial venules, and lymphocyte may also enter lymph nodes by migrating from the blood to peripheral tissues such as skin or gut, and then enter lymph nodes with afferent lymph. Thus, lymphocytes have several different pathways for recirculation in the body.

Mathematical modeling of lymphocyte recirculation

By definition recirculation of lymphocytes is a dynamic process and therefore mathematical modeling is likely to be a useful tool for understanding of lymphocyte recirculation. Mathematical modeling is required to accurately quantify the kinetics of lymphocyte recirculation and to estimate the rates of lymphocyte migration from the blood to tissues and lymphocyte residence times in tissues. There have been many uses of mathematical models to understand cell migration. For example, mathematical models have been used to gain insights of how lymphocytes move in lymphoid and nonlymphoid tissues and how tissue composition impacts movement patterns of T cells [32, 33]. Here our main focus will be on experimental studies providing quantitative data on lymphocyte migration between tissues, and on mathematical modeling attempts to quantify lymphocyte recirculation kinetics (Table 1).

Kinetic aspects of lymphocyte recirculation has been studied since 1950th using multiple mammalian species such as mice, rats, sheep, and pigs (see more below). Prior to pioneering experiments by Gowans the role of small lymphocytes found in the blood was unknown. Collecting lymphocytes from the thoracic duct lymph of rats over several days led to a decline in the number of lymphocytes found in the lymph. However, return of cells collected by thoracic duct cannulation back into circulation prevented loss of lymphocytes from the lymph [34]. These key experiments thus established that lymphocytes are able to recirculate between blood and thoracic duct lymph. By labeling lymphocytes collected from the blood or lymph (e.g., by thoracic duct (rats) or individual lymph node (sheep, pigs) cannulation) with radioactive labels further experiments demonstrated that indeed lymphocytes migrate from the blood to the efferent lymphatics of lymph nodes [8, 35–40].

Spleen

Spleen is a large secondary lymphoid organ and it was previously estimated that about 20% of all lymphocytes in a human are found in the spleen [25, 31]. In mice, around 50% of all lymphocytes from secondary lymphoid tissues are found in the spleen [41, 42]. While it has been estimated that many lymphocytes travel via the spleen of mice, rats, or pigs [28, 30, 43] the exact time lymphocytes spend in the spleen has not been accurately quantified. Previous studies documented accumulation of radioactively labeled lymphocytes in the spleen after i.v. infusion of such cells [43–45] or dynamics of labeled cells in the blood in normal or splenectomized pigs [29]. However, data from these studies have not been analyzed using mathematical models, and thus, these previous data did not lead to estimates of lymphocyte residence (or dwell) times in the spleen.

An interesting approach was taken by Ford who designed an apparatus allowing maintenance of viable rat spleens for an extended period of time (up to 10 days, [46]). By creating an artificial circulation system connecting the spleen’s blood vessels, the author could monitor concentration of lymphocytes that exit the spleen during the perfusion of labeled thoracic duct lymphocytes that had been injected into circulation. Similar experiments with isolated pig spleens was done later by another group [28]. Verbal analysis of the data on migration thoracic duct lymphocytes via isolated perfused spleen led to estimate of lymphocyte residence time in the rat spleen of 4-5 hours [46, 47] and in the pig spleen of 2-4 hours [28]. A relatively complex mathematical model was proposed and fitted to the data of Ford [46]. This mathematical model-based analysis predicted that only about 10-25% of lymphocytes migrating via spleen pass via the marginal zone of the spleen (i.e., enter the spleen parenchyma). Cells entering the marginal zone of the spleen had a residency time of 50 minutes in the tissue. About 10% of cells existing the marginal zone migrated to the white pulp where the residency time was 4.6 hours. In contrast, remaining 90% of lymphocytes exited marginal zone into the red pulp with the average residency time of 5 minutes in that subcompartment of the spleen [48]. While Hammond [48] did not calculate the average time lymphocytes spend in the spleen, given the estimates provided, the average residency time of TDLs in the spleen is $50 + 0.1 \times 4.6 \times 60 + 0.9 \times 5 = 82$ min or 1.4 hours. This is a significantly shorter residency time than concluded by Ford [46] by visual analysis of the data.

Ganusov and Auerbach [49] used another set of data on migration kinetics of thoracic duct lymphocytes (TDLs) in rats. In these experiments (see also below), lymphocytes collected via thoracic duct cannulation were transferred into syngenic rats and the accumulation and loss of transferred cells (labeled with a radioactive label) were measured in multiple tissues of rats [50]. By fitting a mathematical model to these experimental data, Ganusov and Auerbach [49] estimated recirculation kinetics of TDLs including residence of these cells in major secondary lymphoid tissues of rats. The model fits predicted that average residence time of TDLs in the spleen is 2.4 hours which is also shorter than a previous estimate given in Ford [46] but longer than the estimate obtained from parameters of Hammond [48].

Lymph nodes

Migration of lymphocytes from blood to lymph nodes and then back to the blood has been the main focus of many studies on lymphocyte recirculation. In part, this stems from the fact that in small animals such as rats cells that had migrated via lymph nodes can be easily collected via the thoracic duct cannulation, and, thus, the tempo of lymphocyte movement from the blood to thoracic duct

can be easily recorded. In larger animals an efferent lymphatic vessel exiting a given lymph is wide enough to allow cannulation and collection of cells exiting that specific lymph node.

As described above, following the classical studies by Gowans illustrating ability of lymphocytes to recirculate between blood and thoracic duct lymph, there have been multiple studies investigating details of lymphocyte migration from blood to efferent lymph (reviewed in [30, 51–53]). Transit times via a particular lymph node (e.g., inguinal or cervical) have been inferred by transferring labeled lymphocytes into the animal and measuring rate of exit of labeled lymphocytes via a cannulated node in large animals such as sheep and pigs (e.g., [8, 9, 16, 54–57, Figure 1A]). Alternatively, kinetics of lymphocyte migration from the blood to the thoracic duct (i.e., via the whole lymphatic system) have been measured in smaller animals such as mice and rats (e.g., [36, 50, 58–65, Figure 1B]).

Figure 1: Typical examples of experimental data on lymphocyte migration from blood to lymph. In such studies, lymphocytes were isolated from a tissue (e.g., efferent lymph, lymph nodes, blood, spleen) and labeled with a label. In older studies labels were radioactive while newer studies involved fluorescent labels. Then cells were transferred intravenously into the same (e.g., sheep, panel A) or syngenic (e.g., rat, panel B) host. The accumulation of labeled cells in the efferent lymph of a given lymph node (panel A) or in the thoracic duct (panel B) were then followed. Different measures of the number of labeled cells have been used such as the number of labeled cells per fixed number of total isolated cells (e.g., estimated by measuring counts-per-minute (cpm) from the sample, panel A) or as the number or percent of injected cells (measured by total radioactivity) per unit of time (panel B). Data were digitized from previous publications (panel A: Frost et al. [54], panel B: Smith and Ford [50]).

While the data on the migration kinetics of lymphocytes via individual lymph nodes or from the blood to thoracic duct have been collected (e.g., Figure 1) very few studies attempted to estimate the lymphocyte residence (or dwell) time in lymph nodes from these cannulation experiments. For example, by simply looking at the data it is unclear which characteristic of the distribution observed in Figure 1 represents the average residence time. Mode, median, and average could all potentially be good estimates of the average residency time, and intuitively the average of the distribution has been treated in experimental studies as an estimate for residence time of lymphocytes in lymph nodes, e.g., about 48 hours in sheep or 24 hours in rats [30, 58]. However, to accurately estimate the average residence time one need to use mathematical modeling that takes physiology of the recirculatory and lymphatic system of mammals into account.

As far as we know the first mathematical modeling-based attempt to quantify lymphocyte migration via lymph nodes was in a series of papers by Stekel et al. [66–68]. The main idea of the mathematical model considered in these papers was the ability of lymphocytes to attach to and

deattach from the lymphoid tissues while in lymph nodes or the spleen [66]. Deattached cells move through the tissue and this “movement” was described by a transport equation. Attached cells, however, would not move, and thus the process of “attachment – detachment” generated a skewed distribution and matched data on thoracic duct cannulation in rats [66, 67]. This work suggested 20h residency time of lymphocytes in the lymph nodes and 6 h residency time in the spleen of rats [66]. The model was further used to explain different kinetics of lymphocyte migration in irradiated or thymectomized rats [67, 68]. It is now well understood that the basic assumption of the model that cell attachment to structures in lymph nodes prevent lymphocyte exit is incorrect; rather, lymphocyte use structures including fibroblastic reticular cells to move around and to exit lymph nodes although precise mechanisms regulating lymphocyte exit from lymph nodes remain to be fully defined [69]. By using intravital imaging of T lymphocytes moving in murine lymph nodes and by mathematical modeling of T cell movement in the nodes Grigorova et al. [69] estimated the half-life time of T cells in the lymph nodes of mice to be 4-5 hours. Another study analyzed importance of directional movement of T cells in lymph nodes using digital reconstruction of a rat lymph node but the actual residency times of lymphocytes in the nodes was not estimated [70].

An important study quantified the residence time of antigen-specific naive and memory CD8 T cells using a novel “transfer-and-block” technique [71]. In this approach, naive or memory CD8 T cells, specific to the GP33 epitope of LCMV were transferred into congenic hosts and 24 hours after cell transfer further entry of lymphocytes into LNs was blocked by using anti-CD62L antibodies. CD62L is expressed on T cells and is generally required for cell entry into LNs [72]. The declining number of LCMV-specific T cells remaining in the LNs at different times after the blockade was used to infer the lymphocyte residence time. Interestingly, the authors found a non-monotonic rate of loss of T cells from LNs; naive and memory CD8 T cell populations had initial residency times of 5-6 h in LNs which increased significantly at later times for both naive and memory T cell populations to 15-16 hours [71]. Re-analysis of these data in another study revealed that the data could be accurately explained by a model in which residency time of lymphocytes is density-dependent and declines with time since blockade (if blockade was 100% efficient) [49]. In this reanalysis, naive and memory CD8 T cells had different residency times (16 and 9 hours for naive and memory T cells, respectively) [49]. Thus, the use of the same data but different assumptions on lymphocyte migration may result in different estimates of lymphocyte recirculation kinetics.

Mandl et al. [73] extended the study of Harp et al. [71] by transferring polyclonal naive CD4 and CD8 T cells into congenic mice and by blocking entry of new cells 2 hours after cell transfer by using a combination of antibodies to CD62L and VLA4. The authors found that a residence time of naive T cells was dependent on the type of the cell (CD4 vs. CD8 T cell). In contrast with previous work [71] the authors observed that after the blockade the percent of transferred cells declined exponentially over time. By fitting a line to log-transformed cell frequencies the authors estimated that naive CD4 and CD8 T cells spend on average 12 and 21 hours, respectively, in lymph nodes in mice [73].

It is interesting to note that we only know of one study that estimated residence time of lymphocytes from data generated by cannulating individual lymph nodes in sheep [74]. The authors proposed that lymphocyte migration within a lymph node can be described as a Markov process with n states and probability of jump from one state to another forward ($i \rightarrow i + 1$) or backward ($i \rightarrow i - 1$). Reaching the n^{th} state implied exit of a lymphocyte from the lymph node. The model was fitted to the cannulation data similar to that in Figure 1A and predicted the average residence time of lymphocytes in sheep lymph nodes of 31 hour [74]. Using a different mathematical model, which incorporated lymphocyte recirculation kinetics in the whole body, we have analyzed similar data on lymphocyte migration via individual lymph nodes in sheep (McDaniel and Ganusov (in preparation)).

216 Depending on the dataset we found the average residence time of blood-derived lymphocytes in ovine
 217 lymph nodes to be 18-22 hours. This is another demonstration that the estimate of the lymphocyte
 218 residence time may not be robust to the choice of a model [75].

219 We recently performed mathematical modeling-assisted analysis of experimental data on recir-
 220 culation of thoracic duct lymphocytes (TDLs) in rats [49]. In these experiments [50], lymphocytes
 221 collected in rats via thoracic duct cannulation were transferred into a series of syngenic hosts and
 222 accumulation and loss of the transferred cells in multiple lymphoid tissues was followed over time.
 223 By fitting a series of mathematical models to experimental data we estimated the TDL residence
 224 time in multiple tissues including lymph nodes. Interestingly, in contrast with previous studies that
 225 found differences in T lymphocyte residence times in different lymph nodes (e.g., mesenteric vs. sub-
 226 cutaneous LNs) [71, 73] we found the average residence time in subcutaneous or mesenteric LNs or
 227 PPs to be 10 hours [49, see more below].

Figure 2: Schematic of the mathematical model describing loss of photoconverted cells from a LN of Kaede mice. We assume that cells entering a LN must undergo k exponentially distributed transitions before they can exit the node. In essence, this assumes that residency time of lymphocytes in a lymph node follows a gamma distribution. In the model X_i is the number of lymphocytes residing in the i^{th} compartment and m is the rate of transition of cells between subcompartments (see also eqns. (1)–(2)). Cells exiting the last subcompartment exit the lymph node into efferent lymph.

228 A novel experimental technique, the Kaede mice, allows to accurately track exit of lymphocytes
 229 from a given tissue such skin or a lymph node [17, 76]. Kaede mice express a photoconvertable
 230 protein which upon exposure to violet light changes color from green to red [76]. This unique system
 231 allows to label cells in one location, for example, an inguinal lymph node (iLN) or skin, and track the
 232 movement of labeled cells to other tissues in the body [17, 77]. We developed a simple mathematical
 233 model to track the dynamics of photoconverted (red) lymphocytes in the iLNs of Kaede (or other
 234 photoconvertable mice, e.g., KikGR [78, 79]) mice. The model assumes that cells exit the LN and are
 235 not able to re-enter the same LN. Experiments have shown that labeling of cells in the iLN distribute
 236 between all LNs and the spleen with approximately of 2-3% of cells in LNs being red [76]. Therefore,
 237 until the percent of photoconverted cells in the iLN is above 8-10%, re-entry of such cells into the
 238 node can likely be neglected if we assume that photoconversion and surgery associated with it do not
 239 induce strong inflammation impacting cell migration. In experiments, the lack of inflammation was
 240 recorded by similar size of iLNs prior and after the photoconversion.

241 Because our previous study suggested that distribution of residence times of TDLs in LNs was
 242 best described by a gamma distribution (and not by exponential distribution) [49], for experiments
 243 with Kaede we describe exit of lymphocytes from a LN as cell “migration” via multiple (k) subcom-
 244 partments in the LN:

$$\frac{dx_1(t)}{dt} = -mx_1(t), \quad (1)$$

$$\frac{dx_i(t)}{dt} = m(x_{i-1}(t) - x_i(t)), \quad i = 2 \dots k, \quad (2)$$

where m is the migration rate via a given subcompartment. The average residence time of lymphocytes in the lymph nodes is then given by $T = k/m$. Given that following photoconversion cells in all subcompartments at the steady state have equal densities, $x_i(0) = 1/k$, the mathematical model (eqns. (1)–(2)) has a unique solution for the total number of cells in the LN $x(t) = \sum_{i=1}^k x_i(t)$:

$$x(t, m) = e^{-mt} \sum_{i=1}^k \frac{(mt)^{i-1}}{(i-1)!}. \quad (3)$$

By fitting this solution (eqn. (3)) to experimental data from photoconversion experiments (Figure 3) we found that this simple model describes well the data for the loss of photoconverted CD8 T cells and B cells from the iLN (Figure 3B&C). However, the model was inadequate at describing the data for CD4 T cells as judged by the lack of fit test [80, results not shown]. Visually, this is likely because the loss of photoconverted CD4 T cells from the iLN is not exponential (compare Figure 3A & 3B). These results were independent of the number of subcompartments tested. While it is clear that CD4 and CD8 T cell populations most likely consist of subpopulations with perhaps different rates of exit from the iLNs, e.g., naive and memory T cells, why we were not able to detect such heterogeneity for CD8 T cells was unclear. It is possible that heterogeneity in CD4 T cells may come from a more diverse sets of cell types present in this group, for example, naive, memory, and regulatory T cells. Previously it was noted that naive and memory phenotype CD4 T cells have different exit kinetics from LNs [17]. In contrast, LCMV-specific naive and memory CD8 T cells appear to exit iLNs with similar kinetics [71].

Figure 3: Estimated residence time of lymphocytes depends on the shape of residence time distribution. Lymphocytes in inguinal lymph nodes (iLNs) of Kaede mice were photoconverted [76] and the percent of CD4 T cells (panel A), CD8 T cells (panel B), or B cells (panel C) remaining in the iLNs at different times after photoconversion was recorded; the data from individual mice are shown by dots and horizontal lines denote average percent per time point. We fit a series of mathematical models assuming one (eqn. (3)) or two (eqn. (4)) subpopulations of cells with a different number of subcompartments ($k = 1 \dots 4$) to these data. Fits of the models with 2 subpopulations are shown by lines. To normalize residuals we used $\log_{10}$ transformation. The model with two cell subpopulations only improved the fit of the data for CD4 T cells (F-test, $p < 0.005$). Data for CD8 T and B cells were well described by a model with one, homogenous population (F-test, $p > 0.05$). Parameters of the model with 2 subpopulations and $k = 2$ compartments and their 95% confidence intervals (found by bootstrapping residuals with 1000 simulations) for different lymphocyte types are: CD4 T cells: $m_1 = 0.30$ (0.30, 0.31)/h, $m_2 = 0.066$ (0.065, 0.067)/h, $f = 0.53$ (0.52, 0.53), $T = 17.9$ (17.9, 18.0) h; for CD8 T cells: $m_1 = 0.46$ (0.43, 0.48)/h, $m_2 = 0.10$ (0.10, 0.10)/h, $f = 0.16$ (0.15, 0.15), $T = 17.9$ (17.9, 18.0) h; B cells: $m_1 = 0.071$ (0.071, 0.071)/h, $m_2 = 0.07$ (0.07, 0.07)/h, $f = 0.37$ (0.37, 0.37), $T = 28.2$ (28.2, 28.2) h.

To more accurately describe the kinetics of loss of photoconverted CD4 T cells from iLNs we extended the simple model (eqn. (3)) by allowing 2 subpopulations with relative frequencies f and

1 – f and with different exit kinetics determined by the rates m_1 and m_2 , respectively. In this model, the total number of photoconverted (red) cells in the iLN is then given by

$$X(t) = fx(t, m_1) + (1 - f)x(t, m_2), \quad (4)$$

where $x(t, m_i)$ is given in eqn. (3). It is straightforward to extend this model to n subpopulations. The average residence time in this model is defined as $T = fk/m_1 + (1 - f)k/m_2$ where k is the number of subcompartments in each of the subpopulations. This 2 subpopulation model can well describe experimental data on the loss of photoconverted CD4 T cells from iLNs (Figure 3A). Interestingly, for models that fit the data well (e.g., data for CD8 T cells or B cells) the estimated average residence time was not strongly dependent on the number of subpopulations assumed (1 or 2 subpopulations, results not shown). However, the estimate of the average residence time was strongly dependent on the number of subcompartments assumed, and, as the consequences, on the shape of the distribution of residence times. For example, the model fits predicted average residence time for CD8 T cells to be $T = 13.3$ h for $k = 1$ or $T = 22.9$ for $k = 5$ (eqn. (3)) with moderate reduction in the quality of the model fit to data at higher k as judged by AIC (results not shown). Therefore, it appears that the estimate of the average residence time from photoconversion data is not fully robust. Given our previous observation that best description of TDL recirculation kinetics via LNs in rats is given by a gamma distribution with shape parameter $k = 2$ our results suggest that average residence times in mouse iLNs are 18 h for CD4 and CD8 T cells, and 28 h for B cells.

Whole body recirculation kinetics

An important limitation of many of the previously listed studies is that they considered migration of lymphocytes only via individual secondary lymphoid tissues such as spleen or individual lymph nodes. To study lymphocyte migration in the whole organism, Smith and Ford [50] adoptively transferred ^{51}Cr -labeled TDLs and measured the percent of transferred lymphocytes in different organs of the recipient rats including the blood, lung, liver, spleen, skin-draining (subcutaneous) and gut-draining (mesenteric) lymph nodes. These data were initially analyzed with the use of a mathematical model [81] but TDL residence times in different tissues were not estimated. We developed a simple yet large mathematical model describing TDL dynamics, and by fitting the model to Smith and Ford [50] data, for the first time estimated the kinetics of TDL recirculation in the whole body [49, Figure 4].

The model fits predicted that TDLs spend very short time in the main blood vessels (about 30 sec) after which the vast majority of lymphocytes (about 95%) gets trapped in vasculature of the lung or the liver. This trapping is short-lived, however, and within 1 min trapped lymphocytes re-enter circulation (Figure 4). Only 5% of lymphocyte enter secondary lymphoid tissues per one passage of lymphocytes via circulatory system, with half of these entering the spleen, and half entering lymph nodes and Peyer's patches (PPs). Lymphocytes reside for 10 h in LNs/PPs but only for 2.5 h in the spleen (Figure 4).

Because there is a good understanding of the kinetics of blood recirculation in rats, we would like to provide another interpretation of the kinetics at which TDLs pass via major tissues in rats. Previous studies found that the total blood volume in rats is proportional to the rat weight [82], and for 300 g rats, blood volume is $V_b \approx 20$ mL. Heart volume is dependent on the animal size and age, and for 6-10 week old rats, $V_h = 0.5$ mL [83]. Given the high heart rate in rats (462/min)

Figure 4: Estimated residence times of thoracic duct lymphocytes (TDLs) in rats [49]. We developed a mathematical model describing migration of lymphocytes in the whole organisms and by fitting the model to experimental data [50], estimated TDL residence times in major nonlymphoid and all major secondary lymphoid tissues of rats. Predicted residence times were short in the blood, lung and liver ($T \leq 1\text{min}$) with 95% of lymphocytes recirculating between these tissues. Residence time in the spleen was 2.5 h with about 2.5% of lymphocytes entering the tissue. The remaining $\sim 2\%$ of cells were entering lymph nodes and Peyer's patches (PPs), with the average residence time of TDLs in these tissues was 10 hours [49].

and stroke volume 0.3 mL per heart beat, the total cardiac output in rats is approximately $c = 140$ mL/min [84]. This in turn suggests the rate of blood recirculation in rats of $m_0 = c/V_b = 7/\text{min}$ or residency time of 9 seconds (i.e., on average in 9 seconds all blood passes through the heart). Given previously estimated rates at while TDLs enter various tissues [49], we predict that per one circulation of whole rat blood 27% and 6% of TDLs are attaching to the lung and liver vasculature, respectively. This suggests that the majority of TDLs pass via lung and liver vasculatures without attachment! Importantly, only 0.8% of TDLs in the blood will migrate to the spleen and 0.6% will migrate to LNs and PPs in one blood recirculation cycle lasting 9 seconds suggesting that the process of entering secondary lymphoid tissues by recirculating TDLs is not very efficient.

Differential probability of lymphocyte migration via lung and liver vs. secondary lymphoid organs is interesting but perhaps not unexpected given that these two organs are large and are expected to collect large volumes of blood. Amount of blood going to any specific tissue (cardiac output) has been measured in multiple species, for example, by measuring accumulation of labeled small microspheres after injection into the blood [40, 56, 84]. While we did not find a single study in rats measuring cardiac output to the same tissues as in our analysis (Figure 4), we found the cardiac output does not accurately predict the hierarchy of lymphocyte entry into lung, liver, and spleen. In rats, 0.7%, 3.3%, and 0.6% of cardiac output goes to lung, liver, and spleen, respectively [84], which is in contrast to 78%, 17%, and 2.5% of lymphocyte entry probability for these tissues [49]. Thus, migration and retention of lymphocytes in the whole body, while likely is dependent on the blood flow, is not strictly determined by the amount of blood going to any specific tissue.

Lymphocyte type	Tissue/organ	Animal species	Residency time T , h	Reference
TDL	Spleen	Rats	4.5	[46]
TDL	Spleen	Rats	6.0	[66]
TDL	Spleen	Rats	1.4	[48]
TDL	Spleen	Rats	2.4	[49]
Blood lymphocytes	Spleen	Pigs	3.0	[28]
Naive T cells	Lymph nodes	Mice	4.5	[69]
naive CD8 T cells	Lymph nodes	Mice	6.0	[71]
memory CD8 T cells	Lymph nodes	Mice	6.0	[71]
naive CD8 T cells	Lymph nodes	Mice	16.0	[49]
memory CD8 T cells	Lymph nodes	Mice	9.0	[49]
naive CD4 T cells	Lymph nodes	Mice	12.0	[73]
naive CD8 T cells	Lymph nodes	Mice	21.0	[73]
CD4 T cells	Lymph nodes	Mice	18.0	this work
CD8 T cells	Lymph nodes	Mice	18.0	this work
B cells	Lymph nodes	Mice	28.0	this work
TDL	Lymph nodes	Rats	20.0	[66]
TDL	Lymph nodes	Rats	10.0	[49]
ELL	Lymph nodes	Sheep	31.0	[74]
TDL	Peyer's patches	Rats	10.0	[49]

Table 1: Summary of published estimates of residency times of resting lymphocytes in secondary lymphoid tissues such as spleen, lymph nodes, and Peyer's patches. Here ELL are efferent lymph lymphocytes (lymphocytes isolated by cannulation of individual lymph nodes, e.g., in sheep), TDL are thoracic duct lymphocytes (lymphocytes isolated by thoracic duct cannulation). In most other cases lymphocytes were isolated from lymph nodes and/spleen. Listed values for the lymphocyte residency time are as has been reported by authors and in some cases, half-life time of lymphocytes in the tissue ($T_{1/2}$) was converted to the residency time using formula $T = \ln 2 \times T_{1/2}$. In some studies, estimates for the residence time of lymphocytes in lymph nodes were dependent on the lymph node type (e.g., [73]), therefore, the presented estimates are for the pooled data. For time-dependent residency times the initial value was used (e.g., [71]).

Recirculation of activated lymphocytes in mice

The vast majority of previous studies focused on quantifying migration of naive and memory lymphocytes via secondary lymphoid organs. While such lymphocytes are likely to represent the majority of cells in an organism in the absence of infection, infections will result in activation of lymphocytes. Yet, pattern and kinetics of migration of activated lymphocytes remain poorly defined. Cancer immunotherapy, involving *in vitro* expansion of populations of cancer-specific CD8 T cells and transfer of these cells into patients, is one of the novel ways to treat patients [85–87]. Therefore, deeper understanding of migration kinetics of activated T cells may help to improve the efficacy of T cell-based cancer therapies.

To determine the pattern and kinetics of recirculation of activated T lymphocytes we analyzed data from an old set of experiments [61]. In these experiments, Sprent [61] injected thymocytes (cells from the thymus) from CBA (H-2^k) mice into irradiated CBA $\times$ C57Bl/6 (H-2^k $\times$ H-2^b) F₁ mice and isolated activated T cells via the thoracic duct cannulation [60]. Activated T cells were specific to the H-2^b antigen of the donor. Collected T cells were labeled with ¹²⁵IUdR *in vitro* after 1 hour incubation and then injected intravenously into a series of syngenic CBA mice [61]. ¹²⁵IUdR is incorporated into

Figure 5: Schematic of assumed recirculation pathways of activated T lymphocytes [61]. In these experiments, lymphocytes were injected into the blood (B) and from the blood lymphocytes may enter lung ($i = 2$), liver ($i = 3$), spleen (S , $i = 4$), mesenteric LNs (mLNs, $i = 5$), intestine (I , $i = 6$) at rates m_{1i} with $i = 2, \dots, 6$, respectively. Lymphocytes in these tissues may return to circulation at rates m_{i1} with $i = 2, \dots, 6$. Lymphocytes may also leave the blood to other unsampled compartments and/or die at rate d (see eqns. (5)–(8)).

newly synthesized DNA, and therefore, only lymphocytes that were actively dividing *in vitro* became labeled.

Following adoptive transfer recipient mice were sacrificed at different times after cell transfer and the percent of labeled lymphocytes was measured in several major organs of mice including blood, lung, liver, spleen, mesenteric lymph nodes (mLNs), thymus, kidney, and intestine [61, Figure 6]. Because very few cells migrated to thymus and kidney, we ignored these tissues in our following analysis; inclusion of these tissues did not influence significantly estimates of other parameters (results not shown).

To estimate the rates of activated T cell migration to major tissues of mice we adopted a mathematical model from our previous study [49]. In this model we assume that lymphocytes in the blood can migrate to multiple tissues such as lung, liver, spleen, , and intestine, and following passage via the tissue, the cells would return back to the blood. The rate of lymphocyte entry into i^{th} tissue from the blood is denoted as m_{1i} and the rate of exit from the i^{th} tissue into the blood is then m_{i1} where $i = 2, \dots, 6$. Following our previous work and some initial analyses in the model we assume that T cell migration via the lung and liver follows 1st order kinetics (i.e., is described by an exponential distribution), but residence in the spleen, mLNs and intestine is gamma-distributed. Gamma distribution of T cell residence in these tissues was modelled by assuming k subcompartments with migration rate m_{i1} between subcompartments. With these assumptions the mathematical model is given by a set of differential equations:

$$\frac{dx_1}{dt} = -x_1 \left(d + \sum_{i=2}^6 m_{1i} \right) + \sum_{i=2}^3 m_{i1} x_i + \sum_{i=4}^6 m_{i1} x_{ik}, \quad (5)$$

$$\frac{dx_i}{dt} = m_{1i} x_1 - m_{i1} x_i, \quad i = 2, 3, \quad (6)$$

$$\frac{dx_{i1}}{dt} = m_{1i} x_1 - m_{i1} x_{i1}, \quad i = 4, 5, 6, \quad (7)$$

$$\frac{dx_{ij}}{dt} = m_{i1} x_{ij-1} - m_{i1} x_{ij}, \quad i = 4, 5, 6, \quad j = 2, \dots, k, \quad (8)$$

where x_i is the percent of labeled cells found in the blood ($i = 1$), lung ($i = 2$), liver ($i = 3$), and x_{ij} is the percent of labeled cells found in the j^{th} sub-compartment of spleen ($i = 4$), mesenteric LNs ($i = 5$), or intestine ($i = 6$), and $j = 1 \dots k$, d as the rate of removal of lymphocytes from circulation (due to death or migration to unsampled tissues such as other lymph nodes). Note that in this model we assume that cells migrating to the intestine return directly back to circulation without migrating via afferent lymph to mLNs. This assumption was justified by the lack of accumulation of labeled cells in the mLNs (Figure 6).

Figure 6: Experimental data and predictions of a mathematical model on the migration kinetics of *in vivo* activated T lymphocytes in mice. Thymocytes from CBA mice were transferred i.v. into CBA×C57BL/6 mice and activated TDLs were collected 4 to 5 days after cell transfer from the recipient mice. Replicating T cells were labeled with $^{125}\text{IUdR}$, adoptively transferred into CBA mice, and the distribution of transferred cells in different murine organs was followed over time (see Sprent [61] for more experimental detail). The percent of labeled lymphocytes recovered from different organs of the recipient mice are shown by markers. We fit the mathematical model of lymphocyte recirculation (eqns. (5)–(8)) to these experimental data; model fits are shown as lines. Plots are shown on the linear (panel A) or a log-scale (panel B). Parameter estimates are given in Table 2.

The model was fit to experimental data using least squares. A number of interesting observations emerged. First, the model predicted a very short average residence time of activated lymphocytes in the blood, $T \sim 1.2$ min (Table 2). Nearly 65% of activated lymphocytes in the blood migrated to the lung and liver where they spent on average 35 min and 2.2 hours, respectively. These residence times are substantially higher than those for resting TDLs [49, Figure 4]. It is interesting to note that trapping of activated lymphocytes in the liver after the i.v. injection has been observed previously in other experiments [22]. The spleen and intestine took another 20% of activated lymphocytes in the blood, and the average residence times were 8 and 22 hours in the spleen and intestine, respectively

(Table 2). In these experiments very few cells migrated to the mesenteric lymph nodes and we could not reliably estimate the residence times of these *in vivo* activated lymphocytes in the mLNs (e.g., note large confidence intervals in Table 2 for this tissue compartment). We also found that about 15% of transferred cells in the blood per circulation cycle migrated to tissues/organs which have not been sampled in these experiments, e.g., skin or other lymph nodes, or these cells have been dying at a high rate. Indeed, it is expected that activated lymphocytes undergo programmed cell death following clearance of the antigen which has been mimicked by the adoptive transfer experiments. In 24 hours post-transfer, only 34% of injected radioactivity could be recovered from recipient mice [61, Figure 6 and results not shown]. Nevertheless, this analysis highlights a similar hierarchy of migration of naive and activated lymphocytes *in vivo* with the majority of cells entering lung and liver vasculature but residing there for a relatively short period of time in these organs as compared to other tissues.

Organ	Rate of entrance from blood m_{1i} , h^{-1}	Percent cells going to organ from blood	Rate of exit from organ to blood m_{i1} , h^{-1}	Residence time in organ, h
Lung	23.64 (13.3–44.3)	46.1 (37.1–61.6)	1.7 (1.0–3.6)	0.6 (0.3–1.)
Liver	8.61 (5.0–12.8)	16.8 (12.1–20.6)	0.4 (0.3–0.7)	2.2 (1.3–3.3)
Spleen	5.4 (2.9–7.7)	10.5 (7.3–12.8)	0.2 (0.2–0.4)	8.0 (5.7–11.)
MesLN	0.7 (0.3–1.6)	1.3 (0.6–2.9)	0.1 (0.0–0.8)	16.8 (2.5–> 50)
Intestine	5.5 (2.9–7.3)	10.7 (7.2–12.8)	0.1 (0.1–0.1)	22.0 (15.4–31.8)

Table 2: Parameter estimates of the mathematical model that was fitted to the data on migration of *in vivo* activated TDLs in mice [61]. We list i) the rate of TDL entrance into a particular organ from the blood m_{1i} (second column), ii) the percent of cells leaving the blood into a particular organ ($m_{1i}/(d + \sum_{i=1}^5 m_{1i})$, third column), iii) the rate of exit of TDLs from an organ to the blood m_{i1} (fourth column), and iv) the average residence time of TDLs in the organ (fifth column). The rate of migration of TDLs from the blood to all organs, $d + \sum_{i=1}^5 m_{1i}$, is 51.2 h^{-1} or the average residence time of cells in the blood is 1.2 min. The average residence time of cells in a particular organ is calculated as $1/m_{i1}$ (for blood, lung, and liver) and $2/m_{i1}$ for spleen, mLNs, and intestine. We assume $k = 2$ subcompartments in these latter organs since this allowed for the best description of the data based on AIC (results not shown). The rate of cell migration from the blood to other organs in the body $d = 7.50$ (3.68 – 10.02) h^{-1} . No dying cells were assumed to migrate to the liver as including this process did not improve the quality of the model fit to data (not shown). In brackets we show 95% confidence intervals calculated by bootstrapping the residuals with 1000 simulations [88].

Recirculating and non-recirculating lymphocytes

Data and analyses presented so far may create an impression that all (or nearly all) lymphocytes of the immune system recirculate. This is not likely to be the case in general. Multiple factors are likely to influence ability of lymphocyte to recirculate following adoptive transfer [50, 89]. For example, it was noted that handling of lymphocyte *in vitro* at low temperatures dramatically impedes lymphocyte recirculation kinetics; passaging of lymphocytes via an intermediate host before final transfer into definite hosts restores the ability of lymphocytes to recirculate [89]. Accurate counting of lymphocytes entering isolated perfused spleens allowed to conclude that a large fraction of spleen lymphocytes never exit the tissue during 7–10 days of perfusion [46]. More recently, experiments involving surgical joining of syngenic mice (parabiosis) allowed accurate tracking recirculation kinetics of memory CD8 T cells in mice [90]. Initial studies showed limited ability of memory T cells in tissues such as brain to recirculate between two parabiotic mice [90]. A more thorough follow up revealed an inability of LCMV-specific memory CD8 T cells residing in most nonlymphoid tissues to recirculate between

parabiotic mice [91]. Interestingly, some of such tissue-resident memory T cells were also found in secondary lymphoid tissues such as lymph nodes and spleen [92]. Because many of such non-recirculating T cells reside in peripheral, non-lymphoid tissues, perhaps, it is not totally surprising that they are not able to recirculate. However, non-recirculating cells could even be found in the blood, e.g., crawling along sinusoids in the liver [93, 94]. It is important to realize, however, that many of studies documenting “non-recirculatory” nature of lymphocytes have been performed for a relatively short-time period and deeper, mathematical modeling-assisted analyses of the data from such parabiosis experiments are needed in order to accurately quantify the residency time of such tissue-resident lymphocytes.

Summary

There have been multiple studies documenting lymphocyte migration kinetics via secondary lymphoid organs of mice, rats, pigs, and sheep. Many studies involved simple or complex mathematical models to estimate residence times of lymphocytes in different tissues and in different conditions. A quick comparison reveals that these estimates while being approximately similar still vary dramatically in absolute values (Table 1). It is unclear at present if such variability in estimates is simply due to differences in experimental methodologies involved, in mathematical modeling approaches, or both. Future studies should attempt to determine whether estimates of the residence times are robust to the choice of mathematical model. It is often expected (and was found in this analysis) that conclusions arising from models being fitted to data, for example, estimates of model parameters, can be model-dependent [75]. It is also possible that there may not be such a universal parameter such as lymphocyte (e.g., naive CD8 T cell) residency time in a LN or spleen. Residency time may depend on the environment lymphocyte is in (resting vs. inflamed LN), previous history of the lymphocyte, or other factors. Future studies will have to become more mechanistic and instead of simply measuring/estimating lymphocyte migration kinetics, should attempt to determine why some lymphocytes spend 10 hours in tissues while other lymphocytes only 5 hours. Fundamental understanding of lymphocyte recirculation kinetics (or absence of thereof for tissue-resident lymphocytes) should allow to improve therapies that involve lymphocytes such as cancer immunotherapies.

Acknowledgments

We would like to thank multiple people contributing to our discussions on lymphocyte recirculation including Reinhard Pabst, Jurgen Westermann, Dave Masopust, Gudrun Debes, Rob De Boer, Johannes Textor, Judith Mandl. We thank Jeremy Auerbach for the cartoon of the recirculation kinetics of thoracic duct lymphocytes. This work was supported by the NIH grant to VVG (R01 GM118553).

References

1. Janeway CA, Travers P, Walport M, Shlomchik M (2004) Immunobiology 5th edition. Garland Publishing.

2. Kindt TJ, Osborne B, Goldsby RA (2007) *Kuby Immunology*, 6th edition. W. H. Freeman, New York.
3. Jenkins MK, Moon JJ (2012) The role of naive T cell precursor frequency and recruitment in dictating immune response magnitude. *J Immunol* 188:4135–4140.
4. Kotturi MF, et al. (2007) The CD8+ T-cell response to lymphocytic choriomeningitis virus involves the L antigen: uncovering new tricks for an old virus. *J Virol* 81:4928–4940.
5. Polman CH, et al. (2006) A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *N Eng J Med* 354:899–910.
6. Schwab N, Schneider-Hohendorf T, Wiendl H (2015) Therapeutic uses of anti- α 4-integrin (anti-VLA-4) antibodies in multiple sclerosis. *Int Immunol* 27:47–53.
7. Li H, et al. (2018) Clinical adverse effects of natalizumab: Protocol for a meta-analysis of randomized double-blind placebo-controlled clinical trials. *Medicine* 97:e11507.
8. Ford W, Gowans J (1969) The traffic of lymphocytes. *Semin Hematol* 6:67–83.
9. Mackay CR, Marston WL, Dudler L (1990) Naive and memory T cells show distinct pathways of lymphocyte recirculation. *J Exp Med* 171:801–817.
10. Westermann J, Pabst R (1996) How organ-specific is the migration of 'naive' and 'memory' T cells? *Immunol Today* 17:278–82.
11. Westermann J, Bode U, Pabst R (1998) Migration of naive and memory T cells in vivo. *Immunol Today* 19:143–4.
12. Luettig B, et al. (1999) Naive and memory T lymphocytes migrate in comparable numbers through normal rat liver: activated T cells accumulate in the periportal field. *J Immunol* 163:4300–4307.
13. Westermann J, Bode U (1999) Distribution of activated T cells migrating through the body: a matter of life and death. *Immunol Today* 20:302–306.
14. Westermann J, Ehlers E, Exton M, Kaiser M, Bode U (2001) Migration of naive, effector and memory T cells: implications for the regulation of immune responses. *Immunol Rev* 184:20–37.
15. Liu L, Fuhlbrigge RC, Karibian K, Tian T, Kupper TS (2006) Dynamic programming of CD8+ T cell trafficking after live viral immunization. *Immunity* 25:511–520.
16. Bimczok D, Rothkotter H (2006) Lymphocyte migration studies. *Vet Res* 37:325–338.
17. Tomura M, Itoh K, Kanagawa O (2010) Naive CD4+ T lymphocytes circulate through lymphoid organs to interact with endogenous antigens and upregulate their function. *J Immunol* 184:4646–4653.
18. Gebhardt T, et al. (2011) Different patterns of peripheral migration by memory CD4+ and CD8+ T cells. *Nature* 477:216–219.
19. Westermann J, Engelhardt B, Hoffmann J (2001) Migration of T cells in vivo: molecular mechanisms and clinical implications. *Ann Intern Med* 135:279–95.

20. Galkina E, et al. (2005) Preferential migration of effector CD8+ T cells into the interstitium of the normal lung. *J Clin Invest* 115:3473–3483.
21. Jennrich S, Lee MH, Lynn RC, Dewberry K, Debes GF (2012) Tissue exit: a novel control point in the accumulation of antigen-specific CD8 T cells in the influenza A virus-infected lung. *J Virol* 86:3436–3445.
22. Zatz MM, Lance EM (1971) The distribution of ⁵¹Cr-labeled lymphocytes into antigen-stimulated mice. Lymphocyte trapping. *J Exp Med* 134:224–241.
23. Kawashima Y, Sugimura M, Hwang YC, Kudo N (1964) The lymph system in mice. *Jap J Vet Res* 12:69–78.
24. Qatarneh SM, Kiricuta IC, Brahme A, Tiede U, Lind BK (2006) Three-dimensional atlas of lymph node topography based on the visible human data set. *Anat Rec* 289:98–111.
25. Trepel F (1974) Number and distribution of lymphocytes in man. A critical analysis. *Klin Wschr* 52:511–515.
26. Qatarneh S (2006) Development of a whole body atlas for radiation therapy planning and treatment optimization. Division of Medical Radiation Physics Department of Oncology-Pathology Stockholm University & Karolinska Institutet.
27. Pabst R, Trepel F (1975) Quantitative evaluation of the total number and distribution of lymphocytes in young pigs. *Blut* 31:77–86.
28. Pabst R, Trepel F (1975) The predominant role of the spleen in lymphocyte recirculation. I. Homing of lymphocytes to and release from the isolated perfused pig spleen. *Cell Tissue Kinet* 8:529–541.
29. Pabst R, Trepel F (1976) The predominant role of the spleen in lymphocyte recirculation. II. Pre- and postsplenectomy retransfusion studies in young pigs. *Cell Tissue Kinet* 9:179–189.
30. Pabst R (1988) The spleen in lymphocyte migration. *Immunol Today* 9:43–45.
31. Ganusov VV, De Boer RJ (2007) Do most lymphocytes in humans really reside in the gut? *Trends Immunol* 28:514–8.
32. Beltman J, Maree A, Lynch J, Miller M, de Boer R (2007) Lymph node topology dictates T cell migration behavior. *J Exp Med* 204:771–80.
33. Ariotti S, et al. (2012) Tissue-resident memory CD8+ T cells continuously patrol skin epithelia to quickly recognize local antigen. *Proc Natl Acad Sci U S A* 109:19739–19744.
34. Gowans JL (1957) The effect of the continuous re-infusion of lymph and lymphocytes on the output of lymphocytes from the thoracic duct of unanaesthetized rats. *Br J Exp Pathol* 38:67–78.
35. Gowans JL (1959) The recirculation of lymphocytes from blood to lymph in the rat. *The Journal of physiology* 146:54–69.
36. Gowans JL, Knight EJ (1964) The route of re-circulation of lymphocytes in the rat. *Proc R Soc Lond B Biol Sci* 159:257–282.

37. Hall JG, Morris B (1965) The origin of the cells in the efferent lymph from a single lymph node. *J Exp Med* 121:901–910.
38. Ford WL, Gowans JL (1968) Lymphocyte circulation in the rat. *Nouv Rev Fr Hematol* 8:509–518.
39. Cahill RN, Frost H, Trnka Z (1976) The effects of antigen on the migration of recirculating lymphocytes through single lymph nodes. *J Exp Med* 143:870–888.
40. Hay JB, Hobbs BB (1977) The flow of blood to lymph nodes and its relation to lymphocyte traffic and the immune response. *J Exp Med* 145:31–44.
41. Moon J, et al. (2007) Naive CD4(+) T cell frequency varies for different epitopes and predicts repertoire diversity and response magnitude. *Immunity* 27:203–13.
42. Moon JJ, et al. (2009) Tracking epitope-specific T cells. *Nat Protoc* 4:565–581.
43. Christensen BE, Jnsson V, Matre R, Tnder O (1978) Traffic of T and B lymphocytes in the normal spleen. *Scand J Haematol* 20:246–257.
44. Pabst R, Geisler R (1981) The route of migration of lymphocytes from blood to spleen and mesenteric lymph nodes in the pig. *Cell Tissue Res* 221:361–370.
45. van Ewijk W, Nieuwenhuis P (1985) Compartments, domains and migration pathways of lymphoid cells in the splenic pulp. *Experientia* 41:199–208.
46. Ford WL (1969) The kinetics of lymphocyte recirculation within the rat spleen. *Cell Tissue Kinet* 2:171–191.
47. Ford W (1979) Lymphocytes. 3. Distribution. Distribution of lymphocytes in health. *J Clin Pathol Suppl (R Coll Pathol)* 13:63–69.
48. Hammond BJ (1975) A compartmental analysis of circulatory lymphocytes in the spleen. *Cell Tissue Kinet* 8:153–169.
49. Ganusov VV, Auerbach J (2014) Mathematical modeling reveals kinetics of lymphocyte recirculation in the whole organism. *PLoS Comp Biol* 10:e1003586.
50. Smith M, Ford W (1983) The recirculating lymphocyte pool of the rat: a systematic description of the migratory behaviour of recirculating lymphocytes. *Immunology* 49:83–94.
51. Pabst R, Rosenberg YJ (1998) Interpreting data on lymphocyte subsets in the blood of HIV patients - organ distribution, proliferation and migration kinetics are critical factors. *Pathobiology* 66:117–122.
52. Westermann J, et al. (2003) Analyzing the migration of labeled T cells in vivo: an essential approach with challenging features. *Lab Invest* 83:459–69.
53. Di Rosa F, Pabst R (2005) The bone marrow: a nest for migratory memory T cells. *Trends Immunol* 26:360–366.
54. Frost H, Cahill R, Trnka Z (1975) The migration of recirculating autologous and allogeneic lymphocytes through single lymph nodes. *Eur J Immunol* 5:839–843.

55. Reynolds J, Heron I, Dudler L, Trnka Z (1982) T-cell recirculation in the sheep: migratory properties of cells from lymph nodes. *Immunology* 47:415–421.
56. Young AJ (1999) The physiology of lymphocyte migration through the single lymph node in vivo. *Semin Immunol* 11:73–83.
57. Thielke KH, Pabst R, Rothkötter HJ (1999) Quantification of proliferating lymphocyte subsets appearing in the intestinal lymph and the blood. *Clin Exp Immunol* 117:277–284.
58. Ford W, Simmonds S (1972) The tempo of lymphocyte recirculation from blood to lymph in the rat. *Cell Tissue Kinet* 5:175–189.
59. Howard JC (1972) The life-span and recirculation of marrow-derived small lymphocytes from the rat thoracic duct. *J Exp Med* 135:185–199.
60. Sprent J, Miller JF (1976) Fate of H2-activated T lymphocytes in syngeneic hosts. II. Residence in recirculating lymphocyte pool and capacity to migrate to allografts. *Cell Immunol* 21:303–313.
61. Sprent J (1976) Fate of H2-activated T lymphocytes in syngeneic hosts. I. Fate in lymphoid tissues and intestines traced with 3H-thymidine, 125I-deoxyuridine and 51chromium. *Cell Immunol* 21:278–302.
62. Ford W, Smith M (1982) Experimental approaches to lymphocyte traffic: pitfalls of the tracer sample method. *Adv Exp Med Biol* 149:139–145.
63. Fossum S, Smith M, Ford W (1983) The recirculation of T and B lymphocytes in the athymic, nude rat. *Scand J Immunol* 17:551–557.
64. Westermann J, Puskas Z, Pabst R (1988) Blood transit and recirculation kinetics of lymphocyte subsets in normal rats. *Scand J Immunol* 28:203–210.
65. Westermann J, Persin S, Matyas J, van der Meide P, Pabst R (1994) Migration of so-called naive and memory T lymphocytes from blood to lymph in the rat. The influence of IFN-gamma on the circulation pattern. *J Immunol* 152:1744–1750.
66. Stekel DJ (1997) The role of inter-cellular adhesion in the recirculation of T lymphocytes. *J Theor Biol* 186:491–501.
67. Stekel DJ, Parker CE, Nowak MA (1997) A model of lymphocyte recirculation. *Immunol Today* 18:216–221.
68. Stekel DJ (1998) The simulation of density-dependent effects in the recirculation of T lymphocytes. *Scand J Immunol* 47:426–430.
69. Grigorova IL, Panteleev M, Cyster JG (2010) Lymph node cortical sinus organization and relationship to lymphocyte egress dynamics and antigen exposure. *Proc Natl Acad Sci U S A* 107:20447–20452.
70. Textor J, et al. (2011) Defining the quantitative limits of intravital two-photon lymphocyte tracking. *Proc Natl Acad Sci U S A* 108:12401–12406.
71. Harp JR, Gilchrist MA, Onami TM (2010) Memory T cells are enriched in lymph nodes of selectin-ligand-deficient mice. *J Immunol* 185:5751–5761.

- 579 72. von Andrian U, Mempel T (2003) Homing and cellular traffic in lymph nodes. *Nat Rev Immunol*
580 3:867–78.
- 581 73. Mandl JN, et al. (2012) Quantification of lymph node transit times reveals differences in antigen
582 surveillance strategies of naive CD4+ and CD8+ T cells. *Proc Natl Acad Sci U S A* 109:18036–
583 18041.
- 584 74. Thomas N, Matejovicova L, Sriksalanukul W, Shawe-Taylor J, Chain B (2012) Directional
585 migration of recirculating lymphocytes through lymph nodes via random walks. *PLoS One*
586 7:e45262.
- 587 75. Ganusov VV (2016) Strong Inference in Mathematical Modeling: A Method for Robust Science
588 in the Twenty-First Century. *Front Microbiol* 7:1131.
- 589 76. Tomura M, et al. (2008) Monitoring cellular movement in vivo with photoconvertible fluorescence
590 protein "Kaede" transgenic mice. *Proc Natl Acad Sci U S A* 105:10871–10876.
- 591 77. Tomura M, et al. (2010) Activated regulatory T cells are the major T cell type emigrating from
592 the skin during a cutaneous immune response in mice. *J Clin Invest* 120:883–893.
- 593 78. Tomura M, et al. (2014) Tracking and quantification of dendritic cell migration and antigen
594 trafficking between the skin and lymph nodes. *Sci Rep* 4:6030.
- 595 79. Tomura M (2018) New tools for imaging of immune systems: Visualization of cell cycle, cell
596 death, and cell movement by using the mice lines expressing Fucci, SCAT3.1, and Kaede and
597 KikGR. *Meth Mol Biol* 1763:165–174.
- 598 80. Bates DM, Watts DG (1988) Nonlinear regression analysis and its applications. John Wiles &
599 Sons, Inc., Hoboken, NJ, 365 p .
- 600 81. Farooqi ZH, Mohler RR (1989) Distribution models of recirculating lymphocytes. *IEEE Transc*
601 *Biomed Eng* 36:355–362.
- 602 82. Lee HB, Blafox MD (1985) Blood volume in the rat. *J Nucl Med* 26:72–76.
- 603 83. Edgren J, von Knorring J (1973) Radiological determination of heart volume in rats. *Experientia*
604 29:1174–1176.
- 605 84. Miller ED, Kistner JR, Epstein RM (1980) Whole-body distribution of radioactively labelled
606 microspheres in the rat during anesthesia with halothane, enflurane, or ketamine. *Anesthesiology*
607 52:296–302.
- 608 85. June CH (2007) Adoptive T cell therapy for cancer in the clinic. *J Clin Invest* 117:1466–1476.
- 609 86. Svane IM, Verdegaal EM (2014) Achievements and challenges of adoptive T cell therapy with
610 tumor-infiltrating or blood-derived lymphocytes for metastatic melanoma: what is needed to
611 achieve standard of care? *Cancer Immunol Immunother* 63:1081–1091.
- 612 87. Houot R, Schultz LM, Marabelle A, Kohrt H (2015) T-cell-based Immunotherapy: Adoptive Cell
613 Transfer and Checkpoint Inhibition. *Cancer Immunol Res* 3:1115–1122.
- 614 88. Efron B, Tibshirani R (1993) An introduction to the bootstrap. Chapman & Hall, New York,
615 436 p .

- 616 89. Smith ME, Ford WL (1983) The migration of lymphocytes across specialized vascular endothe-
617 lium. VI. The migratory behaviour of thoracic duct lymphocytes retransferred from the lymph
618 nodes, spleen, blood, or lymph of a primary recipient. *Cell Immunol* 78:161–173.
- 619 90. Klonowski K, et al. (2004) Dynamics of blood-borne CD8 memory T cell migration in vivo.
620 *Immunity* 20:551–62.
- 621 91. Steinert EM, et al. (2015) Quantifying Memory CD8 T Cells Reveals Regionalization of Immuno-
622 surveillance. *Cell* 161:737–749.
- 623 92. Schenkel JM, Fraser KA, Masopust D (2014) Cutting edge: resident memory CD8 T cells occupy
624 frontline niches in secondary lymphoid organs. *J Immunol* 192:2961–2964.
- 625 93. Fernandez-Ruiz D, et al. (2016) Liver-resident memory CD8(+) T cells form a front-line defense
626 against malaria liver-stage infection. *Immunity* 45:889–902.
- 627 94. McNamara HA, et al. (2017) Up-regulation of LFA-1 allows liver-resident memory T cells to
628 patrol and remain in the hepatic sinusoids. *Science Immunology* 1–10.