Divergence in the face of gene flow in two *Charadrius* plovers along the

Chinese coast

Xuejing Wang^{1§}, Pinjia Que^{2§}, Gerald Heckel^{3,5}, Junhua Hu⁴, Xuecong Zhang¹,
Chung-Yu Chiang⁶, Qin Huang¹, Simin Liu¹, Jonathan Martinez⁷, Nan Zhang¹, Emilio
Pagani-Núñez¹, Caroline Dingle⁸, Leung Yu Yan⁸, Tamás Székely^{1,9}, Zhengwang
Zhang², Yang Liu^{1*}

1. State Key Laboratory of Biocontrol, Department of Ecology, School of Life
Sciences, Sun Yat-sen University, Guangzhou 510275, China

2. Ministry of Education Key Laboratory for Biodiversity and Ecological
Engineering, College of Life Sciences, Beijing Normal University, Beijing,
100875, China

3. Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012
Bern, Switzerland

4. Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041,
China

5. Swiss Institute of Bioinformatics, Genopode, 1015 Lausanne, Switzerland

6. Department of Environmental Science, Tunhai University, Taichun, Taiwan,
China

7. 14, bis rue des Temples 45240 La Ferté Saint Aubin, France

8. School of Biological Sciences, The University of Hong Kong, Hong Kong
S.A.R., China

9. Milner Center for Evolution, Department of Biology and Biochemistry,
University of Bath, Bath BA1 7AY, UK

Running title: Incipient speciation in *Charadrius* plovers

Total words (without reference): 7930

References □ 133

30 ***Correspondence:** Yang Liu, School of Life Sciences, Sun Yat-sen University,
Guangzhou, China. Email: liuy353@mail.sysu.edu.cn;

32 [§] These authors contributed equally to this work.

34

Abstract

36 Speciation with gene flow is an alternative to the nascence of new taxa in strict
allopatric separation. Indeed, many taxa have parapatric distributions at present. It is
38 often unclear if these are secondary contacts, e.g. caused by past glaciation cycles or
the manifestation of speciation with gene flow, which hampers our understanding of
40 how different forces drive diversification. Here we studied genetic, phenotypic and
ecological aspects of divergence in a pair of incipient species, the Kentish (*Charadrius*
42 *alexandrinus*) and the white-faced Plovers (*C. dealbatus*), shorebirds with parapatric
breeding ranges along the Chinese coast. We assessed divergence based on
44 molecular markers with different modes of inheritance and quantified phenotypic and
ecological divergence in aspects of morphometric, dietary and climatic niches. These
46 analyses revealed small to moderate levels of genetic and phenotypic distinctiveness
with symmetric gene flow across the contact area at the Chinese coast. The two
48 species diverged approximately half a million years ago in dynamical isolation and
secondary contact due to cycling sea level changes between the Eastern and
50 Southern China Sea in the mid-late Pleistocene. We found evidence of character
displacement and ecological niche differentiation between the two species, invoking
52 the role of selection in facilitating divergence despite gene flow. These findings imply
that the ecology can indeed counter gene flow through divergent selection and thus
54 contribute to incipient speciation in these plovers. Furthermore, our study highlights
the importance of using integrative datasets to reveal the evolutionary history and
56 underlying mechanisms of speciation.

KEY WORDS: allopatric speciation, character displacement, gene flow, hybridization,

58 stable isotope analysis, ecological niche

1 INTRODUCTION

60 Understanding how strongly the evolutionary processes, i.e. selection, gene flow and
genetic drift shape the divergence of closely related species has been a long-standing
62 interest in evolutionary biology (Coyne & Orr 2004; Feder et al. 2013; Nosil & Feder
2012; Wagner et al. 2012; Wolf & Ellegren 2017). Allopatric speciation is
64 conventionally considered as the prevalent mode of speciation in which physical
barriers completely restrict gene flow between two populations, facilitating the initiation
66 of divergence through genetic drift or selection (Bernardi et al. 2008; Carson & Clague
1995; Mayr 1963). If populations remain isolated for a period long enough after
68 divergence has been established, then this divergence could be maintained even in
the presence of gene flow after secondary contact (Le Moan et al. 2016; Zhou et al.
70 2016).

72 An increasing number of studies have shown that divergence can arise and be
maintained due to selection imposed by heterogeneous environments despite the
74 constraining effect of gene flow (Martin et al. 2013; Morales et al. 2017). Under this
scenario, divergent selection or ecologically-mediated sexual selection operating on
76 certain (“magic”) traits can lead to reproductive isolation, and incompatibilities in a few
“speciation genes” may be enough to constrain the homogenizing effect caused by
78 gene flow even at an early stage of speciation (Fitzpatrick et al. 2015; Nosil & Schluter
2011; Slatkin 1973). For instance, different host plant preferences result in different
80 digestive and physiological traits in *Timema* walking-stick insects (Nosil 2007).
Locally-adapted phenotypes in body size are related to swimming and foraging
82 ecology in guppies (*Poecilia reticulata*), which can maintain population diversification
in the face of extensive gene flow (Fitzpatrick et al. 2015). Though evidence of
84 speciation with gene flow is accumulating (Martin et al. 2013; Morales et al. 2017;
Schluter 2009; Seehausen et al. 2014), to what extent a balance between selection
86 and gene flow can drive divergence remains largely unknown. This might be
interesting in populations with high propensity of range-wide gene flow such as in

88 waterfowl (e.g. Liu et al. 2012) and shorebirds (Eberhart & Phillips et al. 2015; Küpper et
al. 2012).

90

Shorebirds are a group of migratory species with remarkable movement ability.
92 Seasonal migration may increase the probability of individual dispersal between
populations (Arguedas & Parker 2000; Procházka et al. 2011) and consequently drive
94 frequent gene flow between geographically distant populations (Peters et al. 2012;
Winker et al. 2013). Direct evidence using tracking approaches confirmed that
96 extremely long-distance gene flow can be mediated through individual movements
among remote breeding colonies of pectoral sandpipers (*Calidris melanotos*)
98 (Kempenaers & Valcu 2017). Range-wide phylogeographic studies also revealed
extensive gene flow in several migratory shorebird species (e.g. Trimbos et al. 2011;
100 Küpper et al. 2012; Miller et al. 2014). This may result in weak genetic structure across
a species and consequently prevent population-level divergence (Eberhart & Phillips et
102 al. 2015; Jackson et al. 2017).

104 In this study, we test the role of ecology and gene flow on the divergence of two
shorebird species, the Kentish Plover (*Charadrius alexandrinus*) and the white-faced
106 Plover (*C. dealbatus*) (Figure 1) . *C. alexandrinus* breeds in coastal areas and inland
lakes in Europe, Asia and North Africa (del Hoyo et al. 2016). A previous study
108 uncovered low genetic differentiation of *C. alexandrinus* across the Eurasian continent,
and also between continental and island populations in East Asia (Küpper et al. 2012).
110 *C. dealbatus* was formerly regarded as a subspecies of Kentish plover (Hartert &
Jackson 1915; Swinhoe 1870). Breeding sites of *C. dealbatus* were documented along
112 the coast of China from Fujian to Hainan Island, and in south middle Vietnam (BirdLife
International 2016), yet its geographic range is uncertain (Kennerley *et al.* 2008).
114 Previous studies have found subtle but diagnosable differences in morphometric,
plumage and behavioral traits between the two taxa (Figure 1d), leading to the erection
116 of *C. dealbatus* as a full species (Bakewell & Kennerley 2008; Kennerley et al. 2008;
Rheindt et al. 2011). The first genetic investigation on the relationship between these

118 two taxa provided no evidence of genetic differentiation and concluded that these
species may only differ in a few genomic regions (Rheindt et al. 2011). Although based
120 on a handful of genetic loci, this finding is noteworthy as it raises a probability of
divergence in the face of gene flow between these two young species.

122

Here we provide comprehensive data from multiple sources to explore divergence
124 patterns between these two species of plovers. We carried out extensive sampling on
breeding populations covering the potential area of contact along the Chinese coast.
126 We obtained genetic polymorphism data from multiple markers with various mutation
rates, i.e. mitochondrial DNA, exons and autosomal microsatellites. Using these new
128 datasets, it is possible to estimate the intensity and direction of gene flow (Hey &
Nielsen 2004; Won & Hey 2005) as well as other important demographic parameters
130 such as the effective population size (N_e) and the timing of divergence (reviewed in
Hey 2006) between *C. alexandrinus* and *C. dealbatus*. Because it is difficult to directly
132 quantify divergent selection, inference from the comparisons of traits which are
potentially targets of selection are necessary to offer hints on selective mechanisms in
134 maintaining divergence (Hoskin & Higgie 2010; Lande 1982; Nosil 2007). By collecting
data on morphology, diet, and environmental niche, we attempt to 1) characterize
136 geographic variation in genetic and phenotypic traits in the two plovers, 2) estimate
demographic history and intensity of gene flow, and 3) infer the role of multiple factors,
138 i.e. geographical premating isolation and gene flow during divergence between these
two taxa. Taken together, our investigation provides new insights into the evolutionary
140 history of two incipient bird species.

142 **2 MATERIALS AND METHODS**

DNA sample collection

144 We mainly collected samples along the eastern coastal area of China (Fig 1a), and
also obtained samples from the two biggest continental islands, Taiwan and Hainan,
146 as well as two small islands that are close to the Chinese coastal line (Zhoushan and

Jinmen). Sampling sites were separated by approximately 200-250 km along a 2,300 km transect, spanning almost the entire Chinese coastline. Further, one high altitude population (breeding at 3 350 m a.s.l.) near Qinghai Lake was sampled as an inland outgroup. Breeding individuals were captured using a funnel-trap method described in Székely et al. (2008) between March and July in 2014-2015. For each nest, blood samples of the breeding pair or a chick were collected. Tissue samples were also obtained from dead individuals found in the field. Species identification during sampling was based on the summary of plumage characters, morphometric, and ecological differences between *C. alexandrinus* and *C. dealbatus*, as described in Kennerley et al. (2008). Overall, we collected samples from 454 individuals from 19 breeding sites. All bird-capture and sampling were performed with permission from the respective authorities (Mainland China: Beijing Normal University to PJQ and Sun Yat-sen University to YL; Taiwan: Changhua County Government, New Taipei City Government and Kinmen County Government to ZYJ).

Morphometric analysis

For each adult individual, four basic morphometric measurements were taken: body mass, wing length, tarsus length and bill length. Body mass was measured with an electronic scale (± 0.01 g). Wing length (flattened) was measured with a wing ruler (± 1 mm). Bill length to skull and tarsus lengths were measured using vernier callipers (± 0.01 mm). Measurements were taken by QH, PJQ and ZYJ following the standard described in Redfern & Clark (2001). To avoid potential biases, some individuals were measured twice or three times among these authors to make sure there was no significant difference ($p < 0.05$, between 20 – 30 trials) among them at the beginning and end of the fieldwork in the year 2014-2015.

We carried out principle component analysis (PCA) for the four traits to visualize the variation of morphometric data. Analyses of covariance (ANCOVAs) were conducted based on the PCA results to test statistical significance. We carried out t-tests between species for each measurement. Pairwise morphological difference Q_{ST} based on PC1

scores from PCA were calculated following Storz (2002) and plotted against linear distances between coastal populations. All aforementioned analyses were performed in PAST 3.12 (Hammer et al. 2001).

DNA Sequencing and microsatellite genotyping

We extracted genomic DNA using Tiangen Blood & Tissue Genome DNA Kits, following manufacturer protocols (Tiangen, Beijing, China). DNA quality was measured with NanoDrop 2000 (Thermo Scientific, USA). Three mtDNA loci, partial ATPase subunit 6/8, partial D-Loop of the mitochondrial control region (CR) and NADH dehydrogenase subunit 3 fragment (ND3), were amplified for all samples using primers from Rheindt *et al.* (2011). PCR reactions for mtDNA amplification were carried out in 20 µl volumes containing 1X PCR buffer (Takara Shuzo, Japan), 10-50ng DNA template, 0.5 unit Taq DNA Polymerase (Takara Shuzo, Japan), 1.0 µM of each primer, 2.0 mM of each dNTP and 1.0 µM MgCl₂. We also sequenced 14 autosomal and two Z-linked exonic loci (designed from Liu *et al.* 2018) for a subset of 40 individuals representing both species and several populations within species. PCR for nuclear loci had higher concentration of dNTP at 4.0 mM and 2.5 µM for MgCl₂ and PCR cycling profiles are listed in Table S3. Each PCR product was checked on a 1% agarose gel and then sequenced on ABI3730XL (Applied Biosystems, USA) served by MajorBio, Shanghai, China.

We genotyped all individuals at 22 autosomal microsatellite markers (mst) mostly from Küpper *et al.* (2007) but with C204 from Funk *et al.* (2007) and Hru2 from Primmer *et al.* (1995). Microsatellite loci were amplified using three multiplex PCRs with respective cycling profiles (Table S3). Each PCR was in a total volume of 15 µl containing 1X HotStart buffer (Tiangen), 10ng DNA template, 1 unit Multi HotStart DNA Polymerase (Tiangen), 0.4 µM of each fluorescently labeled primer, 2.0 mM of each dNTP and 2.0 µM MgCl₂. Multiplex PCR products and associated genotypes were isolated on ABI3730XL (Applied Biosystems, USA) served by Invitrogen, Shanghai, China and

206 their length was determined using GeneMapper software v.3.7 (Applied Biosystems)
against an internal size standard (GeneScan-500LIZ; Applied Biosystems).

208

Genetic diversity and population structure analyses

210 For DNA sequence data, we aligned each mitochondrial or nuclear locus using the
CLUSTAL W algorithm implemented in MEGA v.6.06 (Tamura et al. 2013), and the
212 alignment was checked by eye and manually edited if needed. For nuclear DNA
sequence data, we first used PHASE 2.1.1 (Stephens et al. 2001) to reconstruct
214 haplotypes of nuclear sequences with heterozygous sites. Each run was set to 10,000
iterations, 100 burn-in and 10 thinning intervals. For both mtDNA and nuclear loci,
216 basic genetic polymorphism statistics such as haplotype number h , haplotype diversity
 Hd , number of segregating sites S , nucleotide diversity π and Tajima's D (Tajima 1989)
218 of each locus and each population were calculated in DnaSP 5.10.1 (Librado & Rozas
2009). Haplotype networks of each locus were constructed using a median-joining
220 algorithm (Bandelt et al. 1999) in PopART 1.7.2 (Leigh & Bryant 2015).

222 We used FreeNA (Chapuis & Estoup 2007) to check for the frequency of null alleles at
each microsatellite locus. Further tests for Hardy-Weinberg equilibrium and pairwise
224 linkage disequilibrium (LD) were carried out with Arlequin 3.1.1 (Excoffier et al. 2005).
Hardy-Weinberg equilibrium tests were run with 10,000 permutations. LD tests were
226 run for 100,000 steps of Markov chain. To obtain genetic diversity estimates, we
calculated observed heterozygosity (H_o) and expected heterozygosity (H_E) in GenAlEx
228 6.5.1 (Peakall & Smouse 2012).

230 We estimated population structure between species and among sampling sites within
species using several approaches. First, for mtDNA and microsatellites, we performed
232 analyses of molecular variance implemented (AMOVAs) in Arlequin to assess the
proportion of genetic variance explained by the different partition settings, e.g. (1) the
234 two species, (2) *C. alexandrinus* continental populations and Taiwan island
populations, *C. dealbatus*, (3) coastal populations (including Hainan Island), Qinghai

236 and Taiwan. Second, we also calculated pairwise Φ_{ST} and F_{ST} between breeding sites
for mtDNA and mst, respectively, and we derived significance levels using 10,000
238 permutations in Arlequin.

240 For microsatellite genotypes only, we carried out assignment analyses with two
model-based Bayesian approaches. First, we performed a Bayesian clustering
242 analysis using the admixture model with correlated allele frequencies implemented in
STRUCTURE 2.3.4 (Hubisz et al. 2009; Pritchard et al. 2000). Ten independent
244 analyses were run from $K=1$ to $K=8$ for 500,000 Markov chain Monte Carlo (MCMC)
generations with 100,000 burn-in. Replicate runs were combined using STRUCTURE
246 Harvester 0.6.94 (Earl 2012) and CLUMPP 1.1.2 (Jakobsson & Rosenberg 2007). The
most likely number of genetic clusters was also determined using Structure Harvester
248 using the criteria described in Evanno et al. (2005). Second, we used a Bayesian
clustering algorithm that takes the geographical coordinates of each sampling site into
250 account using the R package GENELAND 4.0 (Guillot et al. 2005). 1,000,000 MCMC
iterations were run with thinning set to 100 from 1 to 10 populations, with maximum
252 rate of Poisson process set to 100, uncertainty of spatial coordinates set to 0,
maximum number of nuclei in the Poisson–Voronoi tessellation to 300, and
254 independent Dirichlet distribution model for allele frequencies. With the same package,
the most likely number of clusters was determined based on their posterior density. To
256 confirm the consistency of the results, we repeated the MCMC simulation 10 times.

258 Furthermore, to determine the probability that an individual was a hybrid or a migrant,
we estimated the posterior probability of each individual based on multilocus
260 genotypes. First we used HYBRIDLAB 1.0 (Nielsen et al. 2006) to simulate 100
individuals of each species, F1, F2 and back-crosses in each direction from
262 microsatellite genotypes of 260 individuals with q-value higher than 95% in
STRUCTURE analysis. 100 simulated parents from each species and 100 F1 were
264 used to calculate the threshold for individual hybrid assignment in STRUCTURE
following Vähä and Primmer (2006). Finally, NewHybrid 1.1 (Anderson & Thompson

2002) was used to identify potential hybrid individuals. 100,000 burn-in followed by 400,000 sweeps were performed for both simulated and real data.

Demographic analysis

To infer the demographic histories of the two plover species, we applied Isolation with Migration model (IM) (Hey & Nielsen 2007) analyses using the combined sequence data set of 16 nuclear and three mtDNA loci based on 20 individuals of each species. IM analysis allows the inference of genealogies under different demographic scenarios and estimation of population genetic parameters, such as divergence times, effective population sizes and migration rates between species since their divergence from the common ancestor. The homologues Killdeer *Charadrius vociferus* sequences from GenBank were used as outgroup. The substitution rate of each locus was calculated using the method in Li et al. (2010). The ratio of net genetic distance of each locus across ingroup–outgroup was calculated, compared with net distance of mitochondrial cytochrome b (*cytb*) and then multiplied by the substitution rate for *cytb* (0.0105 ± 0.0005 substitution/site/mya, Weir & Schluter 2008). We used the Phi test (Bruen et al. 2006) for recombination within nuclear loci and no recombination was detected. We implemented models in IMA2p (Sethuraman & Hey 2016), the parallel version of IMA2 (Hey 2010). For each analysis, we ran 48 MCMC chains for 2,500,000 steps of burn-in followed by 500,000 genealogies saved, each recorded after 100 steps. Because IM analysis only provides estimates of average gene flow since the divergence of the two species, we used a Bayesian analysis in BayesAss 3.0.4 (Stephens & Donnelly 2003) based on microsatellite genotypes to characterize the level of gene flow within recent generations. We performed 10,000,000 iterations and 1,000,000 burn-in.

Ecological niche modeling and niche overlaps

To infer potential past range shifts induced by climatic changes, we carried out ecological niche modeling (ENM) using Maxent 3.3.3k (Phillips et al. 2009). The occurrence records of the two species of plovers were obtained from online databases of the Global Biodiversity Information Facility (GBIF, <http://www.gbif.org/>), China Bird

296 Report (<http://www.birdreport.cn>) and our records during the sampling expeditions.
Further, eight bioclimatic variables (i.e., the mean diurnal range, isothermality,
298 minimum temperature of the coldest month, mean temperature of the warmest quarter,
annual precipitation, precipitation of the driest month and precipitation seasonality)
300 were obtained from the WorldClim database v.1.4 (Hijmans et al. 2005).

302 To explore niche similarity between the two species, we performed an ordination null
test of PCA-env in environmental (E)-space (Broennimann et al. 2012; Hu et al. 2016).
304 The PCA-env calculates the occurrence density and environmental factor density
along environmental (principal component) axes for each cell using a kernel smoothing
306 method and then uses the density of both occurrences and environmental variables to
measure niche overlap along these axes (Broennimann et al. 2012). An unbiased
308 estimate of Schoener's D metric was calculated for our data using smoothed densities
from a kernel density function to measure niche overlap between the two species that
310 is ensured to be independent of the resolution of the grid. Statistical confidence in
niche overlap values was then tested through a one-sided niche-similarity test
312 (Broennimann et al. 2012). All statistical analyses were performed in R 3.0.2 (R
Development Core Team 2013) using scripts available in Broennimann et al. (2012).
314 Details of ENM construction and niche-similarity tests are available in Appendix S1.

316 **Dietary niche differentiation inferred by stable-isotope analysis**

Because stable isotopic compositions of consumer tissues can be used to estimate the
318 relative contribution of assimilated dietary sources (DeNiro & Epstein 1978),
stable-carbon(C) and nitrogen(N) isotope analysis is widely used as a tool to study
320 avian dietary patterns (Hobson & Clark 1992; Pagani-Núñez et al. 2017). Carbon
isotope ratios differ between C3, C4 and CAM plants due to differences in the
322 photosynthetic pathways, and these differences are incorporated into an animal when
the plants are consumed and so can be used to infer information about dietary niches
324 (Hobson & Clark 1992). N isotopes are useful for identifying species/individuals which
occupy different trophic positions (high $\delta^{15}\text{N}$ implies higher trophic level; Chen et al.
326 2017). In order to compare dietary differences based on differences in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$
between the two species, we collected the outer pair of rectrices from seven adults per
328 site at eight sites: Qinghai Lake, Tangshan, Lianyungang and Rudong for C.

alexandrinus; Fuzhou, Shanwei, Zhanjiang, Dongfang for *C. dealbatus*. Since both species perform a complete post-breeding molt within their breeding grounds (Ginn & Melville 1983, personal observation), isotope ratios represent trophic level and habitat preferences during the breeding period. We estimated niche width and overlap per species using an isotopic Bayesian approach based on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ profile. Detailed information on lab procedures and statistical analyses can be found in Appendix S2.

3 RESULTS

Morphological differentiation

The two plover species showed subtle but significant differences for most morphological traits (ANCOVA; $p < 0.001$, Figure 1) even though there was considerable overlap in morphology between the species at the level of the individual (Figure 1f). On average, *C. dealbatus* had a longer bill (17.91 ± 0.94 mm vs. 16.69 ± 1.06 mm in *C. alexandrinus*, $p < 0.001$, Figure 1b), longer wings (118.58 ± 2.90 mm vs. 115.59 ± 3.02 mm in *C. alexandrinus*, $p < 0.001$, Figure 1c), and larger body mass (48.99 ± 3.26 g vs. 47.56 ± 3.73 g in *C. alexandrinus*, $p = 0.001$, Figure 1d, Table S5) than those of *C. alexandrinus*. There was no difference in tarsus length between the two species ($p = 0.962$). Individuals of *C. alexandrinus* from Taiwan Island were heavier than continental populations of *C. alexandrinus* and *C. dealbatus* (both $p < 0.001$). Q_{ST} between coastal populations of *C. alexandrinus* and *C. dealbatus* was negatively correlated with geographic distance ($R^2 = 0.293$, $p < 0.001$, Figure 1g).

Genetic diversity and population structure

We sequenced 357 individuals at all three mtDNA loci (224 *C. alexandrinus* and 133 *C. dealbatus*, GenBank Accession No. xxxx-xxxx). For each site, sample size ranged from 11 to 30 individuals. For each individual, we obtained in total 1729 base pairs (bp) of mtDNA sequence, including 846 bp ATPase6/8, 505 bp D-loop and 378 bp ND3. *C. alexandrinus* showed higher intraspecific genetic diversity than *C. dealbatus* (Table 1). Haplotype networks show that in all loci, most individuals were sorted into two major

haplogroups, corresponding to the two species of plovers. One non-synonymous substitution separated the two haplogroups at both ATPase6/8 and ND3 loci (Figure 2a). Moreover, a subset of samples containing 20 individuals of each species were sequenced at 16 loci (range 440-902 bp for each locus; Table 1 and Table S1, GenBank Accession No. xxxx-xxxx) for a total of 11,209 bp nuclear DNA sequence. The haplotype networks from autosomal and Z-linked loci did not show strong patterns of lineage sorting like mtDNA. The most common haplotypes were shared by both species of plover (Figure 2b and S1). Moreover, both species showed the signal of recent demographic expansion as detected by significant Tajima's D values (Table 1).

Overall genetic differentiation was significant and high in mtDNA data ($\Phi_{ST} = 0.506$, $p < 0.001$) and low at microsatellite loci ($F_{ST} = 0.036$, $p < 0.001$). For nuclear sequence data, genetic differentiation between species was also significant at autosomal loci ($\Phi_{ST} = 0.100$, $p < 0.001$) and particularly high at the Z-linked loci Z4 ($\Phi_{ST} = 0.726$, $p < 0.001$) and Z6 ($\Phi_{ST} = 0.309$, $p < 0.001$, Table S3). In the AMOVA analysis, we observed the largest difference between groups when the data were partitioned by species (i.e. *C. alexandrinus* and *C. dealbatus*), with this grouping explaining 49.7% of the variance in mtDNA and 2.4% in the microsatellites ($p < 0.001$; Table 2). These were significantly higher than the values of genetic variation when we partitioned samples as coastal vs. island populations or coastal population vs. Qinghai vs. Taiwan populations. Within *C. alexandrinus*, minor genetic differentiation was found between the inland population (Qinghai Lake) and coastal populations (mtDNA $\Phi_{SC} = 0.042$, $p = 0.022$; mst $F_{SC} = 0.021$, $p < 0.001$). *C. alexandrinus* populations on Taiwan Island also shared haplotypes with other coastal populations (mtDNA $\Phi_{SC} = 0.021$, $p = 0.146$); but were significantly differentiated in microsatellites (microsatellite $F_{SC} = 0.016$, $p < 0.001$).

For microsatellite loci, 18 out of 22 markers were successfully genotyped. However, four markers (Calex-04, 08, 19, C204) showed a large proportion of missing data, while another four loci (Calex-11, 24, 26, 43) showed an estimated frequency of null alleles over 10% and moreover one locus (Calex-35) showed H-W disequilibrium (Table S2).

388 Consequently, the aforementioned nine loci were removed from the dataset, making
genotypes at 13 loci of 402 (271 *C. alexandrinus* and 131 *C. dealbatus*) individuals for
390 the final microsatellite dataset for further analysis.

392 The results of the STRUCTURE analysis clearly showed two genetic clusters
representing each species (Figure 2d) and no obvious gradual transition along the
394 coastline that could be expected from a hybrid zone. The average *Delta K* value when
K = 2 was much higher than other options (Figure S2). Using georeferenced data in
396 GENELAND, our results corroborated the genetic clustering patterns inferred from
STRUCTURE. The posterior probability was 0.70 when *K* = 2 in contrast with the
398 posterior probability of 0.25 when *K* = 3. We visualized the GENELAND results in a
map with the probability distribution of posterior mode of class membership, which
400 further supports the separation between *C. alexandrinus* and *C. dealbatus* along the
China coast (Figure 2c). The divide between these two species was located between
402 Wenzhou and Fuzhou according to the GENELAND results, but it is unclear if these
two species come into contact or forms a putative hybrid zone in this region. Again, the
404 individuals from the sites in Taiwan were assigned to the cluster of *C. alexandrinus*
(Figure 2c).

406

Demographic history in two plovers

408 Isolation with migration analyses suggested that *C. alexandrinus* and *C. dealbatus*
diverged 0.56 (0.41 – 5.19) million years ago. Estimated migration rates in both
410 directions were significant ($p < 0.001$) with slightly higher gene flow from *C.*
alexandrinus into *C. dealbatus* (2.69 migrants per generation) than the vice versa (ca.
412 two migrants per generation). The estimated effective population size of *C.*
alexandrinus ($N_e \approx 1.59 - 4.44$ million) was about 8-10 times higher than for *C.*
414 *dealbatus* ($N_e \approx 0.15 - 0.52$ million).

416 The estimation of recent gene flow between the two species with BayesAss using
microsatellites also suggested bidirectional gene flow. Gene flow from *C. dealbatus* to

418 *C. alexandrinus* was slightly higher (0.013, $p = 0.028$) than that in the other direction
(0.010, $p = 0.028$). In *C. alexandrinus* populations, one likely migrant from *C. dealbatus*
420 and two hybrid F1 individuals were identified (probability higher than 0.5). With the
same threshold value, two migrants and one F1 individual were identified in *C.*
422 *dealbatus* populations.

424 **The detection of hybrid individuals**

Analyses with NewHybrids provided no evidence of a large number of hybrids or a
426 higher frequency of these near the potential contact area between the species. Based
on the STRUCTURE results and simulations, the optimal threshold for distinguishing
428 hybrids from non-hybrids was $q=0.836$. Based on this threshold, 81.9% of the
individuals (204 out of 246) collected from sites at the northern Chinese coastline,
430 Qinghai Lake and Taiwan Island were assigned to *C. alexandrinus* (Figure 2c-d).
Individuals with intermediate q -values, which were possibly hybrids, were found at
432 most northern Chinese sampling sites (Figure 2d). For the southern coastline in China,
145 out of 156 Individuals (92.9%) belonged to *C. dealbatus* with a probability larger
434 than 0.836 (Figure 2c-d). Only one individual of each species was assigned to the
genetic cluster representing the other species with high probability (Figure 2d).

436
The result of NewHybrids was highly concordant with STRUCTURE results and
438 suggested migrants moving in both directions. NewHybrids consistently identified
three migrants mentioned above with probability higher than 99%. However,
440 NewHybrids failed to identify simulated hybrid individuals and recognized them as
migrants. Thus, the result of NewHybrids was not used for hybrid identification (Fig
442 S3).

444 **Niche modeling, projections and comparisons**

Our ecological niche models effectively captured the current distribution of both *C.*
446 *alexandrinus* and *C. dealbatus* (Figure 4) with a high discrimination capacity (AUC
values > 0.88 for training and test data). Jackknife tests on variable importance for *C.*

448 *alexandrinus* revealed that isothermality, precipitation seasonality and mean
temperature of the warmest quarter were the three highest ranked variables when
450 used in isolation. For *C. dealbatus*, mean diurnal range and annual precipitation were
the most important variables. The simulation of the three periods, i.e. Last Interglacial
452 (LIG, 120-100Ka), Last Glacial Maximum (LGM, 21Ka, MIROC model) and current
times, respectively, showed range shifts in both species. In particular, these results
454 suggest that ranges of both species shrank during the LIG in the Chinese coastal area
accompanied by an increase of climatic suitability in the inland region for *C.*
456 *alexandrinus*. In contrast, suitable habitats expanded for both species during the LGM
in the coastal area, the East China Sea and the northern part of the South China Sea
458 (sea area between Hainan and Taiwan), probably due to the fall of the sea level.

460 The ordination approach using PCA-env suggested that the overlap of the current
climatic niches of the two species is relatively low (Figure S4a-b). The two species can
462 be separated based on the first two PCs with an accumulative 81.5% of the total
variance explained (Figure S4c). The niche equivalency test rejected the null
464 hypothesis that the species pair is distributed in identical environmental space ($p =$
0.019; Figure S4d).

466

Stable-isotope analysis

468 Our stable-isotope analysis showed that *C. dealbatus* exhibited significantly higher
 $\delta^{15}\text{N}$ values than *C. alexandrinus* ($p < 0.001$, Figure 5a, Figure S5). In contrast, we did
470 not find significant differences in $\delta^{13}\text{C}$ between the two species ($p = 0.161$). Further
isotope space overlap analysis showed that *C. alexandrinus* individuals had a high
472 probability to be found within the isotopic niche space of *C. dealbatus* (95.3%), while *C.*
dealbatus individuals showed a relatively low probability to be found in the isotopic
474 niche space of *C. alexandrinus* (46.5%, Figure 5b). Moreover, we found that *C.*
alexandrinus showed higher variability in $\delta^{15}\text{N}$ profile across breeding populations than
476 *C. dealbatus*.

478 4 DISCUSSION

Our genetic data show that *C. alexandrinus* and *C. dealbatus* have diverged to a level of advanced sorting of alleles between the two species particularly in mtDNA and Z-linked genes with lower effective population sizes (Figure 2a-b). For autosomal microsatellite data, we also found that genetic differentiation is low at intraspecific level but substantially high between the species (Figure 2c-d, Table 2). Though it is unclear whether a narrow hybrid zone exists in the contact area on the Chinese coast, a considerable level of symmetric gene flow occurred between the two plovers (Table 3). We find diagnosable differences in morphometric traits and ecological characters between the two plovers along the coast. At odds with these results, a previous work found no evidence of genetic differentiation between the two plovers (Rheindt et al. 2011). The present dataset comprises of systematically sampling along the Chinese coast, a larger suite of genetic markers and well-characterized ecological traits (Figure 1) while Rheindt et al. (2011) analyzed samples collected outside the breeding season. Potentially erroneous species assignment of non-breeding birds may underestimate genetic differentiation, because the distinguishing features in non-breeding plumage are relatively subtle (Bakewell & Kennerley 2008; Kennerley *et al.* 2008; Rheindt et al. 2011) and non-breeding *C. alexandrinus* and *C. dealbatus* mix frequently on the coast of the South China sea (own observation and Kennerley et al. 2008).

Phylogeographic patterns in *C. alexandrinus* and *C. dealbatus*

Two genetic lineages were found among breeding *Charadrius* plovers along the Chinese coast, corresponding the northern lineage to *C. alexandrinus* and the southern one to *C. dealbatus*, respectively (Figure 1a, Table 2). The sharp genetic break between the two lineages lies between Wenzhou and Fuzhou (latitude 26-27 °N) north of the Taiwan Strait (Figure 2c-d). Furthermore, samples from Taiwan Island belong to *C. alexandrinus* but the population in Jinmen Island is affiliated with *C. dealbatus* (Figure 2c-d). This pattern likely reflects historical isolation of the two species through separation by some geographical barriers at the Chinese coast during the Pleistocene climatic fluctuation periods. Divergence between the two plovers 560,000 years ago was inferred based on IM for the two species (Table 3, Figure 3).

This period falls into the Marine Isotope Stage 16 of the mid-late Pleistocene, during which was one of the Ice Age (Railsback et al. 2015). Accordingly the divergence between the plovers was probably linked to vicariance when the coastline was separated by a land-bridge raised due to the shallow sills between the East and the South China Sea (Ni et al. 2014). *Charadrius* plovers have originated in the Northern hemisphere and then radiated to the Southern hemisphere as suggested by a recent molecular phylogeny (Dos Remedios et al. 2015). It is thereby likely that *C. dealbatus* originated from *C. alexandrinus* and could have been diverging along the east China coast since the mid-late Pleistocene.

Phylogeographic patterns have been relatively well characterized in shorebirds breeding at high latitudes (Trimbos et al. 2011; Küpper et al. 2012; Miller et al. 2014). Panmixia or weak genetic differentiation was usually suggested, most likely driven by extensive gene flow. However, it seems that islands can act as a barrier to natal dispersal for the continental Kentish plovers (Küpper et al. 2012). In contrast, population structures in temperate and subtropical shorebird populations are poorly documented, probably due to a low level of species diversity. To the best of our knowledge, our study shows the first documented phylogeographic break in a bird species in the coastline in China. Interestingly, concordant phylogeographic patterns have been described in several coastal marine taxa, such as plants (Wang et al. 2015), fishes (Ding et al. 2018; Liu et al. 2007), shellfishes (Ni et al. 2012), and crustaceans (Wang et al. 2008). Though the exact splitting times are not congruent, the observed split line is at approximately 25°N latitude between the East China Sea and the Southern China Sea (reviewed in Ni et al. 2014). This consensus pattern resulted in the hypothesis that historical factors, i.e. sea level fluctuation during the Pleistocene, caused a convergent phylogeographic pattern in multiple coastal marine fauna in the marginal northwestern Pacific Ocean (Ni et al. 2014; Ding et al. 2018). Thus, our study contributes a vertebrate case to the accumulating literature about this species divergence hotspot. Apart from the major role of physical barriers, comparative phylogeographic studies also revealed that other abiotic factors, like ocean current and

hydrothermal conditions, as well as species ecological characters, i.e. dispersal ability,
habitat preference, life-history, and population demography can also play a role in
contributing to divergence of coastline fauna (Liu et al. 2007; Ni et al. 2014).

Morphological and ecological differentiation along a latitudinal gradient

The two plovers, *C. alexandrinus* and *C. dealbatus* are distributed along the Chinese coastline across a latitudinal and associated environmental gradients from temperate to tropical zones. Within each species, we found a general trend that northern populations have larger values than the southern counterparts in morphometry (Figure 1). This pattern may be related to Bergmann's rule which states that body size of homeothermic animals is larger in colder climates than in warmer ones (Bergmann 1848). Populations of both plover species distributed in such a north-south gradient may benefit from an optimal temperature control (Salewski & Watt 2017). A mutually non-exclusive explanation is that the difference in morphological traits, i.e. body mass and bill length link to potential differences in resource exploitation and life history. Our data show that *C. dealbatus* has a larger average body mass than *C. alexandrinus*. Body mass is a comprehensive trait reflecting nutrition assimilation, energy reservation and expenditure (McNab 2009). The difference in body mass might be related to the difference in migration behavior where a lighter body mass in *C. alexandrinus* is favored by decreased transport cost of fuel storage during migration (Alerstam et al. 2003; McWilliams & Karasov 2001). In contrast, larger body mass in *C. dealbatus*, a short-distance migrant or resident, may be beneficial in multiple reproduction within a single breeding season while *C. alexandrinus* produces usually only a single clutch (Lin et al. in prep.). Difference in bill length may be driven by the difference in the use of food resources (Badyaev et al. 2008) but recently, the function of the bill as temperature regulator also started to attract scientific attention (Tattersall et al. 2016). On the other hand, an increasing number of studies have demonstrated that birds at higher latitude and in cooler environments have shorter bills (e.g. European sparrows, Johnston 1969; Australian shorebirds, Nebel et al. 2013), consistent with Allen's Rule (Allen 1877). Nevertheless, we find that populations of *C. alexandrinus* in Taiwan had

similarly large body mass and bill lengths as *C. dealbatus* populations (Figure 1b-d, g), which are significantly larger than mainland conspecific populations. This is probably caused by phenotypic plasticity in the Taiwan population to a sub-tropical ecology and resident life history, similar to the populations of *C. dealbatus* in the South China sea coast.

Climatic niche modelling indicates that the climatic conditions of the two species are spatially separated (Figure S4). *C. dealbatus* is restricted to breeding sites close to the coast, particularly in warmer tropical climate. In contrast, *C. alexandrinus* has a wider climatic niche, as represented by a broader climatic zone (Figure 4). This species can breed not only on temperate coasts (Que et al. 2015) but also in inland saline lake shore (Cramp & Perrins 1983), just as Qinghai Lake. In addition, our isotope analysis revealed that *C. alexandrinus* covered a wider range of isotope ratios than *C. dealbatus*, but *C. alexandrinus* exploited a lower trophic range ($\delta^{15}\text{N}$). This indicates that *C. dealbatus* probably feeds on a higher energy diet than its sister species (Hobson & Clark 1992). However, it is unclear whether such difference may result from diet preference divergence or due to differences in food resource availability (McCormack et al. 2010). Taken together, these results suggest that the ecological niches of the two plovers are significantly different in several aspects, and support a role for ecology in constraining range limits and perhaps habitat preference for the two shorebird species.

Incipient speciation with ongoing gene flow

We find evidence of incipient speciation with ongoing gene flow between the two plovers. The IM results indicate a relatively young split with considerable gene flow between the two species (Figure 3 and Table 3). We suggest this is probably due to gene flow caused by secondary contact during the Pleistocene, because our niche modeling analysis reveals that the ranges of the two species were expanded and consequently overlapped between the Eastern and Southern China Sea during the Last Glacial Maximum (Figure 4). While our niche modeling demonstrates a cycle of

range dynamics caused by sea-level changes, one should bear in mind that the last
600 million years witnessed multiple glaciation cycles (Taylor et al. 2014, 2015). In
particular, marine oxygen isotope records designate 28 isotope stages corresponding
602 to glacio-climatic cycles in the last million years (Railsback et al. 2015). For ocean
marginal and coastline taxa, this may imply that the fluctuations in sea level can lead to
604 alternation between population isolation and contact throughout their evolutionary
histories. This can in turn yield significant consequences on population divergence and
606 historical demography because speciation can proceed through cycles of allopatric
stages interspersed by parapatry, resulting in a diversification process characterized
608 isolation with migration (Rheindt & Edwards, 2011).

Under this premise, a key question arises: after the two plovers diverged, how was
their divergence maintained in the presence of ongoing gene flow? Speciation theory
612 predicts that divergence is initiated either by genetic drift or divergent selection (Lynch
et al. 2016; Rundle & Nosil 2005; Seehausen et al. 2014; Wolf & Ellegren 2017).
614 However, genetic drift is unlikely to have been the only force to initiate divergence in
the plovers for two reasons: the large effective population sizes (N_e) and a relatively
616 high level of gene flow (Table 3). Moreover, large N_e and recent divergence make it
unlikely that genetic drift could have led to diagnosable differences between
618 populations (Ellstrand & Elam 1993; Lanfear et al. 2014), and gene flow directly
counteracts the diverging effects of drift (García-Navas et al. 2015; Poelstra et al.
620 2014). Another possibility is that sufficient divergent selection could contribute to
overcome the homogenizing effects of gene flow (Coyne & Orr 2004; Feder et al. 2013;
622 Nosil & Feder 2012; Wagner et al. 2012; Wolf & Ellegren 2017). In light of this view, the
“genomic islands of speciation” model was proposed to illustrate that divergent
624 selection can have effects in specific regions across the genome leading to divergence
with gene flow (Feder et al. 2013; Nosil & Feder 2012; Wagner et al. 2012; Wolf &
626 Ellegren 2017). Data from several organisms suggest that incipient speciation can be
maintained with divergence at a small number genomic regions (Martin et al. 2013;

Morales et al. 2017; Nadachowska-Brzyska et al. 2013; Towes et al. 2016 but see Curickshan et al. 2014, Burri 2017).

The geographical boundary between the two plovers was ambiguously defined in previous studies (Kennerley et al. 2008; Rheindt et al. 2011). Our results show that the discontinuity in genetic structuring (Figure 2c) and morphometric values (Figure 1b-d) between the two species is situated at coastline between Wenzhou and Fuzhou, indicating a contact zone (Figure 2c). For incipient species, individuals may show a clinal pattern of allele frequency and morphology at their contact and form a hybrid zone (Bastos & Silveira et al. 2012; Harrison 1986; Taylor et al. 2014). Our STRUCTURE results revealed no obvious signs of a hybrid zone (Figure 2d), and rather indicated sporadic migrants in the respective range of the other species. One possibility is that our sampling transect was not fine-grained enough to be able to discover the potential hybrid zone, which would thus be narrower than the 200 km distance between the two neighboring sampling sites. Another possibility for an apparent lack of a hybrid zone is assortative mating between the two plovers at the contact (Bearhop et al. 2005). The latter explanation is supported by the fact that populations of the two species close to this region were much more divergent in morphological traits than ones farther apart (Figure 1b-d). Reproductive character displacement between the two species (Brown & Wilson 1956; Robinson & Wilson 1994; Grant and Grant 2006) in the contact regions would constrain interbreeding (Rybinski et al. 2016; Winkelmann et al. 2014). Obviously, genetic data alone cannot demonstrate this and detailed experiments related to assortative mating in the contact zone are required.

Taxonomic implications

This study offers several taxonomic implications for the *C. alexandrinus* complex in East Asia. Kennerley et al. (2008) recommended that the tropical breeding population, previously defined as the subspecies *dealbatus* warranted species status, based on differentiation in morphology, behavior and distribution from *C. alexandrinus*.

658 Uncertainty arouse about the taxonomic status because *dealbatus* was not
distinguished from *C. alexandrinus* in a first genetic evaluation (Rheindt et al. 2011).
660 Our results provide further support for a pair of incipient species, indicating that *C.*
dealbatus is the youngest species within the genus *Charadrius* (Barth et al. 2013).
662 Although we did not apply explicit Bayesian species delimitation analyses (e.g. Yang &
Rannala 2014), our results clearly demonstrate multiple diagnosable morphological
664 characters and distinct ecological niches, which are likely key factors to maintain
species limits. Beyond the detection of differences between these plovers, this study
666 forms a basis for conservation considerations of *C. dealbatus* because of its restricted
range and adaptation to subtropical climate, as well as a risk of population decline
668 caused by habitat loss in the China coastline (Ma et al. 2014).

670 **Conclusions and perspectives**

Resolving the balance between diverging selection and gene flow is of fundamental
672 importance to understand speciation processes. Here, we show that *C. alexandrinus*
and *C. dealbatus* represent a case of incipient species in which divergent selection
674 associated with ecological differences likely works as efficient mechanism for the
maintenance of divergence in the face of gene flow. While the full species status of *C.*
676 *dealbatus* may be justified, it remains untested whether a strong level of reproductive
isolation has been established between *C. alexandrinus* and *C. dealbatus*. Further, we
678 have shown low genetic divergence between the two plovers, so it would be of
particular importance to explore the patterns of divergence at the genome level and
680 determine whether specific regions are related reproductive isolation and adaptation.
In addition, the IM analysis presented here estimated average historical gene flow, but
682 it would be interesting to evaluate more realistic (and complex) demographic models in
order to estimate changes in *N_e* and the span of isolation and secondary contact better.
684 Genome-wide population genetic approaches promise the power for better resolved
evolutionary history of the two *Charadrius* plovers but also open an avenue to
686 characterize the genetic architecture associated with phenotypic trait divergence and

local adaptation (Freedman et al. 2014; Nadachowska-Brzyska et al. 2013; Towes et al.
2016).

ACKNOWLEDGEMENTS

We thank Qiaoyi Liang, Xiaoyan Long, Derong Meng, Demeng Jiang, Li Tian and Hebo
Peng help with field sampling, Guoling Chen, Yangkai Zhou for their assistance with
laboratory works. Special thanks are due Shaochong Peng for preparing sketches of
plovers in Figure 1.

Funding information:

This study was supported by National Natural Science Foundation of China,
Grant/Award Number: 31301875 and 31572251 to YL, 31600297 to PJQ and
31572288 to ZWZ; and the Open Grant of the State Key Laboratory of Biocontrol of
Sun Yat-sen University to YL and TZ (SKLBC13KF03), Youth Innovation Promotion
Association CAS (2015304) to JHH. Computational machinery time work was granted
by Special Program for Applied Research on Super Computation of the
NSFC-Guangdong Joint Fund (the second phase) under Grant No. U1501501 to YL.

DATA ACCESSIBILITY

Sequences deposited at GenBank: accession number xxxxx-xxxxx
Phenotype, distribution, stable isotope and microsatellite genotypes available at:
Dryad Doi: xxx

AUTHOR CONTRIBUTIONS

Y.L., Z.W.Z. and T. S. conceived and designed the study, P.J.Q., C-Y.C., Q.H., J.M.,
N.Z., X.J.W., and T. S. collected data in the field, X.J.W. and S.M.L conducted
molecular genetic laboratory work, X.J.W. P.J.Q. analyzed genetic data with input from
Y.L. and G.H.. J.H.H. performed niche modeling analysis, C.C.Z. carried out stable
isotope analysis with input from E.P.-G. and C.D.. X.J.W P.J.Q. and Y.L. wrote the

manuscript with input from J.H.H., G.H., X.C.Z. and C.D. All authors gave final approval for publication.

REFERENCES

- Alerstam T, Hedenström A and Åkesson S (2003) Long-distance migration: evolution and determinants. *Oikos*, **103**, 247-260.
- Allen JA (1877) The influence of physical conditions in the genesis of species. *Radical Review* **1**, 108-140.
- Anderson E, Thompson E (2002) A model-based method for identifying species hybrids using multilocus genetic data. *Genetics* **160**, 1217-1229.
- Arguedas N, Parker PG (2000) Seasonal migration and genetic population structure in house wrens. *The Condor* **102**, 517-528.
- Badyaev AV, Young RL, Oh KP, Addison C (2008) Evolution on a local scale: developmental, functional, and genetic bases of divergence in bill form and associated changes in song structure between adjacent habitats. *Evolution* **62**, 1951-1964.
- Bakewell D, Kennerley P (2008) Field characteristics and distribution of an overlooked Charadrius plover from South-East Asia. *Birding Asia* **9**, 46-57.
- Bandelt H-J, Forster P, Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16**, 37-48.
- Barth JM, Matschiner M, Robertson BC (2013) Phylogenetic position and subspecies divergence of the endangered New Zealand Dotterel (*Charadrius obscurus*). *PLoS ONE* **8**, e78068.
- Bearhop S, Fiedler W, Furness RW, *et al.* (2005) Assortative mating as a mechanism for rapid evolution of a migratory divide. *Science* **310**, 502-504.
- Beaumont MA, Zhang W, Balding DJ (2002) Approximate Bayesian computation in population genetics. *Genetics* **162**, 2025-2035.
- Bergmann C (1848) *Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse*.
- Bernardi G, Alva-Campbell YR, Gasparini JL, Floeter SR (2008) Molecular ecology, speciation, and evolution of the reef fish genus Anisotremus. *Molecular Phylogenetics and Evolution* **48**, 929-935.
- Broennimann O, Fitzpatrick MC, Pearman PB, *et al.* (2012) Measuring ecological niche overlap from occurrence and spatial environmental data. *Global Ecology and Biogeography* **21**, 481-497.
- Brown WL, Wilson EO (1956) Character displacement. *Systematic Zoology* **5**, 49-64.
- Bruen TC, Philippe H, Bryant D (2006) A simple and robust statistical test for detecting the presence of recombination. *Genetics* **172**, 2665-2681.
- Burri R (2017) Linked selection, demography and the evolution of correlated genomic landscapes in birds and beyond. *Molecular Ecology* **26**, 3853-3856.
- Carson H, Clague D (1995) Geology and biogeography of the Hawaiian Islands. *Hawaiian Biogeography: Evolution on a Hot Spot Archipelago*, 14-29.
- Chapuis M-P, Estoup A (2007) Microsatellite null alleles and estimation of population differentiation. *Molecular Biology and Evolution* **24**, 621-631.
- Chen Q, Liu Y, Ho WT, Chan SK, Li QH, & Huang JR (2017) Use of stable isotopes to understand food webs in Macao wetlands. *Wetlands Ecology and Management* **25**, 59-66.

- 758 Coyne JA, Orr HA (2004) *Speciation* Sinauer Associates Sunderland, MA.
- Cramp S, Perrins C (1983) The Birds of Europe the Middle East and North Africa, Vol. III (Waders to
- 760 Gulls). *Oxford University Press, Oxford*. pp 153-166.
- Cruickshank TE, Hahn MW. (2014) Reanalysis suggests that genomic islands of speciation are due to
- 762 reduced diversity, not reduced gene flow. *Molecular Ecology* **23**,3133-3157.
- del Hoyo J, Collar N, Kirwan GMS, C.J. (2016) *White-faced Plover (Charadrius dealbatus)*. In: *del Hoyo,*
- 764 *J., Elliott, A., Sargatal, J., Christie, D.A. & de Juana, E. (eds.). Handbook of the Birds of the*
- World Alive*. Lynx Edicions, Barcelona., retrieved from <http://www.hbw.com/node/467300> on
- 766 16 June 2018
- DeNiro MJ, Epstein S (1978) Carbon isotopic evidence for different feeding patterns in two hyrax
- 768 species occupying the same habitat. *Science* **201**, 906-908.
- Ding S, Mishra M, Wu H, et al (2018) Characterization of hybridization within a secondary contact
- 770 region of the inshore fish, *Bostrychus sinensis*, in the East China Sea. *Heredity*, **120**, 51-62.
- Dos Remedios N, Lee PL, Burke T, Székely T, Küpper C (2015) North or south? Phylogenetic and
- 772 biogeographic origins of a globally distributed avian clade. *Molecular Phylogenetics and*
- Evolution* **89**, 151-159.
- 774 Earl DA (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and
- implementing the Evanno method. *Conservation Genetics Resources* **4**, 359-361.
- 776 Eberhart-Phillips LJ, Hoffman JI, Brede EG, et al. (2015) Contrasting genetic diversity and population
- structure among three sympatric Madagascan shorebirds: parallels with rarity, endemism,
- 778 and dispersal. *Ecology and Evolution* **5**, 997-1010.
- Ellstrand NC, Elam DR (1993) Population genetic consequences of small population size: implications
- 780 for plant conservation. *Annual Review of Ecology and Systematics* **24**, 217-242.
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the
- 782 software STRUCTURE: a simulation study. *Molecular Ecology* **14**, 2611-2620.
- Excoffier L, Laval G, Schneider S (2005) Arlequin (version 3.0): An integrated software package for
- 784 population genetics data analysis. *Evolutionary Bioinformatics* **1**, 47-50.
- Feder JL, Flaxman SM, Egan SP, Comeault AA, Nosil P (2013) Geographic mode of speciation and
- 786 genomic divergence. *Annual Review of Ecology, Evolution, and Systematics* **44**, 73-97.
- Fitzpatrick SW, Gerberich JC, Kronenberger JA, Angeloni LM, Funk WC (2015) Locally adapted traits
- 788 maintained in the face of high gene flow. *Ecology Letters* **18**, 37-47.
- Freedman AH, Gronau I, Schweizer RM, et al. (2014) Genome sequencing highlights the dynamic early
- 790 history of dogs. *PLoS Genetics* **10**, e1004016.
- Funk WC, Mullins TD, Haig SM (2007) Conservation genetics of snowy plovers (*Charadrius alexandrinus*)
- 792 in the Western Hemisphere: population genetic structure and delineation of subspecies.
- Conservation Genetics* **8**, 1287-1309.
- 794 García-Navas V, Bonnet T, Waldvogel D, et al. (2015) Gene flow counteracts the effect of drift in a
- Swiss population of snow voles fluctuating in size. *Biological Conservation* **191**, 168-177.
- 796 Ginn H, Melville D (1983) Molt in birds. BTO Guide 19. *British Trust for Ornithology, Tring*.
- Grant PR, Grant BR (2006). Evolution of character displacement in Darwin's finches. *Science* **313**,
- 798 224-226.
- Guillot G, Mortier F, Estoup A (2005) GENELAND: a computer package for landscape genetics.
- 800 *Molecular Ecology Notes* **5**, 712-715.
- Hammer Ø, Harper D, Ryan P (2001) PAST: Paleontological Statistics Software Package for education

and data analysis. *Palaeontologia Electronica* 4.

Harrison RG (1986) Pattern and process in a narrow hybrid zone. *Heredity* **56**, 337.

Hartert E, Jackson AC (1915) Notes on some waders. *Ibis* **57**, 526-534.

Hey, J. (2006). Recent advances in assessing gene flow between diverging populations and species. *Current Opinion in Genetics & Development* **16**, 592-596.

Hey J (2010) Isolation with migration models for more than two populations. *Molecular Biology and Evolution* **27**, 905-920.

Hey J, Nielsen R (2004) Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics* **167**, 747-760.

Hey J, Nielsen R (2007) Integration within the Felsenstein equation for improved Markov chain Monte Carlo methods in population genetics. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 2785-2790.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**, 1965-1978.

Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes I: turnover of ^{13}C in tissues. *Condor* **94**, 181-188.

Hoskin CJ, Higgie M (2010) Speciation via species interactions: the divergence of mating traits within species. *Ecology Letters* **13**, 409-420.

Hu J, Broennimann O, Guisan A, *et al.* (2016) Niche conservatism in *Gynandropaa* frogs on the southeastern Qinghai-Tibetan Plateau. *Scientific Reports* **6**, 32624.

Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with the assistance of sample group information. *Molecular Ecology Resources* **9**, 1322-1332.

BirdLife International (2016) *IUCN Red List for birds*. Downloaded from <http://www.birdlife.org> on 8/09/2016.

Jackson DU, dos Remedios N, Maher KH, *et al.* (2017) Polygamy slows down population divergence in shorebirds. *Evolution* **71**, 1313-1326.

Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**, 1801-1806.

Johnston RF (1969) Character variation and adaptation in European sparrows. *Systematic Biology* **18**, 206-231.

Küpper C, Edwards SV, Kosztolanyi A, *et al.* (2012) High gene flow on a continental scale in the polyandrous Kentish plover *Charadrius alexandrinus*. *Molecular Ecology* **21**, 5864-5879.

Küpper C, Horsburgh GJ, Dawson DA, *et al.* (2007) Characterization of 36 polymorphic microsatellite loci in the Kentish plover (*Charadrius alexandrinus*) including two sex-linked loci and their amplification in four other *Charadrius* species. *Molecular Ecology Notes* **7**, 35-39.

Kempenaers B, Valcu M (2017) Breeding site sampling across the Arctic by individual males of a polygynous shorebird. *Nature* **541**, 528.

Kennerley PR, Bakewell DN, Round PD (2008) Rediscovery of a long-lost *Charadrius* plover from South-East Asia. *Forktail* **24**, 63-79.

Lande R (1982) Rapid origin of sexual isolation and character divergence in a cline. *Evolution* **36**, 213-223.

Lanfear R, Kokko H, Eyre-Walker A (2014) Population size and the rate of evolution. *Trends in Ecology*

846 & *Evolution* **29**, 33-41.

847 Le Moan A, Gagnaire PA, Bonhomme F (2016) Parallel genetic divergence among coastal-marine

848 ecotype pairs of European anchovy explained by differential introgression after secondary

 contact. *Molecular Ecology* **25**, 3187-3202.

850 Leigh JW, Bryant D (2015) Popart: full-feature software for haplotype network construction. *Methods*

in Ecology and Evolution **6**, 1110-1116.

852 Li JW, Yeung CK, Tsai P-W, *et al.* (2010) Rejecting strictly allopatric speciation on a continental island:

 prolonged postdivergence gene flow between Taiwan (*Leucodioptron taewanus*,

854 Passeriformes Timaliidae) and Chinese (*L. canorum canorum*) hwameis. *Molecular Ecology* **19**,

 494-507.

856 Liu JX, Gao TX, Wu SF, Zhang YP (2007) Pleistocene isolation in the Northwestern Pacific marginal seas

 and limited dispersal in a marine fish, *Chelon haematocheilus* (Temminck & Schlegel, 1845).

858 *Molecular Ecology* **16**, 275-288.

 Liu Y, Keller I, Heckel G. (2012). Breeding site fidelity and winter admixture in a long-distance migrant,

860 the tufted duck (*Aythya fuligula*). *Heredity*, **109**, 108-116.

 Liu Y, Liu S, Yeh CF, Zhang N, Chen G, Que P, ... & Li SH. (2018). The first set of universal nuclear

862 protein-coding loci markers for avian phylogenetic and population genetic studies. *bioRxiv*,

 272732.

864 Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data.

Bioinformatics **25**, 1451-1452.

866 Lynch M, Ackerman MS, Gout J-F, *et al.* (2016) Genetic drift, selection and the evolution of the

 mutation rate. *Nature Reviews Genetics* **17**, 704-714.

868 Ma Z, Melville D S, Liu J, *et al.* (2014) Rethinking China's new great wall : Massive seawall construction

 in coastal wetlands threatens biodiversity. *Science*, **346**, 912-914.

870 Martin SH, Dasmahapatra KK, Nadeau NJ, *et al.* (2013) Genome-wide evidence for speciation with

 gene flow in *Heliconius* butterflies. *Genome Research* **23**, 1817-1828.

872 Mayr E (1963) *Animal species and evolution* Belknap Press of Harvard University Press Cambridge,

 Massachusetts.

874 McCormack JE, Zellmer AJ, Knowles LL (2010) Does niche divergence accompany allopatric divergence

 in *Aphelocoma* jays as predicted under ecological speciation?: insights from tests with niche

876 models. *Evolution* **64**, 1231-1244.

 McNab BK (2009) Ecological factors affect the level and scaling of avian BMR. *Comparative*

878 *Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **152**, 22-45.

 McWilliams SR, Karasov WH (2001) Phenotypic flexibility in digestive system structure and function in

880 migratory birds and its ecological significance. *Comparative Biochemistry and Physiology Part*

A: Molecular & Integrative Physiology **128**, 577-591.

882 Miller MP, Haig SM, Mullins TD, Ruan L, *et al.* (2015) Intercontinental genetic structure and gene flow

 in Dunlin (*Calidris alpina*), a potential vector of avian influenza. *Evolutionary Applications*, **8**,

884 149-171.

 Morales AE, Jackson ND, Dewey TA, O'Meara BC, Carstens BC (2017) Speciation with gene flow in

886 North American *Myotis* bats. *Systematic Biology* **66**, 440-452.

 Nadachowska-Brzyska K, Burri R, Olason PI, *et al.* (2013) Demographic divergence history of pied

888 flycatcher and collared flycatcher inferred from whole-genome re-sequencing data. *PLoS*

Genetics **9**, e1003942.

890 Nadeau NJ, Whibley A, Jones RT, *et al.* (2012) Genomic islands of divergence in hybridizing *Heliconius*
butterflies identified by large-scale targeted sequencing. *Phil. Trans. R. Soc. B* **367**, 343-353.

892 Nebel S, Rogers KG, Minton CD, Rogers DI (2013) Is geographical variation in the size of Australian
shorebirds consistent with hypotheses on differential migration? *Emu* **113**, 99-111.

894 Ni G, Li Q, Kong LF, Zheng XD (2012) Phylogeography of bivalve *Cyclina sinensis*: testing the historical
glaciations and Changjiang River outflow hypotheses in northwestern Pacific. *PLoS ONE* **7**,
896 e49487.

Ni G, Li Q, Kong LF & Yu Hong (2014) Comparative phylogeography in marginal seas of the
898 northwestern Pacific, *Molecular Ecology*, **23**, 534-548.

Nielsen EE, Bach LA, Kotlicki P (2006) HYBRIDLAB (version 1.0): a program for generating simulated
900 hybrids from population samples. *Molecular Ecology Notes* **6**, 971-973.

Nosil P (2007) Divergent host plant adaptation and reproductive isolation between ecotypes of
902 *Timema cristinae* walking sticks. *The American Naturalist* **169**, 151-162.

Nosil P, Feder JL (2012) Genomic divergence during speciation: causes and consequences. *Phil. Trans.*
904 *R. Soc. B*, **367**, 332-342.

Nosil P, Schluter D (2011) The genes underlying the process of speciation. *Trends in Ecology &*
906 *Evolution* **26**, 160-167.

Pagani-Núñez E, Renom M, Mateos-Gonzalez F, Cotín J, Senar JC (2017) The diet of great tit nestlings:
908 Comparing observation records and stable isotope analyses. *Basic and Applied Ecology* **18**,
57-66.

910 Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic software for
teaching and research--an update. *Bioinformatics* **28**, 2537-2539.

912 Peters JL, McCracken KG, Pruett CL, *et al.* (2012) A parapatric propensity for breeding precludes the
completion of speciation in common teal (*Anas crecca*, sensu lato). *Molecular Ecology* **21**,
914 4563-4577.

Phillips SJ, Dudik M, Elith J, *et al.* (2009) Sample selection bias and presence-only distribution models:
916 implications for background and pseudo-absence data. *Ecological Applications* **19**, 181-197.

Poelstra JW, Vijay N, Bossu CM, *et al.* (2014) The genomic landscape underlying phenotypic integrity in
918 the face of gene flow in crows. *Science* **344**, 1410-1414.

Primmer C, Møller A, Ellegren H (1995) Resolving genetic relationships with microsatellite markers: a
920 parentage testing system for the swallow *Hirundo rustica*. *Molecular Ecology* **4**, 493-498.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus
922 genotype data. *Genetics* **155**, 945-959.

Procházka P, Stokke BG, Jensen H, *et al.* (2011) Low genetic differentiation among reed warbler
924 *Acrocephalus scirpaceus* populations across Europe. *Journal of Avian Biology* **42**, 103-113.

Que P, Chang Y, Eberhart-Phillips L, *et al.* (2015) Low nest survival of a breeding shorebird in Bohai Bay,
926 China. *Journal of Ornithology* **156**, 297-307.

R Development Core Team (2013) *R: a language and environment for statistical computing*. R
928 Foundation for Statistical Computing, Vienna, Austria. <http://www.rproject.org/>

Railsback LB, Gibbard PL, Head MJ, Voarintosa NRG, Toucanne S (2015) An optimized scheme of
930 lettered marine isotope substages for the last 1.0 million years, and the climatostrati- graphic
nature of isotope stages and substages. *Quaternary Science Reviews* **111**, 94-106.

932 Redfern CP, Clark JA (2001) *Ringers' Manual*. British Trust for Ornithology.

Rheindt FE, Edwards SV (2011) Genetic introgression: an integral but neglected component of speciation in birds. *Auk* 128:620–632

Rheindt FE, Szekely T, Edwards SV, *et al.* (2011) Conflict between genetic and phenotypic differentiation: the evolutionary history of a 'lost and rediscovered' shorebird. *PLoS ONE* 6, e26995.

Robinson BW, Wilson DS (1994) Character release and displacement in fishes: a neglected literature. *The American Naturalist* 144, 596-627.

Rundle HD, Nosil P (2005) Ecological speciation. *Ecology Letters* 8, 336-352.

Rybicki J, Sirkkälä PM, McFarlane SE, *et al.* (2016) Competition driven buildup of habitat isolation and selection favoring modified dispersal patterns in a young avian hybrid zone. *Evolution* 70, 2226-2238.

Salewski V, Watt C (2017) Bergmann's rule: a biophysiological rule examined in birds. *Oikos* 126.

Schluter D (2009) Evidence for ecological speciation and its alternative. *Science* 323, 737-741.

Seehausen O, Butlin RK, Keller I, *et al.* (2014) Genomics and the origin of species. *Nature Reviews Genetics* 15, 176-192.

Sethuraman A, Hey J (2016) IMa2p—parallel MCMC and inference of ancient demography under the isolation with migration (IM) model. *Molecular Ecology Resources* 16, 206-215.

Slatkin M (1973) Gene flow and selection in a cline. *Genetics* 75, 733-756.

Stephens M, Donnelly P (2003) A comparison of bayesian methods for haplotype reconstruction from population genotype data. *American Journal of Human Genetics* 73, 1162-1169.

Stephens M, Smith NJ, Donnelly P (2001) A new statistical method for haplotype reconstruction from population data. *The American Journal of Human Genetics* 68, 978-989.

Storz JF (2002) Contrasting patterns of divergence in quantitative traits and neutral DNA markers: analysis of clinal variation. *Molecular Ecology* 11, 2537-2551.

Swanson HK, Lysy M, Power M, *et al.* (2015) A new probabilistic method for quantifying n-dimensional ecological niches and niche overlap. *Ecology* 96, 318-324.

Swinhoe R (1870) On the plovers of the genus Aegialites found in China. *Proceedings of the Zoological Society London* 1870, 136–142.

Székely T, Kosztolányi A, Küpper C (2008) Practical guide for investigating breeding ecology of Kentish plover. (available at http://www.bath.ac.uk/bio-sci/biodiversity-lab/pdfs/KP_Field_Guide_v3.pdf).

Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123, 585-595.

Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30, 2725-2729.

Tattersall GJ, Arnaout B, Symonds MR (2016) The evolution of the avian bill as a thermoregulatory organ. *Biological Reviews* 92, 1630-1652.

Taylor SA, White TA, Hochachka WM, *et al.* (2014) Climate-mediated movement of an avian hybrid zone. *Current Biology* 24, 671-676.

Taylor SA, Larson EL, Harrison RG (2015) Hybrid zones: windows on climate change. *Trends in Ecology and Evolution* 30, 398–406.

Trimbos K B, Musters C J M, Verkuil Y I, *et al.* (2011) No evident spatial genetic structuring in the rapidly declining Black-tailed Godwit *Limosa limosa limosa*, in The Netherlands. *Conservation*

976 *Genetics* **12**, 629-636.

977 Toews DPL, Taylor SA, Vallender R, Brelsford A, Butcher BG, Messer PW, Lovette IJ. (2016) Plumage

978 genes and little else distinguish the genomes of hybridizing warblers. *Current Biology* **26**,

 2313–2318.

980 Vähä J, Primmer C (2006) Efficiency of model-based Bayesian methods for detecting hybrid

 individuals under different hybridization scenarios and with different numbers of loci.

982 *Molecular Ecology* **15**, 63-72.

 Wagner CE, Harmon LJ, Seehausen O (2012) Ecological opportunity and sexual selection together

984 predict adaptive radiation. *Nature* **487**, 366-369.

 Wang C, Li C, & Li S. (2008) Mitochondrial DNA-inferred population structure and demographic history

986 of the mitten crab (*Eriocheir sensu stricto*) found along the coast of mainland

 China. *Molecular Ecology* **17**, 3515-3527.

988 Wang J, Tsang LM, Dong YW (2015) Causations of phylogeographic barrier of some rocky shore species

 along the Chinese coastline. *BMC Evolutionary Biology* **15**, 114.

990 Weir JT, & Schluter D. (2008). Calibrating the avian molecular clock. *Molecular Ecology*, **17**, 2321-2328.

 Winkelmann K, Genner MJ, Takahashi T, Rüber L (2014) Competition-driven speciation in cichlid fish.

992 *Nature Communications* **5**, 3412.

 Winker K, McCracken KG, Gibson DD, Peters JL (2013) Heteropatric speciation in a duck, *Anas crecca*.

994 *Molecular Ecology* **22**, 5922-5935.

 Wolf JB, Ellegren H (2017) Making sense of genomic islands of differentiation in light of speciation.

996 *Nature Reviews Genetics* **18**, 87.

 Won YJ, Hey J (2005) Divergence population genetics of chimpanzees. *Molecular Biology and Evolution*

998 **22**, 297-307.

 Yang Z, Rannala B. (2014) Unguided species delimitation using DNA sequence data from multiple

1000 loci. *Molecular Biology and Evolution* **31**, 3125-3135.

 Zhang G, Li C, Li Q, *et al.* (2014) Comparative genomics reveals insights into avian genome evolution

1002 and adaptation. *Science* **346**, 1311-1320.

 Zhou Y, Duvaux L, Ren G, *et al.* (2016) Importance of incomplete lineage sorting and introgression in

1004 the origin of shared genetic variation between two closely related pines with overlapping

 distributions. *Heredity* **118**, 211-220.

1006

FIGURE LEGENDS

FIGURE 1 Sampling localities, morphology and trophic level differentiation of the Kentish Plover *Charadrius alexandrinus* (blue) and the White-faced Plover *C. dealbatus* (yellow).

(a) Samples were from an inland site (1, Qinghai Lake), 16 Chinese mainland coastal localities (2, Tangshan; 3, Cangzhou; 4, Weifang; 5, Lianyungang; 6, Nantong; 7, Ningbo; 8, Zhoushan; 9, Wenzhou; 12, Fuzhou; 13, Xiamen; 14, Jinmen; 15, Shanwei; 16, Yangjiang; 17, Zhanjiang; 18, Beihai and 19, Dongfang), and two localities on Taiwan Island (10, Xinbei and 11, Zhanghua).

(b-f) Differences between two species in several characters. *C. alexandrinus* has on average shorter bill length (except for Taiwan populations) **(b)**, wing length **(c)** and lower body mass (except for Taiwan populations) **(d)** than *C. dealbatus*. Localities with less than five measured adults were excluded in these analyses. **(f)** Plots of the first two principal components and their associated variance explained which showed the subtle morphometric differences between the two species. **(e)** *C. alexandrinus* has overall more darkish plumage than *C. dealbatus* and males of the latter species have less black tinged on face and neck during breeding season. **(g)** Pairwise morphological difference Q_{ST} plotted against geographical distance between coastal populations of *C. alexandrinus* and *C. dealbatus*. Q_{ST} values within species were marked by blue or yellow dots. Q_{ST} between species (grey circles) declined as the distance increased (regression showed as the dashed line).

FIGURE 2 Haplotype networks based on mitochondrial and nuclear DNA sequences, and population genetic structures based on microsatellite loci. *C. alexandrinus* marked in blue and *C. dealbatus* in yellow. **(a)** Haplotype networks of three mitochondrial loci. For ATPase6/8 and ND3, over 95% of the individuals are sorted into two major haplogroups. **(b)** Examples of haplotype networks based on 55 to 80 individuals of four exonic nuclear loci. **(c)** Genetic clustering inferred with Geneland based on microsatellite genotypes. Blue and yellow show the assignment probability of a

1038 location for alternative genetic clusters. Location numbers are consistent with Figure 1.

(d) Genetic clustering inferred with STRUCTURE using microsatellites.

1040

FIGURE 3 Posterior densities of population demographic parameters estimated using the isolation-with-migration model (IM) implemented in IMa2p. These analyses used the combined data set of three mitochondrial (1729 bp) and 15 nuclear exonic loci (11209 bp). K represents the Kentish Plover *C. alexandrinus*; W represents White-faced Plover *C. dealbatus*. **(a)** Population divergence times (T) of *C. alexandrinus* and *C. dealbatus*. **(b)** Population migration rates ($2NM$) stand for the average number of individuals in each group that were migrants from the other group in the past. Gene flow on both directions was significant. **(c)** Effective population sizes (N_e) of the two species and their most recent common ancestor.

1050

FIGURE 4 Predicted environmental suitability for *C. alexandrinus* (left) and *C. dealbatus* (right), via ecological niche modeling (ENM). ENM results are shown for the Last Interglacial (LIG, 120-100Ka), Last Glacial Maximum (LGM, 21Ka, MIROC model) and current times, respectively.

FIGURE 5 (a) Raw data (bottom-left), density distribution (top-left and bottom-right), and stable isotope related to dietary niche regions (top right) of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of *C. alexandrinus* and *C. dealbatus* feathers generated by “nicheROVER” (Swanson *et al.* 2015). The niche region of *C. alexandrinus* (blue) was broader than and contained that of *C. dealbatus* (yellow). **(b)** Distribution of the posterior probability that an individual from one species is found within the niche region of the other species. Vertical lines for mean and 95% credible intervals are included in the histogram of each overlap metric.

1064

FIGURE 1

1066

1068

FIGURE 2

1070

FIGURE 3

1072

1074

FIGURE 4

1076

1078

FIGURE 5

1080

1082

TABLE 1 Sampling localities and genetic polymorphism of *C. alexandrinus* and *C.*

1084 *dealbatus*. The number of individuals (*n*) analyzed for mtDNA, autosomal
microsatellites and nuclear exonic loci (nuDNA) are given. Site number corresponds to
1086 the numbers in Figure 1. Estimates of *h*, number of haplotypes; *Hd*, haplotype diversity;
 π , nucleotide diversity; *Tajima's D* value, *Ho*, observed heterozygosity; *He*, expected
1088 heterozygosity were calculated for each locality and species.

Site	Latitude	Longitude	mtDNA						Microsatellites			nuDNA	
			<i>n</i>	<i>h</i>	<i>Hd</i>	π	<i>Tajima's D</i>	<i>n</i>	<i>Ho</i>	<i>He</i>	<i>n</i>	<i>Tajima's</i>	
												<i>D</i>	
<i>C. alexandrinus</i>			224	86	0.929	0.00181	-1.863*	219	0.723	0.802	30-40	0.216	
1	Qinghai Lake	36.693	30	13	0.846	0.00119	-1.785	30	0.703	0.756	13-22		
2	Tangshan	39.169	23	16	0.945	0.00171	-1.434	30	0.797	0.800	12-20		
3	Cangzhou	38.471	14	8	0.868	0.00097	-0.871	16	0.774	0.791			
4	Weifang	37.131	21	14	0.895	0.00158	-1.687	22	0.671	0.787			
5	Lianyungang	34.742	28	20	0.963	0.00189	-1.166	30	0.697	0.777			
6	Nantong	32.575	25	12	0.913	0.00185	-0.016	26	0.731	0.796			
7	Zhoushan	29.991	19	11	0.924	0.00111	-0.141	20	0.746	0.779			
8	Ningbo	30.205	25	18	0.963	0.00179	-0.958	29	0.751	0.793			
9	Wenzhou	27.937	14	13	0.989	0.00161	-1.048	16	0.740	0.768			
10	Xinbei	25.862	14	6	0.736	0.00108	-0.971	30	0.672	0.700	2-8		
11	Zhanghua	24.559	11	5	0.709	0.00076	-0.893	22	0.692	0.669	4-8		
<i>C. dealbatus</i>			133	29	0.667	0.00086	-1.989*	131	0.714	0.758	23-28	0.326	
12	Jinmen	24.448	13	7	0.731	0.00110	-1.362	15	0.733	0.703	5-8		
13	Fuzhou	25.190	29	12	0.874	0.00108	-1.445	15	0.718	0.761			
14	Xiamen	24.162	18	6	0.699	0.00075	-1.532	12	0.755	0.748			
15	Shanwei	22.798	15	3	0.590	0.00039	0.221	23	0.669	0.696			
16	Yangjiang	21.589	10	5	0.822	0.00158	-0.144	11	0.713	0.718			
17	Zhanjiang	20.238	18	7	0.569	0.00099	-1.696	20	0.719	0.721			
18	Beihai	21.426	18	4	0.314	0.00044	-1.849*	23	0.692	0.695	18-12		
19	Dongfang	19.234	12	2	0.167	0.00019	-1.141	12	0.763	0.679			
All			357	109	0.922	0.00216	-1.741	402	0.719	0.780	55-80	0.271	

TABLE 2 Hierarchical analyses of molecular variance (AMOVA) based on

concatenated mtDNA data and 13 microsatellite loci for *C. alexandrinus* and *C. dealbatus*. Samples were partitioned in three groupings: 2 groups (*C. alexandrinus* and *C. dealbatus*); 3 groups (*C. alexandrinus* continental populations and Taiwan island populations, *C. dealbatus*); 4 groups (*C. alexandrinus* inland (Qinghai Lake), *C. alexandrinus* coastal, *C. alexandrinus* Taiwan island, and *C. dealbatus*). *Va*, genetic variation among groups; *Vb*, variation among populations within groups; *Vc*, variation within populations. Between-group genetic differentiation was highest when populations were partitioned into two groups (*C. alexandrinus* and *C. dealbatus*).

	Grouping	<i>Va</i> %	<i>Vb</i> %	<i>Vc</i> %
mtDNA	2	49.712	8.491	41.797
	3	45.695	9.269	45.036
	4	41.782	10.296	47.922
microsatellites	2	2.416	1.798	95.786
	3	2.192	1.749	96.059
	4	1.917	1.776	96.307

All $p < 0.001$

1104 **TABLE 3** Posterior mode, mean and range of 95% highest probability distribution
 (HPD) of six demographic parameters inferred with IMA2p between *C. alexandrinus*
 1106 and *C. dealbatus*. Divergence times (*T*) are given in million years ago (Ma). The
 effective population size (*Ne*) of *C. alexandrinus* was about eight times that of *C.*
 1108 *dealbatus*. Migration rate (*2NM*) into each species was about the same. K represents
 Kentish Plover *C. alexandrinus*; W represents White-faced Plover *C. dealbatus*; A is
 1110 the most recent common ancestor of two species.

Parameter	<i>T</i> /Ma	<i>Ne_W</i> /10 ⁶	<i>Ne_K</i> /10 ⁶	<i>Ne_A</i> /10 ⁶	<i>2NM_{K→W}</i>	<i>2NM_{W→K}</i>
Mode	0.557	0.259	2.036	0.495	2.696*	2.005*
Mean	2.160	0.297	2.450	0.589	2.859	4.306
2.5% HPD	0.413	0.154	1.590	0.134	1.581	0.528
97.5% HPD	5.187	0.515	4.442	1.098	4.503	11.500

* *p* < 0.001

1112

1114