1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Genome-wide identification and expression specificity analysis of the DNA methyltransferase gene family under adversity stresses in cotton

Xiaomin Yang¹, Xuke Lu, Xiugui Chen, Delong Wang, Junjuan Wang, Shuai Wang, Lixue Guo,
Chao Chen, Xiaoge Wang, Binglei Zhang, Mingge Han, Wuwei Ye*

¹ Key Laboratory for Cotton Genetic Improvement, Anyang 455000, Henan, China
Institute of Cotton Research of Chinese Academy of Agricultural Sciences
State Key Laboratory of Cotton Biology

* corresponding author, Wuwei Ye
E-mail: yew158@163.com

- 23 WY contributed equally to the experiment.
- 24 XY, XL, XW,BZ and MH participated in the design of the study and performed the statistical analysis.
- 25 JW, CC,LG,XC, DW and SW provided the materials and revised the manuscript.
- 26 XY drafted the manuscript.
- 27

28 Abstract

29 DNA methylation is an important epigenetic mode of genomic DNA modification that is an
30 important part of maintaining epigenetic content and regulating gene expression. DNA
31 methyltransferases (MTases) are the key enzymes in the process of DNA methylation. Thus far,
32 there has been no systematic analysis the DNA MTases found in cotton. In this study, the whole
33 genome of cotton C5-Mtase coding genes was identified and analyzed using a bioinformatics
34 method based on information from the cotton genome. In this study, 51 DNA MTase genes were
35 identified, of which 8 belonged to *G. raimondii* (group D), 9 belonged to *G. arboreum* L. (group
36 A), 16 belonged to *G. hirsutum* L. (group AD₁) and 18 belonged to *G. barbadense* L. (group
37 AD₂). Systematic evolutionary analysis divided the 51 genes into four subfamilies, including 7
38 MET homologous proteins, 25 CMT homologous proteins, 14 DRM homologous proteins and 5
39 DNMT2 homologous proteins. Further studies showed that the DNA MTases in cotton were more
40 phylogenetically conserved. The comparison of their protein domains showed that the C-terminal
41 functional domain of the 51 proteins had six conserved motifs involved in methylation
42 modification, indicating that the protein has a basic catalytic methylation function and the
43 difference in the N-terminal regulatory domains of the 51 proteins divided the proteins into four
44 classes, MET, CMT, DRM and DNMT2, in which DNMT2 lacks an N-terminal regulatory
45 domain. Gene expression in cotton is not the same under different stress treatments. Different
46 expression patterns of DNA MTases show the functional diversity of the cotton DNA
47 methyltransferase gene family. VIGS silenced *Gossypium hirsutum* L. in the cotton seedling of
48 DNMT2 family gene *GhDMT6*, after stress treatment the growth condition was better than the
49 control. The distribution of DNA MTases varies among cotton species. Different DNA MTase
50 family members have different genetic structures, and the expression level changes with different
51 stresses, showing tissue specificity. Under salt and drought stress, *G. hirsutum* L. TM-1 increased
52 the number of genes more than *G. raimondii* and *G. arboreum* L. *Shixiya* 1. The resistance of
53 *Gossypium hirsutum* L. TM-1 to cold, drought and salt stress was increased after the plants were
54 silenced with *GhDMT6* gene.

Introduction

DNA methylation is the process of transferring a methyl (-CH₃) group to a specific base of a DNA molecule and is catalyzed by DNA MTases, with S-adenosine methionine (SAM) as a methyl donor [1, 2]. DNA methylation widely occurs in the epigenetics of bacteria, plants, and animals and is involved in transposons [3-6], the suppression of gene silencing, genomic imprinting [7], X chromosome inactivation [8], cell differentiation [9], and embryo development [10]. However, DNA methylation is not immutable; under conditions of stress, the plant genome can overcome the limitation of genome instability through DNA methylation by rapid modification. This induces the expression of some genes associated with stress to maintain plant growth and development and evolutionary process [11-13]. Therefore, epigenetic modification precedes genomic evolution in response to adversity, and DNA methylation is considered the molecular response mechanism of plants in the face of adverse stress [5, 6, 11].

DNA methylation occurs mostly in CpGs at carbon 5 in cytosine (C5). It primarily occurs in symmetric sequence CGs but also occurs in CHG and CHH (H=A, C or T) sequences [14]. There are two DNA methylation methods in plants: maintenance methylation and denovo methylation [15]. Maintenance methylation refers to the methylation of a chain of double-stranded DNA molecules through semi-reserved replication, which is passed to the offspring by the parent methylation mode of another chain without methylation. Denovo methylation is a type of DNA methylation that occurs when different DNA MTases catalyze two strands of DNA without methylation [16]. C5-Mtases in plants fall into four categories, MTase (MET), chromomethylase (CMT), domains rearranged MTase (DRM), and Dnmt2 [17, 18]. MET is mainly used in methylation of the heterochromatin region of the CG site of the symmetric sequence, which is a very important part of the methyltransferase [19]. CMT is a specific type of DNA MTases that maintains CHG and CHH site methylation and, to a certain extent plays a role in stabilizing the heterochromatin state of the genome [20]. The function of DRM is to catalyze the methylation of cytosine and to maintain the cytosine methylation of non-CpG sites under the guidance of RNA [21]. DRM is homologous to the DNMT3 of animals [22, 23]. The proteins encoded by the plant Dnmt2 family are very similar to those of mice, bacteria and yeast C5-Mtases, but their mechanism of action in the process of C5 methylation has not been clarified [18].

Cytosine-5 DNA MTases were discovered in 1925 by Robert D. Coghill [24]. DNA methylation catalyzed by MTases was observed in bovine thymus in 1948 by Hotchkiss [25]. In 1964, Gold and Hurwitz identified the first DNA MTases in *Escherichia coli* [26]. Besto identified the first plant DNA MTases, and MET was isolated from *Arabidopsis thaliana*. Its encoded protein, AtMET1, had high homology with the methylation enzyme Dnmt1 in mice [27]. The reaction mechanism of DNA methylation was first explained in prokaryotic organisms in 1993 by Finnegan [28]. DNA MTases were identified in *Arabidopsis thaliana*, rice, maize, solanaceae, Tobacco, Legumes and other plants [22, 29-32].

Materials and methods

Identification of cotton DNA MTases family members

The cotton genome information was downloaded from CottonGen (<https://www.cottongen.org/>), and the DNA-methylase structure domain (PF00145) was downloaded from the Pfam (<http://pfam.xfam.org/>) database (IPR001525). The DNA-methylase.hmm Hidden Markov model was constructed with DNA-methylase.hmm as the reference HMMER3.0 (<http://hmmmer.org/download.html>). The cotton genome database was queried to obtain the gene location and name of candidate protein family members containing DNA-methylase structure domains in cotton and to obtain GFF (general feature format) files from genome annotation files. Then, we obtained the gene position on the chromosome and used local BLAST 2.2.31+ to obtain the CDS sequence and protein sequence of the corresponding gene and obtained the whole sequence of the gene corresponding to the genome based on its position on the chromosome. The gene protein sequence is in the Smart software. The Pfam30.0 database is analyzed to ensure that each candidate gene contains a DNA-methylase structure domain. Subcellular location prediction was performed on cello. The ProtParam was obtained by protein analysis.

Cotton DNA MTases gene structure, and evolutionarily conserved protein domain analysis

The CDS sequence of cotton DNA MTase genes and the corresponding genome-wide sequence in GSDS2.0 (<http://gsds.cbi.pku.edu.cn/>) were used to map the gene structure. The software Smart (<http://smart.embl-heidelberg.de/>) was used to determine the protein conserved structural domains.

Phylogenetic analysis

Cotton DNA-methylase (PF00145, IPR001525) was used as the key word in Phytozome v12.1 (<https://phytozome.jgi.doe.gov/pz/portal.html>) in the database rather than the homologous sequences of other species (file 2). Clustal W software was used to analyze the amino acid sequence alignment. MEGA7.0 software was used to construct the phylogenetic tree with the neighbor-joining method. The number of bootstraps was 1000.

Expression pattern analysis of cotton DNA MTases under stresses

The phytotron sand culture cultivation method was used for *G. raimondii* and *G. arboreum* L. *Shixiya* 1 (16h light / 8h dark, day 28°C, night 25°C). *G. hirsutum* L. TM-1 processing salt (200mM) and PEG6000 (20%) were used at the three-leaf stage at 0h, 1h, 3h, 6h, and 12h. Total RNA was extracted from root, stem, and leaf samples and reverse transcribed into cDNA. The primers for the real-time fluorogenic quantitative PCR were designed with the NCBI-line primer design tool primer-BLAST (<https://www.ncbi.nlm.nih.gov/tools/primer-blast/>) (file 1). The RNA

was reverse transcribed to cDNA samples as a template for quantitative PCR experiments using *G. raimondii*, *G. arboreum* L. *Shixiya* 1, and *G. hirsutum* L. TM-1 to determine the gene expression of DNA MTases. The reaction conditions were 94°C for 30s; 40 cycles of 94°C denaturation for 5s, 50°C annealing for 34s, extension at 72°C for 10s, and a final extension for 34s at 72°C.

PYL156:VIGS silencing and Relative expression of GhDMT6 gene

The plasmid pYL156 was digested by EcoR and Xma I, and the digested product was detected by 1.2% agarose gel. The carrier segment was recycled. In-Fusion technique was used to insert the VIGS silencing fragment into the vector pYL156, and then transferred into DH5a. The positive clones were selected. PCR detection and sample sequencing were carried out to obtain the correct monoclonal. The vector pYL156:*GhDMT6* was successfully constructed.

TM-1(*Gossypium hirsutum* L.) was cultured in the artificial climate chamber by sand culture method. After about 5 days of seedling emergence, the preserved *Agrobacterium tumefaciens* were resuscitated and transferred to 60 ml LB liquid medium. The cultured *Agrobacterium tumefaciens* were shaken to OD₆₀₀=1.5, centrifuged for 5 minutes with 5 000 rpm of bacterial liquid, and the supernatant was discarded. Using 45ml(10mM MES+10 mMgCl₂+200 AS) resuspension to culture the thallus. In order to remove a small amount of antibiotics, repeat the operation, then stewing the thallus in 25°C for 4h.

After mixing the suspension solution containing pYL156:*GhDMT6*, pYL156 and pYL156:CLA1 with the suspension solution containing auxiliary carrier pYL192 isopyknic, the cotton was prepared to be infected. A little pore was stabbed at the back of cotyledon with a sterile needle, and suspension solution injected into the cotyledon spread over the whole cotyledon. After inoculation, the cotton seedlings were put back into the artificial climate chamber for dark culture at 23°C for 24h, and then cultured at 23°C with other conditions unchanged. When the albinism symptoms of the positive control seedlings were obvious, the cotton without infection and pYL156:*GhDMT6* were treated with 200 mM NaCl, 4°C.

Results

Genome-wide identification of cotton DNA MTases family members

A total of 51 DNA MTase members were identified from the whole genome of cotton. Group A had 8 DNA MTases and group D had 9 DNA MTases, which were named *GaDMT1*- *GaDMT8* and *GrDMT1*- *GrDMT9*, respectively, according to their sequence on the chromosomes. Similarly, 16 DNA MTases were identified in the AD₁ group, named *GhDMT1*- *GhDMT16*, and 18 DNA

MTases were identified in the AD₂ group, named *GbDMT1*- *GbDMT18*. Most of the DNA MTases in the four cotton species are located on the chromosome. There are 161-1577 different amino acids, and most contain 300-800 amino acid residues. Because of the difference in the regional gene structure of the N-terminal, *GbDMT3* was up to 1577 amino acids, whereas *GbDMT17* contains only 161 amino acids. The structure of the gene is composed of 118-402 different amino acids, and the theoretical pI (PI) ranges from 4.67 to 9.24. The predicted subcellular localization shows that most DNA MTases are located in the cytoplasm, but some are located in the outer membrane. *GbDMT13* is predicted to be periplasmic (table 1).

Multi-sequence alignment and evolutionary analysis

The DNA MTases in cotton have a C-terminal MTase catalytic region structure domain and a specific N-terminal domain, which is consistent with those of Arabidopsis, rice, maize, solanaceae, Tobacco, Legumes and other crops [22, 29-32]. To evaluate the evolution of DNA MTases in A, D, AD₁ and AD₂, multiple sequence alignment was performed and a phylogenetic tree was constructed for 51 members of the DNA MTase family (Fig. 1A). The DNA MTases in cotton are divided into four subfamilies, namely, CMT, MET, DRM, and Dnmt2 [33]. Among these 51 DNA MTases, the CMT subfamily contains 25 members, with 4, 5, 9, and 7 members in the D, A, AD₁, and AD₂ groups, respectively. There are three different types of CMTs: CMTa has 9 members, with 1, 2, 3, and 3 members in the D, A, AD₁, and AD₂ groups, respectively; CMTb has 6 members, with 1, 1, 2, and 2 members in the D, A, AD₁, and AD₂ groups, respectively; and CMTc has 10 members, with 2, 2, 4, and 2 members in the D, A, AD₁, and AD₂ groups, respectively. MET has 7 members in the D, A, AD₁, and AD₂ groups. DRM has 14 members, with 2, 2, 4, and 6 members in the D, A, AD₁, and AD₂ groups, respectively. There are 5 members in Dnmt2, with 1, 1, 1, and 2 members in the D, A, AD₁, and AD₂ groups, respectively.

Genetic structure and protein domains of DNA MTases in cotton

Gene structure analysis is an important method in the study of genetic evolution. The number of introns and exons in DNA MTase family members in groups D, A, AD₁ and AD₂ were analyzed, and a DNA MTase gene structure for cotton was created (Fig. 1B). The results showed that the numbers of exons in different MTase genes in cotton were very different; the *GbDMT14* gene had the fewest exons, with only 2 exons, whereas *GrDMT8*, *GhDMT9*, *GhDMT11* and *GhDMT14* had 24 exons. The Dnmt2 family contains only 10 exons, the Met family contains 10-12 exons, the CMTa family contains 5-22 exons, the CMTb family contains 19-24 exons, the CMTc family contains 9-24 exons, and the DRM family contains only 2-20 exons.

Motif analysis of the 51 DNA MTase proteins in cotton is shown in Fig. 2 The C-terminal catalytic region has 6 highly conserved motifs: motif x and motif i for Sam-binding sites; motif iv, motif vi, motif viii, and motif ix are the c5-Mtase functional catalytic sites, of which motif iv is the active site, motif vi is the target cytosine binding site. motif viii is the DNA neutralization region, and motif ix is the target sequence location identification area, which is consistent with the related literature [22]. The MTases of cotton DNA have different motif orders. The motif order of

the DRM family is X, VI, VIII, IX, I, and IV, and the order for the DNMT2 family is I, x, vi, viii, iv, and ix. There are three motif orders in CMT: The first is I, iv, vi, x, viii, and ix; the second is IX, i, iv, vi, x, and viii; and the third is VI, i, iv, x, viii, and ix.

The relationship between the cotton DNA MTase family and DNA MTases in other crops

Phylogenetic trees are used to reveal the homologous and evolutionary relationships of DNA MTase families from different species. To show the evolutionary relationship between the members of the cotton DNA MTase family and those of Arabidopsis, cocoa, Medicago, rice, Chlamydomonas reinhardtii and other crops, the amino acid sequences of DNA MTase family members in several crops were compared (table 2) to the DNA of the MET, DRM, and CMT family members. In monocotyledons, the differentiation of DNA MTases was separate from the evolution of the dicotyledons, and the Dnmt2 family did not differentiate in the monocotyledons, which showed that the Dnmt2 family was highly conserved during evolution. The DNA MTases in cotton are closer to those in cocoa in various branches, suggesting that they have similar functions. (Note: The VvDMT8, VvDMT9, ZmDMT7, and StDMT5 sequences are shorter and cannot be used in the construction of the inter-species evolutionary tree.)

Cotton gene expression analysis of different DNA MTases under stress

To study the expression patterns of DNA MTase genes of cotton in different tissues under salt and drought stress, the *G. raimondii*, *G. arboreum* L. Shixiya 1, and *G. hirsutum* L. TM-1 were developed to the trefoil stage, and real-time quantitative PCR was performed. The results showed that the three cotton species had different expression patterns under different stress conditions, and *G. raimondii*, *G. arboreum* L. Shixiya 1, and *G. hirsutum* L. TM-1 had obvious tissue differences when treated with PEG6000 and NaCl. *G. arboreum* L. Shixiya 1 expressed more genes in the cut root and stem under salt and drought stress. The leaf also had more genes, and with different treatments, the gene expression was also different. *GaDMT7* and *GaDMT4* decreased in the stem under salt treatment, and drought treatment also decreased their expression. *G. raimondii* also has organizational specificity. Gene expression was decreased in the rhizome and increased in the leaf. *GrDMT5* decreases in the stem but increases in the root performance under the salt treatment, showing that the same gene is expressed differently in various tissues. *GrDMT1* is decreased in leaves under drought treatment; however, salt treatment increases its expression. For *G. hirsutum* L. TM-1 and the above two cotton varieties, the genes are mainly distributed in the root of the rhizome. Additionally, the leaf has more genes. *GhDMT9* is expressed in the root under salt treatment. In the leaf, expression first increases and then decreases.

Functional analysis of GhDMT6 gene in cotton

After 6 days of natural drought, the plant phenotypes were significantly different. The plants

injected with pYL156 and those of the wild type were yellow and withered, seriously dehydrated, and the plants died. The plants injected with pYL156:*GhDMT6* had no true leaf blight and water loss (Fig. 5a). The relative expression level of *GhDMT6* gene in pYL156:*GhDMT6* infected cotton seedlings was detected. The figure shows that the expression level of *GhDMT6* gene in pYL156 infected cotton seedlings did not change under different stresses. Compared with the control, the transcription level of *GhDMT6* gene in pYL156 infected cotton seedlings decreased the most in the stem, followed by the leaves, and decreased the least in the roots (Fig. 5d).

About 15 days after VIGS infection, the positive control plants showed obvious bleaching and other indexes were normal. Using no infection and pYL156 infection of cotton seedlings as control, the silent plants were treated with cold (4°C), drought (natural drought), salt (200 mM NaCl). After 36h of cold treatment, the phenotypic differences were obvious (Fig. 5b). The plants injected with pYL156 and wild-type true leaf wilted and curly, and the plants injected with pYL156:*GhDMT6* gene grew normally without phenotypic changes. The expression level of *GhDMT6* gene in pYL156 infected cotton did not change under different stresses, and the transcriptional level of *GhDMT6* gene in pYL156:*GhDMT6* infected cotton was significantly reduced compared with the control. Leaf blade decreased the most, followed by root, stem the least (Fig. 5e).

The phenotype of cotton seedlings was significantly different after 3 days, which were treated with 200mM NaCl. The seedlings of pYL156 and wild-type cotyledons were exfoliated, the leaf edges of true leaves were severely coked, and the plants with pYL156:*GhDMT6* gene were not withered and dehydrated (Fig. 5c). The results showed that the expression of *GhDMT6* gene in cotton seedlings infected with pYL156:*GhDMT6* did not change under salt stress. The transcription level of *GhDMT6* gene in cotton seedlings infected with pYL156:*GhDMT6* decreased most in stems, followed by roots, and decreased least in leaves (Fig. 5f).

Discussion

With the completion of the genome project of cotton [34], the identification and study of gene family classifications, evolutionary features and function prediction at the whole-genome level is a hotspot of cotton functional gene research. Cotton is one of the pioneer plants in saline-alkali lands. DNA MTases are key enzymes in DNA methylation, which is closely related to resistance to stress. Therefore, the study of genome-wide DNA MTases is of great significance to cotton breeding, the identification of functional genes and the mechanism of cotton resistance.

DNA methylation affects many biological processes, including disease-associated syndromes in humans [35]. Natural variations of epialleles play a role in plant evolution [36], morphological diversity in plants [37], and the selection and breeding of agronomic traits in crops [4, 38]. Increasing evidence from recent studies suggests that DNA methylation plays an important role in regulating the stress response/adaptation in plants. DNA methylation may be an adaptation mechanism of plants in response to adversity. Osmotic stress causes DNA methylation in the chromatin region of tobacco (*Nicotiana* L.) and tissue culture cells [39]. Salt stress can inhibit

zmPP2C expression and induce *zmGST* expression in maize (*Zea mays*) seedlings [40]. Heavy metals, such as Cd and Pb, can increase the methylation level in the genomes of rice, wheat, rapeseed and other crops and produce toxicity [41].

This study, for the first time, systematically analyzed the DNA MTase gene family in the cotton genome, and 51 DNA MTases were obtained, divided into four Asian families. There are 7 members of the Met family with a region rich in glutamic acid and aspartate, similar to those of *Arabidopsis thaliana*. Although the amino acid residues in this region are important for the function of DNA MTases, the specific effect has not been determined [42]. The CMT family has 25 members, and the main sites are CHG sequences. They have particularly high abnormal chromatin content, possibly because CHG methylation maintains the plant genome regional chromatin state [43]. The DRM family, which has 14 members, contains a ubiquitin-related structural domain (UBA) and an interface between seat proteins, which introduces DRM to specific DNA regions to complete methylation of the region. The DNMT2 family has 5 members and is highly homologous in animals, plants, and prokaryotic organisms. This high homology may be due to the evolution of prokaryotic organisms. Its functional substrate is RNA, and the main target is tRNA^{Asp}. DNMT2 family members can specifically methylate the tRNA^{Asp} of the reverse codon ring 38C [44]. There were 8 members in group A, 9 in group A, 16 in the AD₁ group and 18 in the AD₂ group. The results showed that there were more DNA MTase genes in the AD genome than in the A genome or D genome, but the number of genes in the AD genome was not equal to the sum of genes in genome A and genome D. The number of DNA MTases in the AD₁ genome is less than the sum of genomes A and D, which may be associated with genetic loss during the evolution of the twofold ancestral AD₁ genome. The number of DNA MTases in the AD₂ genome is greater than the sum of genomes A and D, which may be related to gene duplication during the evolution of the twofold ancestral AD₁ genome. However, the quantitative difference between the DA₁ genome and the DA₂ genome may be the difference between natural selection and artificial domestication [45].

Gene expression profiling is usually related to gene functional identification. For the three cotton species under salt and drought stress, DNA MTase gene expression is significantly different in different tissues. This differential gene expression may be caused by the corresponding adverse effects of the regulatory pathway. This shows that different DNA methylation enzymes have different functions and participate in many regulatory pathways and that the plant's response to adversity is the result of synergistic effects through multiple pathways and is a complex process. Drought and salt stress can cause osmotic pressure, and the change in the DNA MTase gene transcription level in cotton may be caused by the joint action of these two types of stress. This is similar to the findings for other crops. Different cotton species in the same subfamily have different patterns of expression, and DNA MTase gene may have specific functions.

The function of the GhDMT6 gene was verified by the VIGS silence. After silencing the cotton GhDMT6 gene, it was less sensitive to cold, salt and drought stress than the control group, which indicated that the gene had a certain significance in improving cotton resistance. In summary, the GhDMT6 gene is a gene involved in stress resistance.

Funding information This study was funded by the National Key Research and

318 Development Program of China (2018YFD0100401).

319 **Compliance with ethical standards** The experiments described here comply with
320 the current laws of China.

321 **Conflict of interest** The authors declare that they have no conflict of interest.

322 References

- 323 1. Yoder JA, Walsh CP, Bestor TH. Cytosine methylation and the ecology of intragenomic parasites.
324 Trends in Genetics Tigr. 1997;13(8):335-340.
- 325 2. Bestor TH. The DNA methyltransferases of mammals. Human Molecular Genetics2000;9(16):
326 2395-2402.
- 327 3. Chen W, Kong J, Cheng Q, Sheng Y, Tan J, Chen Y, et al. Requirement of CHROMOMETHYLASE3
328 for somatic inheritance of the spontaneous tomato epimutation Colourless non-ripening. Sci Rep.
329 2015;5(9192):9192-9192.
- 330 4. Manning K, Tör M, Poole M, Hong Y, Thompson AJ, King GJ, et al. A naturally occurring epigenetic
331 mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening. Nature
332 Genetics. 2006;38(8):948-952.
- 333 5. Zhong S, Fei Z, Chen YR, Zheng Y. Single-base resolution methylomes of tomato fruit
334 development reveal epigenome modifications associated with ripening. Nature Biotechnology.
335 2013;31(2):154-159.
- 336 6. Colot V, Rossignol JL. Eukaryotic DNA methylation as an evolutionary device. Bioessays News &
337 Reviews in Molecular Cellular & Developmental Biology. 1999;21(5):402-411.
- 338 7. Macdonald WA. Epigenetic Mechanisms of Genomic Imprinting: Common Themes in the
339 Regulation of Imprinted Regions in Mammals, Plants, and Insects. Genetics Research International.
340 2012;2012(1):585024.
- 341 8. Bala TN, Brahmachary M, Garg P, Borel C, Alnefaie R, Watson CT, et al. DNA methylation profiling
342 in X;autosome translocations supports a role for L1 repeats in the spread of X chromosome
343 inactivation. Human Molecular Genetics. 2014;23(5):1224-1236.
- 344 9. Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, et al. DEMETER, a DNA
345 glycosylase domain protein, is required for endosperm gene imprinting and seed viability in
346 arabidopsis. Cell. 2002;110(1):33-42.
- 347 10. Gehring M, Henikoff S. Extensive demethylation of repetitive elements during seed development
348 underlies gene imprinting. Science. 2009;324(5933):1447-1451.
- 349 11. Song Q, Zhang T, Stelly DM, Chen ZJ. Epigenomic and functional analyses reveal roles of
350 epialleles in the loss of photoperiod sensitivity during domestication of allotetraploid cottons.

Genome Biology. 2017;18(1):99.

12. Finnegan EJ, Kovac KA. Plant DNA methyltransferases: Springer Netherlands; 2000.

13. Dennis ES. DNA methylation, a key regulator of plant development and other processes. Current Opinion in Genetics & Development. 2000;10(2):217.

14. Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, et al. Non-CG methylation patterns shape the epigenetic landscape in Arabidopsis. Nature Structural & Molecular Biology. 2014;21(1):64.

15. Chen T, Li E. Structure and function of eukaryotic DNA methyltransferases. Current Topics in Developmental Biology. 2004;60(60):55.

16. Jullien Pauline E, Susaki D, Yelagandula R, Higashiyama T, Berger F. DNA Methylation Dynamics during Sexual Reproduction in Arabidopsis thaliana. Current Biology Cb. 2012;22(19):1825-30.

17. JA L, SE J. Establishing, maintaining and modifying DNA methylation patterns in plants and animals. Nat Rev Genet. 2010;11(3):204-20.

18. Pavlopoulou A, Kossida S. Plant cytosine-5 DNA methyltransferases: Structure, function, and molecular evolution. Genomics. 2007;90(4):530-41.

19. Cao X, Aufsatz W, Zilberman D, Mette MF, Huang MS, Matzke M, et al. Role of the DRM and CMT3 methyltransferases in RNA-directed DNA methylation. Current Biology Cb. 2003;13(24):2212-7.

20. Finnegan EJ, Kovac KA. Plant DNA methyltransferases. Plant Molecular Biology. 2000;43(2-3):189-201.

21. Hermann A, Gowher H, Jeltsch A. Biochemistry and biology of mammalian DNA methyltransferases. Cellular & Molecular Life Sciences. 2004;61(19-20):2571-87.

22. Garg R, Kumari R, Tiwari S, Goyal S. Genomic Survey, Gene Expression Analysis and Structural Modeling Suggest Diverse Roles of DNA Methyltransferases in Legumes. Plos One. 2014;9(2):e88947.

23. Sharma R, Singh RKM, Malik G, Deveshwar P, Tyagi AK, Kapoor S, et al. Rice cytosine DNA methyltransferases - gene expression profiling during reproductive development and abiotic stress. Febs Journal. 2009;276(21):6301.

24. Johnson TB, Coghill RD. RESEARCHES ON PYRIMIDINES. C111. THE DISCOVERY OF 5-METHYL-CYTOSINE IN TUBERCULINIC ACID, THE NUCLEIC ACID OF THE TUBERCLE BACILLUS1. Jmchemsoc. 1925;47(11):2838-44.

25. Hotchkiss RD. The quantitative separation of purines, pyrimidines, and nucleosides by paper chromatography. Journal of Biological Chemistry. 1948;175(1):315.

26. Gold M, Hurwitz J. THE ENZYMATIC METHYLATION OF RIBONUCLEIC ACID AND DEOXYRIBONUCLEIC ACID. V. PURIFICATION AND PROPERTIES OF THE DEOXYRIBONUCLEIC ACID-METHYLATING ACTIVITY OF ESCHERICHIA COLI. Journal of Biological Chemistry. 1964;239(9):3858.

27. Finnegan EJ, Peacock WJ, Dennis ES. Reduced DNA methylation in Arabidopsis thaliana results in abnormal plant development. Proceedings of the National Academy of Sciences of the United States of America. 1996;93(16):8449-54.

28. Vongs A, Kakutani T, Martienssen RA, Richards EJ. Arabidopsis thaliana DNA methylation mutants. Science. 1993;260(5116):1926.

29. Nakano Y, Steward N, Sekine M, Kusano T, Sano H. A Tobacco NtMET1 cDNA Encoding a DNA Methyltransferase: Molecular Characterization and Abnormal Phenotypes of Transgenic Tobacco Plants. Plant & Cell Physiology. 2000;41(4):448-57.

30. Teerawanichpan P, Chandrasekharan MB, Jiang Y, Narangajavana J, Hall TC. Characterization of two rice DNA methyltransferase genes and RNAi-mediated reactivation of a silenced transgene in rice

callus. *Planta*. 2004;218(3):337-49.

31. Qian Y, Xi Y, Cheng B, Zhu S. Genome-wide identification and expression profiling of DNA methyltransferase gene family in maize. *Plant Cell Reports*. 2014;33(10):1661.

32. Kumar R, Chauhan PK, Khurana A. Identification and expression profiling of DNA methyltransferases during development and stress conditions in Solanaceae. *Functional & Integrative Genomics*. 2016;16(5):1-16.

33. Finnegan EJ, Dennis ES. Isolation and identification by sequence homology of a putative cytosine methyltransferase from *Arabidopsis thaliana*. *Nucleic Acids Research*. 1993;21(10):2383.

34. Yin Z, Wang J, Wang D, Fan W, Shuai W, Ye W. The MAPKKK Gene Family in *Gossypium raimondii*: Genome-Wide Identification, Classification and Expression Analysis. *International Journal of Molecular Sciences*. 2013;14(9):18740-57.

35. Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, et al. Integrative analysis of 111 reference human epigenomes. *Nature*. 2015;518(7539):317-30.

36. Weigel D, Colot V. Epialleles in plant evolution. *Genome Biology*. 2012;13(10):249.

37. Cubas P, Vincent C, Coen E. An epigenetic mutation responsible for natural variation in floral symmetry. *Nature*. 1999;401(6749):157-61.

38. Quadana L, Almeida J, Asís R, Duffy T, Dominguez PG, Bermúdez L, et al. Natural occurring epialleles determine vitamin E accumulation in tomato fruits. *Nature Communications*. 2014;5(4027):3027.

39. Kovarik A, Koukalová B, Bezdek M, Opatrný Z. Hypermethylation of tobacco heterochromatic loci in response to osmotic stress. *Theoretical & Applied Genetics*. 1997;95(1-2):301-6.

40. Tan MP. Analysis of DNA methylation of maize in response to osmotic and salt stress based on methylation-sensitive amplified polymorphism. *Plant Physiol Biochem*. 2010;48(1):21-6.

41. Labra M, Ghiani A, Citterio S, Sgorbati S, Sala F, Vannini C, et al. Analysis of Cytosine Methylation Pattern in Response to Water Deficit in Pea Root Tips. *Plant Biology*. 2002;4(6):694-9.

42. Finnegan EJ, Kovac KA. Plant DNA methyltransferases. *Plant Mol Biol. Plant Molecular Biology*. 2000;43(2-3):189-201.

43. Papa CM, Springer NM, Muszynski MG, Meeley R, Kaeppler SM. Maize chromomethylase *Zea methyltransferase2* is required for CpNpG methylation. *Plant Cell*. 2001;13(8):1919-28.

44. Goll MG, Kirpekar F, Maggert KA, Yoder JA, Hsieh CL, Zhang X, et al. Methylation of tRNA^{Asp} by the DNA methyltransferase homolog Dnmt2. *Science*. 2006;311(5759):395-8.

45. Xu JH, Messing J. Amplification of prolamin storage protein genes in different subfamilies of the Poaceae. *Theoretical & Applied Genetics*. 2009;119(8):1397-412.

Figures

Fig. 1 Phylogenetic analysis of the DNA Mtase gene family in cotton (A) and genetic structure (B). Cotton DNA Mtase family members are divided into four subfamilies, CMT, MET, DRM, Dnmt2.

Fig. 2 Analysis of the distribution of cotton DNA Mtase protein motifs using online software. The rectangular length conforms to the length of the motif. The order and position of the motif correspond to the order and position of the bases in a single protein sequence.

Fig. 3 DNA Mase family phylogenetic tree of cotton and other crops. The neighbor-joining method was used to construct the tree without a root system, and the value of the bootstrap was 1000.

Fig. 4 The expression of DNA Mtase genes in the roots, stems and leaves of three cotton species, namely, *Shixiya 1*, *G. raimondii*, and *G. hirsutum* L. TM-1, under drought and salt treatments at different times. a *G. arboreum* L. *Shixiya 1*, b *G. raimondii*, and c *G. hirsutum* L. TM-1.

Fig. 5 Relative expression of GhDMT6 gene in different tissues of different stress in *Gossypium. hirsutum*. a. Phenotypic differences of cotton after drought stress. b. Relative expression of GhDMT6 gene after drought stress in *G. hirsutum* L. TM-1. c. Phenotypic differences of cotton after cold stress. d. Relative expression of GhDMT6 gene after cold stress in *G. hirsutum* L. TM-1. e. Phenotypic differences of cotton after salt stress. f. Relative expression of GhDMT6 gene after salt stress in *G. hirsutum* L. TM-1

Tables

Table 1 Basic characteristics of DNA Mtase genes in the cotton genome

gene name	Accession number	Location (chromosome)	Position (domain)	CDS (bp)	AA	PI	Predicted subcellular localization
GaDMT1		CA_chr1: 66883217:66889872-	499-871	2724	907	5.09	Cytoplasmic
GaDMT2	Cotton_A_27044	CA_chr2: 55459962..55465882-	480-771	4638	1545	5.66	Cytoplasmic
GaDMT3	Cotton_A_20175	CA_chr3: 24408653..24412255+	13-389	1179	637	6.02	OuterMembrane
GaDMT4	Cotton_A_29334	CA_chr3: 96706846..96713328-	552-692	2082	693	7.44	OuterMembrane
GaDMT5	Cotton_A_29333	CA_chr3: 96716269:96725942-	475-822	2580	859	6.24	Cytoplasmic
GaDMT6	Cotton_A_36034	CA_chr10: 113686352..113690340+	513-631	1914	663	4.8	Cytoplasmic
GaDMT7	Cotton_A_08447	CA_chr12: 1201266..1207466-	534-653	1992	392	4.94	Cytoplasmic
GaDMT8	Cotton_A_27234	CA_chr12: 56291689..56297548-	1110-1539	1494	497	5.82	Cytoplasmic
GaDMT9	Cotton_A_19737	CA_chr13: 63213811..63219313+	155-372	2421	806	5.25	Cytoplasmic
GrDMT1	Cotton_D_gene_10004803	Chr2:8382256..8388371 +	494-785	2463	820	5.67	Cytoplasmic
GrDMT2	Cotton_D_gene_10010304	Chr3:6796286..6800300-	512-630	1911	636	4.8	Cytoplasmic
GrDMT3	Cotton_D_gene_10010121	Chr4:457766..461539 +	13-389	1203	400	5.55	OuterMembrane
GrDMT4	Cotton_D_gene_10027875	Chr4:20122791..20126504-	511-630	1923	640	4.9	Cytoplasmic
GrDMT5	Cotton_D_gene_10009363	scaffold141: 194013:200650-	495-869	2718	905	5.1	Cytoplasmic
GrDMT6	Cotton_D_gene_10000349	scaffold531:66538..72372 -	1128-1557	4692	1563	5.51	Cytoplasmic
GrDMT7	Cotton_D_gene_10002270	scaffold372: 306669:314046-	530-928	2892	963	7.02	OuterMembrane
GrDMT8	Cotton_D_gene_10002271	scaffold372: 317040:326457-	789-1165	3609	1202	8.44	OuterMembrane
GhDMT1	CotAD_37635	At_chr6:11311066..11315214-	512-630	1911	636	4.8	Cytoplasmic
GhDMT2	CotAD_51709	At_chr9:63179674..63185543-	1076-1505	4536	1511	6.34	Cytoplasmic
GhDMT3	CotAD_46796	At_chr9:65748738..65752451+	511-630	1923	640	4.85	Cytoplasmic
GhDMT4	CotAD_10542	Dt_chr1:40856014..40859213-	1-308	1035	344	7.76	Cytoplasmic
GhDMT5	CotAD_49037	Dt_chr2:8190450..8196335-	496-781	2451	816	5.62	Cytoplasmic
GhDMT6	CotAD_04205	Dt_chr5:13787674..13791315+	13-389	1206	401	5.82	OuterMembrane
GhDMT7	CotAD_13275	Dt_chr6:40651384..40655909-	512-611	1878	625	4.72	OuterMembrane
GhDMT8	CotAD_24264	Dt_chr7:37416068..37421133+	230-632	2007	668	4.81	Cytoplasmic
GhDMT9	CotAD_00990	Dt_chr10: 6477855..6487242+	789-1157	3585	1194	8.51	OuterMembrane
GhDMT10	CotAD_00992	Dt_chr10:6495924..6502242+	299-680	2094	697	6.75	Cytoplasmic
GhDMT11	CotAD_14980	scaffold39.1:602870..611908+	667-846	2646	881	6.94	OuterMembrane
GhDMT12	CotAD_18652	scaffold71.1:1581335..1591239-	547-666	2031	676	4.91	Cytoplasmic
GhDMT13	CotAD_41398	scaffold294.1:1053646..1057162-	319-500	1533	510	5.45	Cytoplasmic
GhDMT14	CotAD_41399	scaffold294.1:1063401:1074111-	780-1161	3597	1198	8.71	OuterMembrane
GhDMT15	CotAD_46012	scaffold1041.1:189137..194922+	1112-1541	4644	1547	5.57	Cytoplasmic
GhDMT16	CotAD_40093	scaffold2005.1:29096..35764+	453-828	2595	864	5.02	Cytoplasmic
GbDMT1	Gbscaffold4563.2.0	At01: 75311449..75320269	441-816	2556	851	5.66	Cytoplasmic
GbDMT2	Gbscaffold7611.1.0	At05:80836461..80840887+	564-683	2082	693	5.03	Cytoplasmic
GbDMT3	Gbscaffold10176.3.0	At05:85393045..85400787+	1142-1571	4734	1577	5.49	Cytoplasmic
GbDMT4	Gbscaffold23728.21.0	At07: 5162033..5169156	503-875	2739	912	4.98	Cytoplasmic
GbDMT5	Gbscaffold11439.6.0	At08:92243382..92245846-	278-428	1431	476	5.77	Cytoplasmic

GbDMT6	Gbscaffold11439.5.0	At08: 92252927..92259967	131-508	1641	546	8.13	Cytoplasmic
GbDMT7	Gbscaffold10104.36.0	At08:106991133..106995128+	13-389	1179	392	5.4	OuterMembrane
GbDMT8	Gbscaffold155.2.0	At09:9281821..9292014+	1068-1186	3579	1192	4.78	Cytoplasmic
GbDMT9	Gbscaffold155.3.0	At09:9298124..9301305+	374-492	1497	498	5.56	Cytoplasmic
GbDMT10	Gbscaffold10257.5.0	Dt01: 55556607..55564136	438-672	2370	789	5.78	Cytoplasmic
GbDMT11	Gbscaffold9562.4.0	Dt04:6380024..6388975+	1126-1555	4686	1561	5.6	Cytoplasmic
GbDMT12	Gbscaffold9562.5.0	Dt04:6390322..6394029+	475-834	2523	840	6.49	Cytoplasmic
GbDMT13	Gbscaffold12826.2.0	Dt04:9853138..9856849+	511-630	1923	640	4.93	Cytoplasmic
GbDMT14	Gbscaffold12826.3.0	Dt04:9869411..9871104+	219-338	1047	348	8.6	Cytoplasmic
GbDMT15	Gbscaffold8153.13.0	Dt07: 5581094..5588192	492-866	2709	902	5.2	Cytoplasmic
GbDMT16	Gbscaffold34888.4.0	Dt08:67685650..67693483-	19-642	1962	653	5.21	OuterMembrane
GbDMT17	Gbscaffold8535.6.0	scaffold8535:88760..90489+	7-125	486	161	9.24	Periplasmic
GbDMT18	Gbscaffold12265.1.0	scaffold12265:8693..15154+	576-694	2103	700	4.67	Cytoplasmic

Table 2 Basic information of related species in analyzing the phylogenetic tree

species name	short	classification	species name	short	classification
<i>Ricinus communis</i>	Rc	D	<i>Chlamydomonas reinhardtii</i>	Cr	A
<i>grapevine</i>	Vv	D	<i>Medicago truncatula</i>	Mt	D
<i>Oryza sativa</i>	Os	M	<i>Arabidopsis thaliana</i>	At	D
<i>Glycine max</i>	Gly	D	<i>Solanum lycopersicum</i>	Sl	D
<i>Zea mays</i>	Zm	M	<i>Solanum tuberosum</i>	St	D
<i>cacao</i>	Co	D			

D, M and A represent Dicotyledon, Monocotyledon and Algae, respectively.