The trichotomy of pneumococcal infection outcomes

Alexis Erich S. Almocera, Gustavo Hernandez-Mejia, César Parra-Rojas and
Esteban A. Hernandez-Vargas*

Frankfurt Institute for Advanced Studies
Ruth-Moufang-Straße 1, 60438 Frankfurt am Main, Germany
* Corresponding author: vargas@fias.uni-frankfurt.de

Abstract

The successful elimination of bacteria such as *Streptococcus pneumoniae* from a host involves the coordination between different parts of the immune system. Previous studies have explored the effects of the initial pneumococcal load (bacterial dose) on different representations of innate immunity, finding that pathogenic outcomes can vary with the size of the bacterial dose. However, others yield support to the notion of dose-independent factors contributing to bacterial clearance. In this paper, we seek to provide a deeper understanding of the immune responses associated to the pneumococcus. To this end, we formulate a model that realizes an abstraction of the innate-regulatory immune host response. Stability and bifurcation analyses of the model reveal the following trichotomy of pneumococcal outcomes determined by the bifurcation parameters: (i) *dose-independent clearance*; (ii) *dose-independent persistence*; and (iii) *dose-limited clearance*. Bistability, where the bacteria-free equilibrium co-stabilizes with the most substantial steady-state bacterial load is the specific result behind dose-limited clearance. The trichotomy of pneumococcal outcomes here described integrates all previously observed bacterial fates into a unified framework.

1. Introduction

The pneumococcus (*Streptococcus pneumoniae*) is a bacterial pathogen associated with pneumonia, otitis media (ear infections), and life-threatening conditions such as sepsis or bacteremia (blood poisoning) and bacterial meningitis (brain infection) [1]. The pneumococcus is notably a coinfective pathogen with the influenza virus, contributing to enhanced morbidity and mortality [2, 3, 4, 5]. According to the World Health Organization (WHO), the pneumococcus is the causative agent behind 16% of mortalities in children under five years of age, and the deaths of 920,136 children in 2015. Diseases caused by the pneumococcus are mostly common among children and the elderly, as well as individuals with a compromised immune system [6, 7, 8]. Current immunization programs may have desirable impacts [9, 10]; however, they remain challenged by antibiotic-resistant serotypes [8, 9, 11, 12]. These challenges emphasize the need for more research and development towards the control and eradication of *S. pneumoniae*.

Briefly speaking, the infection of the pneumococcus begins with entry of pneumococcal particles through the nasal cavity, followed by adherence to epithelial cells (colonization), and concluding with invasive disease [1]. An extensive collection of

studies identified a diverse range of bacterial and environmental factors that contribute to the infection, including enzymes, binding regions, and capsule structures—see, *e.g.*, [13] and [14] for a review. However, here we are interested in the kinetics of *S. pneumoniae* and interactions of the pathogen with first responders, comprising innate immune responses and regulatory mechanisms. To this end, mathematical models provide conceptual frameworks for studying immune responses and changes in bacterial load.

Previous mathematical models investigate the effects of initial bacterial load (dosage or inoculum size) on the successful bacterial clearance by the innate responses. Model formulations assume the phagocytes to form a single group [15], separate phagocytes according to whether they actively engulf pneumococcus [16], or differentiate phagocytes into neutrophils and macrophages [17, 18]. However, these models did not consider the cascading effect of immune responses. Smith et al. [19] formulated three models fitted to bacterial load in mice to investigate the coordination of innate immune cells. This coordination is described by the following cascade of innate responses: alveolar macrophages, neutrophils, and monocyte-derived macrophages. Each model predicts that the bacteria will be cleared in small doses and sustain persistent levels in high doses; we refer to this phenomenon as *dose-limited clearance*. Besides the modeling and experimental work in [19], separate studies identified drug-specific effects [20], genetic variations in either the host or the pathogen [21, 22], and biological switches (for Toll-like receptors and bacteremia threshold) [23] as factors in the pneumococcal outcome in addition to inoculation size. In light of the models reviewed here, we may conceptualize a general form of the innate-regulatory response.

The phenomenon of dose-limited clearance is comparable to the behavior of a bistable system. For a broad range of nonlinear dynamical systems, a simple case of bistability follows from the coexistence of two asymptotically stable equilibrium points. Hence, a typical solution approaches one equilibrium point or the other, depending on the initial state [24]. Malka et al. [25] constructed a one-dimensional equation describing bacterial kinetics and clearance by constant densities of neutrophils. In a specific region of the parameter space, the model exhibits bistability under moderate neutrophil densities. That is, three steady states appear, where the bacteria-free value is mutually stable with the maximum. The intermediate value is unstable and serves as a clearance threshold comparable to Smith et al. [19]. This observation indicates the inadequacy of neutrophils to clear more massive bacterial loads. Furthermore, hysteresis accompanies bistability: when neutrophils decrease to critically small levels and then return to the original state, the clearance of sufficiently small bacterial population suddenly changes to an irreversible persistence event. Notably, the bistability phenomenon agrees with published data from a series of bactericidal experiments [26], suggesting that bistability is a plausible mechanism for fulminant infection. In this paper, we formulate a model of three ordinary differential equations, which generalizes the innate-regulatory immune response to *S. pneumoniae*. The generalized immune response unifies associated mechanisms into abstract components.

We organize the paper as follows. In Section 2, we introduce our model and estimate parameters by fitting the model to murine experimental data in [27]. The experiments in this study revealed that some but not all mice reached undetectable pneumococcal levels after 16 hours post-infection (hpi). Our analysis in Section 3

reveals a bistability event comparable to [25]. Moreover, our model predicts three possible outcomes: clearance, persistence, and dose-limited clearance in the bistability case. The first two outcomes are independent of the bacterial dose size. One threshold parameter controls the window of bistability, while another dictates the predicted outcome. Section 4 concludes this paper with a discussion.

2. Mathematical model

We establish a mathematical model to represent the global panorama of the bacterial (B) interaction with the host and the corresponding immune response which in this case is characterized by the innate (M) and regulatory (N) immune responses. The model reads as follows:

$$\dot{B} = rB \left(1 - \frac{B}{K_B}\right) - c_B BM, \quad (1)$$

$$\dot{M} = \delta_M(M_0 - M) + (\eta B - \theta N)M, \quad (2)$$

$$\dot{N} = \gamma B + \delta_N(N_0 - N), \quad (3)$$

where the dot denotes the derivative with respect to a time variable t , *i.e.*, $\dot{x} = dx/dt$, $x = B, M, N$. Here, the bacteria (B) proliferates logistically at a maximum rate r with a tissue carrying capacity of K_B , given in colony forming units per milliliter (CFU/mL). The clearance of free bacteria occurs at a rate c_B per cell and is assumed to result from the innate immune response. M is assumed to evolve with a constant rate η due to bacterial presence. We consider a constant decline rate of M given by θ , which is influenced by the immune regulatory activity. The elimination rate of M is given by δ_M . Furthermore, the regulatory process is favored by bacteria at a rate γ and its inhibition rate is given by δ_N . The initial immune response levels are M_0 (innate) and N_0 (regulatory). We assume a constant replenishment of the innate ($\delta_M M_0$) and regulatory ($\delta_N N_0$) responses. Figure 1 is a conceptual diagram of the model (1)–(3).

Experimental data. In the experiment of Duvigneau *et al.* [27], C57BL/6J wildtype mice were intranasally infected with a sub-lethal dose of 1×10^6 CFU of the *S. pneumoniae* strain TIGR4 (T4). After the infection, the bacterial load in the bronchoalveolar lavage was measured at different time points, namely at 1.5, 6, 18, 26 and 31 hpi. Figure 2 depicts the experimental data of bacterial load in the lungs of the infected mice. Note that the experiment presents two outcomes, bacterial clearance (a) and persistence (b). We explore the interaction of bacteria with the host immune response through the combination of data and the proposed model (1)–(3) for the two scenarios. For each set of data in Figure 2, we separately estimated the majority of parameters. The remaining ones were set as described below.

Parameter estimation According to Smith *et al.* [19] and Duvigneau *et al.* [27], the bacterial growth rate $r = 1.13 \text{ h}^{-1}$ and the carrying capacity $K_B = 2.3 \times 10^8$ CFU/mL stand for single *S. pneumoniae* infection. Also, following the criteria from Smith *et al.* [19], the *S. pneumoniae* inoculum size (B_0) for simulations is set to 1000 CFU/mL. Several elimination rates of immune actors such as alveolar macrophages, cytokines as interleukin-1 and tumor necrosis factor- α , neutrophils, and monocyte-derived macrophages are reported in [19] for the specific dynamical models there developed. In contrast, our model considers all innate response components as

Figure 1: Schematic diagram of the model (1)–(3). The state variables for bacterial load (B), innate (M) and regulatory (N) response levels are depicted in shaded squares. Solid lines with arrowheads indicate bacterial activation of innate and regulatory responses, associated with constant rates η and γ , respectively. Solid lines ending in bars denote the following inhibitory effects: regulatory levels inhibit innate response growth at a constant rate θ , and the innate response controls bacterial growth at a constant clearance rate c_B . Dashed loops indicate the replenishment of a given state variable according to its corresponding growth term.

107 a single global response rather than as separate actors. The same philosophy is
 108 considered for the regulatory response. Under these considerations, we set the innate
 109 and regulatory inhibition rates as 0.1, a general-response value. In addition, for the
 110 global regulatory response, we set $M_0 = 1$ due to the constant supply of innate
 111 agents, such as alveolar macrophages [19, 27]. A constant regulatory action is also
 112 considered assuming $N_0 = 1$. We would like to remark that different assumptions
 113 of initial values and elimination rates would only rescale the fitted parameters, but
 114 not affect the mechanistic insights from the model selection procedures.

Table 1: Fitted parameter values for the different bacterial infection outcomes. The sets of four parameters were independently fitted based on bacterial data from either the persistence or clearance scenarios.

Parameter	Bacteria clearance	Bacterial persistence
c_B	1.13096	0.97000
γ	0.00376	2.85546×10^{-4}
η	1.3461×10^{-4}	8.07453×10^{-6}
θ	1.3982×10^{-13}	3.50118×10^{-13}

115 To fit the remaining parameters, *i.e.*, c_B , η , θ and γ , we minimized the mean
 116 squared difference (MSD) between the model output and the experimental bacteri-
 117 al measurement, both on logarithmic scale. The model equations were solved in
 118 Python using the numerical integration routines of the SciPy library [28]. The mini-
 119 mization of the MSD was also performed with SciPy using the Differential Evolution
 120 algorithm [29]. Separately, we fitted each of the datasets to uncover the parameter
 121 values that yield either the persistence of the bacterium or its elimination. The
 122 fitted values are shown in Table 1, while Figure 2 shows the resulting dynamics
 123 of the model (1)–(3) for each case. Innate and regulatory responses are plotted in
 124 fold-change, Figure 2(c)–(d). The dynamics of the clearance case shows a marked

Figure 2: Dynamics of bacterial load and immune response. **Top:** bacterial data (circles) and fitted dynamics for (a) the clearance (blue, solid line) and (b) the persistence (orange, dashed line) scenarios. Note that in both panels the data at 1.5 and 6 hpi is the same. **Bottom:** (c) Innate and (d) regulatory immune responses for both bacterial behaviors: clearance (blue, solid line) and persistence (orange, dashed line).

bacterial elimination after 16 hpi—see Figure 2(a). Note that for the clearance scenario, both immune responses present a large marginal increase during the first hours post infection; this effect is more pronounced in the regulatory response, reaching a 10-fold increase before 7 hpi—see Figure 2(d). In contrast, the persistent bacteria appears to provoke a sluggish action of the immune response, making the regulatory and innate responses peak after 20 hpi, where the regulatory effect is almost reaching an 8-fold increase—see panels (c) and (d) of Figure 2.

3. Model analysis

We nondimensionalize the model (1)–(3) to simplify computations. Let us introduce the following dimensionless variables

$$b = \frac{B}{K_B}, \quad m = \frac{M}{M_0}, \quad n = \frac{N}{N_0}, \quad \tau = rt,$$

where τ is a rescaled time variable. This transformation scales the bacterial load with the carrying capacity, and the immune response levels with their respective

initial values. Define the following dimensionless parameters:

$$\begin{aligned}\overline{c_B} &= \frac{c_B M_0}{r}, & \overline{\delta_M} &= \frac{\delta_M}{r}, & \overline{\delta_N} &= \frac{\delta_N}{r}, \\ \overline{\eta} &= \frac{\eta K_B}{r}, & \overline{\theta} &= \frac{\theta N_0}{r}, & \overline{\gamma} &= \frac{\gamma K_B}{r N_0}.\end{aligned}$$

Then the model (1)–(3) has the dynamically equivalent dimensionless form

$$b' = b(1 - b) - b m \overline{c_B}, \quad (4)$$

$$m' = \overline{\delta_M}(1 - m) + m(\overline{\eta}b - \overline{\theta}n), \quad (5)$$

$$n' = \overline{\delta_N}(1 - n) + \overline{\gamma}b, \quad (6)$$

where the prime denotes the derivate with respect to the rescaled time τ .

To determine the local stability of system (4)–(6), we denote a point in state-space by its coordinates (b, m, n) . Then (4)–(6) admits the unique equilibrium point

$$E_0^* := (0, m_0^*, 1), \quad m_0^* := \frac{\overline{\delta_M}}{\overline{\delta_M} + \overline{\theta}} = \frac{\delta_M}{\delta_M + \theta N_0},$$

corresponding to steady state values $B = 0$, $M = m_0^* M_0$ and $N = N_0$. Let

$$\lambda := 1 - \overline{c_B} m_0^*, \quad (7)$$

and denote the Jacobian matrix of (4)–(6) by $J(b, m, n)$. Then the following result is evident from the eigenvalues of $J(E_0^*)$, which take values $-\overline{\delta_N}$, $-(\overline{\theta} + \overline{\delta_M})$ and λ .

Theorem 1. *The dimensionless system (4)–(6) admits the unique equilibrium point $E_0^* = (0, m_0^*, 1)$ where $b = 0$. Moreover, E_0^* is asymptotically stable if $\lambda < 0$ and is a saddle point when $\lambda > 0$.*

We consider λ as our bifurcation parameter, and determine equilibrium points of the form $E^* = (b^*, m^*, n^*)$ where $b^* > 0$. To this end, we introduce the map

$$\mu : (-\infty, 1) \times (-\infty, 1) \rightarrow (0, \infty), \quad \mu(b, \lambda) = \frac{m_0^* (b - 1)}{\lambda - 1},$$

and let

$$\begin{aligned}f(b) &:= \frac{\overline{\gamma}b}{\overline{\delta_N}} + 1, \\ g(b, \lambda) &:= \frac{m_0^*}{\overline{\delta_M}(1 - m_0^*)} \left[\overline{\eta}b - \overline{\delta_M} + \frac{\overline{\delta_M}}{\mu(b, \lambda)} \right], \\ h(b, \lambda) &:= f(b) - g(b, \lambda),\end{aligned}$$

for $b < 1$ and $\lambda < 1$. We are interested in the roots of $h(\cdot, \lambda)$ in the open interval $(0, 1)$ which are later determined to be the values of b^* . To this end, we first establish the following monotone and concave properties, where D_j^k denotes the k th partial derivative with respect to the j th variable and $D_j = D_j^1$.

Lemma 1. *The following properties hold:*

$$D_1 f > 0 = D_2 f, \quad D_1^k f = 0 = D_2^k f, \quad (8)$$

$$D_1 \mu < 0 < D_2 \mu, \quad D_1^k \mu = 0, \quad (9)$$

$$D_1 g > 0 > D_2 g, \quad D_1^2 g > 0 = D_2^k g. \quad (10)$$

143 where $k \geq 2$. Thus, f is a strictly increasing linear function, and $g(b, \lambda)$ is strictly
144 increasing and concave upwards in b while being strictly decreasing in λ . Moreover,
145 $D_1^2 h < 0 < D_2 h$ and $D_2^k h = 0$ for $k \geq 2$, i.e., h is concave downwards in b and is
146 a strictly increasing linear function in λ .

Proof. The properties in (8) follow directly the definition of f , from which f is strictly increasing and linear. To establish (9), we have

$$D_1 \mu(b, \lambda) = \frac{m_0^*}{\lambda - 1} < 0, \quad D_2 \mu(b, \lambda) = \frac{m_0^*(1 - b)}{(\lambda - 1)^2} > 0.$$

Consequently, $D_1 \mu(b, \lambda)$ is independent of b , and $D_1^k \mu = 0$ for $k \geq 2$. Thus, we compute $D_1 g$ and $D_1^2 g$ as follows:

$$D_1 g(b, \lambda) = \frac{m_0^*}{\delta_M(1 - m_0^*)} \left\{ \bar{\eta} - \frac{\delta_M D_1 \mu(b, \lambda)}{[\mu(b, \lambda)]^2} \right\} > 0,$$

$$D_1^2 g(b, \lambda) = \frac{2\delta_M m_0^* [D_1 \mu(b, \lambda)]^2}{\delta_M(1 - m_0^*) [\mu(b, \lambda)]^3} > 0.$$

Finally, we have $D_2 h = -D_2 g$ due to $D_2 f = 0$, and we obtain

$$D_2 g(b, \lambda) = \frac{-m_0^* \delta_M D_2 \mu(b, \lambda)}{\delta_M(1 - m_0^*) [\mu(b, \lambda)]^2} < 0.$$

147 Since $(D_2 \mu)/\mu^2$ expands to a function independent of λ , we have $D_2^k g = 0$ for $k \geq 2$.
148 Therefore, the properties in (10) are true; in particular, g is strictly increasing and
149 concave upwards in b , while being strictly decreasing in λ . The results for $h = f - g$
150 follow from (8) and (10). $\square$

The results of h in Lemma 1 yields the following properties. First, the root of $D_1 h(\cdot, \lambda)$ is uniquely given by a certain $\hat{b} \in (-\infty, 1)$ from which

$$h_{\max}(\lambda) := \max_{b < 1} h(b, \lambda) = h(\hat{b}, \lambda),$$

that is, $\hat{b}$ is the unique maximizer of $h(\cdot, \lambda)$. Furthermore, we have

$$D_1 h(b_1, \lambda) > 0 \quad D_1 h(b_2, \lambda) < 0, \quad (11)$$

for arbitrary values $b_1 < \hat{b}$ and $b_2 > \hat{b}$. The function $h_{\max}$ is linear with positive slope because $D_2^k h = 0$ for $k \geq 2$. Hence, we define the unique root of $h_{\max}$ as

$$\hat{\lambda} := \frac{-h_{\max}(0)}{D_1 h_{\max}(0)} = \frac{-h(\hat{b}, 0)}{D_1 h(\hat{b}, 0)} \quad (12)$$

151 from Maclaurin expansion. That is, $h_{\max}(\widehat{\lambda}) = h(\widehat{b}, \widehat{\lambda}) = 0$.
Now, let

$$\xi := \lim_{b \rightarrow -\infty} \frac{h(b, \lambda)}{b} = \frac{\overline{\gamma\theta} - \overline{\eta\delta_N}}{\overline{\theta\delta_N}}, \quad \zeta := \lim_{b \rightarrow -\infty} [h(b, \lambda) - \xi b] = 1 + \frac{\overline{\delta_M}}{\overline{\theta}}. \quad (13)$$

Then the curve $y = h(b, \lambda)$ is asymptotic to the line $y = \xi b + \zeta$ as $b \rightarrow -\infty$. Moreover, the coefficients ξ and ζ of the asymptote line allow us to write

$$D_1 h(b, \lambda) = \xi - \frac{(1 - \lambda)\zeta}{(1 - b)^2}; \quad (14)$$

from this, we derive

$$\widehat{b} = 1 - \sqrt{\frac{\zeta(1 - \lambda)}{\xi}}. \quad (15)$$

152 In the following lemma, we determine some limiting behavior on the function h and
153 its first derivative.

Lemma 2. *For $0 < b < 1$, we have $h(b, \lambda) < b D_1 h(0, 0)$. Moreover,*

$$\lim_{b \rightarrow -\infty} D_1 h(b, \lambda) = \xi, \quad \lim_{b \rightarrow 1^-} D_1 h(b, \lambda) = \lim_{b \rightarrow 1^-} h(b, \lambda) = -\infty.$$

Proof. Recall that h strictly increases in λ (Lemma 1), and note that the curve $y = h(b, 0)$ is tangent to the line $y = h(0, 0) + b D_1 h(0, 0) = b D_1 h(0, 0)$ at $b = 0$. Hence by the concavity of $h(\cdot, \lambda)$, we have

$$h(b, \lambda) \leq h(b, 0) < b D_1 h(0, 0) \quad (16)$$

154 for $b > 0$. Passing the limit to (16) as $b \rightarrow 1^-$, we have $\lim_{b \rightarrow 1^-} h(b, \lambda) = -\infty$. We
155 complete the proof by passing limits to (14) where $b \rightarrow -\infty$ and $b \rightarrow 1^-$. $\square$

156 We establish that the values of b^* are roots of $h(\cdot, \lambda)$ in the interval $(0, 1)$ on
157 which $f > 0$. From now on, we denote the smallest and largest positive roots by b_1^*
158 and $b_2^* > b_1^*$, respectively.

Lemma 3. *The function $h(\cdot, \lambda)$ has at most two distinct roots, namely b_1^* and b_2^* . If b_2^* exists, then $D_1 h(b_2^*, \lambda) \leq 0$. If b_1^* also exists, then $0 < b_1^* < \widehat{b} < b_2^*$, from which*

$$D_1 h(b_2^*, \lambda) < 0 < D_1 h(b_1^*, \lambda) \quad (17)$$

and $\lambda < 0$. Moreover, the following equations are equivalent:

$$D_1 h(b_2^*, \lambda) = 0, \quad b_2^* = \widehat{b}, \quad \lambda = \widehat{\lambda} < 0. \quad (18)$$

159 *Proof.* Rolle's theorem asserts that a real-valued differentiable function with two
160 distinct roots attains a local maximum or minimum at a point between the roots.
161 Thus, a continuously differentiable function with at least three roots has no fixed
162 concavity. Since $h(\cdot, \lambda)$ is concave downwards by Lemma 1, no more than two roots
163 exist for $h(\cdot, \lambda)$, which are b_1^* and b_2^* .

Recalling from Lemma 2 that $h(b, \lambda) \rightarrow -\infty$ as $b \rightarrow 1^-$, suppose that b_2^* exists so that $h(\cdot, \lambda)$ has no root in the interval $(b_2^*, 1)$. Assuming $D_1 h(b_2^*, \lambda) > 0$ implies that $h(b, \lambda)$ is initially positive then approaches $-\infty$, as b increases from $b = b_2^*$ to $b = 1$. However, a contradiction arises with $h(\tilde{b}, \lambda) = 0$ for some $\tilde{b} \in (b_2^*, 1)$. Thus, $D_1 h(b_2^*, \lambda) \leq 0$. If b_1^* additionally exists, then $0 < b_1^* < \hat{b} < b_2^*$ by Rolle's theorem and the uniqueness of $\hat{b}$ as the root of $D_1 h(\cdot, \lambda)$. We obtain (17) by taking $b_k = b_k^*$ for each k in (11). Furthermore, $h(0, \lambda) < h(b_1^*, \lambda) = 0$. Since $h(0, \lambda)$ has the same sign with λ , we have $\lambda < 0$.

Assuming that b_2^* exists, $D_1 h(b_2^*, \lambda) = 0$ if and only if $b_2^* = \hat{b}$ because $\hat{b}$ is the unique root of $D_1 h(\cdot, \lambda)$. In such case, $h(\cdot, \lambda)$ strictly increases over the interval $(-\infty, b_2^*)$, hence $h(0, \lambda) < h(b_2^*, \lambda) = 0$ and $\lambda < 0$. Now, $b_2^* = \hat{b}$ implies

$$h_{\max}(\lambda) = h(\hat{b}, \lambda) = h(b_2^*, \lambda) = 0.$$

Conversely, $h_{\max}(\lambda) = 0$ implies that either $\hat{b} = b_1^*$ or $\hat{b} = b_2^*$. We must have $\hat{b} = b_2^*$ because the existence of b_1^* necessitates $b_1^* < \hat{b}$ as shown above. Finally, $h_{\max}(\lambda) = 0$ if and only if $\lambda = \hat{\lambda}$, by the uniqueness of $\hat{\lambda}$ as the root of $h_{\max}$. Therefore, the equations in (18) are equivalent. $\square$

We now establish the existence of roots for the function $h(\cdot, \lambda)$ in $(0, 1)$. In particular, we show that the existence of both b_1^* and b_2^* depends on the value of $D_1 h(0, 0)$, which from (14) is given by

$$D_1 h(0, 0) = \frac{\bar{\gamma}}{\bar{\delta}_N} - \left(\frac{\bar{\delta}_M + \bar{\eta}}{\bar{\theta}} + 1 \right).$$

We may alternatively write

$$D_1 h(0, 0) = \frac{\bar{\gamma} - \bar{\gamma}^*}{\bar{\delta}_N}, \quad \bar{\gamma}^* := \left(\frac{\bar{\eta} + \bar{\delta}_M}{\bar{\theta}} + 1 \right) \bar{\delta}_N, \quad (19)$$

to frame our results with $\bar{\gamma}$, which is associated with the proliferation response of interferon growth due to bacterial stimuli. The following result establishes the existence of roots for the function $h(\cdot, \lambda)$. This result is illustrated in Figure 3.

Theorem 2. *If $\lambda > 0$, then b_2^* is the unique root of $h(\cdot, \lambda)$ in $(0, 1)$. If $\lambda \leq 0$, then $h(\cdot, \lambda)$ admits roots in $(0, 1)$ only if $D_1 h(0, 0) > 0$. In this case, b_2^* is the unique root whenever $\lambda = 0$. Moreover, the following trichotomy holds:*

- (i) Both b_2^* and b_1^* exist for $\hat{\lambda} < \lambda < 0$;
- (ii) Only $b_2^* = \hat{b}$ exists for $\lambda = \hat{\lambda}$; and
- (iii) Neither b_1^* nor b_2^* exists when $\lambda < \hat{\lambda}$.

Proof. Recall that $h(0, \lambda)$ is equal in sign to λ . If $\lambda > 0$, then b_2^* is a root of $h(\cdot, \lambda)$ in $(0, 1)$ by the intermediate value theorem, because $h(b, \lambda) \rightarrow -\infty$ as $b \rightarrow 1^-$ from Lemma 2. Appealing to Lemma 3 where $\lambda > 0$, we arrive at the uniqueness of b_2^* as the root in $(0, 1)$.

Now, assume that $\lambda \leq 0$. If $D_1 h(0, 0) \leq 0$, then it follows from Lemma 2 that

$$h(b, \lambda) < b D_1 h(0, 0) \leq 0$$

for $b > 0$. Thus, it is necessary that $D_1 h(0, 0) > 0$ for $h(\cdot, \lambda)$ to have a root in the interval $(0, 1)$. In this case, we infer from similar arguments as Lemma 3 that b_2^* is the unique root whenever $\lambda = 0$. Observe that $D_1 h(0, \lambda) \rightarrow -\infty$ as $\lambda \rightarrow -\infty$, so that we may choose an integer $n > |\lambda|$ such that $D_1 h(0, -n) < 0$. The maximizer of $h(\cdot, -n)$ is negative by virtue of (11) where $\lambda = -n$ and $b_2 = 0$, as does the global maximum due to

$$h(b, -n) < h(b, 0) < 0.$$

for $b < 0$. By contrast, $h(b, 0)$ has a positive maximizer and a positive global maximum. Considering the continuity of the maximizer and global maximum of $h(\cdot, \lambda)$ as functions of λ , it follows that $-n < \hat{\lambda} < 0$. Now, recall that $h(\cdot, \hat{\lambda})$ is maximized at $h(\hat{b}, \hat{\lambda}) = 0$. The desired trichotomy holds by comparing $h(b, \lambda)$ with $h(b, \hat{\lambda})$ and $h(b, 0)$, and appealing to the linear increasing property of h in λ (Lemma 1); this can be associated with the shaded region bounded by $h(b, \hat{\lambda})$ and $h(b, 0)$, located in the right panel of Figure 3. $\square$

Figure 3: The function $h(\cdot, \lambda)$ depicted at different values of λ . In the left panel, $\bar{\gamma} < \bar{\gamma}^*$ for which a unique positive root exists if and only if $\lambda > 0$. The right panel considers the case where $\bar{\gamma} > \bar{\gamma}^*$. The shaded region identifies the family of curves generated by h where $\hat{\lambda} < \lambda < 0$, for which two distinct roots exist. The region is bounded by the bifurcation values $\lambda = 0$ and $\lambda = \hat{\lambda}$. No root exists when $\lambda < \hat{\lambda}$ (bottom curve) and exactly one positive root exists for $\lambda > 0$ (top curve).

Corollary. *Given our bifurcation parameter λ , the only positive equilibrium points for the dimensionless model (4)–(6) are of the form*

$$E_k^* := (b_k^*, \mu(b_k^*, \lambda), f(b_k^*)), \quad k = 1, 2,$$

where b_1^* and $b_2^* > b_1^*$ are the smallest and the largest positive roots of $h(\cdot, \lambda)$, respectively. If $\lambda > 0$, then only E_2^* exists. If $\lambda \leq 0$ then positive equilibrium points exist only if $D_1 h(0, 0) > 0$ and $\hat{b} > 0$. In this case, the following trichotomy holds: both E_1^* and E_2^* exist if $\hat{\lambda} < \lambda \leq 0$; only E_2^* exists where $b_2^* = \hat{b}$ if $\lambda = \hat{\lambda}$; and no positive equilibrium point exists when $\lambda < \hat{\lambda}$.

Proof. Consider a positive equilibrium point $E^* = (b^*, m^*, n^*)$. Then the following equations hold:

$$0 = \frac{m^*(\lambda - 1)}{m_0^*} - (b^* - 1), \quad (20)$$

$$0 = m^* \left[n^* \bar{\delta}_M \left(1 - \frac{1}{m_0^*} \right) + \bar{\eta} b^* \right] - \bar{\delta}_M (m^* - 1), \quad (21)$$

$$0 = \bar{\gamma} b^* - \bar{\delta}_N (n^* - 1). \quad (22)$$

201 We rewrite equations (20) and (22) into $m^* = \mu(b^*, \lambda)$ and $n^* = f(b^*)$, respectively.
 202 Thus, solving (21) in n^* after evaluating m^* yields $n^* = g(b^*, \lambda)$. Consequently,
 203 $h(b^*, \lambda) = f(b^*) - g(b^*, \lambda) = 0$ and b^* is either b_1^* or b_2^* by Lemma 3. Thus, E^* is
 204 either E_1^* and E_2^* . For each k , the equilibrium point E_k^* exists if and only if b_k^* exists.
 205 Therefore, the existence E_1^* and E_2^* follows from Theorem 2. $\square$

Now, let

$$\widehat{b}_0 := \frac{1}{2} \left(1 - \frac{\zeta}{\xi} \right), \quad \Delta := 4 \left[(\widehat{b}_0)^2 + \frac{\zeta\lambda}{\xi} \right]. \quad (23)$$

Then we derive the following equation:

$$\frac{h(b, \lambda)}{D_2 h(b, \lambda)} = \frac{\xi}{\zeta} \left[(\widehat{b}_0)^2 - (b - \widehat{b}_0)^2 \right] + \lambda, \quad (24)$$

where $D_2 h > 0$ (Lemma 1). Hence, the roots b_1^* and b_2^* of $h(\cdot, \lambda)$ are given by

$$b_1^* = \widehat{b}_0 - \frac{\sqrt{\Delta}}{2}, \quad b_2^* = \widehat{b}_0 + \frac{\sqrt{\Delta}}{2}. \quad (25)$$

Observe that $\widehat{b}_0 > 0$ if and only if $D_1 h(0, 0) > 0$ according to (14). From the definitions of Δ and equation (15), the following equations are equivalent:

$$\Delta = 0, \quad (1 - \widehat{b}_0)^2 = \frac{\zeta(1 - \lambda)}{\xi}, \quad \widehat{b} = \widehat{b}_0. \quad (26)$$

If one (hence all) of the equations in (26) is true, then $\widehat{b} = \widehat{b}_0 = b_2^*$ by (25) and equivalently $\lambda = \widehat{\lambda}$ by Lemma 3. We may write $\widehat{\lambda}$ in terms of the dimensionless parameters by solving for $\Delta = 0$:

$$\widehat{\lambda} = 1 + \frac{(\overline{\delta_M} \overline{\delta_N} - \overline{\delta_N} \overline{\eta} + \overline{\delta_N} \overline{\theta} + \overline{\gamma} \overline{\theta})^2}{4 \overline{\delta_N} (\overline{\delta_M} + \overline{\theta}) (\overline{\delta_N} \overline{\eta} - \overline{\gamma} \overline{\theta})}. \quad (27)$$

206 **Theorem 3.** Suppose that E_k^* exists for a given k , and that all eigenvalues of $J(E_k^*)$
 207 have nonzero real part. Then E_k^* is a saddle point for $k = 1$ and asymptotically
 208 stable for $k = 2$.

209 *Proof.* Consider the Jacobian matrix $J(E_k^*)$. To obtain a practical expression of
 210 $J(E_k^*)$, we simplify the first and second diagonal entries by application of equa-
 211 tions (20) and (21), respectively; by i th diagonal entry, we mean the (i, i) -entry. For
 212 the off-diagonal entries, we perform the following algebraic manipulations:

- 213 (i) In the top row, write m_0^* in terms of $D_1 \mu(b_k^*, \lambda)$.
- 214 (ii) In the middle row, evaluate $m^* = \mu(b_k^*, \lambda)$ and write $\overline{\eta}$ in terms of $D_1 g(b_k^*, \lambda)$.
- 215 (iii) In the bottom row, write $\overline{\gamma} = \overline{\delta_N} D_1 f(b_k^*)$.

Additionally, we apply the equation $\overline{\delta_M}(m_0^* - 1)/m_0^* = -\overline{\theta}$ wherever simplification is desired. Thus, we arrive at the following expression:

$$J(E_k^*) = \begin{bmatrix} -b_k^* & \frac{b_k^*}{D_1 \mu(b_k^*, \lambda)} & 0 \\ J_{21} & \frac{-\overline{\delta_M}}{\mu(b_k^*, \lambda)} & -\overline{\theta} \mu(b_k^*, \lambda) \\ \overline{\delta_N} D_1 f(b_k^*) & 0 & -\overline{\delta_N} \end{bmatrix},$$

where

$$J_{21} = \frac{\overline{\delta_M} D_1 \mu(b_k^*, \lambda)}{\mu(b_k^*, \lambda)} + \overline{\theta} \mu(b_k^*, \lambda) D_1 g(b_k^*, \lambda).$$

Denoting the trace and determinant by tr and $\det$, respectively, $J(E_k^*)$ must satisfy three Routh-Hurwitz conditions for E_k^* to be asymptotically stable. The first condition holds for both E_1^* and E_2^* , that is:

$$-\text{tr } J(E_k^*) = b_k^* + \frac{\overline{\delta_M}}{\mu(b_k^*, \lambda)} + \overline{\delta_N} > 0, \quad k = 1, 2,$$

given that $\mu > 0$. The second condition requires

$$\det J(E_k^*) = \frac{-b_k^* \overline{\theta} \overline{\delta_N} \mu(b_k^*, \lambda) D_1 h(b_k^*, \lambda)}{D_1 \mu(b_k^*, \lambda)}$$

216 to be negative. Since $D_1 \mu < 0$, the determinant $\det J(E_k^*)$ shares the same sign
 217 with $D_1 h(b_k^*, \lambda)$. Thus, by Lemma 3, the second condition fails for E_1^* because
 218 $D_1 h(b_1^*, \lambda) > 0$. Since we assumed that all eigenvalues have nonzero real part, E_k^*
 219 must be a saddle point for $k = 1$. Meanwhile, Theorem 2 implies that for E_2^* to
 220 exist, it is necessary that either $\lambda > 0$ or E_2^* coexists with E_1^* . In either case,
 221 $D_1 h(b_2^*, \lambda) < 0$ and the second Routh-Hurwitz condition holds for $k = 2$.

We are left to verify the following last Routh-Hurwitz condition for E_2^* , *i.e.*, assuming that $k = 2$:

$$\sigma := -\text{tr } J(E_2^*) (\sigma_1 + \sigma_2) + \det J(E_2^*) > 0,$$

where

$$\sigma_1 := \frac{\overline{\delta_N} (\overline{\delta_M} + b_2^* \mu(b_2^*, \lambda))}{\mu(b_2^*, \lambda)} > 0 \quad \sigma_2 := \frac{-b_2^* \overline{\theta} \mu(b_2^*, \lambda) D_1 g(b_2^*, \lambda)}{D_1 \mu(b_2^*, \lambda)}.$$

Assuming that E_2^* exists, observe that σ_2 has the same sign with $D_1 g(b_2^*, \lambda)$. Since $D_1 h(b_2^*, \lambda) < 0$ and f has a positive first derivative (Lemma 1), we have

$$D_1 g(b_2^*, \lambda) > D_1 f(b_2^*) > 0.$$

and $\sigma_2 > 0$. Moreover, we have

$$\begin{aligned} \sigma &> (b_2^* + \overline{\delta_N}) \sigma_2 + \det J(E_2^*) \\ &= \frac{\sigma_2 [\overline{\delta_N} D_1 f(b_2^*) + b_2^* D_1 g(b_2^*, \lambda)]}{D_1 g(b_2^*, \lambda)} \end{aligned}$$

222 and $\sigma > 0$. Therefore, E_2^* satisfies all three Routh-Hurwitz conditions and is conse-
 223 quently asymptotically stable. $\square$

224 The existence of roots, as well as their stability—corollary of Theorem 2, and
 225 Theorem 3, respectively—are summarized in Figure 4, showing an illustration of the
 226 different stability regions in the $(\overline{\gamma}, \lambda)$ -plane, and numerical bifurcation plots for b
 227 as a function of λ .

Figure 4: **Top:** Regions of stability determined by λ and $\bar{\gamma}$, each highlighting the corresponding stable equilibrium points; E_0^* has $b = 0$, while E_2^* takes the largest steady-state value b_2^* for b . The specific value $\bar{\gamma}^*$ of γ is given in equation (19), while $\hat{\lambda}$ is explicitly given in (27).

Bottom: Bifurcation diagrams of the system (4)–(6) without (left) and with (right) bistability—respectively, $\bar{\gamma} < \bar{\gamma}^*$ and $\bar{\gamma} > \bar{\gamma}^*$. To generate the diagrams, all parameters were fixed except for $\bar{c}_B$, which was obtained for a given λ from (7). Blue, solid lines: stable equilibria; orange, dashed lines: unstable equilibria. The gray diamond appearing in the right panel highlights the bifurcation point $(\hat{\lambda}, \hat{b})$.

228 4. Discussion

229 Our study is centered on the problem of identifying biological factors that con-
 230 tribute to the elimination of the pneumococcus (*Streptococcus pneumoniae*). Ex-
 231 periments by Smith et al. [19] suggested that inoculum size (dosage) determines
 232 the outcome of bacterial clearance or persistence. In this case, groups of mice were
 233 infected with the pneumococcus at different dose sizes. Each group corresponded
 234 to a single bacterial outcome indicated by the titer readings, depending on whether
 235 the dose size is above or below a threshold. In contrast, experiments from Duvig-
 236 neau et al. [27] showed that inoculum size is not the only factor contributing to
 237 bacterial clearance. Here, all mouse subjects were given doses of identical size. To-
 238 wards the end of the experiment, the bacterial load was undetectable in some mice
 239 and sustained in the rest (Figure 2). The murine experiments of Smith et al. [19]
 240 and Duvigneau et al. [27] provide distinct perspectives on the trade-off between
 241 dosage and bacterial fate, restricted by the microbial instances compatible with the
 242 experimental design.

243 By modeling bacterial kinetics with generalized innate-regulatory immune re-
 244 sponses, our mathematical analysis reveals a qualitative trichotomy of this trade-off
 245 that acknowledges the experiments of both Smith et al. [19] and Duvigneau et al. [27].
 246 Indeed, our model, given by system (1)–(3) and the equivalent dimensionless form
 247 (4)–(6), may be considered an abstraction of previous formulations [17, 18, 16, 15, 19]
 248 where overall immune responses are considered instead of specific phagocyte popu-
 249 lations and chemical mediators. The trichotomy is given by the following cases: (i)
 250 *dose-independent clearance*, where the immune response clears the pneumococcus
 251 independent of dose size; (ii) *dose-independent persistence*, where the pneumococ-
 252 cus outgrows immunity regardless of initial dose; and (iii) *dose-limited clearance*,
 253 where the immune system successfully eliminates the pneumococcus only in small
 254 quantities. Cases (i) and (ii) are corroborated by Duvigneau et al. [27], whereas case
 255 (iii) is supported by Smith et al. [19].

256 We remark that successful clearance of the pneumococcus may also be attributed
 257 to empirical characteristics of the infection other than the inoculum size. Mochan
 258 et al. [21] formulated a model validated with murine datasets including [19] to de-
 259 scribe pulmonary and extrapulmonary pneumococcal kinetics with total phagocyte
 260 levels and damage to epithelial cells (cellular debris) and a homogeneous population
 261 of activated phagocytes. The corresponding simulations indicate that phagocyte
 262 clearance efficiency varies between mouse strains. A follow-up study [22] using the
 263 same model shows that mutations in the pneumococcal strain can influence transient
 264 reduction in pneumococcus. In addition, Schirm et al. [20] adapted the monocyte-
 265 derived macrophage murine model in [19] and incorporated inhalation and antibi-
 266 otic effects to pneumococcal growth and elimination. The models of Mochan et
 267 al. [21, 22] and Schirm et al. [20] are directed towards investigating the effects of
 268 treatment and strain variations on pneumococcal clearance. However, their mod-
 269 eling approaches focus on validation with experimental data instead of bifurcation.
 270 While our mathematical model only considers bacterial kinetics with generalized
 271 innate and regulatory levels via three equations, the results of our comprehensive
 272 stability analysis could provide valuable and testable hypotheses regarding pathogen
 273 clearance.

The following quantities were determined to drive the dynamics of our model:

$$\lambda = 1 - \overline{c_B} m_0^*, \quad D_1 h(0, 0) = \frac{\overline{\gamma} - \overline{\gamma}^*}{\overline{\delta_N}},$$

where

$$\overline{\gamma}^* = \left(\frac{\overline{\eta} + \overline{\delta_M}}{\overline{\theta}} + 1 \right) \overline{\delta_N}.$$

See equations (7) and (19). The stability of the bacteria-free steady state E_0^* according to λ (Theorem 1) portrays the effectiveness of the innate immune response to clear small pneumococcal quantities. Now, λ is a linearly decreasing function of $\overline{c_B}$, which is proportional to the clearance rate c_B with the initial innate response level $M_0 = M(0)$ and all other parameters fixed. Since E_0^* is stable when $\lambda < 0$ or $\overline{c_B} > 1/m_0^*$ (Theorem 1), a rapid innate clearance (large c_B) could promote bacterial eradication at small quantities with the goal of making E_0^* stable. Effective clearance may also hold in mild conditions (moderate values of c_B and M_0) for a large innate level m_0^* . By the same token, $\overline{c_B}$ exhibits inverse proportionality with the maximum logistic proliferation rate r of the bacteria. Thus, the outgrowth of the pneumococcus may benefit from rapid proliferation (large r).

The overall dynamics of our model follows from our main results for positive equilibria. The corollary of Theorem 2 determines which of E_1^* and E_2^* exist, and Theorem 3 establishes that E_1^* is unstable and E_2^* is asymptotically stable. Since $D_1 h(0, 0)$ has the same sign with $\overline{\gamma} - \overline{\gamma}^*$, we may frame our discussion in terms of the dimensionless parameter $\overline{\gamma}$. As illustrated in Figure 4, $\overline{\gamma}$ determines which of the three bacterial outcomes (clearance, persistence, dose-limited clearance) are possible while λ decides which outcome the model predicts.

If the model assumes that $\overline{\gamma} \leq \overline{\gamma}^*$, then bistability does not occur and the model only predicts dose-independent clearance and persistence. That is, the stable equilibrium point is uniquely given by E_0^* for $\lambda < 0$ and E_2^* for $\lambda > 0$. Hence, we expect the innate immune system to eliminate the bacteria for $\lambda < 0$, and for the bacteria to persist for $\lambda > 0$, regardless of the initial bacterial concentration. As $\overline{\gamma}$ increases so that $\overline{\gamma} > \overline{\gamma}^*$, the negative values for λ corresponding to dose-independent clearance are restricted to the interval $(-\infty, \hat{\lambda})$ where $\hat{\lambda} < 0$. Moreover, bistability holds for all values of λ in the interval $(\hat{\lambda}, 0)$, where the unstable equilibrium point E_1^* coexists with the stable equilibria E_0^* and E_2^* . The size of the interval $(\hat{\lambda}, 0)$ changes with $\overline{\gamma}$ according equation (27). We emphasize that the monostability at E_2^* when $\lambda > 0$ is independent of $\overline{\gamma}$.

When scaled with $1/\overline{\delta_N}$ (with fixed $\overline{\delta_N}$), we find that $\overline{\gamma}$ is directly proportional to the tissue carrying capacity K_B and the ratio $\gamma/(\delta_N N_0)$ of innate response promotion to constant replenishment rate. Hence, the model predicts that abundant tissue resources (hypothetically large K_B), and rapid activation of regulatory responses ($\gamma > \delta_N N_0$) contribute to the range of parameter values for bistability and dose-limited clearance. In light of the monostability of E_2^* above, we predict that the corresponding dose-independent persistence may neither depend on tissue carrying capacity nor the activation of regulatory responses.

In the bistability case, the bacterial load b_1^* at E_1^* can serve as a threshold for bacterial clearance. Based on our bifurcation diagrams (Figure 4), one could naively deduce that the immune system clears the bacteria for $b(0) = B(0)/K_B < b_1^*$, and the

314 bacteria succeeds in colonizing the host when $b(0) > b_1^*$ (cf. [25]). However, we must
 315 emphasize that our model is a *three*-dimensional system where stable manifolds of a
 316 saddle node may be one-dimensional curves or two-dimensional surfaces. A deeper
 317 analysis requires investigating the stable and unstable manifolds of E_1^* , which delimit
 318 the basins of attraction for E_0^* and E_2^* .

At this point, we discuss generalizations and future directions of our work. The
 aforementioned local stability as dependent on λ may be qualitatively identified with
 compatible systems exhibiting nonlinear interaction terms. This can be achieved
 with the function

$$G : [0, \infty) \times (0, \infty) \rightarrow [0, 1), \quad G(x, a) = \frac{x}{x + a}.$$

319 For a fixed a , the function $G(\cdot, a)$ typically introduces saturation effects on a
 320 growth/decay rate: the value of $G(x, a)$ approaches its upper bound ($G \approx 1$) with
 321 larger values of x . In different biological contexts, G is associated with the Monod
 322 growth term for microorganisms, the Michaelis-Menten equation for enzyme kinetics,
 323 and the Holling Type-II functional response for predator-prey dynamics.

To demonstrate the robustness of our results to nonlinear interaction terms, we
 show in Figure 5 that a modification of the model (1)–(3) incorporating nonlin-
 ear interaction terms yields comparable qualitative dynamics. Moreover, the same
 trichotomy of bacterial outcomes applies here. The modified system is given by

$$\dot{B} = rB \left(1 - \frac{B}{K_B} \right) - c_B B G(M, K_M), \quad (28)$$

$$\dot{M} = \delta_M(M_0 - M) + [\eta B - \theta G(N, K_N)] M, \quad (29)$$

$$\dot{N} = \gamma G(B, \rho) - \delta_N N. \quad (30)$$

324 This model imposes the following effects: (i) saturated bacterial clearance with high
 325 levels of innate immune response; (ii) bounded regulation of innate response; and
 326 (iii) limited increase in regulatory levels for larger bacterial loads.

Proceeding as before, we generate a bifurcation diagram for (28)–(30) based on
 the eigenvalue

$$\lambda = 1 - \frac{\overline{c_B} m_0}{1 + m_0} \quad (31)$$

characterizing the stability of the unique bacteria-free equilibrium. Here, we have
 used the dimensionless quantities $m_0 = M_0/K_M$ and $\overline{c_B} = c_B/r$, and the time has
 again been rescaled as $\tau = t/r$. The bistability region is now determined by

$$\overline{\gamma}^* = \left(\frac{\overline{c_B} \overline{\delta_M}}{\overline{\theta}} + 1 \right) \frac{\overline{\rho} \overline{\delta_N} \overline{\eta}}{\overline{\theta}}, \quad (32)$$

327 with $\overline{\gamma}$, $\overline{\delta_M}$, $\overline{\theta}$, $\overline{\delta_N}$ and $\overline{\eta}$ as defined before, and $\overline{\rho} = \rho/K_B$. As in the original model
 328 (1)–(3), the parameter $\overline{\gamma}$ determines whether bistability exists, and λ determines
 329 which outcome is predicted. However, the formulation of λ in (31) is independent
 330 of parameters pertaining to innate response as opposed to the original formulation
 331 in (7). This observation suggests that, to an extent, bacterial clearance may depend
 332 on other factors aside from innate response when nonlinear cellular responses are in
 333 action.

Figure 5: Bifurcation diagrams generated by the system (28)–(30), where $b = B/K_B$ and λ , given by (31), is the eigenvalue that determines the stability of the unique equilibrium point satisfying $b = 0$. These diagrams are generated via the same procedure as the ones from Fig. 4, and are shown for $\bar{\gamma} < \bar{\gamma}^*$, corresponding to the case without bistability (left), and $\bar{\gamma} > \bar{\gamma}^*$, corresponding to the bistable case (right); here, $\bar{\gamma}^*$ is given by (32). Blue, solid lines: stable equilibria; orange, dashed lines: unstable equilibria.

Acknowledgments

This work was supported by the Boehringer Ingelheim Stiftung (Exploration Grant, VIBA project) and the Alfons und Gertrud Kassel-Stiftung.

References

- [1] B. Henriques-Normark, E. I. Tuomanen, The pneumococcus: epidemiology, microbiology, and pathogenesis, Cold Spring Harb. Perspect. Med. 3 (2013) a010215.
- [2] J. F. Brundage, Interactions between influenza and bacterial respiratory pathogens: implications for pandemic preparedness, Lancet Infect. Dis. 6 (2006) 303–312.
- [3] R. K. Gupta, R. George, J. S. Nguyen-Van-Tam, Bacterial pneumonia and pandemic influenza planning, Emerg. Infect. Dis. 14 (2008) 1187.
- [4] D. E. Morris, D. W. Cleary, S. C. Clarke, Secondary bacterial infections associated with influenza pandemics, Front. Microbiol. 8 (2017) 1041.
- [5] A. M. Smith, J. A. McCullers, Secondary Bacterial Infections in Influenza Virus Infection Pathogenesis, Springer International Publishing, Cham, pp. 327–356.
- [6] E. Varon, J. Mainardi, L. Gutmann, *Streptococcus pneumoniae*: still a major pathogen, Clin. Microbiol. Infect. 16 (2010) 401.
- [7] K. L. O’Brien, L. J. Wolfson, J. P. Watt, E. Henkle, M. Deloria-Knoll, N. McCall, E. Lee, K. Mulholland, O. S. Levine, T. Cherian, Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates, Lancet 374 (2009) 893–902.
- [8] T. Welte, A. Torres, D. Nathwani, Clinical and economic burden of community-acquired pneumonia among adults in Europe, Thorax 67 (2012) 71–79.
- [9] F. Blasi, M. Mantero, P. Santus, P. Tarsia, Understanding the burden of pneumococcal disease in adults, Clin. Microbiol. Infect. 18 (2012) 7–14.
- [10] C. Ghia, M. Wasserman, M. Fletcher, R. Farkouh, G. Rambhad, Modeling possible inclusion of pneumococcal conjugate vaccine into the National Immunization Program for infants in India, Value Health Reg. Issues 15 (2018) 99–105.
- [11] M.-C. C. Brandileone, S. C. Almeida, R. Minamisava, A.-L. Andrade, Distribution of invasive *Streptococcus pneumoniae* serotypes before and 5 years after the introduction of 10-valent pneumococcal conjugate vaccine in Brazil, Vaccine 36 (2018) 2559–2566.
- [12] E. Usuf, A. Bojang, B. Camara, I. Jagne, C. Oluwalana, C. Bottomley, U. D’Alessandro, A. Roca, Maternal pneumococcal nasopharyngeal carriage and risk factors for neonatal carriage after the introduction of pneumococcal conjugate vaccines in The Gambia, Clin. Microbiol. Infect. 24 (2018) 389–395.
- [13] J. N. Weiser, D. M. Ferreira, J. C. Paton, *Streptococcus pneumoniae*: transmission, colonization and invasion, Nat. Rev. Microbiol. 16 (2018) 355–367.
- [14] C. Feldman, R. Anderson, Recent advances in our understanding of *Streptococcus pneumoniae* infection, F1000Prime Rep. 6 (2014) 82. 82[PII].

- 376 [15] A. Reynolds, J. Rubin, G. Clermont, J. Day, Y. Vodovotz, G. B. Ermentrout, A
377 reduced mathematical model of the acute inflammatory response: I. derivation
378 of model and analysis of anti-inflammation, *J. Theor. Biol.* 242 (2006) 220–236.
- 379 [16] S. S. Pilyugin, R. Antia, Modeling immune responses with handling time, *Bull.*
380 *Math. Biol.* 62 (2000) 869–890.
- 381 [17] D. A. Lauffenburger, Mathematical analysis of the macrophage response to
382 bacterial challenge in the lung, Springer Netherlands, Dordrecht, pp. 351–358.
- 383 [18] S. G. Rudnev, A. A. Romanyukha, Mathematical modeling of immune-
384 inflammatory reaction in acute pneumonia, *J. Biol. Syst.* 03 (1995) 429–439.
- 385 [19] A. M. Smith, J. A. McCullers, F. R. Adler, Mathematical model of a three-
386 stage innate immune response to a pneumococcal lung infection, *J. Theor. Biol.*
387 276 (2011) 106–116.
- 388 [20] S. Schirm, P. Ahnert, S. Wienhold, H. Mueller-Redetzky, G. Nouailles-Kursar,
389 M. Loeffler, M. Witzenthath, M. Scholz, A biomathematical model of pneumo-
390 coccal lung infection and antibiotic treatment in mice, *PLoS ONE* 11 (2016)
391 1–22.
- 392 [21] E. Mochan, D. Swigon, G. B. Ermentrout, S. Lukens, G. Clermont, A math-
393 ematical model of intrahost pneumococcal pneumonia infection dynamics in
394 murine strains, *J. Theor. Biol.* 353 (2014) 44 – 54.
- 395 [22] E. Mochan-Keef, D. Swigon, G. B. Ermentrout, G. Clermont, A three-tiered
396 study of differences in murine intrahost immune response to multiple pneumo-
397 coccal strains, *PLoS ONE* 10 (2015) 1–18.
- 398 [23] E. Domínguez-Hüttinger, N. J. Boon, T. B. Clarke, R. J. Tanaka, Mathemat-
399 ical modeling of *Streptococcus pneumoniae* colonization, invasive infection and
400 treatment, *Front. Physiol.* 8 (2017) 115.
- 401 [24] S. Strogatz, Nonlinear Dynamics and Chaos: With Applications to Physics,
402 Biology, Chemistry, and Engineering, CRC Press, 2018.
- 403 [25] R. Malka, E. Shochat, V. Rom-Kedar, Bistability and bacterial infections,
404 *PLoS ONE* 5 (2010) 1–10.
- 405 [26] Y. Li, A. Karlin, J. D. Loike, S. C. Silverstein, Determination of the critical
406 concentration of neutrophils required to block bacterial growth in tissues, *J.*
407 *Exp. Med.* 200 (2004) 613–622. 20040725[PII].
- 408 [27] S. Duvigneau, N. Sharma-Chawla, A. Boianelli, S. Stegemann-Koniszewski,
409 V. K. Nguyen, D. Bruder, E. A. Hernandez-Vargas, Hierarchical effects of pro-
410 inflammatory cytokines on the post-influenza susceptibility to pneumococcal
411 coinfection, *Sci. Rep.* 6 (2016) 37045.
- 412 [28] E. Jones, T. Oliphant, P. Peterson, et al., SciPy: Open source scientific tools
413 for Python, 2001–. [Online; accessed September 17, 2018].
- 414 [29] R. Storn, K. Price, Differential evolution—a simple and efficient heuristic for
415 global optimization over continuous spaces, *J. Glob. Optim.* 11 (1997) 341–359.