1 **Sexual selection rewires reproductive protein networks**

2

3

4 Timothy L. Karr^{1,*§}, Helen Southern², Matthew Rosenow³, Toni I. Gossmann², and Rhonda R.
5 Snook^{4,*}

6

7 ¹Department of Genomics and Genetic Resources, Kyoto Institute of Technology, Kyoto,
8 Japan

9 ²Department of Animal and Plant Sciences, University of Sheffield, UK.

10 ³Caris Life Sciences, Phoenix, AZ

11 ⁴Department of Zoology, Stockholm University, Sweden

12 *Corresponding authors: Email: tkarr@asu.edu & rhonda.snook@zoologi.su.se

13 §Current address: Arizona State University, School of Life Sciences, Tempe, AZ 85287

14

15

16 Running title: Sexual selection and protein interaction networks

17 **The abbreviations used are:**

18 BLAST, Basic Local Alignment Search Tool

19 Dpse, *Drosophila pseudoobscura*

20 PCSS, postcopulatory sexual selection

21 SFPs, seminal fluid proteins

22 Dmel, *D. melanogaster*

23 SDS, sodium dodecylsulfate

- 24 SDS-PAGE, sodium dodecylsulfate polyacrylamide gel electrophoresis
- 25 MS, mass spectrometry
- 26 LC-MS/MS, liquid chromatography-MS/MS
- 27 AcgP, accessory gland proteome
- 28 FDRs, False Discovery Rates
- 29 AcgS, accessory gland secretome
- 30 exoP, exoproteome
- 31 LFQ, label-free quantitation
- 32 P, polyandry
- 33 M, monandry
- 34 GO, gene ontology
- 35 CC, cellular component
- 36 MF, molecular function
- 37 BP, biological process
- 38 STRING, Search Tool for the Retrieval of Interacting Genes/Proteins
- 39 DIOPT, DRSC Integrative Ortholog Prediction Tools
- 40 ER, endoplasmic reticulum

41 Polyandry drives postcopulatory sexual selection (PCSS), resulting in rapid evolution
 42 of male ejaculate traits. Critical to male and female fitness, the ejaculate is known to
 43 contain rapidly evolving seminal fluid proteins (SFPs) produced by specialized male
 44 secretory accessory glands. The evidence that rapid evolution of some SFPs is driven
 45 by PCSS, however, is indirect, based on either plastic responses to changes in the
 46 sexual selection environment or correlative macroevolutionary patterns. Moreover,
 47 such studies focus on SFPs that represent but a small component of the accessory
 48 gland proteome. Neither how SFPs function with other reproductive proteins, nor how
 49 PCSS influences the underlying secretory tissue adaptations and content of the
 50 accessory gland, has been addressed at the level of the proteome. Here we directly test
 51 the hypothesis that PCSS results in rapid evolution of the entire male accessory gland
 52 proteome and protein networks by taking a system-level approach, combining
 53 divergent experimental evolution of PCSS in *Drosophila pseudoobscura* (Dpse), high
 54 resolution mass spectrometry (MS) and proteomic discovery, bioinformatics and
 55 population genetic analyses. We demonstrate that PCSS influences the abundance of
 56 over 200 accessory gland proteins, including SFPs. A small but significant number of
 57 these proteins display molecular signatures of positive selection. Divergent PCSS also
 58 results in fundamental and remarkably compartmentalized evolution of accessory
 59 gland protein networks in which males subjected to strong PCSS invest in protein
 60 networks that serve to increase protein production whereas males subjected to relaxed
 61 PCSS alters protein networks involved in protein surveillance and quality. These results
 62 directly demonstrate that PCSS is a key evolutionary driver that shapes not only
 63 individual reproductive proteins, but rewires entire reproductive protein networks.

64 Polyandry, in which females mate with different males across a reproductive bout, generates
 65 PCSS in which ejaculates compete for fertilization of a limited supply of ova and females
 66 may choose whose sperm will fertilize those limited ova (1). Polyandry also engenders sexual
 67 conflict, in which male and female reproductive interests differ, as a consequence of the
 68 disproportionate costs and benefits of mating between the sexes (2). PCSS operates in
 69 internally fertilizing organisms between the female reproductive tract and the male ejaculate,
 70 which is composed of both sperm and non-sperm components, including SFPs (3). SFPs
 71 were first identified by their canonical signal peptide sequence that direct proteins to the
 72 secretory pathway (4). Cross-species comparative work has found that general classes of
 73 SFPs are conserved (e.g., proteases and protease inhibitors, lectins and prohormones)
 74 suggesting that their mechanisms of action are also conserved. However, individual SFPs
 75 can rapidly evolve with signals of accelerated rates of adaptive molecular evolution found in
 76 studies of coding sequence and male-biased gene expression observed across different
 77 animal taxa (e.g., mammals (5, 6); birds (7); *Drosophila* (8–10)).

78
 79 Rapid evolution of SFPs is attributed to PCSS because this cocktail of proteins, transferred to
 80 females, has dramatic and sometimes antagonistic effects on male and female fitness,
 81 including increased female fecundity, reduced female receptivity, decreased female life span,
 82 and remodelling of female reproductive tract morphology (4, 11–13). Moreover, males can
 83 tailor ejaculate composition, altering the quantity of specific SFPs, via plastic changes in
 84 response to variation in the PCSS environment (5, 14) although a mechanism by which males
 85 accomplish this is unknown. However, while many studies have examined the evolution and
 86 fitness consequences of SFPs, most of them examine the impact of one or just a few of these
 87 proteins. This focus is problematic because SFPs do not work in isolation, but rather as
 88 partners in protein-protein and protein-nucleic acid networks, and not just solely with other
 89 SFPs (15, 16). Therefore, changes in either protein levels or protein function may have
 90 significant up- and/or down-stream consequences on organismal physiology and fitness.
 91 Indirect studies of the role of PCSS on reproductive protein evolution have not considered

92 other proteins also produced by male accessory gland tissue that may interact with SFPs and
93 influence fitness.

94

95 Most of our current knowledge of SFPs comes from traditional protein detection methods and
96 comparisons aimed at one species, *D. melanogaster* (9, 17–19). However, the advent of
97 sophisticated transcriptomic, proteomic and bioinformatics approaches have identified new
98 SFPs and associated reproductive proteins not only in Dmel but also other taxa (20–26).
99 Furthermore, high throughput mass spectrometry based techniques now routinely produce
100 cellular and tissue proteomes consisting of thousands of protein IDs not only useful for protein
101 identification but also the analysis and identification of protein interaction networks.

102

103 Initial application using these techniques to the analysis of the sperm proteome (24, 25) found
104 heterogeneity in the evolutionary rates of genes compared to SFPs which lead to the
105 hypothesis that adaptive compartmentalization can occur both within and between proteomes
106 involved in reproduction. This hypothesis suggests that interacting reproductive proteins would
107 comprise a set of “core” proteins with essential cellular functions (e.g., sperm motility) that are
108 predicted to be under strong purifying selection and another set of proteins under adaptive
109 selection generated from PCSS, e.g., sperm-egg interactions (24). Other than sperm, direct
110 empirical confirmation of adaptive compartmentalization, and a possible role of PCSS in this
111 process, has yet to be undertaken. Likewise, application of high throughput mass spectrometry
112 based techniques have allowed characterization and dynamics of protein interaction networks
113 in a variety of cellular contexts including disease-induced perturbation and or over evolutionary
114 time frames (see (27, 28) for recent reviews). However, the role of PCSS in shaping the
115 architecture of protein interaction networks is unknown.

116

117 Here we address the overall qualitative and quantitative impact of PCSS on the male
118 accessory gland proteome and directly test the hypothesis that PCSS effects SFP production
119 and their reproductive protein networks. We also test for signatures of positive selection and

examine evidence for adaptive compartmentalization. We perform these tests using replicated experimentally evolved lines of *D. pseudoobscura* (Dpse) in which PCSS was manipulated over many generations by either allowing polyandry (“P” lines; 1 female with 6 males) or enforcing monogamy (“M” lines; 1 female with 1 male) (29). Previous studies using these lines have found that divergent sexual selection impacts the evolution of reproductive traits relevant to PCSS and SFP production, including P males altering female investment in both early and total progeny production (30) and P males having larger accessory glands that resulted in greater reproductive success (31). Sex-biased gene expression has also evolved in these lines (32, 33), and, taken together these results strongly motivate using these lines to directly test the role of PCSS in shaping male reproductive proteomes at the microevolutionary scale.

EXPERIMENTAL PROCEDURES

Experimental lines. The establishment and maintenance of the selection lines were previously described in detail (29). Briefly, an ancestral wild-caught population of Dpse from Tucson Arizona USA, a naturally polyandrous species (wild caught females have been shown to be frequently inseminated by at least two males at any given time; (34)), was used to establish the selection lines. From this population, four replicate lines (replicates 1-4) of two different sexual selection treatments were established. To modify the opportunity for sexual selection, adult sex ratio in vials was manipulated by either confining one female with a single male and enforcing monogamy (“monogamy” treatment, M) or one female with six males promoting polyandry (“polyandry” treatment, P (NB: this treatment has also been referred to as E in other publications). Effective population sizes are equalized between the treatments as described previously (35). At each generation, offspring are collected and pooled together for each replicate line, and a random sample from this pool is used to constitute the next generation in the appropriate sex ratios, thus proportionally reflecting the differential offspring production across families. In total, eight selection lines (M1, M2, M3, M4 and P1, P2, P3, P4) are maintained, in standard vials (2.5 × 80 mm), with a generation time of 28 days. All populations are kept at 22°C on a 12L:12D cycle, with standard food media and added live yeast.

148

149 *Experimental flies.* Flies from replicates 1-4 of each of the selection lines were collected from
 150 generations 157, 156, 155 and 153 respectively. We standardized for maternal and larval
 151 environments (29), but in brief, parental flies were collected and housed en-mass in food
 152 bottles, then groups of about 30 were transferred on egg laying plates for 24 hours, removed
 153 and replaced with a fresh egg plate. This second plate was removed after 24 hours, then 48
 154 hours later, first instar larvae were collected in groups of 100 and housed in standard
 155 molasses/agar food vials at 22°C. Males from these vials were collected on the day of eclosion
 156 and housed in vials of 10 individuals, until they were sexually mature (36), and then dissected
 157 when they were 5 or 6 days old.

158

159 *Accessory gland tissue preparation.* For dissections, males were anaesthetized with ether,
 160 placed in a drop of phosphate buffered saline and the reproductive tract removed. Intact
 161 accessory glands were clipped from the rest of the reproductive tract (Supplemental Figure
 162 1A, B) with fine dissection needles, moved to another drop of phosphate buffered saline for
 163 rinsing, and then transferred to a microcentrifuge tube. 30 accessory glands per replicate were
 164 used for subsequent LC-MS/MS as described below. Samples from each replicate were
 165 solubilized at 4°C by addition of 30 µl of RIPA buffer (Sigma), and a HALT protease inhibitor
 166 mixture containing phenylmethylsulfonyl fluoride (Thermo Fisher Scientific). Once all
 167 accessory glands had been dissected, samples were freeze/thawed three times on dry ice (~5
 168 mins) and then thawed at 37°C for 30 seconds. After the freeze/thaw cycles, samples were
 169 vortexed for 30 seconds, centrifuged at 20,000 rpm for 5 minutes at 4°C and then stored at -
 170 80°C until processing.

171

172 *Sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) and in-gel digestion*
 173 *of proteins.* Protein concentration was determined using a Bradford assay and samples were
 174 solubilized in SDS sample buffer containing 10 mM dithiothreitol and proteins separated on 4-
 175 12% SDS-PAGE gels per manufacturer instructions (Invitrogen). Protein bands were

visualized (Supplemental Figure 1C) using Brilliant Blue G Colloidal Concentrate (Sigma). Each gel lane was manually cut into approximately equivalent sized pieces and destained using 200 mM ammonium bicarbonate and 40% acetonitrile. Gel pieces were then reduced in 200 μ l of a 50 mM ammonium bicarbonate buffer containing 10 mM dithiothreitol, followed by alkylation in a similar volume of a 50 mM ammonium bicarbonate containing 55 mM iodoacetamide. Gel pieces were then centrifuged at 13 Kg for 10 seconds and dried using a vacuum concentrator until all samples were dry (~30 min). The dried pieces were then hydrated in a solution containing 20 μ l of trypsin (New England BioLabs) and 50 μ l of acetonitrile and incubated overnight at 37°C. Peptides were extracted the following day using a standard method with a solution of 100% acetonitrile and 5% formic acid and dried down overnight in a vacuum concentrator at 30°C. Resulting peptides were resuspended in 7.5 μ l of 0.1% (v/v) formic acid, 3% (v/v) acetonitrile, sonicated in a water bath for 5 minutes and centrifuged at 13 x g for 10 seconds, before being transferred to a sample vial and loaded into the autosampler tray of the Dionex Ultimate 3000 μ HPLC system. Samples were set to run using the Xcalibur sequence system.

Liquid chromatography-MS/MS (LC-MS/MS) data collection. All MS data were collected using an LTQ Orbitrap Elite hybrid mass spectrometer (Thermo Fisher Scientific) equipped with an Easy-Spray (Thermo Fisher Scientific) ion source. Peptides were separated using an Ultimate 3000 Nano LC System (Dionex). Peptides were desalted on-line using a capillary trap column (Acclaim Pepmap100, 100 μ m, 75 μ m x 2 cm, C18, 5 μ m; Thermo Fisher Scientific) and then separated using 60 min reverse phase gradient (3-40% acetonitrile/0.1% formic acid) on an Acclaim PepMap100 RSLC C18 analytical column (2 μ m, 75 μ m id x 10 cm; Thermo Fisher Scientific) with a flow rate of 0.25 μ l/min. The mass spectrometer was operated in standard data dependent acquisition mode controlled by Xcalibur 2.2. The instrument was operated with a cycle of one MS (in the Orbitrap) acquired at a resolution of 60,000 at m/z 400, with the top 20 most abundant multiply-charged (2+ and higher) ions in a given chromatographic window further subjected to CID fragmentation in the linear ion trap. An FTMS target value of

1e6 and an ion trap MSn target value of 10000 were used. Dynamic exclusion was enabled with a repeat duration of 30 s with an exclusion list of 500 and exclusion duration of 30 s.

Experimental Design and Statistical Rationale

Database construction of the Dpse accessory gland proteome (AcgP). The mass spectra data files were searched individually using Sequest HT within the Proteome Discover suite (Thermo Fisher Scientific, San Jose, CA, USA; version 1.4.1.14) using *Drosophila pseudoobscura pseudoobscura* fasta file, (December, 2015 release). Peptide matches were further analyzed and validated within Scaffold Q+ (Proteome Software; version 3.2.0) using X!Tandem. Sequest HT and X!Tandem searches were set with a fragment ion mass tolerance of 0.60 Da and a parent ion tolerance of 10.0 parts per million. The oxidation of methionine (15.99), carboxyamidomethyl of cysteine (57.02), and acetyl modification on peptide N-terminus (42.01) were set as variable modifications. Files from Sequest HT searches within the same gel lane were merged together as Mudpit using Scaffold which calculated False Discovery Rates (FDRs) using a reverse concatenated decoy database (FDR was set at 1.0%). Peptide identifications were accepted if they could be established at greater than 95.0% probability as specified by the PeptideProphet⁴⁸ and protein identifications were accepted if they could be established at greater than 99.0% probability and contained at least 2 identified peptides. Protein probabilities were assigned by the Protein Prophet Algorithm (37). Proteins that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony. The dataset was filtered so that every protein must be identified by at least two unique peptides in any one of the biological replicates. Although a conservative approach, this procedure ensured a robust dataset devoid of potential misidentifications often caused by use of a single peptide for protein identification. To establish a working list of the AcgP, protein IDs from Scaffold were converted to Dpse Fly Base gene numbers (FBgns) using the Uniprot website (Uniprot.org). The resulting Dpse FBgns were then used to query Flybase (Flybase.org) to retrieve orthologous Dmel genes from the OrthoDB orthology tables as implemented in Flybase (38). A complete listing can be found in Supplemental Table 1.

232

233 *Database construction of the Dpse accessory gland secretome (AcgS) and exoproteome*
 234 (*exoP*). As a secretory organ, the accessory gland is expected to contain the cellular
 235 machinery necessary for efficient and sustained secretory activity throughout the adult
 236 reproductive life cycle. To examine and focus on potential activities related to secretion, we
 237 assembled an *in silico* AcgS from the 3281 FBgns of the AcgP as input into Uniprot resulting
 238 in 5624 UniProtKB IDs (which includes all predicted protein isoforms). Fasta protein sequence
 239 files from each Uniprot entry were downloaded and input into SignalP
 240 (<http://www.cbs.dtu.dk/services/SignalP/>) (39) and Phobius (<http://phobius.sbc.su.se>) (40),
 241 using default settings. The protein IDs were combined, exported into Excel, yielding a final list
 242 of 771 UniProt identifiers. The Uniprot IDs were mapped back to 535 unique Dpse FBgns (via
 243 Uniprot) which, after submission to OrthoDB (via Flybase) resulted in a final list and 506 Dmel
 244 orthologs. Candidate SFPs were identified by first querying the list of 515 AcgS genes in
 245 Flybase for Gene Ontology (GO) terms containing “extracellular”. The resulting list of 151
 246 proteins therefore represents the accessory gland exoproteome (*exoP*) and is considered to
 247 contain a representative sampling of a major fraction of SFPs.

248

249 *Network and protein interaction network analyses.* The finalized datasets were used for
 250 downstream bioinformatic analyses and subsequent visualizations of GO enrichment. The
 251 protein coding sequences of the AcgP were downloaded from Uniprot and submitted to
 252 Blast2go for annotation and tabulation of the three major GO categories, biological process
 253 (BP), molecular function (MF) and cellular component (CC). GO enrichment and network
 254 visualization and analysis was performed with Cytoscape v3.4 (41, 42) and ClueGO plugin
 255 version 2.2.4 (43, 44). Network parameters used were specific for each dataset as detailed in
 256 figure legends and supplemental tables. Protein interaction network analysis was performed
 257 using Search Tool for the Retrieval of Interacting Genes/Proteins (STRING), a program that
 258 calculates the degree of protein-protein network interconnectivity (45). A combined set of

differentially abundant proteins from each M- and P-line (Qspec, see next section below) was uploaded to the STRING website and the analysis run using the “high confidence 0.9” setting and the “experimental” and “databases” selected for “evidence”. Proteins in each M- and P-up groups were distinguished by color-code: red (M>P) and green (P>M).

Label-free quantitation and statistical analysis. To test for differential abundance of male reproductive proteins in response to postcopulatory sexual selection statistical significance was calculated using a Bayesian approach as implemented by Qspec (46), part of the statistical package Qprot (47). Qspec provides a statistics framework that calculates Bayes factors- essentially likelihood measures of significance in the context of a generalized linear mixed effects model. Local implementation of Qprot (<http://sourceforge.net/p/qprot/>) provided command line access to QSpec that calculated Z-statistics, log-fold change estimates and a local FDR for each pair-wise comparison. This approach has been shown effective in capturing a broad range of differentially abundant proteins using LFQ methods (46, 48). Raw spectral counts of the M and P line datasets from Scaffold were input into Qspec and the output imported into Excel. Differential protein abundance differences were calculated using a 1% false-discovery rate (FDR) cut-off, and proteins assigned based on log-fold changes relative to the M-line, i.e., positive fold-change, $M < P$; negative fold-change, $M > P$.

Molecular evolution parameters. We obtained coding sequence information for the genomes of the two close relatives of Dpse, *D. lowei* and *D. affinis*, downloaded from http://popoolation.at/lowei_genome and http://popoolation.at/affinis_genome (49). These two species have the same karyotype as Dpse and show reasonable divergence (median pairwise dS=0.102 versus *D. lowei* and dS=0.26 versus *D. affinis*) thus avoiding substitution rate saturation. To identify orthologs of the identified AcgP Dpse proteins in the two other Drosophila genomes we combined two approaches. First we used gene annotation ignoring isoforms specification (as these are difficult to identify within a proteomic screen). We then

used best BLAST hits (50) of the *Dpse* gene against each of the two other genomes, but excluded gene sets for which annotation was contradictory to the *Dpse* annotation. Using a pipeline we developed earlier (51), sequences were aligned using MUSCLE (52), uncertain sequences filtered out using ZORRO (53) and input files converted with pal2nal (54). Sequences were then analyzed using PAML v4.9 (55) to obtain dN/dS values for each gene set (one-ratio estimates) as well as to conduct direct tests of positive selection as implemented in the site-specific models models M7a and M8a. For the one-ratio estimates, median differences in dN, dS and dN/dS among groups (AcgP, AcgS, exoP and differentially abundant M and P proteins) were tested using a non-parametric two-tailed test (57). For the site tests of positive selection, we obtained significance by conducting LRTs as described in the PAML manual and corrected P-values for multiple testing using an FDR=0.1 using the method of Benjamini and Hochberg (56). Median differences in dN, dS and dN/dS among groups (AcgP, AcgS, ExoP and differentially abundant M and P proteins) were tested using a non-parametric two-tailed test (57). Enrichment between counts per groups were tested using a 2x2 Fisher's exact test.

RESULTS

The Dpse AcgP. A *Dpse* AcgP was constructed from peptide-based ("bottom-up") shotgun MS/MS spectral data obtained from eight independent runs consisting of four replicate samples from each of two *Dpse* experimental sexual selection populations. For the purposes of assembling a proteome with the broadest coverage, data from all runs were pooled together resulting in a total of 3757 UniProt IDs that mapped to 3281 unique FlyBase gene names. The M- and P- datasets were highly correlated with >90% (3534/3757) overlap (Figure 1A). Likewise, proteins with values in all four replicates for each PCSS treatment represented the majority of identified proteins (M-line 2103/3649; 57.6% and P-line, 2235/3642; 61.4%) A complete listing and tabulation of these results can be found in Supplemental Table S1. The small number of proteins unique to each population (M-unique = 115; P-unique = 108; Figure

1A) most likely represent missed protein assignments due to low quantities (as measured by total spectral counts). Indeed, the average total spectral counts for the unique set of proteins (ave. 4.5; n = 223) was 16-fold lower than the average across the entire dataset (ave. 72.8, n = 3874).

GO and pathway analysis of the AcgP

Pathway and functional analyses began by identifying orthologous Dmel genes, (3159/3281, 96.2%; Supplemental Table 1), providing a useful annotated database to evaluate the overall patterns of the functional elements in the AcgP. As an overview of the major GO groupings, Blast2Go returned "signaling", "reproduction" and "localization" among BP- and numerous proteins annotated as extracellular involved in the CC-categories, respectively (Supplemental Figure 2). Statistical analysis for GO category enrichment (Figure 2) identified BP terms related to intracellular transport (n = 363, P = 2.80E-45), translation (n = 257, P = 2.47E-35), establishment of protein localization (n = 418, P = 1.22E-44), vesicle-mediated transport (n = 301, P = 8.47E-42), endocytosis (n = 106, P = 7E-10), and secretion (n = 140, P = 2.71E-16). Likewise, the AcgP contains a significant number of proteins in CC categories annotated as "extracellular region" (n = 266, P = 3.2E-05), "endomembrane system" (n = 583, P = 1.04E-57) and "vesicle" (n = 230, P = 8.72E-25). Finally, the overall known biochemical pathways of the AcgP, analyzed using the Kyoto Encyclopedia of Genes and Genomes (KEGG), revealed a similar enrichment of 20 overview terms curated by KEGG that included ribosome biogenesis, protein export, endocytosis and the Wnt signaling pathways (Table 1). We conclude the AcgP contains features expected of a metabolically active cell enriched for protein production and protein secretion. See Supplemental Table 2 for a complete listing of all enriched GO categories.

The *Dpse* accessory gland secretome (*AcgS*)

Signal peptides are a ubiquitous class of short (20-22 aa) N-terminal sequences, that target proteins for translocation across, and into, the endomembrane system of the cell (58, 59). Collectively proteins containing signal peptide

sequences are considered part of the secretory pathway, and a subclass - those secreted into the extracellular space - are termed the secretome (also referred to as the exoproteome, exoP). Therefore, some or all components of the exoP can be considered as candidate SFPs. Given the secretory nature of the *Drosophila* accessory gland, we therefore queried the AcgP for proteins containing canonical signal sequences using two predictive programs, SignalP and Phobius (see Methods). SignalP (39) is a neural network-based algorithm designed to detect canonical N-terminal signal sequences and discriminate against N-terminal transmembrane regions known to reduce predictive power, and Phobius utilizes a combined model of both transmembrane and signal peptide predictors (40, 60). The combined output of both resulted in 771 Uniprot IDs that mapped to 535 Dpse FBgns (Figure 1B; Supplemental Table 3). Dmel orthologs (OrthoDB via Flybase) subsequently returned 506 Dmel orthologs to the Dpse AcgS including a small percentage (8/511, 1.6%) of "1:many" matches included in the analysis to capture the greatest proteome coverage of the secretome. Thus, the AcgS represents approximately 15% (535/3281; 16.3%) of the entire AcgP consistent with similar calculations for the predicted human secretome (~15%, <http://www.proteinatlas.org/humanproteome/secretome>). Finally, the DRSC Integrative Ortholog Prediction Tools (DIOPT) website (<http://www.flyrnai.org/diopt>) identified 61% (327/535) of the AcgS that corresponded to orthologous human sequences (Supplemental Table 3).

Gene Ontology (GO) Functional Analysis of the AcgS

The high degree of homology between the Dpse AcgS genes and Dmel (506/528, Supplemental Table 3) provided a putative orthologous secretome useful for GO analysis and network visualization. A significant enrichment in BP terms was observed, many related to multicellular organism reproduction (n=113, p=7E-7), reproduction (n=116, p=7E-7), behavior (n=53, p=2.1E-5 and proteolysis (n=66, p=3.5E-6). The secretome is enriched in MF terms related to oxidoreductase activity (n=55, p=7.1E-7) and hydrolase activity (n=138, p=8.5e-14; see Supplemental Table 3 for a complete list of all AcgS enriched BP, CC and MF GO

categories). We also examined the subcellular localization of the AcpS using the Cerebral layout tool implemented in Cytoscape. As expected for functions related to secretion and proteins containing signal sequences targeted to the secretory pathway, the predicted subcellular localization of AcpS proteins were skewed toward extracellular and plasma membrane proteins (Figure 3).

The AcpS contains a robust repertoire of putative SFPs

We used the AcpS to identify a list of 151 putative SFPs representing >25% (151/506) of the AcpS (Figure 1B; Supplemental Table 4). This list is a conservative estimate as 99 genes had no CC functional annotation. Cytoscape and Cluego network analyses returned enriched BP terms of major functional categories including, insemination, sperm competition, and copulation reproduction, and negative regulation of endopeptidase activity (Table 2; Supplemental Table 4). Compared with a list of 212 Dmel SFPs assembled from the literature (9, 20, 21) an overlap of 32.1% (68/212) was observed that included 32 named SFP genes including Acp53Ea, Acp53C14d and Acp53C14c (Supplemental Table 4).

Divergent sexual selection results in differential AcpP protein abundance

To examine the proteome-wide response to sexual selection, differential protein abundance between the M- and P-lines was determined. A total of 250 differentially abundant Uniprot entries were identified using LFQ and Qspec statistical analysis across both treatments (FDR < 1%). The Uniprot IDs were subsequently mapped to 229 orthologous Dmel proteins (Supplemental Table 5). Taken together, network visualization of all 229 differentially abundant proteins revealed tight clustering of GO terms grouped into processes related to the cytoskeleton, protein translational machinery- particularly ribosomal proteins- and a suite of amino-acyl tRNA synthetases, along with a significant enrichment in categories related to endomembrane systems (e.g., Golgi, ER) and elements of the secretory pathway associated with vesicles and the Golgi apparatus (Figure 4A). Moreover, 14.4% (33/229) of the differentially abundant proteins were members of the exoP, including known Acps

(Supplemental Table 6).

M- and P-line differential protein abundance

To further probe the effect of PCSS on our divergent selection lines, we defined differential protein abundance between the sexual selection treatments based according to the sign of the log-fold change relative to the M-line (i.e., positive fold-change, $M < P$; negative fold-change, $M > P$; Supplemental Table 5). Of the 229 differentially abundant proteins, 133 and 96 were more abundant in the M-line in the P-line, respectively (Figure 1C). To examine differences in the function and cellular location of proteins differentially abundant between the M- and P-lines, the overlap of the two with respect to their GO groupings was generated using Cytoscape and Cluego. This comparison revealed a striking separation of functional grouping, with each selection line showing unique sub-groupings within the overall network of functional annotations (Figure 4B). We found differentially abundant proteins related to translational machinery (e.g., ribosomes) and cytoskeleton groupings predominant within the P-line (Figure 4B, green) with bias toward cytoplasmic cellular activities whereas Golgi- and vesicle- related groupings (including the exoP) biased in the M-line (Figure 4B, red) and functioning near or at the cell periphery.

Thirty-three of the differentially abundant proteins (24 up in M, 9 up in P) were annotated as "extracellular" (i.e., putative SFPs; Supplemental Table 6). While a larger number of candidate SFPs were more abundant in M-line compared to P-line males, over 50% of P abundant SFPs overlapped with known Dmel SFPs (5/9) compared to 29% overlap between M abundant and Dmel SFPs (7/24). Differentially abundant P male proteins did not show significant GO BP enrichment whereas significant enrichment was found in M-line males related to "response to fungus" ($P = 2.1E-02$) and "wound healing" ($P = 2.0E-02$) (9/24, 34.5%; Supplemental Table 6; Supplemental Figure 3).

Sexual selection targets protein interaction networks

While the differentially abundant proteins of each selection line was biased towards distinct sets of GO functional groupings (Figure 4B), we also tested whether these relationships extended to specific protein interaction networks of known network topologies. We used the most stringent filters for network interactions on STRING (45), and found a statistically significant ($P = 0.001$) PPI network of 91 proteins containing nine defined subnetworks and including elements of 34 KEGG pathways (Supplemental Table 7). A striking aspect of the STRING-generated PPI networks was the almost complete segregation of M and P differentially abundant proteins into line-specific protein interaction networks (Figure 5). For example, all elements of the “vesicle transport” and “purine biosynthesis” PPI networks were more abundant exclusively in the M-line. Likewise, all proteins under “microtubule organization” and almost all (15/18) in the “ribosome” category are more abundant in the P-line. These results show that the divergent sexual selection treatments used to generate the M- and P-lines resulted in highly focused and interrelated changes in protein abundance that were specific to each treatment.

Molecular evolutionary rates of accessory gland protein genes

We tested for rates of molecular evolution in male reproductive proteins by estimating omega (dN/dS substitution rates) for each set of proteins: candidate SFPs, secretome proteins (minus SFPs), and the remaining accessory gland proteome proteins using orthologs from two closely related species in the obscura group, *D. lowei* and *D. affinis*. We found that putative Dpse SFPs (Exoproteome) are evolving faster than both accessory gland proteome proteins (AcgP; median omega = 0.088 versus 0.052, $P = 1.1 \times 10^{-9}$) and accessory gland secretome minus the Exoproteome (AcgS; median omega = 0.088 versus 0.078, $P = 2.7 \times 10^{-2}$; Figure 6A). AcgS proteins also evolve faster than the AcgP proteins (median omega = 0.052 versus 0.078, $P = 1 \times 10^{-9}$; Figure 6A).

Given that sexual selection is hypothesized to result in rapid protein sequence evolution, we then compared the evolutionary rates of the selection-specific differentially abundant proteins

and accessory gland proteins (excluding selection-specific proteins). We found that the P treatment showed pronounced reduction in dN compared to the M treatment (Figure 6B, dN=0.009 (P) versus 0.018 (M), $P=0.011$). This is perhaps unsurprising considering that cytosolic ribosomal proteins are highly enriched in P, a protein group that has been shown to be slowly evolving (61). However, slowly evolving genes may be indicative of intensified selective pressures which obscure signals of positive selection, so to account for this we employed a direct site-specific test of positive selection across the accessory gland proteome. We identified 19 proteins that showed positive selection, 14 from the AcgP and 5 showing differential abundance in the selection lines, 3 in P and 2 in M (Table 3). An enrichment analysis shows that genes encoding proteins that were differentially abundant following experimental microevolution were more likely to be under positive selection than those in the AcgP set (Figure 6C, Supplemental Table 8) after correction for multiple testing and this is significant for the P-only set as well.

DISCUSSION

Proteomics of the Dpse male accessory gland

The main aim of this work was to directly test the role of postcopulatory sexual selection in the evolution of reproductive proteins and the male accessory gland reproductive protein network. To accomplish this aim, we employed a “bottom up” shotgun proteomics approach, generating a robust accessory gland proteome containing 3281 proteins, representing the first accessory gland proteome to be described in *Drosophila*. The AcgP proteins in Dpse were enriched for cellular components expected from a tissue whose primary function is secretory, including several cellular component GO terms related to membranes, extracellular regions, and peptidase complex. The top biological process GO terms clearly indicated a large investment in processes directly and indirectly related to protein synthesis, protein assembly, transport and secretion. We then *in silico* concatenated this list to include proteins with secretory signal sequences to identify 545 accessory gland secretome proteins, which were enriched for GO

terms related to the biological processes of reproduction, behavior and proteolysis, with these proteins heavily biased towards subcellular localizations in the plasma membrane or extracellular components, as predicted from proteins with secretory signals.

Candidate Dpse SFPs contains many known Dmel SFPs

SFPs are secreted from the accessory gland tissue and transferred to the females upon mating. To generate a list of putative SFPs, we restricted the AcgS to only those proteins that were associated with GO cellular component annotations containing the keyword “extracellular”. The list of these proteins is most probably conservative as not all proteins were annotated. Nonetheless, the approach identified 151 proteins enriched for biological processes associated with reproduction, mating, insemination, and sperm competition, with approximately 30% that shared homology with previously identified Dmel SFPs. We also compared our list with the 29 putative SFPs previously identified via genome comparison between Dmel and Dpse (9). Of the nearly 50% overlap between these two lists (14/29), protein members of the canonical Sex Peptide (SP) network are particularly well-represented. In Dmel, SP binds to sperm in the female seminal receptacle, a sperm storage organ, and is required for both long-term female resistance to remating and for sperm release from storage (62). Our list of SFPs contains several known proteins involved in the Dmel SP network, including the gene duplicate pair lectins CG1652 and CG1656, aquarius (CG14061), intrepid (CG12558), antares (CG30488), seminase (CG10586), CG17575 (a cysteine-rich secretory protein), and CG9997 (a serine protease homolog) (16). CG9997 is processed in the female and males that do not produce this protein are unable to transfer the lectins, which are required to slow the rate at which CG9997 is processed in the female. All SP pathway proteins, except SP, were detected in our putative list of SFPs. Absence of detectable Dpse SP protein is consistent with the lack of a recognized SP ortholog in this species, and raises the interesting possibility that either the Dpse SP ortholog has significantly diverged, or has been replaced by another gene. If indeed a bona fide Dpse SP gene exists, further MS searches using

algorithms to detect amino acid replacements (63) may be useful in the search for this elusive SFP.

Functional compartmentalization results from divergent sexual selection

While our analysis found many known SFPs, 68% had no known orthology to Dmel. This supports the idea that SFPs and related proteins involved in reproduction show rapid evolution as suggested by genome comparisons (9, 16, 17, 21, 64–66) and experimental evolution research on changes in sex-specific gene expression (32, 33, 67). Here we directly measured, using LC-MS/MS and LFQ, the proteome-wide effect of postcopulatory sexual selection on protein abundance in two populations that had undergone over 150 generations of divergent PCSS. This approach identified significant protein abundance differences in over 200 proteins between the selection lines. Knowledge of the changes in protein abundance under these experimental conditions provided a useful database to compare and contrast the effects of selection from a proteome-wide perspective and to assess the impact of selection on the protein-protein interaction networks revealed by our analysis.

Previous comparative work has suggested protein evolution may be adaptively compartmentalized for core processes and undergoing purifying selection and for interactive proteins that undergo positive selection (24, 26). We directly tested this assertion by comparing protein interaction network architecture in the divergent PCSS treatments and testing for molecular signatures of selection. We found remarkably line-specific compartmentalized changes to proteins after more than 150 generations of experimental sexual selection. P males invest in protein production machinery (i.e., protein quantity) and M males invest in protein surveillance (i.e., protein quality). The number of matings a *Drosophila* male can have is limited by SFP supply and how rapidly males can replenish this supply (68). Moreover, males can respond plastically to increased risk of sperm competition by transferring more of some SFPs to females (14, 69). We have previously shown that P males have larger

accessory glands and can mate with more females sequentially than M males (30, 31). The increased investment in protein production machinery in P males may underlie this phenotypic response, and we suggest this may explain how males in polyandrous mating systems can rapidly adjust and replenish SFP quantity. In contrast, M males overall have increased production of proteins that function in downstream processing of secreted proteins, suggestive of increased investment in protein quality. The transfer of high quality proteins to females may be a response to reduce sexual conflict and improve population fitness, as predicted under enforced monogamy in which the reproductive interests of the sexes become aligned.

The SFPs produced in larger quantities in M compared to P males were enriched for chitin catabolic process (including wound healing) and response to fungus. Mating in *Dmel* initiates an immune response in females that has negative fitness consequences (70). The immune system may be activated to combat pathogens in the ejaculate and/or immunogenic sperm and SFPs (response to fungus), and/or wounds inflicted by males (chitin catabolic processes and wound healing). As enforced monogamy relaxes sexual conflict between males and females, M males may produce more of these proteins to limit damage to their mates. Future work could test this by manipulating these protein levels in the male ejaculate and determining the consequences on female fitness following mating. M males also produced more ejaculatory bulb protein (Ebp; CG2668), a known constituent of the mating plug. The mating plug remains in the female uterus until females eject it, along with unstored sperm. Ebp is necessary for ejaculate retention; knocking down Ebp in *Dmel* results in ejaculate loss, and reduced female sperm storage (Avila et al. 2015b). M males may produce more Ebp for better ejaculate retention, reducing the likelihood of their mates requiring additional matings to maintain sperm stores and subsequent fertility. Given the costs of mating for females, such a response is predicted when sexual conflict is relaxed, as in the M treatment.

Differentially abundant P male proteins did not show significant GO BP enrichment but over 50% of these (5/9) are known *Dmel* SFPs. This includes another mating plug protein, Acp53C14b, known to be involved in egg laying and reproductive fitness (71). Our previous

work on the Dpse selection lines described sexual conflict over female reproductive phenology; M females mated to P males oviposit more eggs earlier in their lifespan than when mated to their own males (31). Given that the M treatment removes sexual conflict, alterations to the M female oviposition schedule when mated to P males is likely to be costly. This manipulation of female oviposition may be mediated by Acp53C14b given that this protein – which P males produce more of – influences oviposition.

Rates of molecular evolution on male reproductive proteins

Rapid evolution at the molecular level is common for reproductive proteins including SFPs (e.g., (10, 18, 22, 26)) and this rapid evolution at the macroevolutionary scale is attributed to PCSS. Here we go beyond these traditional tests and ask whether there are differences in the rate of evolution between different types of reproductive proteins arising from the same tissue at the microevolutionary scale following over 150 generations of experimental manipulation of PCSS. We find that genes encoding SFPs showed significantly faster rates of molecular evolution compared to both other secreted protein encoding genes that were not candidate SFPs and overall accessory gland proteome genes. However, we also found that secretome genes showed higher rates of molecular evolution than non-secreted proteins (i.e., the remainder of the AcpP). These results support the interpretation that genes which interact extracellularly evolve more rapidly than those that remain within the cytoplasm (72, 73).

Evolutionary rates of genes encoding differentially abundant proteins between the sexual selection treatments found pronounced reduction in dN of P males compared to either differentially abundant proteins in M or the AcpP. This was likely due to the compartmentalization we identified, in which P males invested more in slowly evolving cytosolic ribosomal proteins (Barreto and Burton 2013). However, after controlling for this, we found that genes encoding for M- and P- differentially abundant proteins were more likely to have been subjected to positive selection compared to the AcpP genes as a whole. This result reinforces the hypothesis that the proteins we identified as responding to divergent sexual

selection at the microevolutionary level have undergone adaptive processes on a macroevolutionary scale. We argue this directly illustrates a major role of PCSS in the evolution of male reproductive proteins.

Conclusions

By employing divergent postcopulatory sexual selection through the use of experimental evolution of mating systems, we were able to directly test the hypothesis that postcopulatory sexual selection results in the rapid evolution of male-specific reproductive proteins. Within 150 generations, substantial microevolutionary changes occurred such as differential abundance of secreted proteins, including previously identified *Dmel* SFPs that have known effects on male and female reproductive fitness. A small number of these proteins showed signatures of positive molecular evolution. We found PCSS mediates remarkable compartmentalization of subcellular function of the secretory tissue, suggesting sexual selection alters fundamental reproductive protein networks, not just individual SFPs. These changes affected protein production, but in different ways in the divergent treatments, suggesting that PCSS selects for increased protein production whereas relaxation of sexual conflict selects for protein surveillance. Some specific changes may be related to previously described phenotypic responses to divergent sexual selection that facilitate greater reproductive capacity in males subject to polyandry and ameliorates sexual conflict in populations subject to monandry. Our novel results suggest wide-spread, previously unappreciated, consequences of postcopulatory sexual selection on reproductive protein evolution. Combining the increasing use of high throughput proteomics and experimental manipulation of mating systems in different taxa will allow broader tests of this pattern in different taxa and better understanding of how sexual selection and sexual conflict impacts male reproductive protein evolution.

Acknowledgements – We thank the many people who have contributed to the maintenance of the Snook experimental evolution lines. Funding of the experimental evolution lines came from

NSF (DEB-0093149), NERC (NE/B504065/1; NE/I014632/1), and EU (ITN-2008-213780 SPECIATION) to RRS. Funding and technical support for the proteomic work came from the University of Sheffield Biological Mass Spectrophotometry Facility (funded by Yorkshire Cancer Research (Shend01) and the Wellcome Trust). TIG was supported by a Leverhulme Early Career Fellowship (Grant ECF-2015-453) and a NERC grant (NE/N013832/1). Funding from the Kyoto Institute of Technology to TLK also supported this research.

DATA AVAILABILITY

Raw files for the entire datasets used in this study have been deposited via ProteomeXchange on the PRIDE database repository.

REFERENCES

1. Snook, R. R. (2014) The evolution of polyandry. *The Evolution of Insect Mating Systems* 159-180
2. Arnqvist, G. and Rowe, L. (2005) *Sexual conflict*, Princeton University Press, Princeton, N.J.
3. Birkhead, T. R., Hosken, D. J. and Pitnick, S. S. (2008) *Sperm Biology*.
4. Avila, F. W., Sirot, L. K., LaFlamme, B. A., Rubinstein, C. D. and Wolfner, M. F. (2011) Insect seminal fluid proteins: identification and function. *Annual review of entomology* **56**, 21-40
5. Ramm, S. A., McDonald, L., Hurst, J. L., Beynon, R. J. and Stockley, P. (2009) Comparative proteomics reveals evidence for evolutionary diversification of rodent seminal fluid and its functional significance in sperm competition. *Mol Biol Evol* **26**, 189-198
6. Dorus, S., Evans, P. D., Wyckoff, G. J., Choi, S. S. and Lahn, B. T. (2004) Rate of molecular evolution of the seminal protein gene SEMG2 correlates with levels of female promiscuity. *Nat Genet* **36**, 1326-1329
7. Harrison, P. W., Wright, A. E., Zimmer, F., Dean, R., Montgomery, S. H., Pointer, M.

- 643 A. and Mank, J. E. (2015) Sexual selection drives evolution and rapid turnover of male gene
644 expression. *Proc Natl Acad Sci U S A* **112**, 4393-4398
- 645 8. Panhuis, T. M., Clark, N. L. and Swanson, W. J. (2006) Rapid evolution of reproductive
646 proteins in abalone and *Drosophila*. *Philosophical transactions of the Royal Society of London*
647 *B: Biological sciences* **361**, 261-268
- 648 9. Mueller, J. L., Ravi Ram, K., McGraw, L. A., Bloch Qazi, M. C., Siggia, E. D., Clark, A.
649 G., Aquadro, C. F. and Wolfner, M. F. (2005) Cross-species comparison of *Drosophila* male
650 accessory gland protein genes. *Genetics* **171**, 131-143
- 651 10. Clark, N. L., Aagaard, J. E. and Swanson, W. J. (2006) Evolution of reproductive
652 proteins from animals and plants. *Reproduction* **131**, 11-22
- 653 11. Mattei, A. L., Riccio, M. L., Avila, F. W. and Wolfner, M. F. (2015) Integrated 3D view
654 of postmating responses by the *Drosophila melanogaster* female reproductive tract, obtained
655 by micro-computed tomography scanning. *Proceedings of the National Academy of Sciences*
656 *of the United States of America* **112**, 8475-8480
- 657 12. Avila, F. W. and Wolfner, M. F. (2009) Acp36DE is required for uterine conformational
658 changes in mated *Drosophila* females. *Proceedings of the National Academy of Sciences of*
659 *the United States of America* **106**, 15796-15800
- 660 13. Perry, J. C., Sirot, L. and Wigby, S. (2013) The seminal symphony: How to compose
661 an ejaculate. *Trends in Ecology and Evolution* **28**, 414-422
- 662 14. Wigby, S., Sirot, L. K., Linklater, J. R., Buehner, N., Calboli, F. C. F., Bretman, A.,
663 Wolfner, M. F. and Chapman, T. (2009) Seminal fluid protein allocation and male reproductive
664 success. *Current biology : CB* **19**, 751-757
- 665 15. Avila, F. W., Mattei, A. L. and Wolfner, M. F. (2015) Sex peptide receptor is required
666 for the release of stored sperm by mated *Drosophila melanogaster* females. *Journal of insect*
667 *physiology* **76**, 1-6
- 668 16. Findlay, G. D., Sitnik, J. L., Wang, W., Aquadro, C. F., Clark, N. L. and Wolfner, M. F.
669 (2014) Evolutionary rate covariation identifies new members of a protein network required for
670 *Drosophila melanogaster* female post-mating responses. *PLoS Genetics* **10**, e1004108

- 671 17. Swanson, W. J., Clark, A. G., Waldrip-Dail, H. M., Wolfner, M. F. and Aquadro, C. F.
672 (2001) Evolutionary EST analysis identifies rapidly evolving male reproductive proteins in
673 *Drosophila*. *Proceedings of the National Academy of Sciences of the United States of America*
674 **98**, 7375-7379
- 675 18. Mueller, J. L., Ripoll, D. R., Aquadro, C. F. and Wolfner, M. F. (2004) Comparative
676 structural modeling and inference of conserved protein classes in *Drosophila* seminal fluid.
677 *Proceedings of the National Academy of Sciences of the United States of America* **101**, 13542-
678 13547
- 679 19. Wagstaff, B. J. and Begun, D. J. (2005) Comparative genomics of accessory gland
680 protein genes in *Drosophila melanogaster* and *D. pseudoobscura*. *Mol Biol Evol* **22**, 818-832
- 681 20. Findlay, G. D., MacCoss, M. J. and Swanson, W. J. (2009) Proteomic discovery of
682 previously unannotated, rapidly evolving seminal fluid genes in *Drosophila*. *Genome research*
683 **19**, 886-896
- 684 21. Findlay, G. D., Yi, X., Maccoss, M. J. and Swanson, W. J. (2008) Proteomics reveals
685 novel *Drosophila* seminal fluid proteins transferred at mating. *PLoS Biol* **6**, e178
- 686 22. Wilburn, D. B. and Swanson, W. J. (2016) From molecules to mating: Rapid evolution
687 and biochemical studies of reproductive proteins. *J Proteomics* **135**, 12-25
- 688 23. Schumacher, J., Rosenkranz, D. and Herlyn, H. (2014) Mating systems and protein-
689 protein interactions determine evolutionary rates of primate sperm proteins. *Proc Biol Sci* **281**,
690 20132607
- 691 24. Dorus, S., Wasbrough, E. R., Busby, J., Wilkin, E. C. and Karr, T. L. (2010) Sperm
692 proteomics reveals intensified selection on mouse sperm membrane and acrosome genes.
693 *Molecular biology and evolution* **27**, 1235-1246
- 694 25. Dorus, S., Busby, S. A., Gerike, U., Shabanowitz, J., Hunt, D. F. and Karr, T. L. (2006)
695 Genomic and functional evolution of the *Drosophila melanogaster* sperm proteome. *Nature*
696 *genetics* **38**, 1440-1445
- 697 26. Vicens, A., Borziak, K., Karr, T. L., Roldan, E. R. S. and Dorus, S. (2017) Comparative
698 Sperm Proteomics in Mouse Species with Divergent Mating Systems. *Mol Biol Evol* **34**, 1403-

699 1416

- 700 27. Athanasios, A., Charalampos, V., Vasileios, T. and Ashraf, G. M. (2017) Protein-
701 Protein Interaction (PPI) Network: Recent Advances in Drug Discovery. *Curr Drug Metab* **18**,
702 5-10
- 703 28. Levy, E. D. and Pereira-Leal, J. B. (2008) Evolution and dynamics of protein
704 interactions and networks. *Curr Opin Struct Biol* **18**, 349-357
- 705 29. Crudgington, H. S., Beckerman, A. P., Brustle, L., Green, K. and Snook, R. R. (2005)
706 Experimental removal and elevation of sexual selection: Does sexual selection generate
707 manipulative males and resistant females? *American Naturalist* **165**, S72-S87
- 708 30. Crudgington, H. S., Fellows, S. and Snook, R. R. (2010) Increased opportunity for
709 sexual conflict promotes harmful males with elevated courtship frequencies. *Journal of*
710 *evolutionary biology* **23**, 440-446
- 711 31. Crudgington, H. S., Fellows, S., Badcock, N. S. and Snook, R. R. (2009) Experimental
712 manipulation of sexual selection promotes greater male mating capacity but does not alter
713 sperm investment. *Evolution; international journal of organic evolution* **63**, 926-938
- 714 32. Immonen, E., Snook, R. R. and Ritchie, M. G. (2014) Mating system variation drives
715 rapid evolution of the female transcriptome in *Drosophila pseudoobscura*. *Ecology and*
716 *evolution* **4**, 2186-2201
- 717 33. Veltsos, P., Fang, Y., Cossins, A. R., Snook, R. R. and Ritchie, M. G. (2017) Mating
718 system manipulation and the evolution of sex-biased gene expression in *Drosophila*. *Nat*
719 *Commun* **8**, 2072
- 720 34. Anderson, W. W. (1974) Frequent multiple insemination in a natural population of
721 *Drosophila pseudoobscura*. *The American naturalist* **108**, 709-711
- 722 35. Snook, R. R., Brüstle, L. and Slate, J. (2009) A test and review of the role of effective
723 population size on experimental sexual selection patterns. *Evolution; international journal of*
724 *organic evolution* **63**, 1923-1933
- 725 36. Snook, R. R. and Markow, T. A. (2002) Efficiency of gamete usage in nature: sperm
726 storage, fertilization and polyspermy. *Proceedings. Biological sciences / The Royal Society*

727 **269**, 467-473

728 37. Nesvizhskii, A. I., Keller, A., Kolker, E. and Aebersold, R. (2003) A Statistical Model for
729 Identifying Proteins by Tandem Mass Spectrometry abilities that proteins are present in a
730 sample on the basis. *Analytical chemistry* **75**, 4646-4658

731 38. Kriventseva, E. V., Tegenfeldt, F., Petty, T. J., Waterhouse, R. M., Simão, F. A.,
732 Pozdnyakov, I. A., Ioannidis, P. and Zdobnov, E. M. (2015) OrthoDB v8: update of the
733 hierarchical catalog of orthologs and the underlying free software. *Nucleic Acids Res* **43**,
734 D250-6

735 39. Petersen, T. N., Brunak, S., von Heijne, G. and Nielsen, H. (2011) SignalP 4.0:
736 discriminating signal peptides from transmembrane regions. *Nature methods* **8**, 785-786

737 40. Käll, L., Krogh, A. and Sonnhammer, E. L. (2004) A combined transmembrane
738 topology and signal peptide prediction method. *J Mol Biol* **338**, 1027-1036

739 41. Cline, M. S., Smoot, M., Cerami, E., Kuchinsky, A., Landys, N., Workman, C.,
740 Christmas, R., Avila-Campilo, I., Creech, M., Gross, B., Hanspers, K., Isserlin, R., Kelley, R.,
741 Killcoyne, S., Lotia, S., Maere, S., Morris, J., Ono, K., Pavlovic, V., Pico, A. R., Vailaya, A.,
742 Wang, P.-L., Adler, A., Conklin, B. R., Hood, L., Kuiper, M., Sander, C., Schmulevich, I.,
743 Schwikowski, B., Warner, G. J., Ideker, T. and Bader, G. D. (2007) Integration of biological
744 networks and gene expression data using Cytoscape. *Nature protocols* **2**, 2366-2382

745 42. Demchak, B., Hull, T., Reich, M., Liefeld, T., Smoot, M., Ideker, T. and Mesirov, J. P.
746 (2014) Cytoscape: the network visualization tool for GenomeSpace workflows.
747 *F1000Research* **3**, 151

748 43. Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A.,
749 Fridman, W.-H., Pagès, F., Trajanoski, Z. and Galon, J. (2009) ClueGO: A Cytoscape plug-in
750 to decipher functionally grouped gene ontology and pathway annotation networks.
751 *Bioinformatics (Oxford, England)* **25**, 1091-1093

752 44. Bindea, G., Galon, J. and Mlecnik, B. (2013) CluePedia Cytoscape plugin: pathway
753 insights using integrated experimental and in silico data. *Bioinformatics* **29**, 661-663

754 45. Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J.,

- Simonovic, M., Roth, A., Santos, A., Tsafou, K. P., Kuhn, M., Bork, P., Jensen, L. J. and von Mering, C. (2015) STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* **43**, D447-52
46. Choi, H., Fermin, D. and Nesvizhskii, A. I. (2008) Significance Analysis of Spectral Count Data in Label-free Shotgun Proteomics. *Molecular & Cellular Proteomics* **7**, 2373-2385
47. Choi, H., Kim, S., Fermin, D., Tsou, C. C. and Nesvizhskii, A. I. (2015) QPROT: Statistical method for testing differential expression using protein-level intensity data in label-free quantitative proteomics. *J Proteomics* **129**, 121-126
48. Blein-Nicolas, M. and Zivy, M. (2016) Thousand and one ways to quantify and compare protein abundances in label-free bottom-up proteomics. *Biochimica et biophysica acta* **1864**, 883-895
49. Palmieri, N., Kosiol, C. and Schlötterer, C. (2014) The life cycle of Drosophila orphan genes. *Elife* **3**, e01311
50. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990) Basic local alignment search tool. *J Mol Biol* **215**, 403-410
51. Gossmann, T. I., Santure, A. W., Sheldon, B. C., Slate, J. and Zeng, K. (2014) Highly variable recombinational landscape modulates efficacy of natural selection in birds. *Genome Biol Evol* **6**, 2061-2075
52. Edgar, R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792-1797
53. Wu, M., Chatterji, S. and Eisen, J. A. (2012) Accounting for alignment uncertainty in phylogenomics. *PLoS One* **7**, e30288
54. Suyama, M., Torrents, D. and Bork, P. (2006) PAL2NAL: robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Res* **34**, W609-12
55. Yang, Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-1591
56. Hochberg, Y. and Benjamini, Y. (1990) More powerful procedures for multiple

783 significance testing. *Stat Med* **9**, 811-818

784 57. Mann, H. B. and Whitney, D. R. (1947) On a Test of Whether One of Two Random
785 Variables Is Stochastically Larger Than the Other. *Ann. Math. Statistics* **18**, 50-60

786 58. Armengaud, J., Christie-Oleza, J. A., Clair, G., Malard, V. and Duport, C. (2012)
787 Exoproteomics: exploring the world around biological systems. *Expert Rev Proteomics* **9**, 561-
788 575

789 59. Tjalsma, H., Bolhuis, A., Jongbloed, J. D., Bron, S. and van Dijk, J. M. (2000) Signal
790 peptide-dependent protein transport in *Bacillus subtilis*: a genome-based survey of the
791 secretome. *Microbiology and molecular biology reviews : MMBR* **64**, 515-547

792 60. Käll, L., Krogh, A. and Sonnhammer, E. L. (2007) Advantages of combined
793 transmembrane topology and signal peptide prediction--the Phobius web server. *Nucleic
794 Acids Res* **35**, W429-32

795 61. Barreto, F. S. and Burton, R. S. (2013) Evidence for compensatory evolution of
796 ribosomal proteins in response to rapid divergence of mitochondrial rRNA. *Mol Biol Evol* **30**,
797 310-314

798 62. Avila, F. W., Ravi Ram, K., Bloch Qazi, M. C. and Wolfner, M. F. (2010) Sex peptide
799 is required for the efficient release of stored sperm in mated *Drosophila* females. *Genetics*
800 **186**, 595-600

801 63. Starkweather, R., Barnes, C. S., Wyckoff, G. J. and Keightley, J. A. (2007) Virtual
802 polymorphism: finding divergent peptide matches in mass spectrometry data. *Anal Chem* **79**,
803 5030-5039

804 64. Clark, N. L. and Swanson, W. J. (2005) Pervasive adaptive evolution in primate
805 seminal proteins. *PLoS Genet* **1**, e35

806 65. Dean, M. D., Clark, N. L., Findlay, G. D., Karn, R. C., Yi, X., Swanson, W. J., MacCoss,
807 M. J. and Nachman, M. W. (2009) Proteomics and comparative genomic investigations reveal
808 heterogeneity in evolutionary rate of male reproductive proteins in mice (*Mus domesticus*).
809 *Mol Biol Evol* **26**, 1733-1743

810 66. Sirot, L. K., Findlay, G. D., Sitnik, J. L., Frasheri, D., Avila, F. W. and Wolfner, M. F.

- (2014) Molecular characterization and evolution of a gene family encoding both female- and male-specific reproductive proteins in *Drosophila*. *Molecular biology and evolution* **31**, 1554-1567
67. Hollis, B., Houle, D., Yan, Z., Kawecki, T. J. and Keller, L. (2014) Evolution under monogamy feminizes gene expression in *Drosophila melanogaster*. *Nat Commun* **5**, 3482
68. Linklater, J. R., Wertheim, B., Wigby, S. and Chapman, T. (2007) Ejaculate depletion patterns evolve in response to experimental manipulation of sex ratio in *Drosophila melanogaster*. *Evolution* **61**, 2027-2034
69. Sirot, L. K., Wolfner, M. F. and Wigby, S. (2011) Protein-specific manipulation of ejaculate composition in response to female mating status in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 9922-9926
70. Mack, P. D., Kapelnikov, A., Heifetz, Y. and Bender, M. (2006) Mating-responsive genes in reproductive tissues of female *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* **103**, 10358-10363
71. Avila, F. W., Cohen, A. B., Ameerudeen, F. S., Duneau, D., Suresh, S., Mattei, A. L. and Wolfner, M. F. (2015) Retention of Ejaculate by *Drosophila melanogaster* Females Requires the Male-Derived Mating Plug Protein PEBme. *Genetics* **200**, 1171-1179
72. Feyertag, F., Berninsone, P. M. and Alvarez-Ponce, D. (2017) Secreted Proteins Defy the Expression Level-Evolutionary Rate Anticorrelation. *Mol Biol Evol* **34**, 692-706
73. Liao, B. Y., Weng, M. P. and Zhang, J. (2010) Impact of extracellularly on the evolutionary rate of mammalian proteins. *Genome Biol Evol* **2**, 39-43

Figure Legends

Figure 1. Dpse accessory gland summary statistics. (A) Venn diagram of combined and total Uniprot IDs from the M- and P-experimental lines used to assemble the AcgP. (B) Secretome and exoproteome from the AcgP derived *in silico* (see Methods). (C) Differential protein abundance between the M- and P- experimental lines. Numbers in parentheses indicate Dmel orthologs from OrthoDB database.

Figure 2. AcgP GO category enrichment analysis. Distribution of major functional category groupings for Biological Process (A) and Cellular Component (B). Categories related to secretory processes and discussed in main text highlighted in bold.

Figure 3. Functional grouping and subcellular localization of the AcgS mapped to subcellular regions (labeled on the right) using the Cerebral layout in Cytoscape. Shaded regions show all annotations that include the keyword “Extracellular”.

Figure 4. Network and predicted cellular distribution of DE proteins. (A) Overview of the DE proteins (M- and P-lines combined) showing major enriched functional groupings for GO Cellular Components (CC) using Cytoscape and Cluego. The four major groups, cytoskeletal part, golgi part, translation machinery and endomembrane system are shaded with different background colors. Dashed line encloses two functionally distinct categories, the ribosome and tRNA synthetase complex of the translation machinery (see also Table 5). (B) Network comparison of functional annotation groupings between DE proteins of the M- and P-lines. DE proteins were segregated into their respective M- or P- lines based on relative protein abundance and both groups compared using Cluego. A heat map (red-to-green) shows the distribution of DE protein between the M- and P-lines. These distributions were then mapped onto the predicted cellular locations of the groupings using the Cluepedia and the Cerebral layout as implemented in Cytoscape.

863

864 Figure 5. PPI network representation of the M (red, M>P) and P (green, P>M) DE proteins
 865 (inset). The PPI network was generated using STRING, a program that determines protein-
 866 protein interactions based on experimental and database criteria and calculated confidence
 867 levels to assign interactions between protein pairs. Stippled lines indicate major functional
 868 categories describing the networks.

869

870 Figure 6. (A) Evolutionary rate calculations of the ratio of the non-synonymous to synonymous
 871 (dN/dS) substitution rates for (left to right) the Acg proteome (AcpP), the secretome (AcpS,
 872 excluding exoproteome and the exoproteome (AcpS-Extracellular). (B) The rate of
 873 nonsynonymous changes (dn) in genes coding for (left to right); the AcgP (minus DE M and P
 874 treatment genes), differentially abundant M-line genes (DE_M_UP) and differentially abundant
 875 P-line genes (DE_P_UP). Number of genes analyzed in each category listed below category.
 876 (C) Summary table of evolutionary rate analysis indicating significant enrichment of differential
 877 abundance proteins under positive selection.

Table 1. Enriched KEGG pathways of the AcgP.

<i>GOID</i>	<i>GO Term</i>	<i>Nr. Genes</i>	<i>Pvalue*</i>
KEGG:03008	Ribosome biogenesis	33	3.5E-12
KEGG:03040	Spliceosome	82	8.0E-06
KEGG:04141	Processing in endoplasmic reticulum	82	5.9E-05
KEGG:03018	RNA degradation	40	4.6E-04
KEGG:03050	Proteasome	35	5.5E-04
KEGG:04144	Endocytosis	74	5.6E-04
KEGG:04310	Wnt signaling pathway	24	1.8E-03
KEGG:00071	Fatty acid degradation	24	2.9E-03
KEGG:04130	SNARE vesicular transport	17	2.9E-03
KEGG:00330	Arginine and proline metabolism	34	4.6E-03
KEGG:00230	Purine metabolism	40	1.0E-02
KEGG:03060	Protein export	17	1.1E-02
KEGG:00280	Valine, leucine and isoleucine degradation	22	1.3E-02
KEGG:00250	Ala, asp and glu metabolism	20	1.4E-02
KEGG:00020	Citrate cycle (TCA cycle)	27	2.5E-02
KEGG:00030	Pentose phosphate pathway	16	2.8E-02
KEGG:04068	FoxO signaling pathway	16	2.9E-02
KEGG:00500	Starch and sucrose metabolism	20	4.2E-02
KEGG:03010	Ribosome	83	4.3E-02
KEGG:03015	mRNA surveillance pathway	41	4.4E-02

*Bonferroni Corrected

Table 2. Enriched BP categories of the AcgS exoproteome (putative SFPs).

<i>BP Category</i>	<i>Nr. Genes</i>	<i>Pvalue*</i>
multicellular organism reproduction	77	7.5E-29
insemination	11	2.8E-13
drug catabolic process	10	1.3E-06
response to biotic stimulus	20	7.3E-06
lipid transport	9	2.2E-05
response to wounding	8	1.9E-03
negative regulation of catalytic activity	9	2.4E-03
reproduction	78	3.3E-27
multi-multicellular organism process	11	2.8E-13
copulation	11	2.7E-12
sperm competition	10	5.6E-12
regulation of female receptivity, post-mating	9	2.0E-11
regulation of female receptivity	9	1.7E-09
female mating behavior	9	2.0E-08
response to external biotic stimulus	20	7.3E-06
response to other organism	20	7.3E-06
carbohydrate derivative catabolic process	9	1.1E-05
mating	12	4.5E-05
reproductive behavior	12	6.2E-05
mating behavior	11	9.4E-05
response to bacterium	13	5.5E-04
lipid localization	9	9.6E-04
aminoglycan metabolic process	8	2.5E-03
negative regulation of molecular function	9	3.5E-03

*Bonferonni corrected

Table 3

19 genes with site-specific test of positive selection (PAML; M7a vs M8a)

<i>Dpse gene</i>	<i>Dpse FBgn ID</i>	<i>Dmel FBgn ID</i>	<i>Symbol name</i>	Diff Abundance†
GA20754	FBgn0080745	FBgn0262559	Mdh2	M-Up
GA16372	FBgn0076387	FBgn0051659	CG31659	M-Up
GA13959	FBgn0073991	FBgn0265434	zip	P-Up
GA18802	FBgn0078802	FBgn0001098	Gdh	P-Up
GA19458	FBgn0079454	FBgn0027582	CG6230	P-Up
GA23027	FBgn0244429	FBgn0264078	Flo2	
GA16248	FBgn0076264	FBgn0038197	foxo	
GA14679	FBgn0074706	FBgn0027872	rdgBbeta	
GA10763	FBgn0070819	FBgn0033160	Dhx15	
GA11564	FBgn0071615	FBgn0037489	CG1234	
GA19689	FBgn0079685	FBgn0037842	CG6567	
GA20526	FBgn0080520	FBgn0261065	Cpsf73	
GA20026	FBgn0080022	FBgn0035121	Tudor-SN	
GA21237	FBgn0081225	FBgn0024238	Fim	
GA17987	FBgn0077996	FBgn0000299	Col4a1	
GA14182	FBgn0074211	FBgn0016075	vkg	
GA13434	FBgn0073470	FBgn0035533	Cip4	
GA16683	FBgn0076698	FBgn0052104	CG32104	
GA25309	FBgn0246690	ND*	ND	

* no ortholog found

†DA proteins (bold) from the indicated selection lines (M or P) with evidence of positive selection

Figure 1**A** **B** **C**

Figure 2

GO Category Enrichment (% terms per group)

A

B

Figure 4

Figure 5

Figure 6

C

Evolutionary Rate Analysis of Accessory Gland Proteome			
Protein Group	(+)ive selection	no (+) selection	P-value *
AcgP	14	2045	NA
P-line	3	62	0.0137
M-line	2	89	0.1449
M+P-line	5	151	0.0084

* 2x2 Fisher's exact test (vs AcgP)