Molecular Epidemiology and Mechanisms of Antibiotic Resistance in Gram-positive Bacteria in Africa: A Systematic Review and Meta-Analysis from a One Health Perspective

John Osei Sekyere^{ac#} and Eric Mensah^b

Department of Pharmaceutics, Faculty of Pharmacy and Pharmaceutical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.^a

Kumasi Centre for Collaborative Research in Tropical medicine, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.^b

Department of Medical Microbiology, School of Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa.^c

#Address correspondence to John Osei Sekyere, jod14139@yahoo.com

Running title: Resistance mechanisms of Gram-positive bacteria

HIGHLIGHTS

- Gram-positive bacteria (GPB) isolated from human, animal and environmental samples were of the same clones and/or shared common resistance genes and mobile genetic (MGEs).
- Multidrug resistant (MDR) clones such as *S. aureus* ST5 and *E. faecium* ST80 were isolated from human, animal and environmental sources.
- *mecA*, *erm(B)*, *erm(C)* *tet(M/K/L)*, and *vanA/B/C* were common in GPB, including VRSA.
- Mean drug resistance rates of isolates from humans, animals and the environment were respectively 62.0% (95% CI: 54.7 – 69.3%), 68.2% (95% CI: 58.0 -78.4%) and 84.6% (95% CI: 69.9 – 99.31%) (P -value<0.0001).
- SCCmec, IS16, and Tn916 mobilized *mecA*, *erm(B)* and *tet(M)* respectively across various GPB species isolated from animals, humans, and the environment.
- A One Health approach to studying antibiotic resistance mechanisms and molecular epidemiology of GPB is warranted.

ABSTRACT

A systematic review and meta-analysis of antibiotic-resistant Gram-positive bacteria in Africa, showing the molecular epidemiology of resistant species from animal, human and environmental sources, is lacking. Thus, the current burden, type, and sources of Gram-positive bacterial resistance and their dissemination routes from farm to fork is absent. To fill this One Health information gap, we systematically searched PubMed, Web of Science and African Journals Online for English research articles reporting on the resistance mechanisms and clonality of resistant Gram-positive bacteria in Africa within 2007 to 2018. The review and all statistical analysis were undertaken with 130 included articles. From our analyses, the same resistant Gram-positive bacterial clones, resistance genes, and mobile genetic elements (MGEs) are circulating in humans, animals and the environment. The resistance genes, *mecA*, *erm(B)*, *erm(C)*, *tet(M)*, *tet(K)*, *tet(L)*, *vanB*, *vanA*, *vanC*, and *tet(O)*, were found in isolates from

humans, animals and the environment. Commonest clones and mobile genetic elements identified from all three sample sources included *Staphylococcus aureus* ST5 (n=208 isolates), ST 8 (n=116 isolates), ST 80 (n=123 isolates) and ST 88 (n=105 isolates), and *IS16* (n=18 isolates), *Tn916* (n=60 isolates) and *SCCmec* (n=202 isolates). Resistance to penicillin (n=4 224 isolates, 76.2%), erythromycin (n=3 552 isolates, 62.6%), ampicillin (n=1 507 isolates, 54.0%), sulfamethoxazole/trimethoprim (n=2 261 isolates, 46.0%), tetracycline (n=3 054 isolates, 42.1%), vancomycin (n=1 281 isolates, 41.2%), streptomycin (n=1 198 isolates, 37.0%), rifampicin (n=2 645 isolates, 33.1%), ciprofloxacin (n=1 394 isolates, 30.5%), clindamycin (n=1 256 isolates, 29.9%) and gentamicin (n=1 502 isolates, 27.3%) (p -value <0.0001) were commonest.

Mean resistance rates of 14.2% to 98.5% were recorded in 20 countries within the study period, which were mediated by clonal, polyclonal and horizontal transmission of resistance genes. A One Health approach to research, surveillance, molecular epidemiology, and antibiotic stewardship to contain ABR should be prioritized.

Keywords: *Staphylococcus spp.*; *Enterococcus spp.*; *Streptococcus spp.*; MRSA; VRE

1. INTRODUCTION

Antibiotic resistance, a threat to public health

Limited research and surveillance data in Africa makes it impossible to track and monitor the true burden of antibiotic resistance (ABR) ¹, particularly the distribution and dissemination of resistance genes between humans, animals and the environment. According to a recent WHO report, the potential for ABR to lead to higher mortalities and morbidities in low- and middle-income countries such as Africa may even be greater as a result of the higher burden of bacterial infections, limited diagnostic capacity and lower access to second-line antibiotics ^{1,2}. This makes it imperative to have a One Health analysis that describes the burden and epidemiology of resistance genes in bacteria isolated from humans, animals and the environment ³.

In a recent review, Gram-positive bacteria (GPB) were responsible for a high proportion of infections among children and showed a high level of resistance to WHO-recommended drugs in Africa⁴. In some African regions, as many as 80% of *Staphylococcus aureus* infections are methicillin-resistant *S. aureus* (MRSA), which show resistance to most standard licensed drugs including quinolones and peptides²⁵. Although *Enterococcus spp.* are mostly not as virulent as *S. aureus*, their multidrug resistance (MDR) propensities restrict drug options for clinicians⁷. Patients infected with MRSA are estimated to be 64% more likely to demise than those infected with methicillin-susceptible *S. aureus* (MSSA)⁶. Reviews addressing GPB in Africa have reported on increasing rates of ABR from blood-stream infections, pneumonia, urinary tract infections and meningitis caused by *Streptococcus agalactiae*, *S. aureus*, *Streptococcus pneumoniae* and *Enterococcus faecium* in both children and adults. Sepsis due to *S. agalactiae* accounts for about 26% of all neonatal deaths and 10% maternal deaths in Sub-Saharan Africa⁸. However, the potential dissemination of these resistant strains from farm (environment and animals) to fork (humans), are less described.

Sources and anthropogenic activities driving resistance

High-level ABR has been reported in humans, animals and the environment, with indiscriminate antibiotic use being fingered as a major contributor in Africa. Resistance genes have been detected in surface water fed with runoff effluents from farms utilizing antibiotics, hospitals, and sewage processing plants as well as in ground water^{9–11}. Furthermore, genes mediating resistance to last-resort GPB antibiotics such as vancomycin have been recovered from raw milk and animal products, pigs, wild animals (buffalo, zebra and cattle), waste water, effluents and patients, implicating veterinary and agricultural use of antibiotics as potential sources of resistance genes in humans^{12–14}. These reports suggest that a larger share of the antibiotics that end up polluting the environment and communities emanate from livestock production^{15–17}. This interconnectivity between animals, humans and the environment, explains the need to adopt a One Health research policy.

Several studies have reported high rates of MDR among GPB isolates from humans, animals and the environment in Africa, mainly as a result of overuse, underuse and wrong choice of antibiotics^{18–24}. Different factors have been implicated in the high rate of ABR to the limited drugs in Africa. These include: unrestricted access to antibiotics over-the-counter without prescription such as selling on the streets; inadequate hygienic practices; uncontrolled usage of antibiotics as growth promoters in food animals production; wrong diagnosis and prescription, off-label use and errors in dosage regimens; use of untreated poultry and cattle manure to fertilize agriculture lands; extensive use of broad-spectrum antibiotics in poultry production; and inefficient chlorination of hospital wastewater effluents before discharge into the environment^{10,18,22,25–29}. Additionally, inadequate knowledge of animals' diseases, misdiagnosis and poor antibiotic handling practices in animal production add up to the overall burden of ABR in Africa¹⁷.

Molecular ABR mechanisms

Selective pressures exerted by various antibiotics used in human and veterinary medicine, as well as in agriculture, have resulted in the emergence and dissemination of numerous mechanisms of resistance in GPB in Africa. Commonly reported mechanisms include *blaZ*, *erm(B)*, *mecA*, *tet(M)*, *vanB* and *vanC*^{30–33}. These resistance genes have been found to be associated with mobile genetic elements (MGEs) such as transposons, conjugative plasmids, integrons, and insertion sequences, which are capable of mobilizing resistance genes across a wide spectrum of bacterial species^{34,35}. *SCCmec*, *Tn916* and *IS16* are notable MGEs that carry major ABR determinants in Africa and are transmissible between clones of the same or different bacteria species by a conjugative mechanism. These MGEs have the potential to thus spread resistance genes from environmental and animal bacterial hosts to human pathogens in Africa; they have therefore been analysed herein^{36–38}.

Purpose of this review

Excellent reviews addressing antimicrobial resistance in some GPB and Gram-Negative ones in Africa have been published^{4,39-44}. However, reviews discussing the molecular epidemiology and mechanisms of ABR in GPB such as *Staphylococcus spp.*, *Streptococcus spp.* and *Enterococcus spp.* in Africa in the context of resistance rates, resistance mechanisms (and MGEs), clonality, and geographical distribution from a One Health perspective are non-existent, to the best of our knowledge. This review sought to fill this gap by analyzing the burden, types, and molecular epidemiology of resistant GPB from a One Health context.

1.1 Search strategy and inclusion criteria

English research articles published within the last ten years (01/01/2007 to 07/08/ 2018) and indexed in PubMed, Web of Science and African Journals Online were searched with the following keywords: “Enterococcus”, and “Streptococcus”, “Staphylococcus”, in permutations and combinations with “resistance AND Africa”. Studies which did not identify the underlying ABR mechanisms/genes as well as the clonality of antibiotic-resistant GPB were excluded. Thus, studies that only reported on antibiotic sensitivity testing (AST) results or undertook ABR surveillance studies without further molecular tests to characterize the ABR mechanisms and/or clonality of the isolates were excluded (Figure 1). In all, 248 studies were excluded because they only had MIC data (See Supplementary data 1). All searches were undertaken independently by both authors in triplicates to ensure replication of the results.

Data extracted from the articles included year of study, country, GPB species, clones, sample sources, sample size/number of isolates, number of resistant isolates, resistance genes and MGEs and antibiotics to which the strains were resistant (Tables 1-6; Supplementary data 2). The mean rate of ABR among GPB per country and in Africa was determined to identify countries with the highest or lowest levels of resistance in Africa (Table 5). As well, the antibiotics to which the isolates were most resistant were determined to evaluate their correlation with the detected/reported resistance mechanisms (Table 6). The resistance mechanisms, as well as MGEs involved in the transmission of resistance genes per species or clone, were determined to assess the means of resistance transfer i.e., horizontal or vertical (through

clonal expansion), per specimen sources (animal, human, and environment) (Figures 2a & 2b). The distribution of clones, resistance genes, and MGEs were considered to identify countries with most resistant clones, resistance genes, and their associated MGEs (Figure 3a).

1.2 Statistical analysis.

The data was analyzed using Microsoft Excel® 2017 and Graph pad prism™ 6 (GraphPad Software, San Diego, CA, USA) (Supplementary data 2). Calculation for the statistical significance of the data was determined using the kolmogorov-smirnov test (with Dallal - wilkinson-Lilliefors p-value) and/or column statistics or one sample t-test, and the confidence intervals determined at 95%. The p-values were two tailed with a Gaussian approximation. A p-value of <0.05 was considered as statistically significant. Only studies that provided the required information were used in the analysis. In all, 130 articles were used for the data analysis (Fig. 1).

2. RESULTS AND DISCUSSION

Of the 1,486 articles returned from the systematic literature search from PubMed, Web of Science and African Journals Online, 130 studies representing 20 out of 54 African countries were included in this review and data analysis (Fig. 1). A total of 249 papers were excluded because they only had MIC data. Tunisia (n=33 studies) recorded the highest number of studies followed by South Africa (n=21 studies), Egypt (n=21 studies), Nigeria (n=13 studies) and Algeria (n=7 studies), Angola (n=6 studies), Uganda (n=5 studies), Democratic Republic of the Congo (n=3 studies), Ghana (n=3 studies), Kenya (n=3 studies), São Tomé and Príncipe (n=3 studies), Gabon (n=2 studies), Tanzania (n=2 studies), Cape Verde (n=1 study), Libya (n=1 study), Namibia (n=1 study), Senegal (n=1 study) and Sudan (n=1 study). Majority of the included studies were undertaken in Northern Africa (n=65 studies, 50%), Southern Africa (n=35 studies, 26.9%) and West Africa (n=18 studies, 13.9%). Different rates of resistance to antibiotics were reported in different countries in Africa (Tables 2-5; Supplementary data 1).

A meta-analysis of published literature confirmed the presence of a high mean rate of drug resistance in GPB isolated from humans (62.0%, 95% CI: 54.7 – 69.3%), animals (68.2%, 95% CI: 58.0 -78.4%) and the environment (84.6%, 95% CI: 69.9 – 99.3%) (P -value<0.0001) in Africa, albeit many studies that did not address the molecular mechanisms of resistance in GPB were excluded. Obviously, the mean rate of resistance would have been higher had all research articles using only phenotypic methods to describe ABR in GPB been included (Supplementary data 1). Interestingly, although a lesser number of GPB were isolated from environmental sources, they expressed higher ABR than those from humans and animals; hence, the higher mean resistance rate of 84.6%. This also underscores the fact that there is increasing ABR genes in the environment, obviously due to antibiotic pollution from human activity. Evidently, ABR is high among GPB in certain regions in Africa (Figures 3a & 3b) (Table 5) and underpins the need to up the ante against this menace through increased molecular surveillance research, education of clinical microbiologists on ABR, and antibiotic stewardship.

Studies describing detailed molecular mechanisms of GPB resistance and molecular epidemiology in Africa are few, making it difficult to paint a vivid comprehensive picture of ABR in Africa. However, this review shows that *S. aureus* ST5, *E. faecium* ST18, ST80 and ST910, *E. faecalis* and *S. agalactiae* harbouring *mecA*, *tet* and *erm* genes, were commonly found in humans, animals and the environment, particularly in Northern, Western, and Southern Africa. Thus, careful use of β -lactams, tetracyclines, and macrolides is warranted to prevent further selection and dissemination of these resistance genes and resistant clones. Furthermore, it will be prudent for countries within these regions to review their recommended antibiotic regimens, guidelines/protocols for infections caused by these species.

erm(B) and *tet*(M) were found in *S. aureus*, *Enterococcus spp.* and *Streptococcus spp.*, with *erm*(B), *tet*(M) and *vanA* genes being mobilized by Tn916 and IS16, indicating horizontal transfer within same clones, different clones and species. The discovery of same clones and resistance genes in specimens from humans, animals and the environment suggest a possible transmission of these clones between humans, animals and the environment, corroborating the need for a One Health approach to infection

control and management of antibiotic-resistant infections. Further molecular epidemiological surveillance in the above-mentioned states is crucial to forestall further spread of these resistant pathogenic clones both within their borders and from their borders to other countries.

Resistance rates per countries and MDR GPB species

High mean resistance rates were reported in Sudan (98.5%), South Africa (82.7%) Nigeria (71.2%), Egypt (70.5%), Angola (66.2%), Tunisia (66.8%), Ghana (65.1%), Algeria (62.2%) etc. (Table 5). Cross-

contamination of multi-drug resistant bacteria between patients and the environment accounted for the high rate of resistance in Algeria ⁴⁵⁻⁴⁹. The high rate of ABR in Tunisia was attributed to cross contamination between hospital patients and hospital environment, immune deficiency ⁵⁰, overconsumption of antibiotics, heavy consumption of sheep meat, which is a reservoir of MRSA, and high consumptions of antibiotics in animal feed ^{51,52}. In Egypt, inappropriate antibiotic prescription practices ²⁹, inadequate hygienic handling and processing of food ¹², and close contact with pet dogs accounted for the high resistance ⁵³. The high rate of drug resistance in Nigeria has been attributed to the exchange of resistance genes between farm animals or their products and man ^{54,55}, existence of MRSA in clinical and community settings ⁵⁶, uncontrolled usage of antibiotics ⁵⁷ and the presence of efflux pumps in coagulase-negative staphylococcus strains ⁵⁸. Expansion of resistant clones ⁵⁹, variability of hospital acquired MRSA clones ⁶⁰, consumption of unpasteurized milk or inefficient thermal processing of milk ²¹, shedding of resistant clones from animals to the environment and heavy consumption of antibiotics to treat TB due to high HIV burden ⁶¹, were incriminated for the high-level resistance in South Africa.

Staphylococcus spp. (*S. aureus*, *S. haemolyticus* and *S. saprophyticus*); *Streptococcus spp.* (*S. pyogenes* and *S. agalactiae*), and *Enterococcus spp.* (*E. faecium*, *E. faecalis*, *E. hirae*, *E. durans*, and *E. gallinarum*) were the antibiotic-resistant GPB widely distributed in Northern, Southern, Western and Central Africa. The high number of *tet*(M/L/K), *erm*(A/B/C), *aph*(3')-III and *vanA/B/C* in *Staphylococcus spp.*, *Enterococcus spp.*, and *Streptococcus spp.* reported in Tunisia, South Africa, Nigeria, Algeria and Egypt accounted for the high rate of resistance to tetracycline, erythromycin, kanamycin and vancomycin

(Figure 3a). Such resistant GPB are known to compromise the safety of invasive medical procedures such as organ transplants, orthopedic surgery, and cancer treatment. In addition, infections such as sepsis, endocarditis, deep wound infections, pneumonia, meningitis and urinary tract infections caused by these resistant pathogens are becoming increasingly fatal due to limited treatment options^{62,63}. The abuse of antibiotics as growth promoters, prophylaxis, and metaphylaxis in food animals in these countries have been implicated in the selection of resistant bacteria that can pass on to humans through food consumption, direct contact with animals and the environment, as well as trade of animals and food products between countries⁶⁴.

Approximately 26, 385 GPB were isolated from humans (n=83 studies), animals (n=32 studies) and the environment (n=14 studies) (Tables 1-4), with mean rates of ABR varying from 14.2% to 98.5% across the 20 included countries (Tables 2-5). The antibiotics to which the isolates were most resistant to were penicillin (n=4 224 isolates, 76.2%), erythromycin (n=3 552 isolates, 62.6%), ampicillin (n=1 507 isolates, 53.9%), sulfamethoxazole/trimethoprim (n=2 261 isolates, 46.0%), tetracycline (n=3 054 isolates, 42.1%), vancomycin (n=1 281 isolates, 41.2%), streptomycin (n=1 198 isolates, 37.0%), rifampicin (n=2 645 isolates, 33.1%), ciprofloxacin (n=1 394 isolates, 30.5%), clindamycin (n=1 256 isolates, 29.9%), and gentamicin (n=1 502 isolates, 27.3%) (p -value <0.0001) (Tables 2-4 & 6). Countries with high number of studies such as Tunisia, South Africa, Egypt and Nigeria recorded high number of ABR (Table 5) and high number of *mecA*, *erm(B)*, *tet(M)*, *drfG* and *vanB* resistance genes (Figure 3a). Vancomycin resistance was reported in seven studies each for animals and the environment, and 12 studies in Humans. Vancomycin-resistant *Enterococcus spp.* (n=102 isolates) and vancomycin-resistant *Staphylococcus spp.* (n=258 isolates) were reported in humans, animals and the environment (Tables 2-4; Figures 2). Vancomycin-resistant *Staphylococcus aureus* (VRSA) was reported in animals (n=238 isolates), the environment (n=15 isolates) and humans (n=5 isolates). A similar situation occurred with vancomycin-resistant *E. faecium*, which was isolated from the environment (n=306 isolates), animals (n=671 isolates) and humans (n=26 isolates) (Supplementary data 1).

Antibiotic-resistant *S. aureus* ST5, *E. faecium* (ST18, ST80 and ST910) and *E. faecalis* harbouring *mecA*, *erm(B)*, *erm(C)*, *tet(M)*, *tet(K)*, *tet(L)* and *vanB* were isolated from humans, animals and the environment, albeit in higher proportion in humans and animals than the environment (Tables 2-4). For instance, Farhat et al. (2014)⁴⁶, van Rensburg et al. (2012)⁵⁹ and De Boeck et al. (2015)⁶⁵ in Algeria, South Africa and Democratic Republic of Congo respectively, reported on resistant *S. aureus* ST5 in humans whilst Fall et al. (2012)⁶⁶ reported on the same clone (*S. aureus* ST5) in pigs from Senegal. Further, Mariem et al. (2013)²⁴ isolated the same clone (*S. aureus* ST5) from the environment in Tunisia, suggesting that this clone is widely distributed in Africa in humans, animals and environment. It is currently not clear whether this clone first emerged from humans, animals or the environment, but its presence in all three spheres shows the possibility of resistant species and clones being disseminated between animals, humans and the environment. Notably, *S. aureus* ST5 is among the frequently reported clones in Asia⁶⁷ and recent evidence suggest that it has spread from hospitals into communities, resulting in community-acquired MRSA⁶⁸.

Similarly, Lochan et al. (2016)³⁰ in South Africa, Dziri et al. (2016)²⁰ and Elhani et al. (2014)⁶⁹ in Tunisia isolated resistant *E. faecium* ST80 from humans. For the first time, *E. faecium* ST80 was isolated from environmental samples in a hospital in Tunisia by Elhani et al. (2013)⁶⁹ and Dziri et al. (2016)⁷⁰. Transmission of this resistant clone to animals is possible, although not yet reported. This implies that these resistant species and clones are circulating between humans and the environment, underpinning the broad host range and transmissibility of these strains between humans and the environment.

mecA was the predominant resistance gene, which corresponded with the higher penicillin resistance recorded (Figure 2aii). MRSA strains were the most commonly isolated strains ($\geq 2,350$)⁷¹⁻⁷⁴. This is consistent with the global report of increasing prevalence of MRSA^{75,76}. MRSA harbours the *mecA* gene, which is carried by the *SCCmec* MGE, and mediates resistance to multiple β -lactam antibiotics⁷⁷. From this review, MRSA showed resistance to eleven different antibiotic classes: aminoglycosides (gentamicin, tobramycin), β -lactams (penicillin, ampicillin, oxacillin, cefoxitin), fluoroquinolones (ciprofloxacin,

levofloxacin, ofloxacin), glycopeptides (vancomycin), lincosamide (clindamycin), macrolides (erythromycin), phenicols (chloramphenicol), rifamycins (rifampicin), streptogramins (pristinamycin), sulfonamides (trimethoprim/sulfamethoxazole), and tetracyclines (tetracycline). MRSA is thus a worrying public health threat as some strains have evolved resistance to almost all licensed drugs (26). Vancomycin-resistant Enterococci (VREs) (≥ 594), which were reported in Northern and South Africa, also pose a serious threat to public health as they are resistant to vancomycin, a glycopeptide that is reserved for fatal or life-threatening Gram-positive infections, and other important antibiotics such as ampicillin, erythromycin, fluoroquinolones (ciprofloxacin, levofloxacin), gentamicin, rifampicin, streptomycin, trimethoprim/sulfamethoxazole and tetracycline. In this study, enterococcus isolates had a resistance rate of 60.1% (95%, CI=32.2 -87.9) (p-value = 0.0005) to vancomycin (Table 6). Multidrug resistance in VREs increases VRE-associated mortality rates, which is likely to increase to 75% compared with 45% from susceptible strains^{13,80}. As well, evolution of macrolide resistance (42.0%, 95% CI: 12.02 – 72.1) (p-value = 0.0129) in drug-resistant streptococci is limiting treatment options and resulting in high mortalities^{81–83}. In this study, MRSA, VRE and drug-resistant streptococci remain major public health threats, calling for measures to contain ABR. Novel antibiotics such as linezolid, synergid, and daptomycin should be used empirically whilst awaiting susceptibility results. The empirical therapy can be changed or maintained based on the susceptibility report⁸⁴.

Resistance rates of species per animals, humans and the environment

The rates of ABR in isolates recovered from the environment was highest, followed by isolates from animal sources. Among environmental isolates, 91.2% (95%, CI=78.8–103.6) were resistant to penicillin, 82% (95%, CI=40.6–123.4) were resistant to sulfamethoxazole/trimethoprim, 68.5% (95%, CI=24.1–100) were resistant to ampicillin, 60.8% (95%, CI=25.0–96.6) were resistant to vancomycin, 56.9% (95%, CI=–40.7–73.2) were resistant to erythromycin, 54.5% (95%, CI=29.49–79.5) were resistant to ciprofloxacin, and 51.3% (95%, CI=21.3–100) were resistant to clindamycin (Table 6). Among animal isolates, 71.8% (95%, CI=54.9–88.73) were resistant to penicillin, 58.9% (95%, CI=36.1–81.7) were resistant to clindamycin, 58.5% (95%, CI=37.6 –79.4) were resistant to ampicillin, 49.6% (95%,

CI=30.1–69.1) were resistant to trimethoprim/sulfamethoxazole, 42.3% (95% CI=17.7–67.0) were resistant to vancomycin, 47.6% (95% CI=34.0–61.2) were resistant to erythromycin, and 38.8% (95% CI=21.3–56.3)(p-value = 0.15) were ciprofloxacin resistant (Table 6; Supplementary file 1).

The rates of resistance were much lower in humans for most of the antibiotics used (Tables 2-4). Among the various species, *Enterococcus spp.* and *Staphylococcus spp.* recorded high rates of resistance for most antibiotics (Figure 3b). *Streptococcus spp.* reported low rates of resistance except for tetracycline to which it recorded a high rate of 55.13% (95%, CI=20.63.18–89.64) (p-value = 0.006). Resistance to vancomycin was not reported in any *Streptococcus spp.* Isolate (Table 6).

Enterococcus spp., mainly *E. faecium* and *E. faecalis*, recorded a resistance rate of 98.5% (95%, CI=94.5–102.6)(p-value = 0.0001) to clindamycin, 81.6% (95%, CI=52.1–110)(p-value = 0.0008) to trimethoprim/sulfamethoxazole, 64.0% (95%, CI=50.0–78.1)(p-value=0.0001) to erythromycin, 60.1% (95%, CI=32.2–87.9)(p-value = 0.0005) to vancomycin, 57.3% (95%, CI=24 -90.7)(p-value=0.0057) to penicillin, 51.7% (95%, CI=35.8–67.6)(p-value=0.0001) to tetracycline, 49.9% (95% CI=31.3–68.5)(p-value = 0.0001) to ciprofloxacin, 48.9% (95% CI=20.6–77.2)(p-value=0.004) to kanamycin, 47.1% (95% CI=26.7–67.7)(p-value=0.0006) to ampicillin, 40.8% (95% CI=24.3–57.4)(p-value=0.0001) to streptomycin and 34.0% (95% CI=19.7–48.4)(p-value=0.0002) to gentamicin (Table 6).

S. aureus showed high resistance (79.6%) to penicillin (95% CI=69.7–89.5)(p-value = 0.0001), 67.8% to erythromycin (95% CI=11.5–147.0)(p-value = 0.0917), 55.5% to ampicillin (95% CI=44.50–88.5)(p-value = 0.0001), 39.3% to trimethoprim/sulfamethoxazole (95% CI=39.3–47.8)(p-value = 0.0001), 36.9% to tetracycline (95% CI=29.3–44.5)(p-value = 0.0001), 35.8 to streptomycin (95% CI=14.7–57.0)(p-value = 0.004), 33.6% to rifampicin (95% CI=20.1–47.03)(p-value = 0.0001), 24.0% to clindamycin (95% CI=14.9–33.1)(p-value = 0.0001), 23.9% to ciprofloxacin (95% CI=17.6-30.2)(p-value= 0.0001), 22.7% to vancomycin (95% CI=4.3–41.2)(p-value = 0.0212) and 22.2% to vancomycin (95% CI=15.7–28.3)(p-value = 0.0001) (Table 6).

Resistance mechanisms, clones, and MGEs

Few studies identified the clones and MGEs in the resistant isolates. Of the 130 included studies, 32 identified the clones whilst 22 described the MGEs, which were used in the statistical analysis. The most dominant gene detected in Africa, which was widespread and responsible for resistance in GPB, was *mecA* (n=3 547), followed by *erm*(B) (n=1 268), *vanC1/2/3* (n=971), *tet*(M) (n=720), *blaZ* (≥ 565), *dfpG* (n=422), *vanB* (≥ 451), *aph*(3')-IIIa (≥ 170) and *aac*(6')-aph(2') (≥ 268) (p-value = 0.0011) (Fig. 2a).

Figure 2b represents MGEs per clone. *S. aureus* clones ST5, ST8, ST 80 and ST88 were highly associated with *mecA*. Resistant *S. aureus*, *E. faecium* and *E. faecalis* clones such as *S. aureus* ST5, and *E. faecium* clones ST18, ST80, and ST16 were widely distributed in humans, animals and the environment. Similarly, *mecA*, *erm*(B), *erm*(C), *tet*(M), *tet*(K), *tet*(L), *vanB*, *vanA*, *vanC* and *tet*(O) were reported in isolates from humans, animals and the environment (Table 1).

IS16 and *Tn916* were found with the resistance genes *erm*(B) and *tet*(M) in *E. faecium* (ST18, ST80 and ST910), *S. agalactiae* (ST612, ST616 and ST617), *E. faecalis* and *S. pyogenes* (*emm*18, *emm*42, *emm*76 and *emm*118) isolated from humans, animals and the environment (Tables 2-4; Figure 2b). *tet*(M) was associated with *Tn916* transposon in tetracycline-resistant *S. agalactiae*⁸⁵ and *S. pyogenes*⁸¹ in humans in Tunisia. Fischer et al. (2013) also reported the association between *Tn916* and *tet*(M) in tetracycline-resistant *S. agalactiae* in camel in Kenya⁸⁶. Similarly, *IS16* was found in vancomycin-resistant *E. faecium* (ST80, ST180 and ST910) in humans and the environment in Tunisia^{69,70}. Investigations into the association between MGEs and resistance genes were limited by few studies (n=22 studies) on MGEs.

From Tables 2-4, majority of the resistance genes namely, *mecA*, *erm*(B), *tet* (M), *vanA* etc. were responsible for drug resistance to antibiotics such as aminoglycosides (gentamicin, streptomycin, kanamycin), β -lactams (penicillins, cephalosporins), fluoroquinolones (ciprofloxacin), macrolide (erythromycin), sulfamethoxazole/trimethoprim, tetracycline and glycopeptides (vancomycin). These resistance genes were widely distributed in Northern Africa (Tunisia, Algeria, Egypt, Morocco, and

Libya) and Southern Africa (South Africa and Namibia). All the three different MGEs (Tn916, SCCmec and IS16) were reported in Tunisia, with two being reported in Kenya (SCCmec and Tn916). IS16 was only reported in an *E. faecium* infection in Tunisia (Figure 3) whilst *mecA* was mostly associated with SCCmec. *erm*(B) and *tet*(M) were highly associated with Tn916 and IS16.

In Africa, different studies have reported SCCmec-borne *mecA* in *S. aureus* in humans, animals and the environment^{23,47,60,66,87} besides the discovery of IS16 and Tn916 in the environment of *erm*(B) and *tet*(M) genes in Enterococcus and Streptococcus. These reports show that MGEs are mediating the dissemination of these (and possibly other) resistance genes across different GPB clones and species. MGEs-mediated mobilization of various resistance genes in different GPB clones and species in humans, animals and the environment (Tables 1-4; Figure 2b) calls for prompt measures to contain ABR as the situation may worsen if additional resistance genes are acquired by the MGEs. Resistance genes on MGEs can be horizontally transferred to susceptible cells or vertically transferred to daughter clones^{37,88,89}, which can easily spread these resistance genes to susceptible pathogens. The higher number of resistant Gram-positive cocci and mean resistance rate in Tunisia may be due to the presence of these three MGEs in this region^{69,70,81,90}

Molecular epidemiology of antibiotic-resistant GPB

Staphylococcus spp. (*S. aureus*, *S. haemolyticus* and *S. saprophyticus*)

North Africa: Algeria, Egypt, Morocco, Tunisia, Libya

Algeria.

S. aureus was recovered from two different studies in Algeria. In assessing the nasal carriage of *S. aureus* in patients with medical conditions including pneumonia, urinary tract infections, osteoarthritis, heart diseases, diabetes and chronic kidney disease, Djoudi *et al.* (2014) isolated MRSA⁴⁶. They also found nasal carriage of *S. aureus* to be significantly associated with cancer and previous hospitalization of patients with kidney failure due to immunological suppression and hemodialysis. The nine MRSA isolates, i.e. ST80 (n=4), ST5 (n=2), ST22 (n=2) and ST535 (n=1), harboured *mecA* and were resistant to

tobramycin (n=6), gentamicin (n=1), trimethoprim/sulfamethoxazole (n=2), tetracycline (n=3) and erythromycin (n=1). MRSA ST80 is a well-known and frequent etiological agent of infections in North Africa and Middle-East countries^{91,92}. Typing of 64 MRSA isolated from human pus (n=47), venous catheters (n=7), tracheal aspirates (n=4), punction fluids (n=3), blood (n=2) and urine (n=1) in 64 Algerian patients revealed that 50 were hospital acquired (HA-MRSA) and 14 community acquired (CA-MRSA), which were all resistant to ceftiofur and oxacillin⁴⁷. *mecA*, mobilized by SCC*mec*, was the only detected mechanism of resistance.

Egypt

MRSA have been respectively isolated in five animal-based and two human-based studies in Egypt between 2011 to 2017. Hashem et.al (2013) isolated 94 *S. aureus* strains from blood and wounds in which 45 were MRSA while 25 were fluoroquinolone-resistant²⁹. Mutations such as C2402T, T2409C, T2460G, T1497C, and A1578G in gyrase enzymes, which leads to fluoroquinolones' target-site alterations, were implicated in resistance to fluoroquinolones (ciprofloxacin, levofloxacin, ofloxacin). The high rate of fluoroquinolone resistance (55.56%) among MRSA infections is rather concerning as patients unable to tolerate vancomycin are treated with other antibiotics such as fluoroquinolones. Vancomycin is often reserved as a last-resort therapy for MRSA infections due to their high resistance to several antibiotics.

Multidrug resistance to drugs such as gentamicin, ampicillin, amoxicillin, cefepime, tetracycline and chloramphenicol in MRSA is mediated by diverse resistance mechanisms including impermeability effects and efflux pumps. Unrestricted access to antibiotics and inappropriate prescriptions were responsible for the high rates of drug resistance in this study²⁹. In a similar study, MRSA was isolated from patients suffering from surgical wound infections, diabetic foot, abscess and burns. Although *mecA* was the only mechanism of resistance, the isolates were multiple-resistant to several antibiotics belonging to the β -lactams, aminoglycosides, fluoroquinolones, macrolides, lincosamides, tetracyclines and

glycopeptides, indicating other mechanisms of resistance⁹³. It therefore implies that administration of such antibiotics will not relieve patients from *S. aureus* infections. The high rate of *S. aureus* isolation confirms it to be the most prevalent Gram-positive pathogen isolated from soft tissue and wound infections.

Al-Ashmawy *et. al.* detected a high rate of MRSA (53%) in milk and dairy products believed to originate from human contamination rather than contamination from animals. Besides being resistant to β -lactams and other antibiotics, thirty-six of the isolates were resistant to vancomycin known to be effective in treating MRSA infections¹², making milk and dairy products a significant source of multidrug-resistant and toxigenic *S. aureus* infections. The occurrence of MRSA in pets such as dogs admitted in a veterinary clinic⁵³ may confirm a possible route in the community transmission of this pathogen, which is emerging as a veterinary pathogen of public health importance.

In 2017, Osman and colleagues detected *Staphylococcus spp.* in imported beef meat. Sixteen of these isolates were MDR and showed resistance to different groups of antibiotics due to resistance mechanisms such as *mecA*, and mutations in *gyrA* and *gyrB*. Indeed, MRSA has made methicillin and other β -lactams antibiotics clinically useless as a result of their high MDR⁹⁴. Imported meat acts as a transmission vector for MRSA and is worrisome as *Staphylococcus spp.* are among the most common foodborne pathogens causing food poisoning outbreaks worldwide. Of 133 *S. aureus* recovered from animal origin, more than 70% were MDR and 30 were MRSA, exhibiting high resistance to clindamycin, co-trimoxazole, tetracycline, oxacillin, cefoxitin, ceftriaxone and erythromycin; four of the isolates were resistant to vancomycin²³. The isolates showed the maximum sensitivity to imipenem, chloramphenicol and rifamycin, which is consistent with similar reports in China and Pakistan^{95,96}, indicating their effectiveness in treating *S. aureus* infections.

MRSA was isolated from chicken products mainly due to poor hygienic handling processes, posing a risk to public health in 2016. The mean *S. aureus* count in the chicken products were beyond the permissible limits of the Egyptian organization for Standardization and Quality Control (EOSQC 2005), coupled with

resistance to different antibiotics classes; thus, retail chicken products could constitute a high health risk to human consumers²⁸

Morocco

In a study to assess *S. aureus* carriage among end-stage renal diseases patients undergoing hemodialysis, 42.9% were carriers, of which only one was MRSA. The methicillin-susceptible *S. aureus* (MSSA) was resistant to many of the local antibiotics, thus limiting the successful treatment of MSSA infections. Moreover 81.8% of the MSSA were penicillin-resistant. The male gender and age 30 or below were identified as risk factors of *S. aureus* nasal carriage (P -value < 0.001)²⁷. Periodic monitoring of patients with hemodialysis is crucial as they are at increased risk of *S. aureus* infection due to periodic hospitalization, immunosuppression and high invasive vascular interventions.

Tunisia

Resistant *S. aureus* was isolated from the environment, animals and humans between 2011 to 2017. Ben Said, et al. recovered 12 MSSA from wastewater samples that were resistant to penicillin (n=12 isolates), erythromycin (n=7 isolates), tetracycline (n=1 isolate) and clindamycin (n=1 isolate) due to the presence of *blaZ* (n=7), *msr(A)* (n= 7) and *tet(K)*(n=1). These resistant strains were of ST3245(n=7) and ST15(n=1)¹⁸, which have been also reported in animals and humans. In an investigation to evaluate the prevalence of coagulase-negative Staphylococcus (CoNS) in the hospital environment, MDR *S. haemolyticus* and *S. saprophyticus* were the most dominant. Methicillin resistance was detected in *S. haemolyticus*, *S. epidermidis* and *S. saprophyticus*. These isolates were resistant to erythromycin, tetracycline, gentamicin, kanamycin, tobramycin and streptomycin due to the presence of *msrA* (32), *erm(C)* (8), *tet(K)* and *tet(M)*, *aac(6')-Ie-aph(2'')-Ia* (16), *aph(3')-IIIa*(19), *ant(4')-Ia* (n=14) and *ant(6')-Ia* (3)⁹⁷. The high prevalence of MDR *Staphylococci spp.* isolates may result from transmission between the staff, patients and the environment. Strict infection controls are needed as infections caused by CoNS are common cause of death, particularly in low-birth-weight children, and are opportunistic infections in immunocompromised patients⁹⁸.

Moreover, nasal swab from sheep detected five MRSA (*mecA*=5), which were all of ST153 and carried *blaZ*, *ant(6)-Ia*, *aph(30)-IIIa*, *erm(C)*, *tet(K)*, and *fusB* genes that respectively encoded resistance to penicillin, streptomycin, kanamycin, erythromycin, tetracycline and fusidic acid. This study shows that the nares of healthy sheep could act as reservoirs of MRSA⁹⁹.

Between 2011 to 2012, 99 MRSA strains were detected from nasal swabs, blood, catheter, wounds, pleural puncture and abscess, among which 39 were tetracycline resistant. These isolates were resistant to aminoglycosides, fluoroquinolones, macrolides and lincosamide, with mechanisms of resistance including *mecA* (n=24), *tet(K)* (n=6), *tet(L)* (n=1) and/or *tet(M)* (n=18), *erm(A)*(n=14), *aph(2')-acc(6')* (n=13). Identified drug-resistant strains included ST247 (n=12), ST239 (n=6), ST728 (n=2), ST241 (n=1), ST398 (n=1), ST5 (n=1) and ST641 (n=1)⁵⁰. For the first time, clonal lineage ST398, which has been reported in pigs from several studies in USA, South America, Asia and Canada^{100–103}, was found in human MRSA isolates in Africa in a nasal swab of a 74-year old patient.

Additionally, 69 MRSA strains were isolated from hospital-acquired and community-acquired infections. Although *mecA* (n=59) was the only mechanism of resistance identified, the isolates were resistant to aminoglycosides, tetracycline, fluoroquinolones, macrolides and rifampicin. The resistant clones were ST1 (n=2), ST5 (n=5), ST22 (n=1), ST80 (n=41), ST97 (n=2), ST153 (n=2), ST239 (n=4), ST241 (n=3), ST247 (n=3), ST256 (n=1), ST1819 (n=3) and ST1440 (n=1)²⁴.

Mezghani Maalej and colleagues (2012) isolated five pristinamycin-resistant *S. aureus* strains from patients with skin infections. These isolates were MDR (Table 2), being the first detection of resistance to streptogramins due to *vat(B)* and *vga(B)* resistance genes¹⁰⁴, which emerged due to selective pressure from the use of pristinamycin. Thirty-six methicillin-resistant *S. haemolyticus* (MRSHae) were isolated from neutropenic patients (suffering from febrile neutropenia) with hematological cancer between 2002 and 2004. These MDR isolates carried SCC*mec*-borne *mecA* (Table 2)¹⁰⁵, which agrees with a report on *S. haemolyticus*' MDR capacity, particularly in immunocompromised patients^{106,107}

Libya

Due to the high risk of MRSA colonization developing into infections in children, nasal samples were collected from children inpatients, their mothers, healthcare workers and outpatients' workers, which yielded a MRSA nasal carriage rate of 8.3%, 11% ,12.3% and 2.2% respectively in Libya ¹⁰⁸. Thus, nasal carriage of MRSA is common in inpatients children, their mothers and health workers in Libya and could be a source of MRSA infections.

West Africa: Ghana, Nigeria, Senegal

Ghana

Among 308 staphylococcus isolates collected across Northern, Central and Southern Ghana in 2013, low prevalence of antibiotic resistance was reported except for penicillin (97%), tetracycline (42%) and erythromycin (6%) ¹⁰⁹. Moreover, *mecA* was detected in only nine isolates, implying the presence of other β -lactam resistance mechanisms. The MRSA clones included ST8 (n=1), ST72 (n=1), ST88 (n=2), ST239 (n=1), ST250 (n=2), ST789 (n=1), and ST2021 (n=1). In a similar study that characterized 30 MRSA isolates resistant to tetracycline, fluoroquinolones and macrolides, *tet(M)* (n=13), *tet(K)* (n=10), *aphA3* (n=7), *aacA-aphD* (n=5) and *erm(C)* (n=4) were detected. Similar and different resistant clones, viz. ST88 (n=8), ST8 (n=5), and ST247 (n=4) were detected ¹¹⁰, indicating high MRSA clonal diversity in Ghana. These studies show a high rate of resistance to non- β lactams that further complicate MRSA treatment. Furthermore, the isolation of USA300 and other epidemic multidrug-resistant MRSA clones calls for MRSA surveillance and adequate control measures.

Nigeria

Five different studies reported drug-resistant *S. aureus* from several human anatomical sites such as throat swabs, soft skin and tissue infection, urinary tract and respiratory infections, wound, vagina, otitis, conjunctivitis, septicemia and bronchitis. Of a total ≥ 602 isolates, ≥ 433 were resistant to several antibiotic classes (Table 1). Of note, 429 of the ≥ 433 drug-resistant isolates were all resistant to cotrimoxazole or trimethoprim/sulfamethoxazole (SXT). Mechanisms of resistance included *mecA* (≥ 54), *blaZ* (n=284),

dfrA (≥ 5) and *dfrG* (≥ 152). *S. aureus*-resistant clones ST8, ST14, ST37, ST39, ST88, ST152, ST241, and ST772 were present. Colonized persons, including immune-compromised individuals, facilitated the spread of *S. aureus* and MRSA ST8 identified as ubiquitous in various geographic areas of Nigeria. High utilization of co-trimoxazole or SXT because of low cost and easy obtainability through lenient medication regulations were implicated for the high resistance⁵⁶. Besides *S. aureus*, *S. haemolyticus* was the major species isolated, and is considered as the second most detected and clinically important *Staphylococci spp.*, particularly in immunocompromised patients¹¹¹. All the *S. haemolyticus* isolates detected were resistant to at least three antibiotics classes (Tables 2-4)¹¹².

Moreover, O. Ayepola *et al.* (2015) reported a higher rate of 20.8% *S. aureus* from UTIs than the reported ranges in Africa (6.3-13.9%), and far exceed the rate reported from Europe and Brazil (1.1%)¹¹³. None of the isolates exhibited resistance to vancomycin, linezolid, daptomycin and mupirocin; indicating their usefulness in treating *S. aureus* infections. Co-trimoxazole, which was previously clinically valuable in treating MRSA infections, demonstrated the highest level of resistance, hence it's not recommendable^{56,57,90,112}. In a study to examine the genetic mechanism(s) of resistance in CoNS in faecal samples, all the 53 isolated CoNS were Penicillin V-resistant and between three to 19 exhibited multidrug resistance (Table 2); *mecA* (n=15), *erm(C)*, *tet(M)* (n=4) and *tet(K)* (n=6) were identified¹¹². CoNS isolates from faeces carrying tetracycline, macrolides and aminoglycosides resistance genes may transfer them inter- and intra-species, disseminating MDR in *Staphylococcus*.

Senegal

A low prevalence of MRSA (10.5%) was reported in Senegalese pigs compared to those reported in developed countries. This might be due to a lesser veterinary antibiotic use as growth promoters and/or for therapy. However, all the isolates were resistant to penicillin, 27 were resistant to co-trimoxazole and 16 were resistant to tetracycline⁶⁶. Five of the MRSA were of ST5⁶⁶, evincing the spread of this clone in

animals, humans^{46,59}, and the environment²⁴; the importance of this clone as a cause of human infections is well-established⁶⁸.

Cape verde

In Cape Verde, a low prevalence of 5.6% (6/107) MRSA nasal carriage was documented in 2015. The predominant MRSA clones was ST5 (n=3), ST8 (n=1) and ST88 (n=2). These isolates showed significant level of resistance to erythromycin (ERY), sulphamethoxazole-trimethoprim (SXT) and penicillin G (PEN)¹¹⁴.

Central Africa: Gabon, D.R. Congo

Gabon

In Gabon, *S. aureus* isolated from colonized persons, blood, as well as soft and skin tissue infections resulted in 49% (104/212) resistance to trimethoprim: *dfrA* (n=1), *dfrG* (n=100), *dfrK+G* (n=1), *dfrB* (n=2), and *mecA* (n=1) were detected in the isolates⁵⁵. Thus, *dfrG* is obviously the most abundant and common trimethoprim resistance mechanism in Africa, refuting *dfrB* mutation as the main mechanism of resistance to trimethoprim^{115–117}.

D.R. Congo (DRC)

A total of 215 (79.3%) drug-resistant *S. aureus* isolates were collected between 2015 to 2017 from nasal swab and bloodstream infections in the D. R. Congo; 70 isolates were MRSA. Other major resistance genes mediating resistance to trimethoprim/sulfamethoxazole, aminoglycoside, macrolides, tetracycline, penicillin, and chloramphenicol were *dfrG* (≥ 120), *tet(K)* (≥ 98), and *femA* (≥ 98). MRSA showed high-level resistance to β -lactams, aminoglycoside, macrolides and tetracycline. The pathogen caused severe infections such as pneumonia, meningitis, complicated urinary tract infections, gynaecological infections and peritonitis. *S. aureus* ST8 (≥ 47) was the dominant clone, followed by ST152 (≥ 17), ST5 (≥ 2) and ST88 (≥ 2). In DRC, MRSA ST8 outnumbers the African MRSA clone ST88, which is dominant in Africa. The high-level oxacillin resistance in DRC was associated with a mutation in *femA* (Y195F) whilst

high-level trimethoprim resistance was due to the detection of *dfrG*, which is consistent with trimethoprim resistance in Africa and Asia. In Africa, SXT or cotrimoxazole is frequently administered as prophylactic to immuno-suppressed patients such as HIV/AIDS patients to prevent opportunistic infections such as *Pneumocystis carinii* pneumonia, toxoplasmosis and bacterial pneumonia¹¹⁸. Hence, prophylactic use of SXT in HIV patients may impact resistance. Additionally, there was high-level MDR among MRSA, which is a great concern as microbiological laboratories/facilities and second-line antibiotics are rare in DRC. Moreover, the detection of nasal carriage among healthcare workers' demands strict infection controls and surveillance^{65,119,120}.

East Africa: Kenya, Tanzania

Kenya

In contrast to earlier studies done in Kenya, Omuse and colleagues (2016) detected a wide genetic diversity of MRSA and well-established epidemic MRSA clones among clinical isolates. MRSA clonal complexes 5, 22 and 30, implicated in several outbreaks were described. These clones included ST5 (n=1 isolates), ST8 (n=2 isolates), ST22 (n=4 isolates), ST88 (n=1 isolates), ST241 (n=12 isolates), ST239 (n=2 isolates) and ST789 (n=1 isolates). Approximately 41% of the MRSA in the study were MDR (Table 2), showing resistance to clindamycin, erythromycin and SXT⁸⁷. Detection of these clones in referral hospitals in Kenya calls for implementation of strict infection control measures to reduce the high morbidities and mortalities associated with HA-MRSA infections.

Tanzania

In a study to investigate the molecular epidemiology of trimethoprim resistance in MSSA causing skin and soft tissues infections, *dfrG* was detected in all 32-trimethoprim resistant isolates. Other reported trimethoprim resistance mechanisms such as *dfrA*, *dfrB* and *dfrK* were missing, confirming *dfrG* as the main trimethoprim resistance mechanism in Sub-Sahara Africa⁵⁵.

Uganda

A MRSA carriage of 56.1% (23/41) was detected in milk from pastoral communities in Uganda, exactly 70% of which were tetracycline-resistant. MRSA clones ST97 and ST1 were identified. Furthermore, over 90% of the isolates carried genes encoding enterotoxin that causes food-borne diseases. The weak veterinary delivery system and the high dependency on animals and animal products for food in Uganda was implicated for the high prevalence of MRSA ¹²¹.

S. aureus isolates, including 24 MRSA and 40 MSSA, were isolated from patients with surgical site infections (SSI). The MRSA isolates were MDR (including resistance to oxacillin, gentamicin, ciprofloxacin and chloramphenicol) compared to the MSSA. Inducible clindamycin resistance was found in 17.2% of the isolates, mostly in MRSA. In a multivariate analysis, inducible clindamycin resistance and cancer were identified as independent predictors of MRSA-SSI ¹²².

Southern Africa: Angola, Malawi, Mozambique, Namibia, South Africa

Angola

Conceição et al (2014) reported a nasal *S. aureus* carriage of 23.7% (n=128 isolates), out of which 58.1% (n=77 isolates) were MRSA. Fifty-seven of the MRSA clones were of ST5, followed by ST88 (n=9), ST8 (n=5) and ST72 (n=3). This study represents the first description of the spread of MRSA ST5 in Africa. All the 77 MRSA strains were resistant to SXT, cefoxitin (FOX) and PEN ¹²³. In a study to identify oxacillin-susceptible *mecA*-positive *S. aureus* (OS-MRSA) for the first time in Africa, a prevalence of 17.7% was detected among healthy healthcare workers in Angola and São Tomé & Príncipe, making them potential OS-MRSA reservoirs ¹²⁴. OS-MRSA have been reported worldwide in humans, animals and food animals ^{125–128}. The OS-MRSA isolates expressed MDR (Table 2) and belonged to ST88 (n=15 isolates) and ST8 (n=9 isolates). In sub-Saharan Africa, the identification of clinically important *S. aureus* is heavily based on phenotypic agar-screening and oxacillin disc-diffusion methods.

Mozambique

The prevalence of HA-MRSA and CA-MRSA in Mozambique was found to be 15.1% and 1%, respectively. MRSA showed high-level resistance to penicillin, cefoxitin, gentamicin, ciprofloxacin, erythromycin, SXT, chloramphenicol and tetracycline, compared to MSSA. Additionally, inducible macrolide–lincosamide–streptogramin B (MLSB) resistance was 41.7% and 10.7% in hospital-acquired *S. aureus* (HA-SA) and community-acquired *S. aureus* (CA-SA) isolates respectively¹²⁹, further limiting therapeutic options for *S. aureus* infections. This study, which is the first to detect the emergence of HA-MRSA within post-operative abdominal wounds and burn wounds in Mozambique, reported that patients with infected burn wounds had a significantly longer hospitalization than patients with post-operated abdominal wounds. Efforts to prevent the transmission of MDR HA-SA, such as education on proper hand-washing techniques, are urgently needed.

Namibia

The dominant resistance gene mediating trimethoprim resistance in MRSA and MSSA in Namibia was *dfrG*. This is similar to reports in other Africa countries⁵⁵. Moreover, *dfrG* was frequently detected in *S. aureus* from SSTIs in travelers returning from other African countries, suggesting that *dfrG* can be transmitted into populations with low antifolate resistance such as North America and Europe^{130,131}.

South Africa

Thirty MDR *S. aureus* were recovered between April 2015 to April 2016 from ten beaches in the Eastern Cape Province, South Africa (Table 2). Notably, the isolates harbored *mecA*, *femA*, *rpoB*, *blaZ*, *erm*(B) and *tet*(M)¹¹, making marine environments and public beaches potential depositaries of MDR *S. aureus* that can be transmitted to animals and humans. Further, the 50% resistance to vancomycin recorded is concerning to global health due to its role as a last-resort antibiotic for treating MRSA infections. *S. aureus* was detected in raw and pasteurized milk at an isolation rate of 75% and 29% respectively, due to inefficient thermal processing and post-process contamination. A high proportion (60%-100%) of these isolates showed resistance to aminoglycosides, β -lactams, vancomycin, tetracycline and erythromycin, albeit only 19 *mecA* genes were present²¹. Evidently, raw and pasteurized milk can harbour MDR *S. aureus*, exposing consumers to colonization and/or infections. Again, *Staphylococcus spp.*, including *S.*

aureus, *S. haemolyticus*, *S. xylosus* and *S. capitis* were isolated from healthy pigs and cattle, of which between 75 to 100% were resistant to penicillin G, tetracycline, sulfamethoxazole and nalidixic acids, due to their use as growth promoters; *mecA* and *mphC* were identified. Additionally, 12% of the isolates were resistant to vancomycin and erythromycin, evincing the important role of animals in the dissemination of resistance determinants and the importance of commensals to public health ⁶¹.

Van Rensburg et al. ⁵⁹ detected 43.4% (1432/3298 isolates) and 3.1% (328/10448 isolates) rifampicin resistance rate among MRSA and MSSA respectively. Similar studies in South Africa have also reported of high rifampicin resistance in MRSA ^{132,133}, obviously due to frequent use of rifampicin among tuberculosis patients, who are highly prevalent in South Africa. MRSA ST5 and ST612 were detected while H481Y/N and I527M mutations in *rpoB* were associated with high-level rifampicin resistance, similar to reports in Italy ¹³⁴. Additionally, novel H481N, I527M, K579R mutations were also detected.

Three studies reported a prevalence of 29.1% ¹³⁵, 45.44% ⁶⁰ and 100% ¹³⁶ MRSA recovered from humans, expressing resistance to macrolides, tetracycline, aminoglycoside, cotrimoxazole and rifampicin. MRSA ST612, ST239, ST36 and ST5 were the dominant strains similar to other findings in Australia and Europe¹³⁷. The study showed that *S. aureus* bacteremia is common and account for high mortality in South Africa. For instance, in a study by Perovic et al., ¹³⁵ 202 patients died from *S. aureus* bacteremia infections, with HIV patients being more likely to acquire HA-MRSA. The isolates were however susceptible to glycopeptides, fluoroquinolones, linezolid, tigecycline, fosfomycin and fusidic acid, confirming their clinical usefulness in treating MRSA infections. In a recent study, a high prevalence and genetic diversity of multi-drug efflux (MDE) resistance genes were found in clinical *S. aureus* isolates, including 81 MRSA and 16 MSSA ¹³⁸. *norA*, *norB*, *mepA*, *tet(38)*, *sepA*, *mdeA*, *imrs* and *sdrM* were present in at least 86% of the isolates, predicting resistance to broad-spectrum biocides and fluoroquinolones, which is disturbing. Efforts to develop efflux pump inhibitors can mitigate such resistance mechanisms.

Sao Tome & Principe

MRSA prevalence of 26.9%¹³⁹ and 25.5%¹¹⁴ was reported in nasal swabs in 2014 and 2015, respectively, in Sao Tome & Principe. Additionally, a high prevalence of oxacillin-susceptible *mecA*-positive *S. aureus* was reported in the same study in Sao Tome & Principe and Angola¹²⁴. The most dominant MRSA clone was ST8 (n=25 isolates), followed by ST5 (n=13 isolates) and ST80 (n=13 isolates). High genetic variability was found in the MSSA strains. Both MRSA and MSSA showed different levels of resistance to SXT, ERY, CIP and TET; however, all the MRSA isolates were resistant to ceftiofur.

Streptococcus spp. (S. pyogenes, S. pneumoniae and S. agalactiae)

Drug resistant *Streptococcus spp.* including *S. agalactiae* and *S. pyogenes* have been identified in Northern, Eastern and Southern Africa. *S. pyogenes* were reported in only humans whilst *S. agalactiae* was reported in both animals (camels) and humans with a high rate of resistance to tetracycline and erythromycin.

North Africa: Algeria, Egypt, Morocco, Tunisia, Libya

Algeria

A sole study has so far detected 44 tetracycline (100%, 44/44 isolates)- and erythromycin-resistant (43.18%, 19/44 isolates) *S. agalactiae* from vaginal swabs; *tet(M)*; and *erm(B)* respectively mediated this resistance. A high diversity of resistant clones viz., ST1, ST19, ST10, ST158, ST166, ST233, ST460, ST521 and ST677 were detected⁴⁵, which have been reported worldwide for causing life-threatening invasive diseases such as meningitis and sepsis^{140,141}.

Egypt

Similarly, Shabayek et al. (2014) detected 98% and between 14-17% *S. agalactiae* resistance to tetracycline and macrolides respectively. *tet(M)* was detected in all the 98 tetracycline-resistant isolates whilst *erm(B)* and *erm(A)* mediated erythromycin resistance. Efflux pump genes such as *tet(K)* (n=12 isolates), *tet(L)* (n=1 isolates) and *mefA/E* (n=1 isolates) were also found³², which reflects the increasing reports of *S. agalactiae* resistance to tetracycline and macrolides¹⁴². This study also showed that

vancomycin and fluoroquinolones are effective replacement for erythromycin and clindamycin, and for patients allergic to penicillin. Although penicillin is the antibiotic of choice for treating *S. agalactiae* infections, reports of penicillin resistance in USA and China calls for increased surveillance in Africa ¹⁴².

Tunisia

S. agalactiae

From January 2007 to December 2009, 226 *S. agalactiae* were isolated from female genitals and gastric fluid of infected newborns. Of these, 97.35% (220/226 isolates), 40% (90/226 isolates) and 19.1% (43/226 isolates) were resistant to tetracycline, erythromycin and rifampicin respectively. Additionally, seven isolates were resistant to aminoglycoside (gentamycin and streptomycin) and chloramphenicol. *tet(M)* (n=205 isolates), encoding a ribosomal protection protein, which protect the ribosome from the action of tetracycline, was the main tetracycline resistance mechanism, and was significantly associated with Tn916 (p-value = 0.0002). Other resistance genes including *erm(B)* (n=79 isolates) and *tet(O)* (n=50 isolates) were detected. All isolates were however susceptible to β -lactams and quinupristin-dalfopristin ⁸⁵. Between 2005 and 2007, 160 erythromycin-resistant *S. agalactiae* were isolated from humans, with a high resistance rate of 84.3% (135/160 isolates) to the constitutive macrolides-lincosamides, streptogramins B (MLSB) ¹⁴³.

S. pyogenes

Hraoui *et al.*, (2011) reported a low macrolide resistance rate (5%, 5/103) and a high tetracycline resistance rate (70%, 72/103) among human isolates, with *tet(M)*, associated with Tn916, being responsible for tetracycline resistance ¹⁴⁴. Increase tetracycline use in food animals was implicated in this instance, leading to selection and dissemination of resistance genes from animals to human. Macrolide resistance was only detected in seven isolates, which is corroborated by the findings of Ksia *et al.* (2010), who detected low-level macrolides resistance among Children ¹⁴⁵.

East Africa: Kenya, Tanzania

Kenya

S. agalactiae

In the horn of Africa, camel plays a significant role in the survival of humans by providing milk, meat and transportation. In 2013, Fischer et al. detected 36% (37/92) tetracycline resistance in *S. agalactiae* isolates from camels' wound infections and mastitis that was mainly mediated by a Tn916-borne *tet*(M). ST616 (n=22) was the major resistant clone, followed by ST612 and ST617¹⁴⁶. Shifting from tetracycline to other antibiotics is evidently necessary for effective treatment outcomes in camel infections in Kenya.

Southern Africa: Angola, Malawi, Mozambique, Namibia, South Africa

South Africa

S. agalactiae

A *S. agalactiae* colonization rate of 30.9% was detected from vaginal and rectal swabs of pregnant women. Similar to other reports in Africa, a high rate of tetracycline (94.5%, 120/128 isolates) and macrolide (21.1%, 27/128) resistance was documented. All the isolates were however sensitive to penicillin, ampicillin, vancomycin and gentamicin. Macrolide and clindamycin resistance were associated with *erm*(B) and *mefA* genes¹⁴⁷. The study highlights the need for research on treatment options for patients allergic to penicillin due to high-level resistance in alternative drugs such as macrolides and lincosamides.

Enterococcus spp. (E. faecium, E. faecalis, E. hirae, E. durans, E. gallinarum)

North Africa: Algeria, Egypt, Morocco, Tunisia, Libya

Algeria

The first study to molecularly characterize *Enterococcus spp.* from urinary tract and wound infections in Algeria revealed a high rate of resistance to erythromycin (86.4%, 108/125 isolates), tetracycline (82.4, 103/125 isolates), levofloxacin (71.2%, 89/125 isolates) and gentamicin (54.4, 68/125 isolates). Only

3.2% (4/125 isolates) were VRE, confirming glycopeptides as ideal antibiotics for treating Enterococcus infections. A mortality rate of 10% was reported due to infections caused by Enterococcus. *E. faecium*, *E. faecalis* and *E. gallinarum* were the main Enterococcus isolated. Majority of these isolates were from females (53%). *erm*(B) (≥ 92) and *vanCI*(≥ 4) were the main mechanisms of resistance. A high genetic diversity among strains was seen in *E. faecium* and *E. faecalis*, with *E. faecium* ST78 being the dominant resistant strain¹⁴⁸, which is also prevalent in Asian (Japan, Taiwan, China and Korea) and European (Italy and Germany) countries^{149–151}. A novel ST317 (n=33) clone was predominant among the *E. faecalis* isolates. Rational use of antibiotics, as well as close monitoring of the epidemiology of the strains are crucial.

Egypt

In a similar study to characterize *E. faecium* and *E. faecalis* from patients, 82% of the isolates were MDR, showing high-level resistance to aminoglycosides, β -lactams and tetracycline. *VanA* was detected in two *E. faecium* isolates, all of which were resistant to all antibiotics tested. Bioinformatic (sequence) analysis revealed that *vanA* was transmitted horizontally to *S. aureus*, showing the importance of horizontal gene transfer in ABR and subsequent management of enterococci infections such as bacteremia, endocarditis and urinary tract infections¹⁵².

Tunisia

Antimicrobial-resistant Enterococcus was found in faeces of pet and camel, irrigation water from farm environments, food vegetables, hospital environments, animal meat and patients in Tunisia^{19,22,31,51,52,69}. High-level resistance to vancomycin, macrolides, aminoglycosides, β -lactams and tetracycline was detected in the environment, animals and humans with majority of the isolates being *E. faecium*, followed by *E. faecalis*. *tet*(M), *tet*(L), *erm*(B), *ant* (6)-*la*, *vanA* and *aph*(3')-IIIa were the major resistance mechanisms, with *IS16* being the main MGE disseminating the resistance genes. *E. faecium* ST80, ST910 and ST16 were the dominant resistant clones in Tunisia. The studies show that meat, animals, pets,

hospital environment and wastewater used for farm irrigation play a crucial role in the spread of antibiotic resistant *Enterococcus*.

West Africa: Cape Verde, Ghana, Nigeria, Senegal

Nigeria

Enterococcus spp. isolated from poultry and cattle as well as their manure demonstrated high-level resistance to tetracycline, erythromycin, gentamicin, ampicillin and streptomycin. Sixty isolates were MDR, showing resistance to three or more antimicrobials¹⁵³. The rate of MDR is a reflection of the substantial use of broad-spectrum antibiotics in Nigeria, raising major public health concerns as practices such as the use of untreated poultry and cattle manure for fertilizing agricultural soils, particularly vegetables, are a common practice in Africa. This could transfer MDR *Enterococci* to humans, and cause serious nosocomial infections including endocarditis, bacteremia and urinary tract infections that can result in high morbidities and mortalities.

Ngbede et al. (2017) recently characterized 63 ampicillin- and 37 gentamicin-resistant *E. faecium* from vegetables, soil, farms, animal and manure²⁵. Approximately 95% (35/37 isolates) and 8% (5/63 isolates) of the aminoglycoside- and ampicillin-resistant clones were recognized as high-level aminoglycosides- and ampicillin-resistant *E. faecium* respectively. Modifying enzymes' genes such as *aac(6')-Ie-aph(2'')-Ia*), *aph(2')-Ic*, *aph(3')-IIIa*., and *ant(4')-Ia* accounted for the aminoglycoside resistance.

East Africa: Kenya and Tanzania

Tanzania

In a study to determine if cattle co-grazing with wild life influence ABR, ABR in wild animals such as buffalo, zebra and wildebeest was higher than in cattle, although wildlife is periodically treated with antibiotics. Ten VRE and ampicillin-resistant *Enterococcus* were found in the wild animals but not cattle. Additionally, *Enterococcus* isolates from wildlife were highly resistant to tetracycline, rifampicin, macrolides, aminoglycosides and cotrimoxazole¹⁴. *tet(W)* and *sulI* were the resistance genes identified in the isolates. The practice of co-grazing possibly resulted in transmission of ABR genes from livestock

to wildlife. The high presence of ABR bacteria in wildlife was likely due to contact with more environmental surfaces that have been contaminated with human, birds or animal excreta. Result from this study demonstrates the presence of ABR Enterococci in wild animals without antibiotic pressure.

Southern Africa: Angola, Malawi, Mozambique, Namibia, South Africa

South Africa

Multiple antibiotic-resistant Enterococci were isolated from borehole water, waste water, pigs and humans in South Africa. Notably, a very high-level vancomycin, aminoglycoside, β -lactam, macrolides and fluoroquinolones resistance was detected among the Enterococci isolates compared to other countries. *erm(B)* (≥ 300 isolates), *vanC* 2/3 (162 isolates), *vanB* (≥ 138 isolates), *vanC* (≥ 120 isolates), *strA* (≥ 120 isolates) were the major resistance genes. The vancomycin-resistant isolates were from patients with haematological malignancies, bacteremia, pigs, wastewater and underground water^{9,10,26,30}. Inefficient chlorination to kill bacteria accounted for the high resistance rates in the final effluents' discharge into the environment. Hospital wastewater is therefore a major source of MDR Enterococcus. Sub-therapeutic antibiotic usage in animal feed also accounted for the emergence of ABR in pigs whilst the construction of boreholes near pit toilets resulted in high enterococcal isolation and resistance rates in South Africa.

Experimental procedures used in included studies

The studies included in this review basically used the following experimental procedures. Transport media such as stuart agar, cary-blair medium, and gel transport swabs with charcoal were used to transport the samples to the laboratory^{53,65}. Cotton swabs were used to swab sample specimens, tissues, surfaces, fluids, etc. and cultured on nutrient agar, blood agar, tryptone soya agar, mannitol salt-phenol red agar, brain-heart infusion broth, Slanetz-Bartley mannitol salt agar, and Edwards agar media prior to identifying the 24-hour colonies using Gram-staining and different biochemical tests such as catalase and coagulase tests, latex coagulase test and DNase agar test. Subsequently, antimicrobial susceptibility testing (AST) using disc diffusion (Kirby-Bauer method or E-test) on Mueller Hinton agar plates and a 0.5 McFarland bacterial inoculum was performed. Antibiotics such as ampicillin (AMP), amoxicillin

(AMX), amikacin (AMK), ampicillin-Sulbactam (SAM), amoxicillin-clavulanic acid (AMC), azithromycin (AZI), apramycin (APR), chloramphenicol (CHL), cefoxitin (FOX), ceftazidime (CFZ), clarithromycin (CLR), ciprofloxacin (CIP), cefuroxime (CXM), clindamycin (CLI), cephalexin (LEX), cefoperazone (CFP), cefepime (FEP), cefotaxime (CTX), ceftaroline (CPT), cephalothin (CET), cloxacillin (CLX), doxycycline (DOX), erythromycin (ERY), fusidic acid (FUS), fosfomycin (Fof), gatifloxacin (GAT), gentamicin (GEN), imipenem (IPM), kanamycin (KAN), levofloxacin (LVX), linezolid (LZD), lincomycin (LIN), meropenem (MER), mupirocin (MUP), minocycline (MIC), moxifloxacin (MXF), methicillin (MET), metronidazole (MTZ), nitrofurantoin (NIT), norfloxacin (Nor), nalidixic acid (NAL), netilmicin (NEL), oxacillin (OXA), ofloxacin (OFX), perfloxacin (PF), penicillin (PEN), pristnamycin (PRI), rifampicin (RIF), streptomycin (STR), streptogramin B (SB), sulfamethoxazole (SMZ), tetracycline (TET), teicoplanin (TEC), telithromycin (TEL), tobramycin (TOB), trimethoprim-sulfamethoxazole (SXT), and vancomycin (VAN) were mostly used for the AST. Polymerase chain reaction (PCR) was used to detect the antimicrobial resistance genes and clones (i.e. molecular typing) of the isolates.

3. CONCLUSION AND STUDY LIMITATIONS

We report of high rate of ABR among GPB in several African countries, mediated largely by *S. aureus* ST5, ST8, and ST80, *Enterococcus faecium* and *Enterococcus faecalis* strains, *SCCmec*, *Tn916* and *IS16* MGEs are a major threat to clinical medicine, the economy and socio-economic development. This calls for national as well as international rules and regulations to contain resistance. Heavy consumption of antibiotics in animal feed, exchange of resistance genes between animals and food animal products to man, uncontrolled and inappropriate antibiotics prescription practices, inadequate hygienic handling and processing of food, close contact with pet dogs, shedding of resistant clones from animals to humans and the environment, as well as high consumption of antibiotics in humans, particularly in HIV patients, account for the high rate of ABR in Africa.

Effective surveillance and monitoring of antimicrobial drug usage and licensing, banning or restricting the prescription of reserved, expired and substandard drugs, periodic monitoring of pharmacies and veterinary shops, and antibiotic stewardship are recommended measures to contain ABR. Improving animal health through hygienic practices on farms, avoiding prophylactic or growth-promoting antibiotic usage in veterinary medicine, integrative efforts between human and veterinary medicine as well as environmental health are urgently needed to contain ABR. Implementation of these policies will decrease the high rate of ABR in Africa, reduce longer hospital stays and the resort to expensive but toxic antibiotic alternatives, with a concomitant reduction in morbidity and mortality rates. Few studies reporting on the molecular determinants of ABR in GPB in Africa limited the study to 130 articles. Among these, only few studies reported on MGEs and resistant clones.

Role of Funding Source: Not applicable.

Contributors: JOS conceived, designed and supervised the study, analysed and vetted the results, wrote the paper, edited and formatted it for publication. EM co-conceived and co-designed the study, gathered and analysed the data and drafted the paper. Both authors approved the final version for submission.

Funding: None

Declaration of interests: The authors declare no conflict of interest.

Acknowledgments: None

References

1. World Health Organization. Antimicrobial resistance—global report on surveillance. Geneva, Switzerland. *WHO* 2014.
2. Frean J, Perovic O, Fensham V, McCarthy K, von Gottberg A de GL et al. External quality assessment of national public health laboratories in Africa, 2002-2009. *Bull WHO* 2012; **90**:

- 191–199A. Available at: doi:10.2471/BLT.11.091876.
3. Osei Sekyere J. Current State of Resistance to Antibiotics of Last-Resort in South Africa: a Review From a Public Health Perspective. *Front Public Heal* 2016; **4**: 209. Available at: <https://doi.org/10.3389/fpubh.2016.00209>.
4. Williams PC, Isaacs D BJ. Antimicrobial resistance among children in sub-Saharan Africa. *Lancet Infect Dis* 2017; **18**(2): Pages e33-e44. Available at: [https://doi.org/10.1016/S1473-3099\(17\)30467-X](https://doi.org/10.1016/S1473-3099(17)30467-X).
5. Liu Y-Y, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. . *Lancet Infect Dis* 2016; **16**: 161–8. Available at: doi:10.1016/S1473-3099(15)00424-7.
6. World Health Organization. WHO’s first global report on antibiotic resistance reveals serious, worldwide threat to public health. In Antimicrobial resistance–global surveillance report. Virtual Press Conference 2014 Apr. *WHO* 2014; (Vol. 30). Available at: <http://www.who.int/mediacentre/news/releases/2014/amr-report/en/>.
7. Witte W. ‘Antibiotic resistance in gram-positive bacteria: epidemiological aspects.’ *J Antimicrob Chemother* 1999; **44**, su: 1–9. Available at: https://doi.org/10.1093/jac/44.suppl_1.1.
8. Seale AC, Mwaniki M, Newton CR BJ. Maternal and early onset neonatal bacterial sepsis: burden and strategies for prevention in sub-Saharan Africa. *Lancet Infect Dis* 2009; **9**(7): 428–38. Available at: [https://doi.org/10.1016/S1473-3099\(09\)70172-0](https://doi.org/10.1016/S1473-3099(09)70172-0).
9. Ateba CN, Lekoma KP KD. ‘Detection of vanA and vanB genes in vancomycin-resistant enterococci (VRE) from groundwater using multiplex PCR analysis.’ *J Water Health* 2013; **11**:4: 684–691.
10. Iweriebor, Benson Chuks, Sisipho Gaqavu, Larry Chikwelu Obi, Uchechukwu U. Nwodo and AIO. ‘Antibiotic susceptibilities of Enterococcus species isolated from hospital and domestic wastewater effluents in Alice, eastern Cape Province of South Africa.’ 4231-4246. *Int J Environ Res Public Health* **12**, no. 4. Available at: <https://doi.org/10.3390/ijerph120404231>.

11. Akanbi OE, Njom HA, Fri J, Otigbu AC CA. Antimicrobial Susceptibility of Staphylococcus aureus Isolated from Recreational Waters and Beach Sand in Eastern Cape Province of South Africa. *Int J Environ Res public Heal* 2017; **14**(9): 1001. Available at: doi:10.3390/ijerph14091001.
12. Al-ashmawy MA, Sallam KI, Abd-elghany SM. Prevalence, Molecular Characterization, and Antimicrobial Susceptibility of Methicillin-Resistant Staphylococcus aureus Isolated from Milk and Dairy Products. *FOODBORNE Pathog Dis* 2016; **13**: 156–62. Available at: <https://doi.org/10.1089/fpd.2015.2038>.
13. Iweriebor BC, Obi LC OA. Virulence and antimicrobial resistance factors of Enterococcus spp. isolated from fecal samples from piggery farms in Eastern Cape, South Africa. *BMC Microbiol* 2015; **15**(1): 136. Available at: <https://doi.org/10.1186/s12866-015-0468-7>.
14. Katakweba AA, Møller KS, Muumba J, Muhairwa AP, Damborg P, Rosenkrantz JT, Minga UM, Mtambo MM OJ. Antimicrobial resistance in faecal samples from buffalo, wildebeest and zebra grazing together with and without cattle in Tanzania. *J Appl Microbiol* 2015; **118**(4): 966–75. Available at: <https://doi.org/10.1111/jam.12738>.
15. Osei Sekyere J. Types and selling practices of antibiotics in veterinary shops in ashanti region, GHANA. *Int J Food, Agric Vet Sci* 2014; **4**(2): 87–96. Available at: <http://www.cibtech.org/jfav.htm>.
16. Osei Sekyere, John and FA. ‘Prevalence of Multidrug Resistance among Salmonella enterica Serovar Typhimurium Isolated from Pig Faeces in Ashanti Region, Ghana.’ *Int J Antibiot* 2015. Available at: <http://dx.doi.org/10.1155/2015/898790>.
17. Osei Sekyere J. ‘Antibiotic types and handling practices in disease management among pig farms in Ashanti Region, Ghana.’ *J Vet Med* 2014. Available at: <http://dx.doi.org/10.1155/2014/531952>.
18. Said MB, Abbassi MS, Gómez P, Ruiz-Ripa L, Sghaier S, Ibrahim C, Torres C HA. Staphylococcus aureus isolated from wastewater treatment plants in Tunisia: occurrence of human and animal associated lineages. *J water Heal* 2017; **15**(4): 638–43.

19. Ben Said L, Klibi N, Dziri R, Borgo F, Boudabous A, Ben Slama K TC. ‘Prevalence, antimicrobial resistance and genetic lineages of *Enterococcus* spp. from vegetable food, soil and irrigation water in farm environments in Tunisia.’ *J Sci Food Agric* 2016; **96(5)**: 1627–33. Available at: <https://doi.org/10.1002/jsfa.7264>.
20. Dziri R, Lozano C, Said LB, Bellaaj R, Boudabous A, Slama KB, Torres C KN. Multidrug-resistant enterococci in the hospital environment: detection of novel vancomycin-resistant *E. faecium* clone ST910. *J Infect Dev Countries* 2016; **10(08)**: 799–806. Available at: doi: <https://doi.org/10.3855/jidc.8014>.
21. Akindolire MA, Babalola OO AC. ‘Detection of antibiotic resistant *Staphylococcus aureus* from milk: A public health implication.’ *Int J Environ Res Public Health* 2015; **12**, no. **9**: 10254–10275. Available at: <https://doi.org/10.3390/ijerph120910254>.
22. Ben Said L, Dziri R, Sassi N, Lozano C, Ben Slama K, Ouzari I, Torres C KN. Species distribution, antibiotic resistance and virulence traits in canine and feline enterococci in Tunisia. *Acta Vet Hungarica* 2017; **65(2)**: 173–84. Available at: <https://doi.org/10.1556/004.2017.018>.
23. El-Hamid MI BM. ‘Comparative phenotypic and genotypic discrimination of methicillin resistant and susceptible *Staphylococcus aureus* in Egypt.’ *Cell Mol Biol* 2015; **61**, no. **4**: 101–112. Available at: <http://www.cellmolbiol.com>.
24. Mariem BJ, Ito T, Zhang M, *et al.* Molecular characterization of methicillin-resistant Pantone-valentine leukocidin positive *staphylococcus aureus* clones disseminating in Tunisian hospitals and in the community. *BMC* 2013; **12**: 2. Available at: <https://doi.org/10.1186/1471-2180-13-2>.
25. Ngbede EO, Raji MA, Kwanashie CN, Kwaga JK, Adikwu AA, Maurice NA AA. Characterization of high level ampicillin-and aminoglycoside-resistant enterococci isolated from non-hospital sources. *J Med Microbiol* 2017; **10;66(7)**: 1027–32. Available at: doi: [10.1099/jmm.0.000518](https://doi.org/10.1099/jmm.0.000518).
26. Iweriebor BC, Obi LC OA. Virulence and antimicrobial resistance factors of *Enterococcus* spp. isolated from fecal samples from piggy farms in Eastern Cape, South Africa. *BMC Microbiol* 2015; **15**: 136. Available at: <https://doi.org/10.1186/s12866-015-0468-7>.

27. Oumokhtar B, Elazhari M, Timinouni M, Bendahhou K, Bennani B, Mahmoud M, El Ouali Lalami A, Berrada S, Arrayhani M SHT. Staphylococcus aureus nasal carriage in a Moroccan dialysis center and isolates characterization. *Hemodial Int* 2013; **2**: 542–7. Available at: <https://doi.org/10.1111/j.1542-4758.2012.00759.x>.
28. El Bayomi RM, Ahmed HA, Awadallah MA, Mohsen RA, Abd El-Ghafar AE AM. Occurrence, Virulence Factors, Antimicrobial Resistance, and Genotyping of Staphylococcus aureus Strains Isolated from Chicken Products and Humans. *VECTOR-BORNE ZOONOTIC Dis* 2016; **16**: 157–64. Available at: <https://doi.org/10.1089/vbz.2015.1891>.
29. Hashem RA, Yassin AS, Zedan HH, Amin MA. Fluoroquinolone resistant mechanisms in methicillin-resistant Staphylococcus aureus clinical isolates in Cairo , Egypt. *J Infect Dev Ctries* 2013; **7**, no. **11**: 796–803. Available at: <https://doi.org/10.3855/jidc.3105> .
30. Lochan H, Moodley C, Rip D, Bamford C, Hendricks M, Davidson A EB. ‘Emergence of vancomycin-resistant Enterococcus at a tertiary paediatric hospital in South Africa.’ *SAMJ South African Med J* 2016; **106**, no. **6**: 562–566. Available at: [10.7196/SAMJ.2016.v106i6.10858](https://doi.org/10.7196/SAMJ.2016.v106i6.10858).
31. Said LB, Klibi N, Lozano C, Dziri R, Slama KB, Boudabous A TC. ‘Diversity of enterococcal species and characterization of high-level aminoglycoside resistant enterococci of samples of wastewater and surface water in Tunisia.’ . *Sci Total Environ* 2015; **530**: 11–7. Available at: <https://doi.org/10.1016/j.scitotenv.2015.05.091>.
32. Shabayek S, Abdalla S. Macrolide- and tetracycline-resistance determinants of colonizing group B streptococcus in women in Egypt. *J Med Microbiol* 2014; **63**: 1324–7. Available at: doi: [10.1099/jmm.0.077057-0](https://doi.org/10.1099/jmm.0.077057-0) .
33. Mbelle NM, Maningi NE, Tshisevhe V, Modipane L, Amoako DG, Osei Sekyere J. Draft Genome Sequence of a Clinical Enterococcus faecium Sequence Type 18 Strain from South Africa. *Genome Announc* 2017; **5**: e01381-17. Available at: <https://doi.org/10.1128/genomeA.01381-17>.
34. Malachowa N DF. ‘Mobile genetic elements of Staphylococcus aureus.’ *Cell Mol life Sci* 2010; **67.18**: 3057–3071. Available at: DOI [10.1007/s00018-010-0389-4](https://doi.org/10.1007/s00018-010-0389-4).

- 931 35. Roberts AP MP. 'Tn916-like genetic elements: a diverse group of modular mobile elements
932 conferring antibiotic resistance.' *FEMS Microbiol Rev* 2011; **35**, no. **5**: 856–871. Available at:
933 <https://doi.org/10.1111/j.1574-6976.2011.00283.x>.
- 934 36. Frost LS, Leplae R, Summers AO TA. 'Mobile genetic elements: the agents of open source
935 evolution.' *Nat Rev Microbiol* 2005; **3**, no. **9**: 722–732. Available at: doi:10.1038/nrmicro1235.
- 936 37. Thomas CM NK. 'Mechanisms of, and barriers to, horizontal gene transfer between
937 bacteria.' (2005): . *Nat Rev Microbiol* 2005; **3**, no. **9**: 711–21. Available at:
938 doi:10.1038/nrmicro1234.
- 939 38. Smillie C, Garcillán-Barcia MP, Francia MV, Rocha EP de la CF. 'Mobility of plasmids.'
940 *Microbiol Mol Biol Rev* 2010; **74**, no. **3**: 434–452.
- 941 39. Bernabé KJ, Langendorf C, Ford N, Ronat JB MR. Antimicrobial resistance in West Africa:
942 a systematic review and meta-analysis. *Int J Antimicrob agents* 2017; **50(5)**: 629–39. Available
943 at: <https://doi.org/10.1016/j.ijantimicag.2017.07.002>.
- 944 40. Tadesse BT, Ashley EA, Ongarello S, *et al*. Antimicrobial resistance in Africa□: a
945 systematic review. *BMC Infect Dis* 2017; **17:616**: 1–17. Available at:
946 <https://doi.org/10.1186/s12879-017-2713-1>.
- 947 41. Workneh M, Katz MJ, Lamorde M, Cosgrove SE MY. Antimicrobial Resistance of Sterile
948 Site Infections in Sub-Saharan Africa: A Systematic Review. *Open forum Infect Dis*: ofx209).
949 US: Oxford University Press. Available at: <https://doi.org/10.1093/ofid/ofx209>.
- 950 42. Schaumburg F, Alabi AS, Peters G BK. New epidemiology of Staphylococcus aureus
951 infection in Africa. *Clinical. Microbiol Infect* 2014; **20(7)**: 589–96. Available at:
952 <https://doi.org/10.1111/1469-0691.12690%0A>.
- 953 43. Nejad SB, Allegranzi B, Syed SB, Ellis B PD. Health-care-associated infection in Africa: a
954 systematic review. *Bull World Heal Organ* 2011; **89(10)**: 757–65. Available at:
955 doi:10.2471/BLT.11.088179.
- 956 44. Abdulgader SM, Shittu AO, Nicol MP KM. Molecular epidemiology of Methicillin-resistant
957 Staphylococcus aureus in Africa: a systematic review. *Front Microbiol* 2015; **6**: 348. Available

at: <https://doi.org/10.3389/fmicb.2015.00348>.

45. Bergal A, Loucif L, Benouareth DE, Bentorki AA, Abat C RJ. Molecular epidemiology and distribution of serotypes, genotypes, and antibiotic resistance genes of *Streptococcus agalactiae* clinical isolates from Guelma, Algeria and Marseille, France. *European Journal of Clinical Microbiology & Infectious Dis* 2015; **34**, no. **12**: 2339–48. Available at: DOI 10.1007/s10096-015-2487-6.

46. Djoudi F, Benallaoua S, Aleo A, Touati A, Challal M, Bonura C MC. Descriptive Epidemiology of Nasal Carriage of *Staphylococcus aureus* and Methicillin-Resistant *Staphylococcus aureus* Among Patients Admitted at Two Healthcare Facilities in Algeria. *Microb Drug Resist* 2014; **21**(2): 218–23. Available at: <https://doi.org/10.1089/mdr.2014.0156>.

47. Ouchenane Z, Agabou A, Smati F, Rolain JM RD. Spread of methicillin-resistant *Staphylococcus aureus* between the community and the hospitals in Asian countries: an ANSORP study. *Journal of antimicrobial chemotherapy. Pathol Biol* 2013; **61**: 280–1. Available at: <https://doi.org/10.1016/j.patbio.2013.05.006>.

48. Djahmi N, Boutet-Dubois A, Nedjai S, Dekhil M, Sotto A LJ. Molecular epidemiology of *Enterococcus* sp . isolated in a university hospital in Algeria. *Scand J Infect Dis* 2012; **2011**: 656–62. Available at: <https://doi.org/10.3109/00365548.2012.673232>.

49. Conceição T, Coelho C, de Lencastre H A-SM. Frequent occurrence of oxacillin-susceptible *mecA* -positive *Staphylococcus aureus* (OS-MRSA) strains in two African countries . *J Antimicrob Chemother* 2015; 2015; **70**: 3200–4. Available at: <https://doi.org/10.1093/jac/dkv261>.

50. Elhani D, Gharsa H, Kalai D, Lozano C, Gómez P, Boutheina J, Aouni M, Barguellil F, Torres C SK. ‘Clonal lineages detected amongst tetracycline-resistant meticillin-resistant *Staphylococcus aureus* isolates of a Tunisian hospital, with detection of lineage ST398.’ *J Med Microbiol* 2015; **64**, no. **6**: 623–629. Available at: doi: 10.1099/jmm.0.000066 .

51. Klibi N, Said LB, Jouini A, Slama KB, López M, Sallem RB, Boudabous A TC. ‘Species distribution, antibiotic resistance and virulence traits in enterococci from meat in Tunisia.’ (2013): . *Meat Sci* 2013; **93**, no. **3**: 675–80. Available at:

986 <https://doi.org/10.1016/j.meatsci.2012.11.020>.

987 52. Klibi N, Lagha AB, Slama KB, Boudabous A TC. 'Faecal enterococci from camels in
988 Tunisia: species, antibiotic resistance and virulent genes.' *Vet Rec* 2013; **172**, : 213–213.
989 Available at: doi: 10.1136/vr.100910.

990 53. Abdelmoein KA, ElHariri M SA. Methicillin-Resistant *Staphylococcus aureus*: An
991 Emerging Pathogen of Pets in Egypt with a Public Health Burden. *Transbound Emerg Dis* 2012;
992 **59**: 331–5. Available at: <https://doi.org/10.1111/j.1865-1682.2011.01273.x>.

993 54. Fowoyo PT, Ogunbanwo ST. Antimicrobial resistance in coagulase negative staphylococci
994 from Nigerian traditional fermented foods. *Ann Clin Microbiol Antimicrob* 2017; **16**:4: 1–7.
995 Available at: <https://doi.org/10.1186/s12941-017-0181-5>.

996 55. Nurjadi D, Olalekan AO, Layer F, Shittu AO, Alabi A, Ghebremedhin B, Schaumburg F,
997 Hofmann-Eifler J, Van Genderen PJ, Caumes E FR. Emergence of trimethoprim resistance gene
998 *dfrG* in *Staphylococcus aureus* causing human infection and colonization in sub-Saharan Africa
999 and its import to Europe. *J Antimicrob Chemother* 2014; **27**: 2361–8. Available at:
1000 <https://doi.org/10.1093/jac/dku174>.

1001 56. O'Malley SM, Emele FE, Nwaokorie FO, Idika N, Umezudike AK, Emeka-Nwabunnia I,
1002 Hanson BM, Nair R, Wardyn SE ST. Molecular typing of antibiotic-resistant *Staphylococcus*
1003 *aureus* in Nigeria. *J Infect Public Health* 2015; **8**: 187–93. Available at:
1004 <http://dx.doi.org/10.1016/j.jiph.2014.08.001>.

1005 57. Shittu A, Oyedara O, Abegunrin F, Okon K, Raji A, Taiwo S, Ogunsola F, Onyedibe K EG.
1006 Characterization of methicillin-susceptible and -resistant staphylococci in the clinical setting: a
1007 multicentre study in Nigeria. *BMC Infect Dis* 2012; **12**(1): 286. Available at:
1008 [https://doi.org/10.1016/S0732-8893\(98\)00118-7](https://doi.org/10.1016/S0732-8893(98)00118-7).

1009 58. Vitali LA, Petrelli D, Lamikanra A, Prenna M, Akinkunmi EO. Diversity of antibiotic
1010 resistance genes and staphylococcal cassette chromosome *mec* elements in faecal isolates of
1011 coagulase-negative staphylococci from Nigeria. *BMC Microbiol* 2014; **14**:106. Available at:
1012 <https://doi.org/10.1186/1471-2180-14-106>.

- 1013 59. van Rensburg MJ, Whitelaw AC EB. Genetic basis of rifampicin resistance in methicillin-
1014 resistant *Staphylococcus aureus* suggests clonal expansion in hospitals in Cape. *BMC Microbiol*
1015 2012; **12**: 46. Available at: <https://doi.org/10.1186/1471-2180-12-46>.
- 1016 60. Perovic O, Iyaloo S, Kularatne R, Lowman W. Prevalence and Trends of *Staphylococcus*
1017 *aureus* Bacteraemia in Hospitalized Patients in South Africa , 2010 to 2012□: Laboratory- Based
1018 Surveillance Mapping of Antimicrobial Resistance and Molecular Epidemiology. *PLoS One*
1019 2015: 1–14.
- 1020 61. Adegoke AA, Okoh AI. Species diversity and antibiotic resistance properties of
1021 *Staphylococcus* of farm animal origin in Nkonkobe Municipality , South Africa. *FoliaMicrobiol*
1022 2014; **59**: 133–40. Available at: DOI 10.1007/s12223-013-0275-1.
- 1023 62. WHO. Tackling Antibiotic Resistance from a Food Safety Perspective in Europe. *WHO-*
1024 *Europe, Denmark* 2011: 1–88. Available at: <http://www.euro.who.int/.../e94889.pdf>.
- 1025 63. O'Neill J. 'Tackling drug-resistant infections globally: final report and recommendations.'
1026 *Rev Antimicrob Resist* 2016.
- 1027 64. Cavaco LM AF. 'Resistance in bacteria of the food chain: epidemiology and control
1028 strategies.' *Microb Drug Resist Futur Med Ltd, Brastislava, Slovakia* 2013: 136–158. Available
1029 at: <https://doi.org/10.2217/ebo.12.361>.
- 1030 65. Boeck H De, Vandendriessche S, Hallin M, Batoko B, Alworonga J. *Staphylococcus aureus*
1031 nasal carriage among healthcare workers in Kisangani , the Democratic Republic of the Congo.
1032 *Eur J Clin Microbiol Infect Dis* 2015; **34(8)**: 1567–72. Available at: DOI 10.1007/s10096-015-
1033 2387-9.
- 1034 66. Fall C, Seck A, Richard V, Ndour M, Sembene M, Laurent F BS. Epidemiology of
1035 *Staphylococcus aureus* in Pigs and Farmers in the Largest Farm in Dakar , Senegal. *FOOD*
1036 *BORNE Pathog Dis* 2012; **9**: 962–5. Available at: <https://doi.org/10.1089/fpd.2012.1197>.
- 1037 67. Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H MF. 'Meticillin-
1038 resistant *Staphylococcus aureus* (MRSA): global epidemiology and harmonisation of typing
1039 methods.' *Int J Antimicrob Agents* 2012; **39**, no. **4**: 273–282. Available at:

1040 <https://doi.org/10.1016/j.ijantimicag.2011.09.030>.

1041 68. Song JH, Hsueh PR, Chung DR, Ko KS, Kang CI, Peck KR et al. Spread of methicillin-
1042 resistant *Staphylococcus aureus* between the community and the hospitals in Asian. *J Antimicrob*
1043 *Chemother* 2011; **66**: 1061–9. Available at: <https://doi.org/10.1093/jac/dkr024>.

1044 69. Elhani D, Klibi N, Dziri R, Hassan MB, Mohamed SA, Said LB, Mahjoub A, Slama KB,
1045 Jemli B, Bellaj R BF. ‘vanA-containing *E. faecium* isolates of clonal complex CC17 in clinical
1046 and environmental samples in a Tunisian hospital.’ *Diagn Microbiol Infect Dis* 2014; **79**, no. **1**:
1047 60–63. Available at: <http://dx.doi.org/10.1016/j.diagmicrobio.2014.01.011>.

1048 70. Dziri R, Lozano C, Said LB, Bellaaj R, Boudabous A, Slama KB, Torres C KN. ‘Multidrug-
1049 resistant enterococci in the hospital environment: detection of novel vancomycin-resistant *E.*
1050 *faecium* clone ST910.’ *J Infect Dev Ctries* 2016; **10**, no. **8**: 799–806. Available at: doi:
1051 <https://doi.org/10.3855/jidc.8014>.

1052 71. Udo EE, Pearman JW GW. Genetic analysis of community isolates of methicillin-resistant
1053 *Staphylococcus aureus* in Western Australia. *J Hosp Infect* 1993; **25**: 97–108. Available at:
1054 [https://doi.org/10.1016/0195-6701\(93\)90100-E%0A](https://doi.org/10.1016/0195-6701(93)90100-E%0A).

1055 72. Salgado CD, Farr BM CD. Community-acquired methicillin-resistant *Staphylococcus aureus*:
1056 a meta-analysis of prevalence and risk factors. *Clin Infect Dis* 2003, **36**: 131–139.
1057 Available at: <https://doi.org/10.1086/345436>.

1058 73. Hiramatsu K, Okuma K, Ma XX, Yamamoto M, Hori S et al: New trends in *Staphylococcus*
1059 *aureus* infections: glycopeptide resistance in hospital and methicillin resistance in the
1060 community. *Curr Opin Infect Dis* 2002, **15**: 407–413. Available at:
1061 doi:10.1097/00132980-200201060-0000.

1062 74. Chambers H. The changing epidemiology of *Staphylococcus aureus*? *Emerg Infect Dis* 2001;
1063 **7**: 178–182.

1064 75. Diekema DJ, Pfaller MA, Schmitz FJ, Smayevsky J, Bell J et al. Survey of infections due to
1065 *Staphylococcus* species: frequency of occurrence and antimicrobial susceptibility of isolates
1066 collected in the United States, Canada, Latin America, Europe, and the Western Pacific region

- 1067 for the SENTRY Antimicrobial Surveillanc. *Clin Infect Dis* 2001; **32 Suppl 2**: S114–132.
1068 Available at: <https://doi.org/10.1086/320184>.
- 1069 76. NNIS. National Nosocomial Infections Surveillance (NNIS) System Report, data summary
1070 from January 1992 through June 2004. *Am J Infect Control* 2004; **2**: 470–485.
- 1071 77. Katayama Y, Ito T HK. A new class of genetic element, staphylococcus cassette
1072 chromosome mec, encodes methicillin resistance in Staphylococcus aureus. *Antimicrob agents*
1073 *Chemother* 2000; **44(6)**: 1549–55. Available at: doi: 10.1128/AAC.44.6.1549-1555.2000.
- 1074 78. Akinkunmi E LA. Species distribution and antibiotic resistance in coagulase-negative
1075 staphylococci colonizing the gastrointestinal tract of children in Ile-Ife, Nigeria. *Trop J Pharm*
1076 *Res* 2010; **9(1)**: 35–43. Available at: <http://dx.doi.org/10.4314/tjpr.v9i1.52033>.
- 1077 79. Archer GL CM. Antimicrobial susceptibility of coagulase-negative staphylococci.
1078 *Antimicrob Agents Chemother* 1994; **38**: 2231–2237.
- 1079 80. Hashem YA, Yassin AS AM. Molecular characterization of Enterococcus spp. clinical
1080 isolates from Cairo, Egypt. *Indian J Med Microbiol* 2015; **33(5)**: p.80. Available at: doi:
1081 10.4103/0255-0857.148836.
- 1082 81. Hraoui M, Boubaker IB, Doloy A, Redjeb SB BA. Molecular Mechanisms of Tetracycline
1083 and Macrolide Resistance and emm Characterization of Streptococcus pyogenes Isolates in
1084 Tunisia. *Microb Drug Resist* 2011; **17(3)**: 377–82. Available at:
1085 <https://doi.org/10.1089/mdr.2010.0160>.
- 1086 82. Rachdi M, Boubaker IB, Moalla S, *et al*. Phenotypic and genotypic characterization of
1087 macrolide resistant Streptococcus pneumoniae in Tunisia. *Pathol Biol* 2008; **56**: 125–9.
1088 Available at: <https://doi.org/10.1016/j.patbio.2007.05.005>.
- 1089 83. Wolter N, Gottberg A Von, Gouveia L De, Klugman KP. Molecular basis and clonal nature
1090 of increasing pneumococcal macrolide resistance in South Africa , 2000 – 2005. *Int J Antimicrob*
1091 *Agents* 2008; **32**: 2000–5. Available at: <https://doi.org/10.1016/j.ijantimicag.2008.01.013>.
- 1092 84. Hancock RE. Mechanisms of action of newer antibiotics for Gram-positive pathogens.
1093 *Lancet Infect Dis* 2005; **5**: 209–18. Available at: [https://doi.org/10.1016/S1473-3099\(05\)70051-](https://doi.org/10.1016/S1473-3099(05)70051-)

- 1094 7.
- 1095 85. Hraoui M, Boubaker IB, Rachdi M, Slim A RS. Macrolide and tetracycline resistance in
1096 clinical strains of Streptococcus agalactiae isolated in Tunisia. *J Med Microbiol* 2017; **61**: 1109–
1097 13. Available at: doi: 10.1099/jmm.0.037853-0 .
- 1098 86. Fischer A, Liljander A, Kaspar H, Muriuki C, Fuxelius HH, Bongcam-Rudloff E, de Villiers
1099 EP, Huber CA, Frey J, Daubenberger C BR. Camel Streptococcus agalactiae populations are
1100 associated with specific disease complexes and acquired the tetracycline resistance gene tetM via
1101 a Tn 916 -like element Camel Streptococcus agalactiae populations are associated with specific
1102 disease complex. *Vet Res* 2013, 2013; **44**:86. Available at: [https://doi.org/10.1186/1297-9716-](https://doi.org/10.1186/1297-9716-44-86)
1103 44-86.
- 1104 87. Omuse G, Zyl KN Van, Hoek K, Abdulgader S, Kariuki S, Whitelaw A. Molecular
1105 characterization of Staphylococcus aureus isolates from various healthcare institutions in Nairobi
1106 , Kenya: a cross sectional study. *Ann Clin Microbiol Antimicrob* 2016; **15**:51: 1–9. Available
1107 at: <https://doi.org/10.1186/s12941-016-0171-z>.
- 1108 88. Stokes HW GM. ‘Gene flow, mobile genetic elements and the recruitment of antibiotic
1109 resistance genes into Gram-negative pathogens.’ *FEMS Microbiol Rev* 2011; **35**, no. **5**: 790–819.
1110 Available at: <https://doi.org/10.1111/j.1574-6976.2011.00273.x>.
- 1111 89. Gogarten JP TJ. ‘Horizontal gene transfer, genome innovation and evolution.’ *Nat Rev*
1112 *Microbiol* 2005; **3**, no. **9**: 679–687. Available at: doi:10.1038/nrmicro1204.
- 1113 90. Bouchami O, Hassen AB, De Lencastre H MM. High prevalence of mec complex C and ccrC
1114 is independent of SCC mec type V in Staphylococcus haemolyticus. *Eur J Clin Microbiol Infect*
1115 *Dis* 2012; **31**: 605–14. Available at: 10.1007/s10096-011-1354-3.
- 1116 91. Khalil W, Hashwa F, Shihabi A TS. Methicillin-resistant Staphylococcus aureus ST80-IV
1117 clone in children from Jordan.. *Diagn Microbiol Infect Dis* 2012; **73**: 228–230. Available at:
1118 <https://doi.org/10.1016/j.diagmicrobio.2012.03.012>0AGet .
- 1119 92. Udo EE SE. The dissemination of ST80-SCCmec-IV community-associated methicillin
1120 resistant Staphylococcus aureus clone in Kuwait hospitals. *Ann Clin Microbiol Antimicrob* 2010;

- 1121 9: 1–7. Available at: <https://doi.org/10.1186/1476-0711-9-31>.
- 1122 93. Ahmed EF, Gad GF, Abdalla AM, Hasaneen AM AS. Prevalence of Methicillin Resistant
1123 Staphylococcus aureus among Egyptian Patients after Surgical interventions. *Surg Infect*
1124 (*Larchmt*) 2014; **15**: 404–11. Available at: <https://doi.org/10.1089/sur.2013.212>.
- 1125 94. Osman K, Alvarez-Ordóñez A, Ruiz L, Badr J, ElHofy F, Al-Maary KS, Moussa IM,
1126 Hessain AM, Orabi A, Saad A EM. Antimicrobial resistance and virulence characterization of
1127 Staphylococcus aureus and coagulase-negative staphylococci from imported beef meat. *Ann Clin*
1128 *Microbiol Antimicrob* 2017; **16(1)**: 35. Available at: <https://doi.org/10.1186/s12941-017-0210-4>.
- 1129 95. Perveen I, Majid A, Knawal S, Naz I, Sehar S, Ahmed S RM. Biochemical characters and
1130 antibiotic susceptibility of S. aureus isolates. *Asian Pac J Trop Biomed* 2011; **1**: 212–216.
1131 Available at: doi: 10.1016/S2221-1691(11)60029-4.
- 1132 96. Perveen, I., Majid, A., Knawal, S., Naz, I., Sehar, S. A, S. and Raza MA. Prevalence and
1133 antimicrobial susceptibility pattern of methicillin-resistant Staphylococcus aureus and
1134 coagulasenegative staphylococci in Rawalpindi, Pakistan. *Br J Med Med Res* 2013; **3**: 198–209.
1135 Available at: doi: 10.9734/%0ABJMMR/2013/2109.
- 1136 97. Dziri R, Klibi N, Lozano C, *et al.* High prevalence of Staphylococcus haemolyticus and
1137 Staphylococcus sapro- phyticus in environmental samples of a Tunisian hospital. *Diagn*
1138 *Microbiol Infect Dis* 2016; **85.2**: 136–140. Available at:
1139 <http://dx.doi.org/10.1016/j.diagmicrobio.2016.03.006>.
- 1140 98. Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA ER, Lemons JA, Donovan EF,
1141 Stark AR, Tyson JE, Oh W BC, Korones SB, Shankaran S, Laptook AR, Stevenson DK, Papile
1142 LA P, WK. Late-onset sepsis in very low birth weight neonates: the experience of the NICHD
1143 Neonatal Research Network. *J Pediatr* 2002; **110**: 285–291. Available at:
1144 doi:10.1542/peds.110.2.285.
- 1145 99. Gharsa H, Slama KB, Lozano C, Gómez-Sanz E, Klibi N, Sallem RB, Gómez P, Zarazaga
1146 M, Boudabous A TC. Prevalence, antibiotic resistance, virulence traits and genetic lineages of
1147 Staphylococcus aureus in healthy sheep in Tunisia. *Veterinary microbiol. Vet Microbiol* 2012;
1148 **156(3)**: pp.367-373. Available at: <https://doi.org/10.1016/j.vetmic.2011.11.009>.

- 1149 100. Cui S, Li J, Hu C, Jin S, Li F, Guo Y, Ran LU MY. Isolation and characterization of
1150 methicillin-resistant *Staphylococcus aureus* from swine and workers in China. *J Antimicrob*
1151 *Chemother* 2009; **64**: 680–683. Available at: <https://doi.org/10.1093/jac/dkp275>.
- 1152 101. Jiménez JN, Vélez LA, Mediavilla JR, Ocampo AM, Vanegas JM, Rodríguez EA,
1153 Kreiswirth BN CM. Livestock-associated methicillin-susceptible *Staphylococcus aureus* ST398
1154 infection in woman, Colombia. *Emerg Infect Dis* 2011; **17**: 1970–1971. Available at:
1155 [doi:10.3201/eid1710.110638](https://doi.org/10.3201/eid1710.110638).
- 1156 102. Khanna T, Friendship R, Dewey C WJ. Methicillin resistant *Staphylococcus aureus*
1157 colonization in pigs and pig farmers. *Vet Microbiol* 2008; **128**: 298–303. Available at:
1158 <https://doi.org/10.1016/j.vetmic.2007.10.006>.
- 1159 103. Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, Capuano AW,
1160 Herwaldt LA DD. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present
1161 in midwestern U.S. swine and swine workers. *PLoS One* 2009; **4**,: e4258.
- 1162 104. Maalej SM, Malbruny B, Leclercq R HA. Emergence of *Staphylococcus aureus* strains
1163 resistant to pristinamycin in Sfax (Tunisia). *Pathol Biol* 2012; **60**: e71–4. Available at:
1164 <http://dx.doi.org/10.1016/j.patbio.2011.10.012>.
- 1165 105. Bouchami O, Hassen AB, De Lencastre H MM. High prevalence of mec complex C and
1166 ccrC is independent of SCCmec type V in *Staphylococcus haemolyticus*. *Eur J Clin Microbiol*
1167 *Infect Dis* 2012; **31**(4): 605–614. Available at: DOI 10.1007/s10096-011-1354-3.
- 1168 106. Tabe Y, Nakamura A, Oguri T IJ. Molecular characterization of epidemic multiresistant
1169 *Staphylococcus haemolyticus* isolates. *Diagn Microbiol Infect Dis* 1998; **32**: 177–183. Available
1170 at: [https://doi.org/10.1016/S0732-8893\(98\)00118-7](https://doi.org/10.1016/S0732-8893(98)00118-7).
- 1171 107. Santos Sanches I, Mato R, de Lencastre H, Tomasz A C, Collaborators C and the I. Patterns
1172 of multidrug resistance among methicillin-resistant hospital isolates of coagulase-positive and
1173 coagulase-negative staphylococci collected in the international multicenter study RESIST in
1174 1997 and 1998. *Microb Drug Resist* 2000; **6**: 199–211. Available at:
1175 <https://doi.org/10.1089/mdr.2000.6.199>.

- 1176 108. Al-haddad OH, Zorgani A GK. Nasal Carriage of Multi-Drug Resistant Pantone-Valentine
1177 Leucocidin-Positive Methicillin-Resistant Staphylococcus aureus in Children in Tripoli-Libya.
1178 2014; **90**: 724–7. Available at: doi: <https://doi.org/10.4269/ajtmh.13-0715> .
- 1179 109. Egyir B, Guardabassi L, Sørnum M, Nielsen SS, Kolekang A, Frimpong E, Addo KK,
1180 Newman MJ LA. Molecular epidemiology and antimicrobial susceptibility of clinical
1181 Staphylococcus aureus from healthcare institutions in Ghana. *PLoS One*, 2014; **9**(2): p.e89716.
- 1182 110. Egyir B, Guardabassi L, Monecke S, Kwasi K, Jemima M, Rhod A. Resistance Methicillin-
1183 resistant Staphylococcus aureus strains from Ghana include USA300. *J Glob Antimicrob Resist*
1184 2015; **3**: 26–30. Available at: <http://dx.doi.org/10.1016/j.jgar.2014.11.006>.
- 1185 111. Bouchami O, Ben Hassen A, de Lencastre H MM. High prevalence of mec complex C and
1186 ccrC is independent of SCCmec type V in Staphylococcus haemolyticus. *Eur J Clin Microbiol*
1187 *Infect Dis* 2012; **31**(4): 605–614. Available at: DOI 10.1007/s10096-011-1354-3.
- 1188 112. Shittu A, Oyedara O, Abegunrin F, Okon K, Raji A, Taiwo S, Ogunsola F, Onyedibe K EG.
1189 ‘Characterization of methicillin-susceptible and-resistant staphylococci in the clinical setting: a
1190 multicentre study in Nigeria.’ *BMC Infect Dis* **12**, no. **1**: 286. Available at:
1191 <https://doi.org/10.1186/1471-2334-12-286>.
- 1192 113. Naber KG, Schito G, Botto H, Palou J MT. Surveillance Study in Europe and Brazil on
1193 Clinical Aspects and Antimicrobial Resistance Epidemiology in Females with Cystitis (ARESC):
1194 Implications for Empiric Therapy. *Eur Urol* 2008; **54**: 1164–1178. Available at:
1195 <https://doi.org/10.1016/j.eururo.2008.05.010>.
- 1196 114. Conceição T, Coelho C, Silva IS, de Lencastre H A-SM. Staphylococcus aureus in former
1197 Portuguese colonies from Africa and the Far East: missing data to help fill the world map. *Clin*
1198 *Microbiol Infect* 2015; **21**(9): 842–e1. Available at: <https://doi.org/10.1016/j.cmi.2015.05.010>.
- 1199 115. Jensen SO LB. Genetics of antimicrobial resistance in Staphylococcus aureus. *Futur*
1200 *Microbiol* 2009; **4**: 565–82. Available at: <https://doi.org/10.2217/fmb.09.30>.
- 1201 116. Frey KM, Lombardo MN, Wright DL AA. Towards the understanding of resistance
1202 mechanisms in clinically isolated trimethoprim-resistant, methicillin-resistant Staphylococcus

1203 aureus dihydrofolate reductase. *J Struct Biol* 2010; **170**: 93–7. Available at:
1204 <https://doi.org/10.1016/j.jsb.2009.12.011>.

1205 117. Kadlec K, Fessler AT, Hauschild T SS. Novel and uncommon antimicrobial resistance
1206 genes in livestock-associated methicillin-resistant Staphylococcus aureus. *Clin Microbiol Infect*
1207 2012; **18**: 745–55. Available at: <https://doi.org/10.1111/j.1469-0691.2012.03842.x>.

1208 118. Badri M, Ehrlich R, Wood R MG. Initiating co-trimoxazole prophylaxis in HIV-infected
1209 patients in Africa: an evaluation of the provisional WHO/UNAIDS recommendations. *AIDS*
1210 2001; **15**(9): 1143–8. Available at: [doi:10.1097/00002030-200106150-00009](https://doi.org/10.1097/00002030-200106150-00009).

1211 119. Phaku P, Lebughe M, Strauß L, *et al*. Unveiling the molecular basis of antimicrobial
1212 resistance in Staphylococcus aureus from the Democratic Republic of the Congo using whole
1213 genome sequencing. *Clin Microbiol Infect* 2016. Available at:
1214 <http://dx.doi.org/10.1016/j.cmi.2016.04.009>.

1215 120. Vandendriessche S, De Boeck H, Deplano A, Phoba MF, Lunguya O, Falay D, Daully N,
1216 Verhaegen J, Denis O JJ. Characterisation of Staphylococcus aureus isolates from bloodstream
1217 infections, Democratic Republic of the Congo. *Eur J Clin Microbiol Infect Dis* 2017; **36**(7):
1218 1163–71. Available at: DOI 10.1007/s10096-017-2904-0.

1219 121. Asiimwe BB, Baldan R, Trovato A CD. Prevalence and molecular characteristics of
1220 Staphylococcus aureus, including methicillin resistant strains, isolated from bulk can milk and
1221 raw milk products in pastoral communities of South-West Uganda. *BMC Infect Dis* 2017; **17**(1):
1222 422. Available at: <https://doi.org/10.1186/s12879-017-2524-4>.

1223 122. Seni J, Bwanga F, Najjuka CF, Makobore P, Okee M, Mshana SE, Kidenya BR, Joloba ML
1224 KD. Molecular Characterization of Staphylococcus aureus from Patients with Surgical Site
1225 Infections at Mulago Hospital in Kampala, Uganda. *PLoS One* 2013; **8**: 1–7.

1226 123. Conceição T, Coelho C, Santos-Silva I, de Lencastre H A-SM. Epidemiology of methicillin-
1227 resistant and-susceptible Staphylococcus aureus in Luanda, Angola: first description of the
1228 spread of the MRSA ST5-IVa clone in the African continent. *Microb Drug Resist* 2014; **20**(5):
1229 441–9. Available at: <https://doi.org/10.1089/mdr.2014.0007>.

1230 124. Conceição T, Coelho C, de Lencastre H A-SM. Frequent occurrence of oxacillin-susceptible
1231 mecA-positive *Staphylococcus aureus* (OS-MRSA) strains in two African countries. *J*
1232 *Antimicrob Chemother* 2015; **70(12)**: 3200–4. Available at: <https://doi.org/10.1093/jac/dkv261>.

1233 125. Hososaka Y, Hanaki H EH et al. Characterization of oxacillinsusceptible mecA-positive
1234 *Staphylococcus aureus*: a new type of MRSA. *J Infect Chemother* 2007; **13**: 79–86. Available at:
1235 <https://doi.org/10.1007/s10156-006-0502-7>.

1236 126. Petinaki E, Kontos F MA. Emergence of two oxacillin-susceptible mecA-positive
1237 *Staphylococcus aureus* clones in a Greek hospital. *J Antimicrob Chemother* 2002; **50**: 1090–1.
1238 Available at: <https://doi.org/10.1093/jac/dkf235>.

1239 127. Pu W, Su Y LJ et al. High incidence of oxacillin-susceptible mecA-positive *Staphylococcus*
1240 *aureus* (OS-MRSA) associated with bovine mastitis in China. *PLoS One* 2014; **9**: e88134.

1241 128. Corrente M, Normanno G, Martella V, Bellacicco AL, Quaglia NC, Dambrosio A,
1242 Buonavoglia D, D'Abramo M BC. Comparison of methods for the detection of methicillin
1243 resistance in *Staphylococcus aureus* isolates from food products. *Lett Appl Microbiol* 2007; **45**:
1244 535–9. Available at: <https://doi.org/10.1111/j.1472-765X.2007.02226.x%0A>.

1245 129. Meeren BT, Millard PS, Scacchetti M, Hermans MH, Hilbink M, Concelho TB, Ferro JJ
1246 WP. Emergence of methicillin resistance and Panton-Valentine leukocidin positivity in hospital-
1247 and community-acquired *Staphylococcus aureus* infections in Beira , Mozambique. *Trop Med Int*
1248 *Heal Vol* 2013; **0**. Available at: <https://doi.org/10.1111/tmi.12221%0A>.

1249 130. De Angelis G, Cipriani M, Cauda R TE. Treatment of skin and soft tissue infections due to
1250 community-associated methicillin-resistant *Staphylococcus aureus* in Europe: the role of
1251 trimethoprim-sulfamethoxazole. *Clin Infect Dis author reply 2* 2011; **52**: 1471–2. Available at:
1252 <https://doi.org/10.1093/cid/cir247>.

1253 131. Chua K, Laurent F CG et al. Antimicrobial resistance: not community-associated
1254 methicillin-resistant *Staphylococcus aureus* (CA-MRSA)! A clinician's guide to community
1255 MRSA—its evolving antimicrobial resistance and implications for therapy. *Clin Infect Dis* 2011;
1256 **52**: 99–114. Available at: <https://doi.org/10.1093/cid/ciq067>.

1257 132. Marais E, Aithma N, Perovic O, Oosthuysen WF, Musenge E DA. Antimicrobial
1258 susceptibility of methicillin-resistant *Staphylococcus aureus* isolates from South Africa. . *SAMJ*
1259 2009; **99**: 170–3.

1260 133. Shittu AO LJ. Antimicrobial susceptibility patterns and characterization of clinical isolates
1261 of *Staphylococcus aureus* in KwaZulu-Natal province, South Africa. *BMC Infect Dis* 2006; **6**:
1262 125. Available at: <https://doi.org/10.1186/1471-2334-6-125>.

1263 134. Wichelhaus TA, Schafer V, Brade V BB. Molecular characterization of *rpoB* mutations
1264 conferring cross-resistance to rifamycins on methicillin-resistant *Staphylococcus aureus*.
1265 *Antimicrob Agents Chemother* 1999; **43**: 2813–2816.

1266 135. Perovic O, Singh-Moodley A, Govender NP, Kularatne R, Whitelaw A, Chibabhai V,
1267 Naicker P, Mbelle N, Lekalakala R, Quan V SC. A small proportion of community-associated
1268 methicillin-resistant *Staphylococcus aureus* bacteraemia, compared to healthcare-associated
1269 cases, in two South African provinces. *Eur J Clin Microbiol Infect Dis* 2017; **36(12)**: 2519–32.
1270 Available at: DOI 10.1007/s10096-017-3096-3.

1271 136. Amoako DG, Bester LA, Somboro AM, Baijnath S, Govind CN, Essack SY. Plasmid-
1272 mediated resistance and virulence mechanisms in the private health sector in KwaZulu-Natal ,
1273 South Africa□: An investigation of methicillin resistant *Staphylococcus aureus* (MRSA)
1274 clinical isolates col. *Int J Infect Dis* 2016; **46**: 38–41. Available at:
1275 <https://doi.org/10.1016/j.ijid.2016.03.019>.

1276 137. Sit PS, Teh CS, Idris N, Sam IC, Syed Omar SF SH, Thong KL, Kamarulzaman A PS.
1277 Prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) infection and the molecular
1278 characteristics of MRSA bacteraemia over a two-year period in a tertiary teaching hospital in
1279 Malaysia. *BMC Infect Dis* 2017; **17**: 274. Available at: [https://doi.org/10.1186/s12879-017-2384-](https://doi.org/10.1186/s12879-017-2384-y)
1280 [y](https://doi.org/10.1186/s12879-017-2384-y).

1281 138. Antiabong JF, Kock MM, Bellea NM EM. Diversity of Multidrug Efflux Genes and
1282 Phenotypic Evaluation of the In vitro Resistance Dynamics of Clinical *Staphylococcus Aureus*
1283 Isolates Using Methicillin; a Model β -lactam. *open Microbiol journal* 2017; **11**: 132–41.
1284 Available at: doi: 10.2174/1874285801711010132.

- 1285 139. Conceição T, Santos Silva I, de Lencastre H A-SM. Staphylococcus aureus nasal carriage
1286 among patients and health care workers in Sao Tome and Principe. *Microb Drug Resist* 2014;
1287 **20(1)**: 57–66. Available at: <https://doi.org/10.1089/mdr.2013.0136>.
- 1288 140. Sadowy E, Matynia B HW. Population structure, virulence factors and resistance
1289 determinants of invasive, noninvasive and colonizing Streptococcus agalactiae in Poland. *J*
1290 *Antimicrob Chemother* 2010; **65(9)**: 1907–1914.
- 1291 141. Bohnsack JF, Whiting A, GottschalkM, DunnDM, Weiss R A, PH, Philips JB 3rd, Weisman
1292 LE, Rhoads GG LF. Population structure of invasive and colonizing strains of Streptococcus
1293 agalactiae from neonates of six U.S. Academic Centers from 1995 to 1999. *J Clin Microbiol*
1294 2010; **46(4)**: 1285–1291.
- 1295 142. Verani, J. R., McGee, L., Schrag SJ. Prevention of perinatal group B streptococcal
1296 disease— revised guidelines from CDC, 2010. *MMWR Recomm Rep* 2010; **59 (RR-10)**: 1–36.
- 1297 143. Rachdi M, Boubaker IB, Hraoui M RS. High rates of macrolide resistance among clinical
1298 isolates of Streptococcus agalactiae in Tunisia. *Arch Inst Pasteur Tunis* 2010; **87(1/2)**: p.35.
- 1299 144. Hraoui M, Boubaker IB, Doloy A, Redjeb SB BA. ‘Molecular mechanisms of tetracycline
1300 and macrolide resistance and emm characterization of Streptococcus pyogenes isolates in
1301 Tunisia.’ *Microb Drug Resist* 2011; **17**, no. **3**: 377–382. Available at:
1302 <https://doi.org/10.1089/mdr.2010.0160>.
- 1303 145. Ksia S, Smaoui H, Hariga D KA. Biotypes and antimicrobial susceptibility of Streptococcus
1304 pyogenes strains isolated in children in Tunis. *Bull la Société Pathol Exot* 2010; **103(2)**: pp.69-
1305 74. Available at: <http://www.springerlink.com/.../>.
- 1306 146. Fischer A, Liljander A, Kaspar H, Muriuki C, Fuxelius HH, Bongcam-Rudloff E, de
1307 Villiers EP, Huber CA, Frey J, Daubenberger C BR. ‘Camel Streptococcus agalactiae
1308 populations are associated with specific disease complexes and acquired the tetracycline
1309 resistance gene tetM via a Tn 916-like element.’ *Vet Res* 2013; **44**, no. **1**: 86. Available at:
1310 <https://doi.org/10.1186/1297-9716-44-86>.
- 1311 147. Bolukaoto JY, Monyama CM, Chukwu MO, Lekala SM, Nchabeleng M, Maloba MR,

1312 Mavenyengwa RT, Lebelo SL, Monokoane ST, Tshepuwane C MS. Antibiotic resistance of
1313 *Streptococcus agalactiae* isolated from pregnant women in Garankuwa , South Africa. *BMC Res*
1314 *Notes* 2015: 6–12. Available at: <https://doi.org/10.1186/s13104-015-1328-0>.

1315 148. Djahmi N, Boutet-Dubois A, Nedjai S, Dekhil M, Sotto A LJ. Molecular epidemiology of
1316 *Enterococcus* sp. isolated in a university hospital in Algeria. *Scand J Infect Dis* 2012; **44(9)**,.:
1317 pp.656-662. Available at: <https://doi.org/10.3109/00365548.2012.673232>.

1318 149. Matsushima A, Takakura S, Yamamoto M, Matsumura Y, Shirano M, Nagao M, Ito Y,
1319 Inuma Y, Shimizu T, Fujita N IS. Regional spread and control of vancomycin-resistant
1320 *Enterococcus faecium* and *Enterococcus faecalis* in Kyoto, Japan. *Eur J Clin Microbiol Infect*
1321 *Dis* 2011. Available at: DOI 10.1007/s10096-011-1412-x.

1322 150. Fallico L, Boldrin C, Grossato A, Franchin E DCE, Tommasini T et al. Molecular
1323 epidemiology of *Enterococcus faecium* isolates from an Italian hospital. *Infection* 2011; **39**: 127
1324 – 33. Available at: DOI 10.1007/s15010-011-0086-8.

1325 151. Hsieh YC, Lee WS, Ou TY HP. Clonal spread of CC17 vancomycin-resistant *Enterococcus*
1326 *faecium* with multilocus sequence type 78 (ST78) and a novel ST444 in Taiwan. *Eur J Clin*
1327 *Microbiol Infect Dis* 2010; **29**: 25 – 30. Available at: DOI 10.1007/s10096-009-0810-9.

1328 153. Iweriebor, Benson C., Larry C. Obi and AIO. ‘Virulence and antimicrobial resistance
1329 factors of *Enterococcus* spp. isolated from fecal samples from piggery farms in Eastern Cape,
1330 South Africa.’ *BMC Microbiol* **15**, no. **1**: 136. Available at: [https://doi.org/10.1186/s12866-015-](https://doi.org/10.1186/s12866-015-0468-7)
1331 [0468-7](https://doi.org/10.1186/s12866-015-0468-7).

1332 154. Djahmi N, Messad N, Nedjai S, Moussaoui A, Mazouz D, Richard JL, Sotto A LJ.
1333 Molecular epidemiology of *Staphylococcus aureus* strains isolated from inpatients with infected
1334 diabetic foot ulcers in an Algerian University Hospital. *Clin Microbiol Infect* 2013; **19(9)**: E398–
1335 404. Available at: <https://doi.org/10.1111/1469-0691.12199>.

1336 155. Conceição T, Coelho C, Santos Silva I, de Lencastre H A-SM. Methicillin-resistant
1337 *Staphylococcus aureus* in the community in Luanda, Angola: blurred boundaries with the
1338 hospital setting. *Microb Drug Resist* 2016; **22**: 22–7. Available at:
1339 <https://doi.org/10.1089/mdr.2015.0236>.

- 1340 156. Abbassi MS, Znazen A, Mahjoubi F, Hammami A BA. Emergence of vancomycin-resistant
1341 Enterococcus faecium in Sfax: clinical features and molecular typing. *Med Mal Infect* 2007; **37**:
1342 240–1. Available at: doi: 10.1016/j.medmal.2006.09.011 .
- 1343 157. Abdelkareem MZ, Sayed M, Hassuna NA, Mahmoud MS AS. Multi-drug-resistant
1344 Enterococcus faecalis among Egyptian patients with urinary tract infection. *J Chemother* 2017;
1345 **29**: 74–82. Available at: <https://doi.org/10.1080/1120009X.2016.1182358>.
- 1346 158. Abouelfetouh A, Kassem M, Naguib M E-NM. Investigation and Treatment of Fusidic Acid
1347 Resistance Among Methicillin-Resistant Staphylococcal Isolates from Egypt. *Microb Drug*
1348 *Resist* 2017; **23**: 8–17. Available at: <https://doi.org/10.1089/mdr.2015.0336>.
- 1349 159. Bendary MM, Solyman SM, Azab MM, Mahmoud NF HA. Genetic diversity of multidrug
1350 resistant Staphylococcus aureus isolated from clinical and non clinical samples in Egypt. *Cell*
1351 *Mol Biol* 2016; **62**: 55–61. Available at: doi: 10.14715/cmb/2016.62.10.9.
- 1352 160. Bendary MM, Solyman SM, Azab MM, Mahmoud NF HA. Characterization of Methicillin
1353 Resistant Staphylococcus aureus isolated from human and animal samples in Egypt. *Cell Mol*
1354 *Biol* 2016; **62**: 94–100. Available at: doi: 10.14715/cmb/2016.62.2.16.
- 1355 161. El-Sharif A AH. Community-acquired methicillin-resistant Staphylococcus aureus (CA-
1356 MRSA) colonization and infection in intravenous and inhalational opiate drug abusers. *Exp Biol*
1357 *Med* 2008; **233**: 874–80. Available at: <https://doi.org/10.3181/0711-RM-294>.
- 1358 162. Okuda KV, Toepfner N, Alabi AS, Arnold B, Blard S, Falke U, Menschner L, Monecke S,
1359 Ruppelt-Lorz A BR. Molecular epidemiology of Staphylococcus aureus from Lambarene,
1360 Gabon. *Eur J Clin Microbiol Infect Dis* 2016; **35**: 1963–73. Available at: DOI 10.1007/s10096-
1361 016-2748-z.
- 1362 163. Egyir B, Guardabassi L, Nielsen SS, Larsen J, Addo KK, Newman MJ LA. Prevalence of
1363 nasal carriage and diversity of Staphylococcus aureus among inpatients and hospital staff at
1364 Korle Bu Teaching Hospital, Ghana. *J Glob Antimicrob Resist* 2013; **1**: 189–93. Available at:
1365 <https://doi.org/10.1016/j.jgar.2013.05.006>.
- 1366 164. Maina EK, Kiiyukia C, Wamae CN, Waiyaki PG KS. Characterization of methicillin-

- 1367 resistant *Staphylococcus aureus* from skin and soft tissue infections in patients in Nairobi,
1368 Kenya. *Int J Infect Dis* 2013; **17**: e115–9. Available at:
1369 <https://doi.org/10.1016/j.ijid.2012.09.006>.
- 1370 165. Al-haddad OH, Zorgani A GK. Nasal Carriage of Multi-Drug Resistant Pantone-Valentine
1371 Leucocidin-Positive Methicillin-Resistant *Staphylococcus aureus* in Children in Tripoli-Libya.
1372 *Am J Trop Med Hyg* 2014; **90**: 724–7. Available at: doi: <https://doi.org/10.4269/ajtmh.13-0715> .
- 1373 166. Zriouil SB, Bekkali M ZK. Epidemiology of *Staphylococcus aureus* infections and nasal
1374 carriage at the Ibn Rochd University hospital center, Casablanca, Morocco. *Brazilian J Infect Dis*
1375 *2012 May 1;16(3)*279-83 2012; **16**: 279–83. Available at: [https://doi.org/10.1016/S1413-](https://doi.org/10.1016/S1413-8670(12)70324-0)
1376 [8670\(12\)70324-0](https://doi.org/10.1016/S1413-8670(12)70324-0).
- 1377 167. Ayepola OO, Olasupo NA, Egwari LO, Becker K. Molecular Characterization and
1378 Antimicrobial Susceptibility of *Staphylococcus aureus* Isolates from Clinical Infection and
1379 Asymptomatic Carriers in Southwest Nigeria. *PLoS One* 2015; **2304**: 4–11. Available at:
1380 <http://dx.doi.org/10.1371/journal.pone.0137531>.
- 1381 168. Kolawole DO, Adeyanju A, Schaumburg F, Akinyoola AL, Lawal OO, Amusa YB, Köck R
1382 BK. Characterization of colonizing *Staphylococcus aureus* isolated from surgical wards' patients
1383 in a Nigerian university hospital. *PLoS One* 2013; **8(7)**: e68721.
- 1384 169. Shittu AO, Okon K, Adesida S, Oyedara O, Witte W, Strommenger B, Layer F NU.
1385 Antibiotic resistance and molecular epidemiology of *Staphylococcus aureus* in Nigeria. *BMC*
1386 *Microbiol* 2011; **11**: 92. Available at: <https://doi.org/10.1186/1471-2180-11-92>.
- 1387 170. Okon KO, Basset P, Uba A, Lin J, Oyawoye B, Shittu AO BD. Cooccurrence of
1388 predominant Pantone-Valentine leukocidin-positive sequence type (ST) 152 and multidrug-
1389 resistant ST 241 *Staphylococcus aureus* clones in Nigerian hospitals. *J Clin Microbiol* **47**: 3000–
1390 3. Available at: doi: [10.1128/JCM.01119-09](https://doi.org/10.1128/JCM.01119-09).
- 1391 171. Ghebremedhin B, Olugbosi MO, Raji AM, Layer F, Bakare RA, König B KW. Emergence
1392 of a community-associated methicillin-resistant *Staphylococcus aureus* strain with a unique
1393 resistance profile in Southwest Nigeria. *J Clin Microbiol* 2009; **47**: 2975–80. Available at: doi:
1394 [10.1128/JCM.00648-09](https://doi.org/10.1128/JCM.00648-09).

172. Mbelle NM, Maningi NE, Tshisevhe V, Modipane L, Amoako DG SJ. First Report of a Whole-Genome Shotgun Sequence of a Clinical *Enterococcus faecalis* Sequence Type 6 Strain from South Africa. *Genome Announc* 2017; **5(50)**: e01382-17. Available at: doi: 10.1128/genomeA.01382-17.
173. Mbelle NM, Maningi NE, Tshisevhe V, Modipane L, Amoako DG SJ. Draft Genome Sequence of a Clinical *Enterococcus faecium* Sequence Type 18 Strain from South Africa. *Genome Announc* 2017; **5(48)**: e01381-17. Available at: <https://doi.org/10.1128/genomeA.01381-17>.
175. Shittu AO, Udo EE LJ. Phenotypic and molecular characterization of *Staphylococcus aureus* isolates expressing low-and high-level mupirocin resistance in Nigeria and South Africa. *BMC Infect Dis* 2009; **9**: 10. Available at: <https://doi.org/10.1186/1471-2334-9-10>.
176. Shittu A, Lin J MD. Molecular identification and characterization of mannitol-negative methicillin-resistant *Staphylococcus aureus*. *Diagnostic Microbiol Infect Dis* **57(1)**: 93–5. Available at: <https://doi.org/10.1016/j.diagmicrobio.2006.05.004>.
177. Elhassan MM, Ozbak HA, Hemeg HA, Elmekki MA AL. Absence of the *mecA* gene in methicillin resistant *Staphylococcus aureus* isolated from different clinical specimens in shendi city, Sudan. *BioMed Res Int* 2015; 2015. Available at: <http://dx.doi.org/10.1155/2015/895860>.
179. Nejma MB, Mastouri M, Jrad BB NM. Characterization of ST80 Panton-Valentine leukocidin-positive community-acquired methicillin-resistant *Staphylococcus aureus* clone in Tunisia. *Diagnostic Microbiol Infect Dis* 2013; **77**: 20–4. Available at: <https://doi.org/10.1016/j.diagmicrobio.2008.02.010>.
180. Bouchami O, Achour W, Mekni MA, Rolo J HA. Antibiotic resistance and molecular characterization of clinical isolates of methicillin-resistant coagulase-negative staphylococci isolated from bacteremic patients in oncohematology. *Folia Microbiol* **56**: 122. Available at: DOI 10.1007/s12223-011-0017-1.
181. Slama KB, Gharsa H, Klibi N, Jouini A, Lozano C, Gómez-Sanz E, Zarazaga M, Boudabous A TC. Nasal carriage of *Staphylococcus aureus* in healthy humans with different levels of contact with animals in Tunisia: genetic lineages, methicillin resistance, and virulence

- 1423 factors. *Eur J Clin Microbiol Infect Dis* 2011; **30**: 499–508. Available at: DOI 10.1007/s10096-
1424 010-1109-6.
- 1425 182. Bouchami O, Achour W HA. ‘Prevalence of resistance phenotypes and genotypes to
1426 macrolide, lincosamide and streptogramin antibiotics in Gram-positive cocci isolated in Tunisian
1427 Bone Marrow Transplant Center.’ *Pathol Biol* 2011; **59**, no. **4**: 199–206. Available at:
1428 <https://doi.org/10.1016/j.patbio.2009.03.010>.
- 1429 183. Bouchami O, Achour W HA. Typing of staphylococcal cassette chromosome mec encoding
1430 methicillin resistance in *Staphylococcus aureus* strains isolated at the bone marrow transplant
1431 centre of Tunisia. *Curr Microbiol* 2009; **59**: 380–5. Available at: DOI 10.1007/s00284-009-
1432 9448-1.
- 1433 184. Klibi N, Ben Slama K, Sáenz Y, Masmoudi A, Zanetti S, Sechi LA, Boudabous A TC.
1434 Detection of virulence factors in high-level gentamicin-resistant *Enterococcus faecalis* and
1435 *Enterococcus faecium* isolates from a Tunisian hospital. *Can J Microbiol* 2007; **53**(3). Available
1436 at: <https://doi.org/10.1139/W06-136>.
- 1437 185. Achour W, Bouchami O, Galopin S, Leclercq R BHA. Analysis of pristinamycin-resistant
1438 *Staphylococcus epidermidis* isolates in the Tunisian Bone Marrow Transplant Center. *Lett Appl*
1439 *Microbiol* **46**: 358–63. Available at: <https://doi.org/10.1111/j.1472-765X.2007.02318.x>.
- 1440 186. Bouchami O, Achour W HA. Prevalence and mechanisms of macrolide resistance among
1441 *Staphylococcus epidermidis* isolates from neutropenic patients in Tunisia. *Clin Microbiol Infect*
1442 2007; **13**: 103–6. Available at: <https://doi.org/10.1111/j.1469-0691.2006.01567.x>.
- 1443 187. Seni J, Bwanga F, Najjuka CF, Makobore P, Okee M, Mshana SE, Kidenya BR, Joloba ML
1444 KD. ‘Molecular characterization of *Staphylococcus aureus* from patients with surgical site
1445 infections at Mulago Hospital in Kampala, Uganda.’ *PLoS One* 2013; **8**, no. **6**: e66153.
- 1446 188. Okee MS, Joloba ML, Okello M, Najjuka FC, Katabazi FA, Bwanga F, Nanteza A KD.
1447 Prevalence of virulence determinants in *Staphylococcus epidermidis* from ICU patients in
1448 Kampala, Uganda. *J Infect Dev Countries* 2011; **6**: 242–50. Available at:
1449 <https://doi.org/10.3855/jidc.2007>.

- 1450 189. Kitara LD, Anywar AD, Acullu D, Odongo-Aginya E, Aloyo J FM. Antibiotic
1451 susceptibility of Staphylococcus aureus in suppurative lesions in Lacor Hospital, Uganda.
1452 *African Heal Sci* **11**: 34–9. Available at: <http://dx.doi.org/10.4314/ahs.v11i3.70068>.
- 1453 190. Ojulong J, Mwambu TP, Joloba M, Bwanga F K-MD. Relative prevalence of methicilline
1454 resistant Staphylococcus aureus and its susceptibility pattern in Mulago Hospital, Kampala,
1455 Uganda. *Tanzania J Heal Res* 2009; **11**. Available at:
1456 <http://dx.doi.org/10.4314/thrb.v11i3.47703>.
- 1457 191. Martins E, Novais C, Freitas AR, Dias AR, Ribeiro TG, Antunes P PL. Filling the map for
1458 antimicrobial resistance in sub-Saharan Africa: ampicillin-resistant Enterococcus from non-
1459 clinical sources in Angola. *J Antimicrob Chemother* **70**: 2914–6. Available at:
1460 [doi:10.1093/jac/dkv172](https://doi.org/10.1093/jac/dkv172).
- 1461 192. Awad A, Ramadan H, Nasr S, Ateya A AS. Genetic Characterization Antimicrobial
1462 Resistance Patterns And Virulence Determinants of Staphylococcus aureus Isolated Form Bovine
1463 Mastitis. *Pak J Biol Sci* 2017; **20**: 298–305. Available at:
1464 <https://scialert.net/abstract/?doi=pjbs.2017.298.305> .
- 1465 193. Ammar AM, Attia AM, Abd EH, MI ES. IM and Abd El-Kader S. Genetic basis of
1466 resistance waves among methicillin resistant Staphylococcus aureus isolates recovered from milk
1467 and meat products in Egypt. *Cell Mol Biol Noisy-le-Grand* 2016; **62**: 7–15.
- 1468 194. Sallam KI, Abd-Elghany SM, Elhadidy M TT. Molecular characterization and antimicrobial
1469 resistance profile of methicillin-resistant Staphylococcus aureus in retail chicken. *J food Prot*
1470 2015; **78**: 1879–84. Available at: <https://doi.org/10.4315/0362-028X.JFP-15-150>.
- 1471 195. Fischer A, Liljander A, Kaspar H, Muriuki C, Fuxelius HH, Bongcam-Rudloff E, de
1472 Villiers EP, Huber CA, Frey J, Daubenberger C BR. Camel Streptococcus agalactiae populations
1473 are associated with specific disease complexes and acquired the tetracycline resistance gene tetM
1474 via a Tn 916-like element. *Vet Res* 2013; **44**(1): 86. Available at: [https://doi.org/10.1186/1297-](https://doi.org/10.1186/1297-9716-44-86)
1475 [9716-44-86](https://doi.org/10.1186/1297-9716-44-86).
- 1476 196. Ngbede EO, Raji MA, Kwanashie CN KJ. Antimicrobial resistance and virulence profile of
1477 enterococci isolated from poultry and cattle sources in Nigeria. *Trop Anim Heal Prod* 2017;

- 1478 **49(3)**: 451–8. Available at: DOI 10.1007/s11250-016-1212-5.
- 1479 197. Chah KF, Gómez-Sanz E, Nwanta JA, Asadu B, Agbo IC, Lozano C, Zarazaga M TC.
1480 Methicillin-resistant coagulase-negative staphylococci from healthy dogs in Nsukka, Nigeria.
1481 *Brazilian J Microbiol* 2014; **45**: 215–20. Available at: [http://dx.doi.org/10.1590/S1517-](http://dx.doi.org/10.1590/S1517-83822014005000034)
1482 83822014005000034.
- 1483 198. Iweriebor BC, Obi LC OA. Macrolide, glycopeptide resistance and virulence genes in
1484 Enterococcus species isolates from dairy cattle. *J Med Microbiol* 2016; **65**: 641–87.
- 1485 199. Soyegbe AO, Ajayi AO, Ngwenya E, Basson AK OA. Vancomycin and Oxacillin Co-
1486 Resistance of Commensal Staphylococci. *Jundishapur J Microbiol* 2014; **7**. Available at: doi:
1487 10.5812/jjm.9310.
- 1488 200. Freitas AR, Elghaieb H, León-Sampedro R, Abbassi MS, Novais C, Coque TM, Hassen A
1489 PL. Detection of optrA in the African continent (Tunisia) within a mosaic Enterococcus faecalis
1490 plasmid from urban wastewaters. *J Antimicrob Chemother* 2017; **72(12)**: 3245–51. Available at:
1491 <https://doi.org/10.1093/jac/dkx321>.
- 1492 201. Chairat S, Gharsa H, Lozano C, Gómez-Sanz E, Gómez P, Zarazaga M, Boudabous A,
1493 Torres C BSK. Characterization of Staphylococcus aureus from raw meat samples in Tunisia:
1494 detection of clonal lineage ST398 from the African continent. *Foodborne Pathog Dis* 2015; **12**:
1495 686–92. Available at: <https://doi.org/10.1089/fpd.2015.1958>.
- 1496 202. Gharsa H, Slama KB, Gómez-Sanz E, Lozano C, Zarazaga M, Messadi L, Boudabous A
1497 TC. Molecular characterization of Staphylococcus aureus from nasal samples of healthy farm
1498 animals and pets in Tunisia. *Vector-Borne Zoonotic Dis* 2015; **15**: 109–15. Available at:
1499 <https://doi.org/10.1089/vbz.2014.1655>.
- 1500 203. Gharsa H, Sallem RB, Slama KB, Gómez-Sanz E, Lozano C, Jouini A, Klibi N, Zarazaga
1501 M, Boudabous A TC. High diversity of genetic lineages and virulence genes in nasal
1502 Staphylococcus aureus isolates from donkeys destined to food consumption in Tunisia with
1503 predominance of the ruminant associated CC133 lineage. *BMC Vet Res* 2012; **8**: 203. Available
1504 at: <https://doi.org/10.1186/1746-6148-8-203>.

204. Akanbi OE, Njom HA, Fri J, Otigbu AC CA. Antimicrobial Susceptibility of Staphylococcus aureus Isolated from Recreational Waters and Beach Sand in Eastern Cape Province of South Africa. *Int J Environ Res public Heal* 2017; **14**(9): 1001. Available at: doi:10.3390/ijerph14091001.

205. Molale LG BC. Antibiotic resistance, efflux pump genes and virulence determinants in Enterococcus spp. from surface water systems. *Environ Sci Pollut Res* **23**: 21501–10. Available at: DOI 10.1007/s11356-016-7369-7.

206. Gharsa H, Dziri R, Klibi N, Chairat S, Lozano C, Torres C, Bellaaj R SK. Environmental Staphylococcus aureus contamination in a Tunisian hospital. *J Chemother* 2016; **28**: 506–9. Available at: <https://doi.org/10.1179/1973947815Y.00000000036>.

Table 1. Frequency distribution of Gram-positive bacterial species, resistance genes and MGEs isolated from animals, humans and environmental specimens.

Bacteria species, ARGs and MGEs		Human (n) ¹	Animal(n) ²	Environment(n) ³
Species	<i>E. faecalis</i>	225	129	66

¹ Total number of species or ARGs or MGEs in human isolates

² Total number of species or ARGs or MGEs in animal isolates

³ Total number of species or ARGs or MGEs in environmental isolates

	<i>E. faecium</i>	299	577	523
	<i>S. agalactiae</i>	658	92	0
	<i>S. aureus</i>	25559	1609	65
	<i>S. haemolyticus</i>	96	43	38
	<i>S. pyogenes</i>	296	0	0
ARGs	<i>mecA</i>	3057	462	28
	<i>erm(B)</i>	551	520	197
	<i>erm(C)</i>	102	23	8
	<i>tet(M)</i>	524	115	81
	<i>tet(K)</i>	179	80	22
	<i>tet(L)</i>	25	57	37
	<i>vanB</i>	4	387	60
	<i>vanA</i>	23	0	23
	<i>vanC1/2/3</i>	8	862	101
	<i>dfrA/G</i>	422	0	0
MGEs	<i>aph(3')-IIIa</i>	50	5	115
	<i>aac(6')-aph(2')</i>	178	17	73
	<i>ant(6)-Ia</i>	5	24	38
	<i>blaZ</i>	403	127	35
	<i>IS16</i>	3	0	5
	<i>SCCmec</i>	2471	27	8
	<i>Tn916</i>	62	37	0

1526 **Table 2. Geographical distribution, species, clones, and resistance mechanisms of antibiotic-resistant Gram-positive bacteria isolated from**
1527 **humans in Africa from 2007-2018**

Country (n) ⁴	Year	Organism/Species (n) ⁵	Specimen Sources (n) ⁶	Sample size (Resistant isolates)	Resistance rate (%)	Clones (n) ⁷	Resistance genes/mechanisms (n) ⁸	Antibiotics to which strains were resistant(n) ⁹	MGEs (n) ¹⁰	Refer ence
Algeria (6)	2015	<i>S. agalactiae</i> (44)	Vaginal swab (44)	(44)	100	ST1(9), ST19(14), ST10(4), ST158, ST166, ST233, ST460, ST521, ST677	<i>tet(M)</i> (44), <i>erm(B)</i> (19), <i>mefA/E</i> (1), <i>erm(A)</i> (1)	TET (44) ERY (13)	ND	⁴⁵
	2014	<i>S. aureus</i> (159)	Nasal swab (159)	159 (9)	5.66	ST80 (4), ST5 (2), ST22 (2), ST535 (1)	<i>mecA</i> (9)	GEN ((3), TET (3), TOB(6) SXT(2)	SCCm ec (9)	⁴⁶
	2013	<i>S. aureus</i> (85), <i>E. faecalis</i> (7), <i>C. ONs</i> (31)	Human(123)	123(NS)	NS	NS	<i>mecA</i> (73), <i>aphA</i> (70), <i>aacA-aphD</i>	Methicillin(73)	ND	¹⁵⁴
	2012	<i>E. faecium</i> (80), <i>E. faecalis</i> (39) <i>E. gallinarum</i> (4), <i>E. raffinosus</i> (1), and <i>E. durans</i> (1).	Urinary (85), cutaneous (24), blood (14), pus (2)	125 (108)	87	ST 317 (33), ST51(20), ST52(11), ST175 (8), ST78(25), ST578(4), ST81(2), ST16(2)	<i>erm(B)</i> (92), <i>vanC1</i> (4)	AMP (38), GEN (68), TET (103), ERY (106), CAM (18), LVX ((89), NIT (24), VAN (4).	ND	¹⁴⁸
	2012	<i>S. aureus</i> (64)	Pus (47), venous catheters (7)	(64)	100	ND	<i>mecA</i> (64)	MET (64), OXA (64), FOX (64)	SCCm ec	⁴⁷

⁴ Total number of studies per country

⁵ Total number of isolates

⁶ Total number of specimen source

⁷ Total number of resistant clones

⁸ Total number of resistant genes

⁹ Total number of antibiotics to which strains were resistant to.

¹⁰ Total number of MGEs

			tracheal aspirates (4), puncture fluids (3), blood (2), urine (1)						(46)	
	2010	<i>S. aureus</i> (221)	Skin and soft tissue(158),bone and joint (infection(25),bacteraemia(20),pneumonia(12),eye infection(7),meningitis(3), UTI(2)	221(41)	18.55	ST80(13),ST241(9)	<i>mecA</i> (97)	KAN(29),TET(25),ERY(25),FUS(41)	SCCmec(97)	
Angola (4) and Sao Tome principe (1)	2015	<i>S. aureus</i> (164)	Nasal swab (164)	164 (29)	17.68	ST88(15), ST8(9)	<i>mecA</i> (NS)	FOX (29), SXT (26), TET (18), ERY (16), CIP (9) and CLI (8)	SCCmec (NS)	49
	2015	<i>S. aureus</i> (203)	Nasal (203)	203(128)	63.05	ST8(16), ST5(83), (ST88(19), ST72(5), ST789(1), ST5/2629(2), ST30(2), ST22(1)	<i>mecA</i> (127)	SXT (136), FOX (128), TET (39), PEN (200), RIF (156), CLI (4), ERY (14), CIP (20), GEN (43), CHL (18)	SCCmec (128)	114
	2015	<i>S. aureus</i> (70)	Nasal swab(70)	70(61)	87.14	ST5(13),ST88(6),ST601(1)	<i>mecA</i> (20)	PEN(67),FOX(20),RIF(61),SXT(15),CHL(6),GEN(3),TET(7),FUS(1),CIP(1)	ND	155
	2014	<i>S. aureus</i>	Nasal swab (128)	128(124)	96.88	ST8(57), ST88(9), ST8(5), ST72(3), ST789(1)	<i>mecA</i> (77)	PEN (124), FOX (77), SXT (80), GEN (24), RIF (97), CHL (11), CIP (10), TET (16), ERY (8)	SCCmec (128)	123
Cape verde (1)	2015	<i>S. aureus</i>	Nasal swab (113)	113(16)	14.16	ST88(2), ST8(1), ST5(3)	<i>mecA</i> (6)	FOX (5), TET (5), PEN (109), CIP (2), CLI (3), SXT (12), ERY (16), (FUS (5), MUP (6)	SCCmec (6)	
Democratic Republic of Congo (3)	2017	<i>S. aureus</i> (108)	blood(108)	108(27)	25	ST5(11),ST8(30),ST88(1),ST152(17)	<i>dfgG</i> (24), <i>aac</i> (6')- <i>aph</i> (2'')(25), <i>tet</i> (K)(23), <i>erm</i> (C)(20)	TET(61),LIN(20),CIP(20),PEN(87),CHL(5),SXT(4),	ND	120

	2016	<i>S. aureus</i> (100)	Nasal swab (100)	100 (97)	97	ST8 (9)	<i>dfrG</i> (72), <i>tet</i> (K) (44), <i>femA</i> (98), <i>mecA</i> (33)	TMP(72), PEN (97), TET(45),GEN(25),OXA(24),ERY (20),LUV(16),RIF(7),CHL(7),CLI(4)	ND	119
	2015	<i>S. aureus</i> (63)	Nasal swabs (63)	63(10)	15.87	ST8 (8), ST5 (1), ST88 (1)	<i>mecA</i> (10)	TET(21),ERY(12),CLI(8),PG(60), CHL(9),KAN(12),GEN(12),TOB(12), SXT(6)	SCCmec (10)	65
Egypt (10)	2017	<i>S. aureus</i> (20), <i>S. haemolyticus</i> (9), <i>S. schleifer</i> (3), <i>S. warnei</i> (2), <i>S. lugdunensis</i> (4)	Urine(NS), Blood(NS)	58(38)	65.52	ND	<i>mecA</i> (19)	FOX(25),CIP(21),CLI(21), SXT(21),ERY(38),GEN(32),RIF(14), TET(27)	SCCmec	156
	2016	<i>E. faecalis</i> (57)	Urine(57)	57(52)	91.23	ND	<i>acc</i> (6) <i>la-aph</i> (2) <i>la</i> (21), <i>erm</i> (B)(51), <i>mef</i> (A/E) (1)	AMX(14),VAN(2),FoF(36),GEN(20), AMK(52)	ND	157
	2016	<i>Staphylococcus spp</i>	Urine(3),blood(10), pus(7),sputum(4),bronchoalveolar lavage(2)	81(26)	32.1	ND	<i>fusB</i> (8), <i>fusC</i> (9)	GEN(14),RIF(5),AMP(17)	ND	158
	2016	<i>S. aureus</i> (60)	Human(60)	60(NS)	NS	ST22(1),ST239(1)	<i>mecA</i> (14), <i>erm</i> (C)(14)	CLI(NS),CIP(NS),GEN(NS),SXT(NS), VAN(NS),OXA(NS),ERY(NS)	ND	159
	2016	<i>S. aureus</i> (64)	Sputum(18),pus(35), urine(10),CSF(1)	64(45)	69.23	ND	<i>mecA</i> (NS)	CRO(45),ERY(38),OXA(38),SXT(31), GEN(22),CIP(19),CLI(17),VAN(3)	ND	160
	2015 H	<i>E. faecium</i> (26), <i>E. faecalis</i> (47)	Urine (100)	(73)	100	ND	<i>vanA</i> (2)	PEN(17), AMP(38), CIP(22), GEN(41), STR(73), CHL(12), TET(50), VAN(2)	ND	80

	2014	<i>S. agalactiae</i> (100)	Vaginal swab (100)	100 (98)	98	ND	<i>erm</i> (B) (9), <i>erm</i> (A) (1), <i>mefA</i> /E(1), <i>tet</i> (M) (99), <i>tet</i> (L)(12), <i>tet</i> (K)(1), <i>tet</i> (O) (1)	ERY(17), CLI(14), AZI(16), TET(98) and CHL(1)	ND	32
	2014	<i>S. aureus</i> (127)	Diabetic foot ulcers (39), surgical site infection (48) and abscess infections (25), burn discharges (15).	127 (111)	87.40	ND	<i>mecA</i> (29)	AMP(111), AMX(104), OXA(31), LEX(83), CXM(67), CFP(43), FEP(56), CTX(32), SAM(37), AMC(41), AMK(3) CIP(32), NOR(37), OFX(31), LVX(11), GAT(5), ERY(59), Cll(34), TET(66), VAN(2), CHL(44), RIF(35)	ND	93
	2013	<i>S. aureus</i> (94)	Blood and wound	94 (45)	47.87	ND	<i>gyrA</i> (C2402T, T2409C, T2460G) (60), <i>gyrB</i> (T1497C, A1578G) (5)	CIP(26), LUX(26), AMC(26), FEP(24), GEN(11), TET(17), CHL(5)	ND	29
	2008	<i>S. aureus</i> (60)	Sputum(13),throat swabs(11), nasal swabs(31), blood(9)	60(31)	51.67	ND	<i>mecA</i> (18)	MET(31)	ND	161
Gabon (2)	2016	<i>S. aureus</i> (103)	Throat swab(79),skin lesions(24)	103(61)	59.22	ND	<i>mecA</i> (3), <i>blaZ</i> (90), <i>mrs</i> (A)(8), <i>aphA3</i> (1), <i>df rA</i> (2), <i>tet</i> (K)(56), <i>tet</i> (M)(6), <i>qacC</i> (4)	PEN(90),OXA(1), CXM(1),ERY(8),TET(61),SXT(5 1),CIP(3)	ND	162
	2014	<i>S. aureus</i> (212)	Skin and soft tissue (100) and bloodstream (12)	212 (104)	49.06	ND	<i>df rA</i> (1), <i>df rG</i> (100), <i>df rK</i> +G (1), <i>df rB</i> (2) <i>mecA</i> (1)	TMP;(104), SXT(100), SMZ(6)	ND	55
Ghana (3)	2015	<i>S. aureus</i> (30)	Skin and Soft Tissue Infections (16) , bacteraemia (5), nasal swab (9)	(30)	100	ST88 (8),ST8 (5), ST247 (4)	<i>tet</i> (M) (13) , <i>tet</i> (K) (10), <i>aphA3</i> (7), <i>aacA-aphD</i> (5), <i>erm</i> (C) (4).	TET(20), NOR(12), MXF(11), ERY(11), CLI(9), KAN(9),GEN(9) and CPT (6)	ND	110
	2014	<i>S. aureus</i> (308)	Blood (112), SST1(173), others (23)	308 (208)	67.53	ST88 (2), ST8 (1), ST789 (1), ST72 (1), ST2021 (1), ST250 (2), ST239 (1)	<i>mecA</i> (9)	PEN(208), TET(129), and ERY(18)	ND	109

	2013	<i>S. aureus</i> (105)	Nasal swab(105)	105(29)	27.62	ST88(4),ST8(1),ST 172(1)	<i>mecA</i> (6)	PEN(98),FUS(13),TET(29),FOX(6),SXT(3),ERY(5),CLI(3),NOR(2),GEN(2),RIF(1),MUP(1)	SCCmec(6)	163
Kenya (2)	2016	<i>S. aureus</i> (93)	Blood(93)	93 (32)	34.41	ST22(4),ST88(1),ST789(1),ST5(1),ST8(2),ST241(12),ST239(2)	<i>mecA</i> (32)	CLI(10), ERY(9) and SXT(9),MXF(1), RIF(3), TET(6),LUX(5)	SCCmec (32)	87
	2013	<i>S. aureus</i> (82)	Boil(39),abscess(14),cellulitis(18),ulcer(11),	82(69)	84.12	ND	<i>mecA</i> (52)	ERY(56),CLI(31),CIP(55),OXA(69),FOX(69),SXT(51),GEN(69)	SCCmec	164
Libya (1)	2014	<i>S. aureus</i> (208)	Nasal swab (44)	208(70)	33.69	ND	<i>mecA</i> (35)	CIP(22), GEN(24), FUS(49)	ND	165
Morocco (2)	2013	<i>S. aureus</i> (30)	Nasal swab (30)	30 (25)	83.33	ND	<i>mecA</i> (1)	PEN(25), GEN(1), TOB(1), KAN(1), PF(1), TET(1), ERY(1), SXT(1)	ND	27
	2012	<i>S. aureus</i> (79)	Human(79)	79(43)	54.43	ND	<i>mecA</i> (28)	PEN(74),KAN(29),TOB(27),GEN(27),ERY(21),FUS(25),PF(30),TET(43),MIC(34),RIF(25),SXT(19)	ND	166
Mozambique (1)	2013	<i>S. aureus</i> (24)	Wound (24)	24 (9)	19.15	ND	<i>mecA</i> (9)	FOX(9), OXA(8)	ND	129
Namibia (1)	2014	<i>S. aureus</i> (116)	skin and soft tissue (31), urinary tract(19), respiratory tract (37), ear (7), eye (4) and bloodstream (3)	116 (34)	29.31	ND	<i>dfrA</i> (14), <i>dfrG</i> (20) <i>mecA</i> (11)	SXT(20), TMP(34) SMZ(20)	ND	55
Nigeria (9)	2015	<i>S. aureus</i> (38)	throat (40), nasal (23), wound (10)	38 (32)	84.21	ST8 (5), ST152 (1), ST772 (1), ST14(1)	<i>mecA</i> (16)	TET(32),LUX(7), GEN(5), ERY(5), PEN, SXT(29)	ND	167
	2015	<i>S. aureus</i>	Skin and nasal		72.76	ND	<i>mecA</i> (7), <i>blaZ</i>	PEN(284), SXT(233),	SCCm	58

		(290)	swab (120), wounds, blood	290 (211)			(284))	TET(51),OXA(7),GEN(11),TOB(11),LUX(23),MXF(21),TGC(51),	ec (7)	
	2014	<i>S. epidermidis</i> (20), <i>S. haemolyticus</i> (10), <i>S. saprophyticus</i> (5), <i>S. capitis</i> (5), <i>S. lugdunensis</i> (2), <i>S. warneri</i> (4), <i>S. xylosus</i> (n4), <i>S. cohnii</i> (3).	Stool (53)	(53)	100	ND	<i>mecA</i> (15), <i>aac(6)</i> – <i>aph(2)</i> (3), <i>erm(C)</i> (4), <i>msrA</i> (1), <i>tetK</i> (6), <i>tet(M)</i> (4)	PEN(53), OXA(15), GEN(3), ERY(5), TET(7), SXT(19), CHL(4),AMC (31),CIP(1)	SCCmec(15)	58
	2014	<i>S. aureus</i> (183)	Skin and soft tissue (32), urinary tract (9), ear (7), unknown site (4), oropharynx (3), eye (3) and bloodstream (1)	183 (154)	84.15	ND	<i>dfrA</i> (2), <i>dfrG</i> (152), <i>mecA</i> (16)	(TMP)(154), SXT(83),SMZ(85)	ND	55
	2013	<i>S. aureus</i> (61)	Human(61)	61(27)	44.26	ST39(1),ST5(2),ST241(1),ST250(1),ST88(2)	<i>mecA</i> (7)	PEN(45),TET(26),CLI(2),GEN(10),LVX(6), SXT(27)	SCCmec(2)	168

	2012	<i>S. aureus</i> (51) <i>S. haemolyticus</i> (21), <i>S. sciuri</i> (9), <i>S. saprophyticus</i> (5), <i>S. warneri</i> (3), <i>S. epidermidis</i> (1) and <i>S. hominis</i> (1),	wounds, (11) skin and soft tissues (12), osteomyelitis (5), burns (1), urinary tract infection (6), septicaemia (17), urinary tract infection (10), otitis media (2), bronchitis (2)	91 (36)	39.56	ST241 (1), ST8 (1), ST152 (1), ST37 (37), ST39, ST88	<i>mecA</i> (15), <i>dfrA</i> (3)	SXT(13), PEN(15), OXA(15), GEN(6), CIP(7), MXF(1), ERY(5), CLI(4), TET(13), SXT(13), RIF(2)	SCCmec (15)	112
	2011	<i>S. aureus</i>	Human(68)	68(49)	72	ND	<i>mecA</i> (11), <i>ermA</i> (6), <i>msrA</i> (2), <i>aacA-aphD</i> (10), <i>tetM</i> (11), <i>tetK</i> (27)	PEN(60), OXA(11), GEN(10), TET(38), CIP(20), MXF(7), SXT(49), ERY(8), CLI(6)	SCCmec(11)	169
	2009	<i>S. aureus</i> (96)	Human(96)	96(12)	12.5	ST241(12)	<i>mecA</i> (12)	PEN(12), OXA(12), FOX(12), GEN(12), ERY(12), CLI(9), SXT(12), CIP(12)	ND	170
	2009	<i>S. aureus</i> (346)	Human(346)	346(206)	59.54	ST5 (72), ST7 (44), ST121 (38), ST250(28), ST88 (33), ST30(26), ST8(18), ST1(20), ST15(8), ST80 (8), ST241 (7), ST25 (5), ST72 (3)	<i>mecA</i> (70)	PEN(316), SXT(206), TET(182), CIP(58), ER7(26), GEN(42)	SCCmec(70)	171

Saõ Tome Príncipe (3)	2015	<i>S. aureus</i> (114)	Nasal swab (114)	114(29)	25.5	ST5(2),ST88(11), ST8(13),ST1(2),ST105(1)	<i>mecA</i> (29)	FOX(29),PEN(114),TET(30),CIP(28),RIF(6),GEN(20),CLIN(20),SXT(58),ERY(25),CH	SCCmec (29)	114
Sao Tome principe and Angola	2015	<i>S. aureus</i> (164)	Nasal swab (164)	164 (29)	17.68	ST88(15),ST8(9)	<i>mecA</i> (NS)	FOX(29), SXT(26), TET(18), ERY(16), CIP (9) and CLI(8)	SCCmec (NS)	49
		<i>S. aureus</i> (52)	Nasal swab (52)	52(27)	51.92	ST8(3), ST88(2),ST5(1),ST105(1)	<i>mecA</i> (14)	SXT(27),ERY(11), CIP(11),TET(12),FOX(14),RIF(2)	SCCmec	139
South Africa (11)	2017	<i>S. aureus</i> (1914)	Blood (1914)	1914(557)	29.10	ST239(8),ST612(8),ST4121(1),ST36(4),ST5(4),ST33(3)	<i>mecA</i> (483)	β-lactams(557),TET(NS),aminoglycoside(NS),SXT(NS)	SCCmec (482)	135
	2017	<i>S. aureus</i> (97)	Human	97(96)	99	ND	<i>norA</i> (96), <i>norB</i> (96), <i>mepA</i> (95), <i>tet</i> (38)(96), <i>sepA</i> (94), <i>mdeA</i> (93), <i>imrs</i> (86), <i>sdrM</i> (83), <i>norC</i> (77), <i>qacA/B</i> (34), <i>smr</i> (42)	NS	ND	138
	2017	<i>E. faecalis</i> (1)	Urine (1)	1	100	ST6(1)	<i>aph</i> (3')-III(1), <i>ant</i> (6)-Ia (1), <i>aac</i> (6')- <i>aph</i> (2'') (1), <i>isa</i> (A)(1), <i>mphd</i> (1), <i>tet</i> (M)(1)	GEN(1),STR(1),ERY(1),CLI(1),TET(1),CLI(1),TET(1),CIP(1)	ND	172
	2017	<i>E. faecium</i> (1)	Urine (1)	1	100	ST18(1)	<i>aph</i> (3')-III(1), <i>ant</i> (6)-Ia (1), <i>tet</i> (M)(1), <i>erm</i> (B)(1), <i>msr</i> (C)(1), <i>tet</i> (L)	GEN(1),STR(1),ERY(1),CLI(1),TET(1),CLI(1),TET(1),CIP(1)	ND	173

	2016	<i>S. aureus</i> (27)	Blood (5), nasal (2), CVP(2), Endotracheal tube (2), pus (2), sputum (1), wound (20), Eye (1), humerus (1), bone (1), cheek (1), buttock (1), head (1)	(27)	100	ND	<i>mecA</i> (27) and <i>blaZ</i> (27), <i>aac</i> (6')- <i>aph</i> (2") (25), <i>erm</i> (C) (13)	CIP(23), GEN(20), RIF(19), TET(18), ERY(17), CLI(3)	ND	174
	2016	<i>E. faecium</i> (120) <i>E. faecalis</i> (40)	Blood (4)	(4)	100	ST80 (1),ST203 (1),ST18 (1),ST817(1)	<i>vanA</i> (3), <i>vanB</i> (1)	VAN (4)	ND	147
	2015	<i>S. agalactiae</i> (128)	vaginal and rectal swabs (128)	128 (121)	94.53	ND	<i>erm</i> (B), (28), <i>linB</i> (48) <i>mefA</i> (48)	ERY(27), CLI(32), CHL(32),TET(111),CIP(24)	ND	60
	2015	<i>S. aureus</i> (2709)	Blood (2709)	2709 (1231)	45.44	ND	<i>mecA</i> (1160)	TET(NS), RIF (NS),MUP(NS), CIP(NS) and SXT(NS) MET(1231)	SCCmec (1160)	59
	2012	<i>S. aureus</i> (13746)	Human (13746)	13746(3298)	24	ST5 (1), ST612 (44),	<i>RpoB</i> (H481Y, H481N, I527M) (NS)	RIF(1760)	ND	59
	2009	<i>S. aureus</i> (17)	Human(17)	17(13)	76.47	ND	<i>mupA</i> (3)	ERY(12),CIP(10),RIF(4),CHL(4)	ND	175
	2007	<i>S. aureus</i> (3), <i>S. lugdunensis</i> (2)	Wound(4),blood(1)	5(5)	100	ND	<i>mecA</i> (5)	PEN(5), OXA(5),GEN(5),ERY(4),TET(5), SXT(5),RIF(5)	SCCmec(5)	176
Sudan(1)	2015	<i>S. aureus</i> (200)	Wound(49),ear swab(57),urine(47), nasal swab(47)	200(197)	98.5	ND	<i>mecA</i> (111)	PEN(197), AMP(197),GEN(122),KAN(136),I PM(89),AMO(87),CIP(123),CLI(113),SXT(105)	ND	177

Tanzania (1)	2014	<i>S. aureus</i> (87)	Skin and soft tissue (39) and bloodstream (2)	87 (32)	36.78	ND	<i>dfrG</i> (32)	SMZ(5), TMP (32)	ND	50
Tunisia (18)	2015	<i>S. aureus</i> (99)	Human (99)	(99)	100	ST247 (12), ST239 (6), ST728 (2), ST241 (1), ST398 (1), ST5 (1) and ST641 (1)	<i>mecA</i> (24), <i>tet(K)</i> (6), <i>tet(L)</i> (1), <i>tet(M)</i> (18), <i>erm(A)</i> , <i>aph(2'')-acc(6')</i> (13)	TET(24), GEN(18), ERY(15), FOF(1), CLI(14), OFX(16), TOB(20), FUS(5)	ND	69
	2014	<i>E. faecium</i> (13), <i>E. gallinarum</i> (3)	blood (8), pus (3), urine (2) and rectal swabs (3).	(16)	100	ST18 (1) and ST80 (2)	<i>vanA</i> (13), <i>vanC1</i> (3), <i>erm(B)</i> (16), <i>tet(M)</i> (15), <i>tet(L)</i> (1), <i>aac(6')-aph(2'')</i> (13) <i>aph(3')-IIIa</i> (16), <i>ant(6)</i> (3)	VAN(16), TEC(13), AMP(16), CIP(16), ERY, TET(16), KAN(13), STR(13), SXT(16), GEN(8),	<i>IS16</i> (3)	178
	2013	<i>S. aureus</i> (69)	Human (69)	(69)	100	ST80 (41), ST1440 (1), ST1 (2), ST5 (5), ST22 (1), ST97 (2), ST239 (4), ST241 (3), ST247 (3), ST1819 (3), ST153 (2), ST256 (1)	<i>mecA</i> (59)	KAN(62), AMK(62(18), TETs(61), OFX(20), CIP(31), ERY(38), CLI(12), RIF(22)	<i>SCCmec</i> (59)	85
	2013	<i>S. aureus</i> (64)	Pus(53)pus, blood culture (6), articular Puncture (4), venous catheter r(1).	(64)	100	ST80(64)	<i>mecA</i> (64)	PEN(64), OXA(64), FOX(64), AMK (64), KAN(63), ERY(13), TET(3), LI N(3)	<i>SCCmec</i> (64)	179
	2012	<i>S. agalactiae</i> (226)	Female genital (120), gastric fluid (106)	226 (220)	97.34	ND	<i>erm(B)</i> (79), <i>mef(A)</i> (2), <i>tet(M)</i> (205), <i>tet(L)</i> (10), <i>tet(O)</i> (5), <i>tet(T)</i> (1)	CHL(7), RIF(43), ERY(90) and TET(220), STR(7), GEN(7)	<i>Tn916</i>	105

	2012	<i>S. haemolyticus</i> (46)	Blood (19), intravascular catheters (14), others (13)	46 (36)	78.26	ND	<i>mecA</i> (28)	PEN(36), OXA(36), GEN(34), KAN(34), and TOB(34), ERY(33), SXT(32), OFX(32), CIP(32), STR(25), fusidic acid(14), TET(11), RIF(9), LIN(6), CHL(1), FOF(1)	SCC <i>mec</i> (28)	104
	2011	<i>S. aureus</i> (1463)	Skin (1463)	160 (5)	3.13	ND	<i>erm</i> (C)(3), <i>erm</i> (A) (1), <i>vat</i> (B) (5), <i>vga</i> (B) (5)	PEN(5), OXA(4), GEN(4), KAN(5), TOB(4(5) and RIF(5), LIN(5)	ND	144
	2011	<i>S. pyogenes</i> (103)	skin (43), respiratory tract (41), blood (12), fluids (4), endometrium (1), vagina (1), and urine (1).	103 (72)	70	<i>emm</i> 18 (4), <i>emm</i> 42 (9), <i>emm</i> 76 (6), <i>emm</i> 118(10)	<i>erm</i> (B) (5), <i>tet</i> (M) (63), <i>tet</i> (O)(3)	ERY(5), CLI (5), and TET(72),	<i>Tn916</i> (62)	145
	2011	<i>S. epidermidis</i> (34), <i>S. haemolyticus</i> (10), <i>S. hominis</i> (1)	Blood(45)	45(42)	93.33	ND	<i>mecA</i> (43), <i>mrsA</i> (13), <i>erm</i> (C)(7), <i>erm</i> (B)(2), <i>erm</i> (A)(6), <i>aac</i> (6')- <i>le-aph</i> (2'')(35), <i>ant</i> (4')- <i>la</i> (18), <i>aph</i> (3')- <i>lla</i> (4), <i>tet</i> (K)(6), <i>tet</i> (M)(1)	PEN(45), OVA(43), GEN(35), KAN (42), TOB(40), ERY(25), CLI(11), TET(5), CHL(3), RIF(15), SXT(31), CIP(25), FUS(27), FOF(18)	SCC <i>mec</i> (43)	180
	2010	<i>S. pyogenes</i> (193)	throat (63), pus (89), punctures (30), blood (4), other sources (7)	193 (13)	6.74	ND	<i>ermB</i> (6), <i>mefA</i> (2)	ERY(7) and TET(6)	ND	143
	2010	<i>S. aureus</i> (55)	Nasal swab(55)	55(19)	35.55	ST80(1)	<i>mecA</i> (1), <i>ant</i> (6)- <i>la</i> (3), <i>tet</i> (K)(7), <i>aph</i> (3')- <i>lla</i> (4), <i>dfrA</i> (1), <i>tet</i> (M)(1), <i>tet</i> (L)(1)	PEN(54), OXA(19), FOX(1), TET(11), STR(5), KAN(3) CIP(8)	SCC <i>mec</i> (1)	181
	2010	<i>S. agalactiae</i> (160)	Urinary tract (160)	(160)	100	ND	<i>erm</i> (B) (132), <i>erm</i> (TR) (13), <i>mef</i> (A) (3)	ERY(160), LIN(135) and SB (135)	ND	182

	2010	<i>S. aureus</i> (13)	Pus(32),blood(16), catheter(12)	72(42)	58.33	ND	<i>mecA</i> (13)	PEN(65),STR(11),GEN(4),KAN(11), OXA(13),TOB(4),LIN(3),TET(42)ERY(11),RIF(6),CHL(2),CIP(5),FUS(8),FOF(1)	ND	183
	2009	<i>S. epidermis</i> (77), <i>S. mitis</i> (50), <i>E. faecium</i> (45)	blood cultures (55), central venous catheters, (22),stool cultures (40), respiratory tract (2) and different sites (3), systematic nasopharyngeal specimens (42), upper respiratory tract(5)	172(95)	55.23	ND	<i>erm</i> (C) (18), <i>erm</i> (B) (6), <i>erm</i> (A)(11), <i>msrA</i> (5)	OXA(39), AMP(28),PEN(90),ERY(119),LIN(97),PRI (3),GEN(71),RIF(78),TEC(50),	ND	82
	2007	<i>E. faecalis</i> (34), <i>E. faecium</i> (12)	Blood(10), pus(26),catheter(7) ,plural aspirate(2)	46(46)	100	ND	<i>aac</i> (6')- <i>aph</i> (2'')(46)	GEN(46),KAN(46),PEN(12),ERY(45),CHL(25),TET(32),STR(26)	ND	184
	2007	<i>E. faecium</i> (2)	Urine(2)	2	-	ND	<i>vanA</i> (2)	STR(2), ERY(2),CIP(2),VAN(2)	ND	156
	2007	<i>S. epidermidis</i> (346)	Human(346)	346(7)	2.02	ND	<i>erm</i> (A)(6), <i>erm</i> (C)(1), <i>vga</i> (7)	PRI(7),OXA(7),GEN(7),ERY(7),LIN(7),RIF(7),SXT(7)TEC(1)	ND	185
	2007	<i>S. epidermidis</i> (34)	Blood(55), urine(22)	(34)	100	ND	<i>icaA</i> (26), <i>erm</i> (C)(18), <i>erm</i> (A)(11), <i>mrsA</i> (5), <i>vga</i> (3),	ERY(34),OXA(28),GEN(34),LIN(33),OFX(33),RIF(28)	ND	186
Uganda (4)	2013	<i>S. aureus</i> (64)	Nasal swab (64)	64(24)	37.5	ND	<i>mecA</i> (24)	OXA(22), GEN(8), CIP(12), CHL(9)	SCCmec (24)	187
	2012	<i>S. epidermidis</i> (50)	Nasal swab(20),catheter(14),blood(9),wound(3)	50(26)	52	ND	<i>aph</i> (')- <i>lla</i> (28), <i>blaZ</i> (2), <i>mecA</i> (3), <i>vanA</i> (3), <i>vanB1</i> (3),	ERY(20),GEN(26),PEN(32),TET(15),SXT(17),OXA(6)	IS256(33)	188
	2011	<i>S. aureus</i> (122)	pus	122(48)	39.34	ND	<i>mecA</i> (2)	AMP(48),CHL(42),CIP(1),ERY(5),TET(29),SXT(320)	ND	189
	2009	<i>S. aureus</i> (54)	Human(54)	54(15)	27.78	ND	<i>mecA</i> (17)	CIP(12),GEN(10),SXT(15),CHL(15),ERY(15)	NG	190

1529 **Table 3. Geographical distribution, species, clones, and resistance mechanisms of antibiotic-resistant Gram-positive bacteria isolated from**
1530 **animals in Africa from 2007-2018.**

Country (n) ¹¹	Year	Organism/Species (n) ¹²	Specimen Sources (n) ¹³	Sample size (Resistant isolates)	Resistance rate (%)	Clones (n) ¹⁴	Resistance genes/mechanisms (n) ¹⁵	Antibiotics to which strains were resistant (n) ¹⁶	MGEs (n) ¹⁷	Reference
Angola(1)	2015	<i>E. faecium</i> (3)	Pig faeces(1), Chicken faeces(2)	3	-	ST971, ST245(2)	<i>tet</i> (L)(1), <i>tet</i> (M)(2), <i>erm</i> (B)(2)	CIP(1),TET(3),ERY(2),STR(2),NIT(2),Q/D(2)	ND	191
Egypt (10)	2017	<i>S. aureus</i> (3), <i>S. hycus</i> (6), <i>S. intermedius</i> (3), <i>S. epidermis</i> (1), <i>S. hemolyticus</i> (1), <i>S. hominis</i> (1), <i>S. l ugdunensis</i> (3), <i>S. simulans</i> (1), <i>S. scuri</i> (4)	imported beef meat (23)	23(16)	69.57	ND	<i>mecA</i> (5), <i>gyrA</i> (12), <i>griA</i> (10), <i>gyrB</i> (6),	AMP((6),CHL(1),CIP(8),CLI(15), ERY(6),GEN(14),MET(8),OXA(13), PEN(22), TET(6)	ND	94
	2017	<i>S. aureus</i> (84)	Milk(84)	84(80)	95.23	ND	<i>mecA</i> (42), <i>blaZ</i> (67)	AMX(54), SXT(66),GEN(20),CIP(12),CHL(58),PEN(70), RIF(32),AMK(14), VAN(64),STR(50),TET(44),ERY(40), AMP(80),OXA(42)	ND	192
	2016	<i>S. aureus</i> (73)	Animal(73)	73(NS)	NS	ST113(1), ST80(1)	<i>mecA</i> (14), <i>erm</i> (C)(14)	CLI(NS),CIP(NS),GEN(NS),SXT(NS),OXA(NS), ERY(NS)	ND	159

¹¹ Total number of studies per country

¹² Total number of bacteria isolated

¹³ Total number of Specimen source

¹⁴ Total number of resistant clones

¹⁵ Total number of resistant genes

¹⁶ Total number of isolates resistant to antibiotics

¹⁷ Total number of mobile genetic elements: plasmids, transposons, integrons

	2016	<i>S. aureus</i> (30)	raw chicken breast fillet (40), sliced luncheon meat (20), and chicken nuggets (20), Human (18)	40 (21)	33.33	ND	<i>mecA</i> (10)	DOX(31), AMX(29), OFX(10), CFP(23), CLI(21), GEN(20), APR(16), ERY(21), SXT(23), LUX(18), NAL(20), OFX(10), CIP(16).	ND	28
	2016	<i>S. aureus</i> (70)	Bovine(70)	70(41)	58.57	ND	<i>mecA</i> (NS)	CRO(41), ERY(35), OXA(41), SXT(14), GEN(14), CIP(11), CLI(8), VAN(1)	ND	160
	2016	<i>S. aureus</i> (40)	Milk(30), meat(10)	40(22)	55	ND	<i>erm</i> (A)(18), <i>mrs</i> (A)(4), <i>mphC</i> (6), <i>erm</i> (B)(3)	ERY(22), CLI(4), TET(24), CIP(4), CHL(5), AMX(26), FOX(22), SXT(1), RIF(5), GEN(4), CRO(14)	ND	193
	2016	<i>S. aureus</i> (200)	Raw milk (40), Damietta Cheese (40), Kareish cheese (40), ice cream (40), and yogurt (40)	200 (106)	53	ND	<i>mecA</i> (106)	TET(270), NEL(78), AMX(230), CLX(314), STR(186), SXT(58), GEN(114), PEN(364), RIF(152), CHL(128), AMK(146), VAN(36)	ND	12
	2015	<i>S. aureus</i> (133)	cow milk samples (61), various origins (14), minced meat (6), sausage (4) and burger (7), pus (22), sputum (17), urine (1), cerebrospinal fluid (1)	133 (96)	72.18	ND	<i>mecA</i> (30)	CRO(96), TET(90), OXA(70), FOX(65), ERY(81), VAN(4), IPM(7), CRO(96), CHL(12), GEN(36), CLI(29), CIP(31), RIF (18)	SCCmec (25)	23
	2015	<i>S. aureus</i> (288)	Chicken(288)	288(256)	88.89	ND	<i>mecA</i> (76)	PEN(269), AMP(256), CLX(240), AMX(224), ERY(212), TET(197), STR(150), RIF(113), AMK(99), CH	ND	194

								L(91),GEN(70),CIP(39),NEL(48),SXT(39),VAN(17)		
	2011	<i>S. aureus</i> (4)	dogs swab (70), cats swab (48), human nasal and oral swabs (50).	(4)	100	ND	<i>mecA</i> (4)	OXA(4), FOX(4), AMP(3),FOX(4),RIF(3),GEN(2),CLI(2),RIF(2),CIP(2),TET(1)	ND	53
Kenya (1)	2013	<i>S. agalactiae</i> (92)	Camel(92)	92 (37)	36	ST617 (8), ST-612 (1),ST-616 (22)	<i>tet</i> (M) (37)	TET(37)	Tn916 (37)	195
Nigeria(3)	2017	<i>S. aureus</i> (30), <i>S. epidermidis</i> (16), <i>S. saprophyticus</i> (2), <i>S. sciuri</i> (1), <i>S. xylosus</i> (1)	Pork(26),beef(14),chicken(10)	50(48)	96	ND	<i>mecA</i> (49)	PEN(48),CLI(48),CHL(46),SXT(46),KAN(46),AMX(460)	ND	156
	2016	<i>E. faecium</i> (108), <i>E. gallinarum</i> , (30), <i>E. faecalis</i> (5), <i>E. hirae</i> . (5) <i>E. mundtii</i> (12)	Cattle (130), chickens (130),manure (130)	167 (102)	61.0	ND	<i>tet</i> (K) (NS), <i>tet</i> (L) (NS), <i>tet</i> (M) (NS), <i>tet</i> (O) (NS) and <i>erm</i> (B) (NS)	TET (102), ERY (102), CHL (13), GEN(55), STR(47),AMP(75)	ND	196
	2014	<i>Coagulase negative staphylococcus</i> (16)	Groin swab of dogs(16)	(16)	100	ND	<i>mecA</i> (16), <i>blaZ</i> (1), <i>tetK</i> (12), <i>tet</i> (M)(8), <i>erm</i> (B)(3), <i>aacA-aphD</i> (11)	PEN(16),OXA(16),FOX(16),TET(13),ERY(9),CLI(9),GEN(5),KAN(12),TOB(1),SXT(10),CHL(7)	ND	197
South Africa (6)	2017	<i>E. faecium</i> (180), <i>E. durans</i> (80), <i>E. hirae</i> (29), <i>E. casseliflavus</i> (20)	Cattle (241)	100	100	ND	<i>vanB</i> (67), <i>vanC1</i> (85), <i>vanC2/3</i> (137), <i>erm</i> (B)(137)	ERY(338),CLI(330),VAN(341),PEN(310),CET(300),STR(320),CLX(100),AMK(252),CIP(41)	ND	198
	2017	<i>S. aureus</i> (104)	Chicken(104)	(104)	100	ND	<i>mecA</i> (45), <i>blaZ</i> (12), <i>tet</i> (K)(32)	AMP(46),GEN(29),ERY(64),FOX(71),KAN(52),STR(57),TET(82),VAN(43)		156
	2015	<i>S. aureus</i> (211)	Milk (211)	211 (124)	58.77	ND	<i>mecA</i> (19)	PEN (124), AMP(99), OXA (93), VAN(47), TEC(116), TET(56),ERY(56),STR(89),KAN(55),GEN(47),SXT (37)	ND	21

	2015	<i>E. faecalis</i> (40), <i>E. hirae</i> (100), <i>E. durans</i> (60), <i>E. faecium</i> (120)	Pigs (320)	(320)	100	ND	<i>vanB</i> , (320), <i>vanC1</i> (320), <i>vanC2/3</i> (320), <i>erm</i> (B)(300)	VAN(320), STR(320) and CLX(320), STR(320), CET(286), PEN(292), CIP(248), AMO(64), AMK(272), CLI(316), ERY(280), IPM (52),	ND	13
	2014	<i>Staphylococcus spp</i> (120)	Pigs(30),cattle(30),cows(30),goats(30)	120(53)	44.17	ND	<i>mecA</i> (12)	VAN(12), CRO(12), CFZ(37), CTX(19), SAM(13), PEN(53), MER(4)	ND	199
	2013	<i>S. xylosus</i> (18), <i>S. aureus</i> (28), <i>S. haemolyticus</i> (42), <i>S. capitis</i> (18), and other <i>Staphylococcus spp.</i> (14)	Animals (120)	(120)	100	ND	<i>mecA</i> (NS), <i>mphC</i> (NS)	PEN (90), MER(3), VAN(14), CTX(14), CFZ(48), OXA(46), MIC(19), TET(100), ERY(14), CLI(19), NAL(120), CIP(5), OFX(6), LUX(2)	SCCmec (NS)	61
Senegal (1)	2012	<i>S. aureus</i> (57)	Swabs from pigs (300) and farmers	57(35)	61.40	ST5 (5)	<i>mecA</i> (6)	PEN(57), SXT(35), TET(20)	SCCmec (6)	66
Tanzania (1)	2014	<i>E. faecium</i> (95) <i>E. faecalis</i> (9) <i>E. gallinarum</i> (7) <i>E. Hirae</i> (9)	Faecal samples of buffalo (35), wildebeest (40), zebra (40) and cattle (20)	120 (42)	35	ND	<i>tet</i> (W) (NS), <i>sull</i> (NS)	VAN(10), AMP(10), TET(40), SXT(32), RIF(53), ERY(42), GEN(35), AMP(31)	ND	14
Tunisia (8)	2017	<i>E. faecium</i> (31), <i>E. faecalis</i> (14), <i>E. durans</i> (6), <i>E. casseliflavus</i> (2), <i>E. gallinarum</i> (2)	Faecal sample of cats(20), dogs(50)	58(31)	53.45	ND	<i>erm</i> (B) (22), <i>tet</i> (M)(5), <i>tet</i> (M), <i>tet</i> (L)(16), <i>tet</i> (L)(4), <i>ant</i> (6')- <i>la</i> (11), <i>aac</i> (6')- <i>le-aph</i> (2'')- <i>la</i> (16), <i>aph</i> (3')- <i>lla</i> (11), <i>catA</i> (1)	AMP(1), ERY(26), CIP(30), PRI(9), STR(12), KAN(12), GEN(9), TET(21), CHL(7)	ND	22
	2017	<i>E. faecalis</i> (2), <i>E. faecium</i> (NS), <i>Enterococcus spp</i> (NS)	Urban wastewater (5)	5(2)	40	ST86(2)	<i>optrA</i> (2), <i>erm</i> (A), <i>erm</i> (B), <i>tet</i> (M)(1), <i>tet</i> (L)(1), <i>aac</i> (6')- <i>aph</i> (2''),	CHL(2), CIP(2), ERY(2), TET(1), GEN(1), STR(2)	ND	200

	2015	<i>S. aureus</i> (43)	Chicken(19), Veal (9), sheep(14), horse(1)	43(13)	30.23	ST30(1), ST398(1)	<i>tet</i> (M)(2), <i>erm</i> (C)(4), <i>erm</i> (A)(2), <i>erm</i> (T)(1), <i>tet</i> (K)(6), <i>tet</i> (L)(3), <i>tet</i> (M)(2), <i>aph</i> (3')-IIIa(4), <i>ant</i> (4)-Ia(1), <i>mrsA</i> (4)	PEN(41), OXA(2), FOX(2), KAN(4), TOB(1)	SCCmec(2)	201
	2015	<i>S. aureus</i> (17)	Goat, cats dogs(17)	17(7)	41.18	ST45(1), ST15(1), ST6(1), ST21(1), ST188(1)	<i>blaZ</i> (7), <i>tet</i> (M)(1), <i>erm</i> (A)(1), <i>ant</i> (6)-Ia(1)	PEN(6), TET(1), ERY(1), STR(1), CIP(1)	ND	202
	2013	<i>E. faecalis</i> (49), <i>E. faecium</i> (30), <i>E. gallinarum</i> (12), <i>E. hirae</i> (12), <i>E. casseliflavus</i> (2), <i>E. durans</i> (2)	Meat (199)	(119)	78.5	ST260(1), ST454(1), ST452(1), ST22(1), ST300(1), ST455(1), ST453(1), ST456(1)	<i>tet</i> (M) (36), <i>tet</i> (L) (32), <i>erm</i> (B) (33), <i>aac</i> (6')-aph(2") (1), <i>ant</i> (6) (7)	TET(57), ERY(43), STR(17), CHL(4), GEN (1)	ND	51
	2013	<i>E. mundtii</i> , (23) <i>E. casseliflavus</i> (20), <i>E. hirae</i> (19), <i>E. faecalis</i> (10), <i>E. faecium</i> (10), <i>E. durans</i> (7), <i>E. gallinarum</i> (7), <i>E. dispar</i> (2)	Cattle (92)	92 (72)	78	ND	<i>erm</i> (B) (7), <i>tet</i> (M) (4), <i>tet</i> (L)(4)	ERY(10), TET(4) and SXT(72)	ND	52
	2012	<i>S. aureus</i> (73)	nasal swab from sheep (73)	73 (5)	6.85	ST153(5)	<i>mecA</i> (5), <i>blaZ</i> (28), <i>ant</i> (6)-Ia (5), <i>erm</i> (C) (5), <i>tet</i> (K) (30)	PEN(5), STR(5), KAN(5), ERY(5), TET (5), FUS(5)	ND	99
	2012	<i>S. aureus</i> (50)	Nasal swab of donkey(50)	50(30)	60	ST133(15), ST1738(4), ST1(2), ST6(4), ST2057(4), ST2110(1), ST2181(1), ST1660(1)	<i>baZ</i> (12), <i>erm</i> (A)(8), <i>erm</i> (C)(2), <i>tet</i> (M)(1), <i>fusC</i> (1)	PEN(12), ERY(8), TET(1), Fusic acid(12),	ND	203

Uganda (1)	2017	<i>S. aureus</i> (41)	milk(30),sour milk sample(11)	41(30)	73.17	ST97(1),ST1(2)	<i>mecA</i> (23)	TET(30),RIF(1),SXT(2),ERY(1),GEN((1),CLI(1)	ND	121
------------	------	-----------------------	-------------------------------	--------	-------	----------------	------------------	---	----	-----

1531

1532 **Table 4. Geographical distribution, species, clones, and resistance mechanisms of antibiotic-resistant Gram-positive bacteria isolated from**
1533 **the environment in Africa from 2007-2017.**

Country (n) ¹⁸	Year	Organism/Species (n) ¹⁹	Specimen Sources (n) ²⁰	Sample size (Resistant isolates)	Resistance rate (%)	Clones (n) ²¹	Resistance genes/mechanisms (n) ²²	Antibiotics to which strains were resistant(n) ²³	MGEs (n) ²⁴	Reference
Angola	2015	<i>E. faecium</i> (5)	Chicken farm facility(4), water from hospital and community(1)	5(4)	80	ST245(1), ST650(2)	<i>tet</i> (M)(4), <i>erm</i> (B)(4), <i>tet</i> L(2)	TET(4),ERY(4),STR(4),NIT(2),Q/D	ND	191
Egypt	2016	<i>S. aureus</i> (23)	Food sample(23)	23(NS)	NS	ST689(1)	<i>mecA</i> (3), <i>vanA</i> (1), <i>vanB</i> (1)	VAN(NS), CIP(NS), GEN(NS),SXT(NS),OXA(NS),ERY(NS)	ND	159
Nigeria (1)	2017	<i>E. faecium</i> (100)	Vegetables soil, farm, Cloacal swabs (25), Manure (8), Rectal swabs(2)	(100)	100	ND	<i>aac</i> (6')-Ie- <i>aph</i> (2')-Ia(35), <i>aph</i> (2')-Ic(31), <i>aph</i> (3')-IIa(32), <i>ant</i> (4')-Ia(14)	AMP (63), GEN(37)	ND	25
South Africa (4)	2017	<i>S. aureus</i>	Recreational waters and beach sand (30)	(30)	100	ND	<i>mecA</i> (5), <i>femA</i> (16), <i>rpoB</i> (11), <i>blaZ</i> (16), <i>ermB</i> (15), <i>tet</i> (M)(8)	AMP (29),PEN (29),RIF(24), CLI(24),OXA	ND	204

¹⁸ Total number of studies per country

¹⁹ Total number of isolates

²⁰ Total number of specimen source

²¹ Total number of resistant clones

²² Total number of resistant clones

²³ Total number of isolates resistant to antibiotics

²⁴ Total number of mobile genetic elements : plasmids,transposons, integrons

								(22), ERY(21), VAN(15), TET(13), SXT(13), CIP(10), GEN(1)		
	2016	<i>E. faecium</i> (30), <i>E. faecalis</i> (37) <i>E. mundtii</i> (36), <i>E. casseliflavus</i> (14), <i>E. gallinarum</i> (5), <i>E. hirae</i> (1), <i>E. sulfureus</i> (1)	Surface water(124)	124(86)	69.35	ND	<i>tet</i> (L)(17), <i>msrC</i> (9)	AMP(59), AMX(53), PEN(87), STR(8), VAN(86), CHL(23), CIP(47) ERY(68), TET(59)	ND	205
	2015	<i>E. faecium</i> (30), <i>E. durans</i> . (15)	waste water (32) and effluent (32)	(45)	100	ND	<i>erm</i> (B) (40), <i>vanB</i> , (42), <i>vanC1</i> (42), <i>vanC2/3</i> (42)	PEN(38), ERY(40), CTX(43), GEN(28), IPM(43), TET(45), KAN(43), CIP(43), VAN(42), CLI(45)	ND	10
	2013	<i>E. faecium</i> (179)	Borehole Water (179)	179 (172)	96.09	ND	<i>vanA</i> (17) and <i>vanB</i> (17)	AMP(158), VAN (166)and PEN(172), CHL(11), KAN(12), GEN(3), AMX (155), ERY(86)	ND	9
Tunisia (7)	2017	<i>S. aureus</i> (12)	Wastewater	12	100	ST3245(7), ST15(1)	<i>blaZ</i> (7), <i>msrA</i> (7), <i>tet</i> (K)(1)	PEN(12), ERY(7), TET(1), C LI(1)	ND	18
	2016	<i>E. faecium</i> (86), <i>E. faecalis</i> (8), <i>E. casseliflavus</i> (6)	Hands (50), inanimate such as beds, treatment tables, toilets, faucets, wrists, sinks (250)	(100)	100	ST910 (13), ST80 (1)	<i>erm</i> (B) (71), <i>tet</i> (M) (18), <i>aph</i> (3')-IIIa (27), <i>ant</i> (6)-Ia (15), <i>cat</i> (A) (4), <i>vanC2</i> (6)	ERY(73), TET(20), STR(27) and KAN(28), VAN(14), CHL(10), SXT(100), CIP(48), PRI(18)	IS16 (14)	20
	2016	<i>S. saprophyticus</i> (30), <i>S. haemolyticus</i> (38), <i>S. epidermidis</i> (NS), <i>S. cohnii</i> (NS), <i>S. warneri</i> (NS), <i>S. sciuri</i> (NS), <i>S. simulans</i>	Inanimate surfaces (83)	83 (32)	38.55	ND	<i>mecA</i> (20), <i>msrA</i> (32), <i>erm</i> (C)(8), <i>tet</i> (K)and/or <i>tet</i> (M)(21), <i>aac</i> (6')-Ie- <i>aph</i> (2')-Ia (16), (<i>aph</i> (3')-IIIa(19), <i>ant</i> (4')-Ia (n=14), <i>ant</i> (6')-Ia (3)	ERY(32), TET(21), GEN(16), KAN(19), TOB(14), STR(3),	ND	97

		(NS)s, <i>S. pasteurii</i> (NS), <i>S. arlettae</i> (NS) and <i>S. xilosus</i> (NS)								
	2015	<i>E. faecium</i> (34), <i>E. hirae</i> (23), <i>E. faecalis</i> (4), and <i>E. casseliflavus</i> (4)	Vegetable food (34), soil and irrigation water (27)	65 (40)	61.54	ST2 (5), ST16 (2), , ST528 (2), ST56 (1), ST885 (1), ST886 (1)	<i>erm</i> (B) (12), <i>tet</i> (M)- <i>tet</i> (L)(10), <i>aph</i> (3')-III, (10) <i>ant</i> (6) (2), <i>van</i> C2(4)	CIP(42), ERY(12), TET(10), KAN(10), CHL(5), STR(2), and GEN(5), VAN(4)	ND	19
	2015	<i>E. faecium</i> (54), <i>E. faecalis</i> (17), <i>E. hirae</i> (8) <i>E. casseliflavus</i> (4), <i>E. durans</i> (2)	waste and surface water (114)	(85)	100	ST480 (1), ST531 (1),ST55 (1),ST532(1),ST202 (1),ST314(1), ST985(1),S T30 (1),ST986 (1),ST12 (1),ST296 (1),ST327(1)	<i>aph</i> (3')-IIIa (22), <i>ant</i> (6)- <i>la</i> (4), <i>erm</i> (B) (34), <i>tet</i> (M) (13), <i>tet</i> (L)(8), <i>aac</i> (6')-Ie- <i>aph</i> (2')(15)	GEN(22), KAN(22), STR(7), ERY(36), TET(13), SXT(79), CIP(6),	ND	31
	2015	<i>S. aureus</i> (12)	Hospital environment(12)	12(6)	50	ST247(2)	<i>bla</i> Z(12), <i>erm</i> (A), <i>tet</i> (M)(2), <i>aac</i> (6')- <i>aph</i> (2')(2),	STR(2),KAN(2),ERY(2),CL I(2),TET(2),FUS(2),TOB(2), GEN(2),AMK(2),OXA(6),PEN(12),FOX(2)	SCCmec(2)	206
	2014	<i>E. faecium</i> (5), <i>E. casseliflavus</i> (7)	Hospital environment((beds, treatment table, toilet, faucet, wrist and sink) (100)	(12)	100	ST80(1)	<i>van</i> A(5), <i>van</i> C2(7), <i>erm</i> B(12), <i>tet</i> M(5), <i>aph</i> (3')-IIa(5), <i>aac</i> (6')- <i>aph</i> (2')(5)	VAN,(12),AMP(5),CIP(12), ERY(12),TET(8), STR(6),KAN(80,SXT(11),GEN(3),TEC(5)	<i>IS</i> 16(1)	69

1534

1535 **Table 5. Mean antibiotic resistance rates per country in Africa**

Country	No. of studies N=130	Mean rate of ABR (%)	95% CI	P-value
Algeria	7	62.24	4.76 - 119.7	0.03
Angola	6	66.19	9.98 – 122.4	0.03
Cape Verde	1	14.16	-	-
Democratic Republic of the Congo	3	45.96	-	-
Egypt	21	70.49	59.30 – 81.68	0.0001
Gabon	2	54.14	-	-
Ghana	3	65.05	-	-
Kenya	3	51.51	-	-
Libya	1	33.69	-	-
Morocco	2	68.88	-	-
Mozambique	1	19.15	-	-
Namibia	1	29.31	-	-
Nigeria	13	71.23	54.81 – 87.65	0.0001
São Tome´ Pri´ncipe	3	31.70	12.87 – 76.27	0.092
South Africa	21	82.72	70.73 – 94.69	0.0001
Sudan	1	98.5	-	-
Tanzania	2	35.89	24.58 – 47.00	0.016
Tunisia	33	66.82	54.73 – 78.91	0.0001
Uganda	5	45.96	24.25 – 67.66	0.0042
Senegal	1	61.40	-	-

Table 6: Antibiotic resistance rates of various Gram-positive bacterial species isolated from humans, animals and the environment in Africa between 2007 and 2018.

Species	Total isolates	Antibiotic resistance rate (%)											
		AMP ²⁵	CIP ²⁶	CL ²⁷	ERY ²⁸	GEN ²⁹	KAN ³⁰	PEN ³¹	RIF ³²	STR ³³	TET ³⁴	SXT ³⁵	VAN ³⁶
Human													
<i>E. faecalis</i>	179	30.4	26.3	-	91.35	77.2	100	26.0	-	56.5	76.0	19.30	52.6
<i>E. faecium</i>	205	56.1	19.0	-	88.0	61.4	90.6	21.5	-	70.3	75.5	100	51.3
<i>S. agalactiae</i>	658	-	18.8	19.5	50.6	3.1			19.0	3.1	68.8	-	-
<i>S. aureus</i>	24160	64.7	24.1	16.3	82.4	20.3	32.9	81.5	31.2	12.2	35.4	40.5	3.13
<i>S. haemolyticus</i>	91	-	62.6	24.4	63.7	75.9	73.9	78.3	31.2	12.2	35.4	69.2	-
<i>S. pyogenes</i>	148	-	-	4.9	5.8	-	-	-	-	-	36.5	-	-
Animal													
<i>E. faecalis</i>	129	24.2	64.6	98.8	43.5	19.5	20.7	16.6	44.17	16.7	32.7	52.5	50.1
<i>E. faecium</i>	577	31.4	43.7	97.8	57.5	23.0	20.7	53.7	44.2	37.9	43.9	52.5	66.7
<i>S. aureus</i>	1601	62.8	23.1	39.3	32.7	28.9	36.8	69.5	33.5	45.3	42.2	37.6	24.4
<i>S. haemolyticus</i>	43	-	-	-	-	-	-	59.6	-	-	-	83.3	-
Environment													
<i>E. faecalis</i>	66	47.6	47.6	-	45.7	20.5	23.1	70.2	-	17.0	23.6	96.5	29.8
<i>E. faecium</i>	523	59.2	69.2	100	64.6	26.1	39.7	83.6	-	29.1	49.3	94.9	62.7
<i>S. aureus</i>	77	96.7	25	35	48.2	10.0	16.67	98.9	80.0	16.7	25.8	43.0	50.0

²⁵ Ampicillin

²⁶ Ciprofloxacin

²⁷ Clindamycin

²⁸ Erythromycin

²⁹ Gentamicin

³⁰ Kanamycin

³¹ Penicillin

³² Rifampicin

³³ Streptomycin

³⁴ Tetracycline

³⁵ Sulphamethoxazole-trimethoprim

³⁶ Vancomycin

1540 **Figure 1.** PRISMA-adapted flow chart showing included and excluded articles. All search were conducted on PubMed, Web of Science and African Journals
1541 Online, and a final number of 130 articles were used for the quantitative analysis.

1542 **Figure 2.** Frequency and distribution of resistance genes, antibiotics, and mobile genetic elements (MGEs) with recorded resistance in Gram-positive bacteria in
1543 Africa. 2ai) Shows the frequency of the various resistance genes found in the drug-resistant Gram-Positive bacterial strains. *mecA* and *erm(B)* were the most
1544 dominant resistance genes detected, followed by *tet(M)*, *dfrG*, *vanB*, *vanC1* etc. 2aii) Shows the antibiotics to which the isolates were most resistant:
1545 erythromycin (ERY) was the least effective drug, followed by rifampicin (RIF), tetracycline (TET), penicillin (PEN), sulphamethoxazole/trimethoprim (SXT),
1546 ciprofloxacin (CIP), gentamicin (GEN), vancomycin (VAN), ampicillin (AMP), clindamycin (CLI), streptomycin (STR), chloramphenicol (CHL), and
1547 kanamycin (KAN). 2b) Shows the MGEs per resistant Gram-positive bacterial clones in Africa. The figure represents resistant clones and the different MGEs
1548 they carry. Each colour represent a particular resistant clone. *S. agalactiae* (ST612, ST616, ST617) and *S. pyogenes* (*emm18*, *emm42*, *emm76*, *emm118*), *E.*
1549 *faecium* (ST18, ST80, ST910) and *S. aureus* (ST5, ST22, ST35) were associated with *Tn916*, *IS16* and *SCCmec* respectively.

1550 **Figure 3.** Frequency distribution of resistant Gram-positive bacterial species, clones and mobile genetic elements (MGEs) per country in Africa. 3a) Shows the
1551 distribution frequencies of the resistant species, clones and MGEs per country in Africa whilst 3b) shows the total frequency per clone in Africa. It is obvious that
1552 *S. aureus* ST5 is predominant in Tunisia, the DRC and Senegal whilst ST22 is highly prevalent in Algeria. *SCCmec* was the commonest MGE in most of the
1553 countries except in Tunisia where *IS16* and *Tn916* were higher in prevalence. *S. aureus* ST8 and ST80 were the most common clones reported, followed by *E.*
1554 *faecium* ST317.

1555 **Supplementary data 1.** List of excluded articles on the basis of only phenotypic (antibiotic sensitivity) tests.

1556 **Supplementary data 2.** Raw data and analysis of extracted information from included articles.

Figure 2ai. Frequency of resistance genes conferring resistance to antibiotics in Gram-positive bacteria in Africa.

Figure 2aii. Frequency of antibiotics to which Gram-positive bacteria were resistant to in Africa.

Figure 2b. Frequency and distribution of mobile genetic elements (MGEs) and resistant clones in Gram-positive bacteria in Africa.

Figure 3a. Frequency distribution of resistant Gram-positive bacterial species, clones and mobile genetic elements (MGEs) per country in Africa.

Figure 3b. Frequency of resistant Gram-positive bacterial species, clones and mobile genetic elements (MGEs) per country in Africa.